

VERMONT'S Wildlife Action Plan

Draft 10/1/2015



protect. respect. enjoy.



Vermont Fish & Wildlife Department

Protecting and conserving our fish, wildlife, plants and their habitats for the people of Vermont.



Vermont Wildlife Action Plan, 2015

Draft 10/1/2015

Acknowledgments

This section is incomplete. It will be completed after the draft Action Plan is reviewed by conservation partners and the general public.

Federal Reviewers Guide to the Wildlife Action Plan

This section is incomplete. It will be completed after the draft Action Plan is reviewed by conservation partners and the general public.

Cover credits, clockwise from top

Vermont Landscape-Vermont Department of Tourism & Marketing
Northeastern Bulrush (*Scirpus ancistrochaetus*)- Bob Popp, VFWD
Bobcat-M Tribble, USFWS
Rainbow Bluet-© Bryan Pfeiffer
Wood Turtle- John Mosesso, NBII
Brook Trout-Photo.com
Wood Thrush- S. Maslowski, USFWS

Recommended Citation

Vermont Wildlife Action Plan Team. 2015. Vermont Wildlife Action Plan 2015. Vermont Fish & Wildlife Department. Montpelier, VT. <http://www.vtfishandwildlife.com>

Funding

Funding for the development of the Vermont Wildlife Action Plan was provided by the State Wildlife Grant Program and the Vermont Fish & Wildlife Department and the many organizations, agencies and individuals who volunteered their time and expertise.

I. Executive Summary: Vermont's Wildlife Action Plan

Congress created the State and Tribal Wildlife Grants Program (SWG) in 2001. To receive SWG funds, each state and territory is required to develop a Wildlife Action Plan. The goal of both the State Wildlife Grants program and the Action Plan is to prevent wildlife from becoming endangered through early, strategic efforts to conserve wildlife and habitat. SWG provides funding and the Action Plan provides the strategic guidance. SWG is now the nation's core program for preventing endangered species listings. Since 2001, Vermont has received or become eligible for more than \$8 million in State Wildlife Grant funds.

According to Congressional mandate, Wildlife Action Plan's must be updated at least every 10 years. This report represents the first revision of Vermont's Wildlife Action Plan, originally created in 2005. This revision has nine chapters and eight appendices as outlined here:

Chapter 1. Introduction describes the Vermont Fish & Wildlife Department's mission and approach to revising the Wildlife Action Plan. It discusses Vermonter's deep and abiding interest in fish and wildlife conservation, and the difficulties of conserving Vermont's natural heritage without stable and secure funding. The Congressional requirements for Wildlife Action Plans are presented along with a summary of the revision process and notable changes and additions to the 2015 Vermont Wildlife Action Plan, including: an in-depth consideration of climate change impacts to wildlife conservation; a design for landscape and habitat connectivity; greater attention to disease and pollinators. This chapter also contains the statewide goals and objectives for Wildlife Action Plan implementation.

Chapter 2. Vermont Overview describes Vermont's landscape and the ecological and land-use factors that contribute to Vermont's natural heritage. It provides both a historic and contemporary perspective on land use history and summarizes the significant threats and problems to Vermont's wildlife. It looks back at conservation successes and at the importance of education, law enforcement and recreation to wildlife conservation efforts—three areas not eligible for SWG funding.

Chapter 3. Climate Change and Wildlife Conservation is an in-depth look at climate change and wildlife conservation. It reviews the historic climate trends and predictions for future climate and summarizes the impacts of a changing climate on Vermont's ecology.

Chapter 4. Conservation at Multiple Scales explains how conservation is organized in this Wildlife Action Plan and how habitat classifications were developed for aquatic species and terrestrial species. From a focus on the status and needs of individual Species of Greatest Conservation Need (Appendix A) to their habitats (Appendix B) to taxon-wide conservation (Chapter 5) and landscape-scale conservation (chapter 6)

Chapter 5. Species of Greatest Conservation Need provides lists of Vermont's Species of Greatest Conservation Need and summarizes SGCN conservation needs by taxonomic group (amphibians and reptiles, birds, fishes, invertebrates, mammals and plants).

Chapter 6. Landscape Conservation describes the condition of Vermont's landscapes (historic, current and desired), provides a framework for identifying and prioritizing landscapes important to SGCN and natural heritage conservation based on six key landscape components (Interior Forest Blocks, Connectivity Blocks, Surface Waters and Riparian Areas, Riparian Areas for Connectivity, Physical Landscape Diversity Blocks, and Wildlife Road Crossings), identifies SGCN benefitting from landscape conservation, and identifies significant threats and priority conservation strategies.

Chapter 7. Monitoring, Implementation & Review outlines plans to track the status of Species of Greatest Conservation Need and their habitats, evaluate the effectiveness of conservation actions, implement the Wildlife Action Plan and keep it up-to-date.

Chapter 8. Glossary and Acronym Key is our best effort to round-up and explain all the terms and acronyms used in this report.

Chapter 9. Action Plan Revision describes the efforts made to review and revise the Wildlife Action Plan. It describes the revision timeline, the planning team structure and outreach and public involvement efforts. It explains how SGCN were selected and how conservation actions were developed.

Appendices A1-A5. Species of Greatest Conservation Need Conservation contains detailed reports on each SGCN exploring species status, distribution, habitat usage, threats and problems, research and monitoring needs and conservation actions.

Appendix B. Habitat & Community Summaries contains detailed reports on the 25 major habitat types found in Vermont. Reports explore habitat characteristics and locations, habitat condition (historic, current and desired future), efforts to benefit these habitats as part of 2005 Action Plan implementation, SGCN typically found in each habitat, threats and information needs, and priority conservation actions.

Appendix C. Threats and Actions defines the categories of threats, problems and conservation actions used in this report.

Appendix D. Vermont Species & Habitat Climate Vulnerability Assessment provides data from a 2013 effort to assess the vulnerability of fish, wildlife, plants and their habitats to climate changes. This Vermont Species & Habitat Climate Vulnerability Assessment investigated 18 species, 20 upland habitats, 11 wetland habitats, and 13 freshwater habitats.

Appendix E. Implementing the 2005 Wildlife Action Plan illustrates how that Action Plan has guided conservation in Vermont over the past decade. It provides multiple examples of how SWG funding has supported habitat and species population restoration, has helped leverage the conservation expenditures of other organizations, and has supported research and monitoring for a wide variety of species and the status of emerging threats such as White-nose Syndrome in bats.

Appendix F. Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape presents a landscape-level conservation design for protecting and enhancing ecological function. The lands and waters identified in this document are the areas of the state that are of highest priority for maintaining ecological integrity. Together, these lands comprise a connected landscape of large and intact forested habitat, healthy aquatic and riparian systems, and a full range of physical features (bedrock, soils, elevation, slope, and aspect) on which plant and animal natural communities depend. When conserved or managed appropriately to retain or enhance ecological function, these lands are expected to sustain Vermont's natural legacy into the future.

Appendix G. Mapping Vermont's Natural Heritage was developed to help municipal partners implement the Wildlife Action Plan. It is an essential guide to understanding natural resource data designed specifically for Vermont towns. It provides an understanding of available datasets, what they mean, and how to access them. It also serves as an important compliment to the 2004 "Conserving Vermont's Natural Heritage" that makes the case for why certain resources are critically important for conservation planning and provides an essential framework for looking at the landscape at different ecological scales. Using this same framework, "Mapping Vermont's Natural Heritage" provides the specific local data to help users connect ecological concepts to land-use planning tools and guides users through the next steps in ecological prioritization.

Appendix H. Vermont Big Game Management Plan is Vermont Fish & Wildlife Department's guide to managing the state's White-tailed Deer, Moose, Black Bear and Wild Turkey populations through 2020.

Table of Contents

I. Executive Summary.....	i
Chapter 1. Introduction to Vermont's Wildlife Action Plan	1.2
VFWD: Mission and Strategic Focus.....	1.2
Vermonters Care about Wildlife	1.3
Problem and Need: Secure Funding	1.3
Revising Vermont's Wildlife Action Plan.....	1.4
The Eight Required Elements of a Wildlife Action Plan.....	1.5
A Decade of Conservation: Implementing the 2005 Wildlife Action Plan.....	1.6
Statewide Goals, Objectives, Actions for Wildlife Action Plan Implementation	1.7
Literature Cited	1.13
Chapter 2. Vermont Overview.....	2.1
The Vermont Landscape—an Overview.....	2.1
Biophysical Regions of Vermont	2.3
Vermont's Landscape—an Historical Perspective.....	2.4
Vermont's Contemporary Land Use	2.4
Contemporary Problems and Threats to Vermont's Wildlife.....	2.5
Conservation Success! Keeping Common Species Common.....	2.10
The Importance of Education, Law Enforcement and Recreation to Wildlife Conservation	2.12
Literature Cited	2.15
Chapter 3. Climate Change & Conservation	3.2
Historic Trends and Future Projections	3.3
Vermont's Changing Seasons	3.4
Impacts of Climate Change on Vermont's Ecology.....	3.5
Conserving Wildlife in the Face of Climate Change.....	3.10
Identifying Climate Adaptation Strategies for Vermont's Species and Habitats.....	3.20
Priority Actions to Address Climate Change Impacts to Species and Habitats.....	3.20
Managing Vermont's Wildlife and Habitats into the Future.....	3.21
Literature Cited	3.22
Chapter 4. Conservation at Multiple Scales	4.2
Selection of Classification Systems.....	4.3
Habitat Succession, Species of Greatest Conservation Need & the Action Plan.....	4.4
Literature Cited	4.7
Chapter 5. Vermont's Species of Greatest Conservation Need	5.1
SGCN Lists	5.4
Conserving Vermont's Amphibians & Reptiles	5.13
Conserving Vermont's Birds.....	5.19
Conserving Vermont's Fishes.....	5.27
Conserving Vermont's Invertebrates	5.34
Conserving Vermont's Mammals	5.45
Conserving Vermont's Plants	5.54
Chapter 6. Landscape Conservation	6.2
Landscape Condition	6.2
Implementing the 2005 Wildlife Action Plan	6.4
SGCN Benefitting from Landscape Conservation	6.6
Landscape Characteristics	6.6

The Ecologically Functional Landscape	6.9
Threats & Information Needs.....	6.11
Priority Conservation Strategies.....	6.11
Literature Cited	6.13
Chapter 7. Monitoring, Adaptive Management, Implementation and Review	7.1
Monitoring, Adaptive Management and Vermont's Wildlife Action Plan.....	7.1
Literature Cited	7.10
Action Plan: Implementation	7.12
Action Plan Review	7.15
Ten Year Review	7.16
Chapter 8. Glossary & Acronym Key	8.1
Chapter 9. Revising Vermont's Wildlife Action Plan.....	9.2
Threats, Problems and Species of Greatest Conservation Need.....	9.4
Outreach and Public Involvement.....	9.7
Identifying Species of Greatest Conservation Need.....	9.8
Conservation of Species of Greatest Conservation Need	9.12
Literature Cited	9.13

Appendices

A. Species of Greatest Conservation Need Assessment Summaries

- A1-Amphibians & Reptiles
- A2-Birds
- A3-Fishes
- A4-Invertebrates
- A5-Mammals

B. Habitat & Community Conservation Summaries

- Conservation at Multiple Scales 2

Terrestrial Habitats

- Northern Hardwood Forest B.9
- Spruce-Fir-Northern Hardwood Forest ... B.16
- Oak-Pine-Northern Hardwood Forest..... B.23
- Young Forest-Old Forest..... B.32
- Floodplain Forest
- Hardwood Swamp..... B.38
- Softwood Swamps..... B.43
- Vernal Pools & Seeps
- Open Peatlands..... B.53
- Marshes and Sedge Meadows
- Wet Shores..... B.62
- Shrub Swamps..... B.66
- Upland Shores..... B.70

- Outcrops & Upland Meadows B.73
- Cliff & Talus..... B.76
- Grassland & Hedgerow
- Mines and Quarries
- Subterranean..... B.87
- Buildings & Other Structures

Aquatic Habitats

- Riparian B.93
- Streams & Rivers
- Lake Champlain Tributaries..... B.106
- Lower Connecticut River
- Lake Champlain..... B.117
- Lakes (excluding Lake Champlain)..... B.122

C. Threat & Action Definitions

D. Vermont Species & Habitat Climate Vulnerability Assessment

F. Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape

G. Mapping Vermont's Natural Heritage

H. Vermont Big Game Management Plan

Vermont's Wildlife Action Plan

Chapter 1 Introduction

Draft September, 2015

Introduction	2
The Vermont Fish & Wildlife Department: Mission and Strategic Focus	2
Vermonters Care about Wildlife	3
Problem and Need: Secure Funding	3
Revising Vermont's Wildlife Action Plan.....	4
The Eight Required Elements of a Wildlife Action Plan	5
A Decade of Conservation: Implementing the 2005 Wildlife Action Plan.....	6
Statewide Goals, Objectives, Actions for Wildlife Action Plan Implementation	7
Literature Cited.....	13

Introduction

Vermont is known for its beautiful juxtaposition of forests and farms, streams and ponds, Lake Champlain and the Green Mountains. Fish and wildlife are an integral part of the Vermont experience and quality of life for Vermonters. White-tailed Deer, Black Bear, Wild Turkey, and geese are thriving and offer sustainable hunting and viewing opportunities. In recent years, and after a lot of effort, Common Loon, Osprey, Peregrine Falcon populations have recovered allowing for their removal from the state threatened and endangered species list, the Bald Eagle is again nesting in Vermont and many of our waters teem with healthy fish communities. The landscape and the wildlife attest to the state's meaningful conservation ethic.

Over the past 10-years Vermont's Wildlife Action Plan helped guide the Vermont Fish & Wildlife Department, partners, stakeholders and others in the conservation of our Species of Greatest Conservation Need and efforts to keep common species common. This 10-year revision to the Action Plan has allowed us to: evaluate our implementation efforts; reassess the status of our Species of Greatest Conservation Need (SGCN) and their habitats; investigate emerging issues; and, renew our vision for fish and wildlife conservation in Vermont.

Notable in this 2015-2025 Wildlife Action Plan are the addition of a climate change chapter; the re-recognition of diseases as significant threats to some species, and of the vital role pollinators play in the environment (a suite of bumble bees are included as SGCN); and, that habitat loss and degradation remain the primary threat to most SGCN.

This Wildlife Action Plan's emphasis on landscapes reflects a doubling-down on habitat conservation as the most effective and efficient strategy for keeping common species common and reducing the need to place species on the endangered species list. By maintaining and enhancing healthy, connected, landscapes we maintain the habitat integrity, help a wide range of species, improve climate change resiliency, and reduce the potential impacts of diseases and non-native invasives. Moreover, healthy landscapes support healthy economies, human communities and our quality of life, and ensure that wildlife and the places they live will be there for future generations to enjoy.

This Wildlife Action Plan has not changed its focus on implementation through collaboration and partnership. Conserving and managing Vermont's wildlife requires everyone's help. From landowners and land managers, to municipalities and regional planning commissions, state and federal agencies, and non-profit organizations and for-profit businesses all need to work together if we are to achieve our goal.

The Vermont Fish & Wildlife Department: Mission and Strategic Focus

The Vermont Fish & Wildlife Department's mission is to protect and conserve our fish, wildlife, plants and their habitats for the people of Vermont. The Department's three planning goals are:

Conserve, enhance, and restore Vermont's natural communities, habitats, and plant and wildlife species along with the ecological processes that sustain them.

Provide a diversity of safe and ethical fish and wildlife-based activities and opportunities that allow hunting, fishing, trapping, viewing, and the utilization of fish, plants, and wildlife resources consistent with the North American Model of fish and wildlife conservation.

Maintain safe fish and wildlife-based activities and limit harmful human encounters with fish and wildlife species, and provide general public safety service incidental to our primary fish and wildlife duties.

Throughout its 100-year history, many of the Fish & Wildlife Department's initiatives focused on game species, in part because of constituent interest, as well as the user-pay system of funding fish and wildlife conservation through license sales and excise taxes on hunting and fishing equipment. As our mission statement suggests, the scope of conservation challenges facing Vermont extends beyond species which people choose to harvest.

Vermonters Care about Wildlife

Wildlife is very important to the people of Vermont. This love of wildlife is more than anecdotal. The 2011 National Survey of Fishing, Hunting, and Wildlife Associated Recreation conducted by the U.S. Fish and Wildlife Service documented that 62 percent of Vermonters went fishing, hunting, or wildlife watching. Vermont ranked second, only two points behind Alaska in participation (U.S. Dept of Interior 2011). When it comes to wildlife watching, however, Vermont was first in the nation with an impressive 53 percent of residents enjoying this activity. This same survey estimates more than \$704 million was spent on fish-and wildlife-based recreation in Vermont.

A July, 2015 public attitude survey measured trends in Vermonters' support for the Department's mission: "The conservation of fish, wildlife, plants, and habitats for the people of Vermont" (Duda et al 2015)." Findings included:

- 83 percent responded, when asked to compare the importance of wildlife with economic development, that the use and development of land should be restricted to protect fish and wildlife; and 81 percent responded that wildlife habitat must be protected even if it reduces the land use options of some landowners and developers.
- 81 percent strongly agreed that endangered species must be protected (only 2% moderately or strongly disagreed). 86 percent of hunters and anglers strongly agree that endangered species should be protected.
- When asked which of the following were big problems, residents listed invasive species (71%), climate change (64%), forest and habitat fragmentation (46%), fish and wildlife habitat loss (46%), loss of Vermont's scenic landscape (41%), and posting land closed to hunting/recreation (26%).

Problem and Need: Secure Funding

Historically, dedicated funds have been available for the conservation and management of game and sportfish species, and to a lesser extent for Threatened and Endangered species. Unfortunately, there has not been a dedicated revenue stream supporting management for the vast majority of wild animals that do not fall within either category. For example, 269 species of birds are found in Vermont. Only about 30 of these are hunted and only a handful (e.g., Common Loon, Osprey, Peregrine Falcon) have benefitted from Endangered Species recovery funding.

In 1985, Vermont's Nongame and Natural Heritage Program (now the Wildlife Diversity Program) was established within the Fish and Wildlife Department. Both an income tax check-off and a conservation license plate were important funding mechanisms for the management and recovery of non-game species, generating approximately \$200,000 per year, but has never adequately met program needs.

Congress responded to the funding need in 2001 with a new annual appropriation, first called the Wildlife Conservation and Reinvestment Act and later the State Wildlife Grants Program (SWG). SWG funds were

to be targeted to fish and wildlife species (plants are not eligible) in need, including those that had not previously benefitted from traditional funding. Since that time, the Vermont Fish and Wildlife Department has received \$8.1 million in SWG appropriations and have vastly improved Vermont's ability to fund new research, inventory, and management initiatives for species such as Timber Rattlesnake, Lake Sturgeon, butterflies, and the Bicknell's Thrush. Education and law enforcement are not eligible activities.

In order to receive SWG funds, Congress requires states to develop Wildlife Action Plans to guide use of the funds. But SWG funds are not sufficient to fully implement the Vermont Wildlife Action Plan. Annual SWG allocations have declined by 40% since 2002 and additional cuts are threatened every year. Currently Vermont receives less than 10% of what would be needed for full Wildlife Action Plan implementation.

Within the State of Vermont, the annual funding picture is rosier. Thanks to the strong support of the Governor and Legislature, general fund allocations to the Fish & Wildlife Department have increased approximately four-fold from \$1.25 million in 2011 to over \$5 million annually in 2015. While only a small portion of these funds are directly invested in Wildlife Action Plan implementation, it all helps the Department meet its broader mission. In 2015 the [Vermont Habitat Stamp](#) was created to provide additional opportunities for Vermonters and friends of Vermont to contribute to the purchase and maintenance of Wildlife Management Areas.

Revising Vermont's Wildlife Action Plan

The revision of Vermont's Wildlife Action Plan followed the [eight required elements](#), revision guidance provided by the U.S. Fish & Wildlife Service and that published in Best Practices for State Wildlife Action Plans (AFWA 2012) and The Northeast Lexicon: Terminology Conventions and Data Framework for State Wildlife Action Plans in the Northeast Region (Crisfield 2014).

Revision began with a close review of the 2005 Action Plan. We found that it had weathered the decade surprisingly well. Much of the plan, in terms of focus, organization and goals, did not need change, only some updating. The foundation of both the 2005 and 2015 plans are the detailed reports on individual SGCN (Appendix A1-4). Both Action Plans focus on habitat conservation as a primary means of conserving SGCN (the 2015 Action Plan provides additional emphasis on landscape conservation and connectivity). And, both stress collaborations and partnerships in implementation.

With this revision we take advantage of the many tools, guidance documents and programs developed since 2005 designed to support the conservation and management of wildlife by partners and the general public—several of which were created as a direct result of the first Wildlife Action Plan. These include the [Community Wildlife Program](#), [Foresters for the Birds](#), [BioFinder](#), [Vermont Invasives](#) the [Landowners Guide - Wildlife Habitat Management for Lands in Vermont](#) and [Community Strategies for Vermont's Forests and Wildlife](#) among many.

Notable changes/additions to the Vermont's Wildlife Action Plan with this revision include:

- An expanded discussion of climate change impacts to SGCN and their habitats and strategies to help wildlife adapt and to improve resiliency (chapter 3).
- A focus on landscape conservation and habitat connectivity (chapter 5) and a design for landscape conservation (appendix F).
- Greater attention to diseases as significant threats to some species.

- Recognition of the important role pollinators play in their ecosystems and the addition of a suite of nine bumble bee species included as SGCN (Appendix A4). The 2005 Action Plan was silent on pollinators.
- A plant conservation summary (chapter 4). The 2005 Action Plan included only a list of plant SGCN.
- Expanded guidance to help municipalities implement the Action Plan (Appendix H).
- Addition of the state's Big Game Management Plan (for Black Bear, Moose, White-tailed Deer, and Wild Turkey) as an appendix (I).
- Revisions to Species of Greatest Conservation Need: SGCN lists have been updated (chapter 4) along with conservation summaries for each SGCN (appendix A). Table 1.1 summarizes these changes.

Table 1.1 Changes to SGCN Lists 2005:2015

Taxon	2005 SGCN	2015 SGCN	Change Notes
Amphibians & Reptiles	19	19	No changes
Birds	58	51	Long-eared Owl, Henslow's Sparrow, Osprey, Cooper's Hawk, Barn Owl, Veery, Blue-winged Teal removed.
Fishes	33	27	Arctic Char, Atlantic salmon (anadromous), Brassy Minnow, Muskellunge, Quillback and the Sea Lamprey (CT River) removed.
Invertebrates	191	200	19 species removed; 28 added, including nine bumble bee species.
Mammals	33	34	Mink removed. Moose and Snowshoe Hare added.
Plants	577	645	

The Eight Required Elements of a Wildlife Action Plan

Wildlife Action Plans are meant to guide the use of SWG funds, so in order to remain eligible for State Wildlife Grants funding Congress requires that states develop Wildlife Action Plans and to revise them at least every ten years. Congress prescribed the following eight required elements for Wildlife Action Plans:

1. Identify wildlife distribution and abundance: Provide information on the distribution and abundance of species of wildlife, including low and declining populations as the State fish and wildlife agency deems appropriate, that are indicative of the diversity and health of the State's wildlife.
2. Describe location and condition of key habitats: Describe the locations and relative condition of key habitats and community types essential to conservation of species identified in (1).
3. Describe key problems and research needs: Describe problems that may adversely affect species identified in (1) or their habitats, and priority research and survey efforts needed to identify factors which may assist in restoration and improved conservation of these species and habitats.
4. Describe and prioritize conservation actions: Describe conservation actions proposed to conserve the identified species and habitats and priorities for implementing such actions.

5. Monitor species, habitats and conservation actions: Describe plans to monitor species identified in (1) and their habitats; monitor the effectiveness of the conservation actions proposed in and, adapt these conservation actions to respond appropriately to new information or changing conditions.
6. Develop a plan review process: Describe procedures to review the Wildlife Action Plan at intervals not to exceed ten years.
7. Coordinate with other plans: Coordinate the development, implementation, review, and revision of the Action Plan with Federal, State, and local agencies and Indian tribes that manage significant land and water areas within the State or administer programs that significantly affect the conservation of identified species and habitats.
8. Include public participation: Describe public participation in the development, revision, and implementation of the Action Plan and projects and programs.

A Decade of Conservation: Implementing the 2005 Wildlife Action Plan

A summary of efforts to implement Vermont's first Wildlife Action Plan can be found in appendix E.

Statewide Goals, Objectives, Actions for Wildlife Action Plan Implementation

- 1. Goal: Conserve, restore, and enhance habitats, natural plant and animal communities, and ecosystem integrity to maintain suitability for SGCN and ecological function and to improve resiliency to climate change.**
 - 1.1. Objective: Prioritize and implement land acquisition based on conservation planning initiatives to protect important SGCN habitats through acquisition of easements or fee title.**
 - 1.1.1. Action: Acquire fee title of lands providing important SGCN habitat and habitat connectivity. Measure: Number of acres/miles protected.
 - 1.1.2. Action: Acquire conservation easements on lands providing important SGCN habitats and habitat connectivity. Measure: Number of acres/miles protected.
 - 1.1.3. Track the number of acres of contiguous forest lost to fragmentation/development. Measure: Number of acres lost to fragmentation/development.
 - 1.2. Objective: Participate in regulatory permitting and enforcement activities to protect significant natural communities, endangered and threatened species, and fish and wildlife habitats.**
 - 1.2.1. Action: Participate in Section 248 and Act 250 development processes to reduce negative impacts from development. Measure: Number of acres of habitat protected from development.
 - 1.2.2. Action: Participate in regulatory review processes to reduce negative impacts of altered flow in streams and rivers. Measure: Number of permits modified or issued that ensure sufficient flows.
 - 1.2.3. Action: Participate in regulatory review processes to reduce negative impacts of shoreline development to prevent increases in improperly hardened shorelines on lakes and streams through regulatory review and enforcement. Measure: Linear feet of shoreline habitat protected from improper development through project review and permitting.
 - 1.2.4. Action: Participate in regulatory review processes to reduce negative impacts of development near wetlands.
 - 1.2.5. Action: support the updating of wetland delineations and maps particularly for Class 1 wetlands. Measure: Acres of wetlands added to the state wetland maps.
 - 1.3. Objective: Maintain, restore and promote management of public and private lands and waters to enhance their suitability for SGCN, integrity and ecological function, and climate change resiliency.**
 - 1.3.1. Action: Implement ecosystem and landscape-level management on lands managed by ANR and other agencies. Measure: Number of state lands with Long Range Management Plans addressing landscape management goals.
 - 1.3.2. Action: Restore or enhance riparian buffers at key locations. Measure: Miles of stream bank re-vegetated.
 - 1.3.3. Action: Restore aquatic organism passage through barrier mitigation in waters critical to SGCN. Measure: Miles of natural passage reconnected in SGCN habitats.

- 1.3.4. Action: Restore natural hydrologic flow regime through barrier mitigation in waters critical to SGCN. Measure: Miles of natural stream flow reconnected in key SGCN habitats.
- 1.3.5. Action: Restore and maintain grassland habitats within grassland focus areas. Measure: Number of acres appropriately managed for grassland SGCN.
- 1.3.6. Action: Implement management to restore and maintain habitat for young forests (early successional) and old forests (late succession) dependent species on state lands to meet objectives of landscape conservation plans. Measure: Percentage of forest meeting target successional stages.
- 1.3.7. Action: Provide technical and financial assistance to private forest owners to support land management to meet targeted successional stages for SGCN. Measure: Acres of private forest land primarily managed for young forest/old forest SGCN; Number of landowners and acres enrolled in the Current Use Ecologically Sensitive Treatment Area (ESTA) program.
- 1.3.8. Action: Provide technical assistance to local state and federal agencies that own or manage lands or set policies affecting land management and SGCN. Measure: the number of municipalities with town plans that address SGCN needs.

1.4. Goal: Integrate SGCN conservation into natural resource planning.

- 1.4.1. Objective: Develop and maintain interagency cooperatives to protect and manage SGCN.
- 1.4.2. Action: Participate in existing cooperatives to include SGCN in program implementation, including the interstate fish and wildlife cooperatives, Lake Champlain Basin Program. Measure: Number of interagency cooperatives incorporating SGCN needs into planning documents.
- 1.4.3. Action: Implement existing cooperative natural resource plans including the Lake Champlain Opportunities for Action Management Plan. Measure: Number of implementation actions conserving SGCN habitats or populations.
- 1.4.4. Action: Continue the collaborative biodiversity conservation work of the northeastern states, including the Regional Conservation Needs Program and the Northeast Fish and Wildlife Diversity Technical Committee. Measure: Number of regional conservation projects.
- 1.4.5. Action: Continue engagement with the, North Atlantic Landscape Conservation Cooperatives (LCC). Measure: Number of LCC projects in Vermont benefitting SGCN.

1.5. Objective: Control problematic native species and invasive species deleterious to SGCN and prevent introduction of these species.

- 1.5.1. Action: Develop and adopt guidelines in cooperation with the Agency of Natural Resources for the management of riparian buffers, vernal pools and other ecologically significant habitats and. Measure: Number of guidelines development and implemented.
- 1.5.2. Action: Control invasive plants in key SGCN habitats to protect ecosystem health and stability. Measure: number of acres with dominant native vegetation restored.
- 1.5.3. Action: Manage and control the spread of insect invasive species in forested SGCN habitats. Measure: Estimate of reduction or prevention of forest damage.
- 1.5.4. Action: Manage white-tailed deer and moose populations to foster forest regeneration

- and understory growth in areas of very high deer or moose population density per VFWD's Big Game Management Plan. Measure: Deer/moose density.
- 1.5.5. Action: Control double-crested cormorant, gull, and problematic mammalian predators at locations where they are negatively impacting nesting birds, turtles and other SGCN. Measure: Nesting success of SGCN species.
 - 1.5.6. Action: Continue efforts to limit the introduction of aquatic invasive species. Measure:
 - 1.5.7. Action: Continue efforts to limit the introduction of emerald ash borer, Asian long-horn beetle, hemlock wooly adelgid and other forest pests. Measure: Number of confirmed infestations.
 - 1.5.8. Action: Work with the Forests, Parks & Recreation Department and researchers to identify examples of hemlock and other forest types to be protected from forest pests and diseases, and strategies to combat infestations while limiting impacts to non-target species.
 - 1.5.9. Action: Prevent introduction of feral swine. Measure: Number of feral swine in the state.

1.6. Objective: Protect habitat viability by developing and maintaining programs to reduce environmental contamination and pollution, atmospheric mercury and acid deposition and the atmospheric discharges of greenhouse gases that are contributing to climate change.

- 1.6.1. Action: Implement the [Lake Champlain restoration plan](#) to reduce phosphorous and other nutrient pollutant inputs into the Lake Champlain watershed. Measure: Progress toward meeting the Total Maximum Daily Load (TMDL) goals for the lake and its tributaries.
- 1.6.2. Action: Reduce the occurrence of combined sewer overflow discharges. Measure: The number of waste water treatment systems that have been updated to reduce combined sewer overflow events.
- 1.6.3. Action: Continue controlling mercury discharges from energy generating and industrial facilities. Measure: Decline in mercury levels in impacted SGCN.
- 1.6.4. Action: Continue controlling acid discharges from combustion sources. Measure: Increase in pH in impacted waters.
- 1.6.5. Action: Continue participating in the Regional Greenhouse Gas Initiative. Measure: Number of carbon pollution credits sold.
- 1.6.6. Work with state and federal agencies, legislators, and partners to reduce water and air pollution and greenhouse gas emissions.

1.7. Objective: Manage human disturbance in SGCN habitats.

- 1.7.1. Action: Restrict public access to bat hibernacula to protect bats and continue closures of cliff areas and turtle nesting beaches to protect peregrine falcons and turtles during the nesting seasons. Measures: Number of hibernacula with restricted access; nesting success of Peregrine Falcons and Spiny Softshell Turtles in protected areas.
- 1.7.2. Action: Manage recreational use of off-road vehicles in important SGCN habitats. Measure: Number of acres protected.
- 1.7.3. Support changes in laws to minimize degradation to fish and wildlife habitat.

1.8. Objective: Promote habitat connectivity for SGCN.

- 1.8.1. Action: Restore and maintain natural habitats in linkage areas between large blocks of

habitat for SGCN. Measure: Number of acres of viable habitat in identified linkage areas.

- 1.8.2. Action: Work with VTrans to identify and mitigate barriers to habitat connectivity. Measure: Number of projects implemented to reduce road mortality and increase connectivity.
- 1.8.3. Action: Remove or replace culverts and dams that impede passage of aquatic SGCN in high priority areas. Measure: Miles of SGCN habitat with restored connectivity.
- 1.8.4. Action: Foster northeast regional habitat connectivity projects for wide-ranging SGCN. Measure: Identification and conservation of important habitat linkages between Vermont and neighboring states and provinces.

1.9. Objective: Promote practices to improve habitat for pollinator species by providing a diversity of flowering plants throughout the active season.

- 1.9.1. Action: Enhance pollinator habitat on state lands. Measure: acres of pollinator habitat enhanced.
- 1.9.2. Action: Provide information on pollinator-friendly landscaping and agricultural practices to the public via the VFWD website. Measure: Number of visits to pollinator habitat management webpage.
- 1.9.3. Action: Develop mowing plans for state lands (e.g., parks, roadsides, rest stops, old fields and rights-of-way) to benefit SGCN pollinators and to limit the spread of invasive plants. Develop similar mowing best management practices for use by partners including VTrans and municipal highway departments. Measure: Completion and implementation of plans and best management practices.

2. Goal: Conserve and manage Vermont's Species of Greatest Conservation Need for self-sustaining populations.

2.1. Objective: Maintain appropriate legal protection for all SGCN.

- 2.1.1. Action: Revise Vermont's Endangered/Threatened statutes to better support habitat protection.
- 2.1.2. Action: Update rules and policies for plant and animal importation, possession and collection/harvest to protect populations and to prevent disease introductions.
- 2.1.3. Through policy and education support the enforcement of existing laws that protect Species of Greatest Conservation Need.

2.2. Objective: Recover populations of Species of Greatest Conservation Need.

- 2.2.1. Action: Continue spiny softshell turtle, spotted turtle, lake sturgeon, spruce grouse, Canada Lynx, American Martin recovery projects. Measure: Status of species populations.
- 2.2.2. Action: Collaborate with neighboring states to manage SGCN populations that cross state boundaries. Measure: Number of collaborative projects with neighboring states.
- 2.2.3. Action: Evaluate the need for recovery plans for additional SGCN and develop such plans where feasible. Measure: the number of recovery plans developed for targeted species.

2.3. Objective: Maintain fish and wildlife health programs and biosecurity to document the

causes of SGCN mortality and occurrence and to track disease outbreaks impacting SGCN.

- 2.3.1. Action: Investigate and report mortality of fish SGCN as appropriate, and report unusual incidences of mortality. Measure: Annual reports on causes of fish mortality in Vermont.
- 2.3.2. Action: Investigate and report mortality of animal SGCN as appropriate, and report unusual incidences of mortality through the Northeast Wildlife Disease Cooperative. Measure: Annual summary reports on causes of wildlife mortality in Vermont.
- 2.3.3. Action: Prevent the transmission of diseases, and invasive species, through the practice of standard disinfection procedures.
- 2.3.4. Action: Monitor chemical contaminants in fish, bird, and mammal top predators. Measure: Number of species monitored for contaminants.

3. Goal: Inventory, monitor, and research SGCN, their habitats and natural communities to provide baselines for conservation and to maintain ecological integrity.

3.1. Objective: Inventory and monitor the abundance, distribution and status of Vermont's SGCN.

- 3.1.1. Action: Identify and prioritize inventory and research needs, and collect information according to priority.
- 3.1.2. Action: Develop abundance and distribution objectives for fish and wildlife species and, where appropriate, prepare species management plans in order to maintain sustainable species population levels within ecological limits while considering social implications.
- 3.1.3. Action: Develop a collaborative, statewide monitoring and adaptive management program to determine SGCN baselines, measure progress toward desired SGCN outcomes, and to evaluate and improve the effectiveness of conservation strategies.
- 3.1.4. Action: Continue and enhance habitat monitoring programs to better track the distribution, abundance and status of SGCN habitats.

3.2. Objective: Maintain data management systems to share data among partnerships and better integrate Vermont data into regional collaborations.

- 3.2.1. Action: Maintain and enhance the Natural Heritage Inventory and other data collection and management programs to inform SGCN and habitat conservation.
- 3.2.2. Action: Continue participation in regional efforts including those with other Northeast states to assess SGCN and their habitats, share data and to implement regional conservation actions. Measure: Number of regional actions implemented.

3.3. Objective: Foster research that improves our understanding of SGCN population dynamics and habitat utilization.

- 3.3.1. Action: Partner with universities fish and wildlife cooperative research units and nongovernmental organizations on research into species and habitat issues.

4. Goal: Maintain and enhance public support for SGCN conservation

4.1. Objective: Inform and engage the public and conservation communities about SGCN conservation and the Wildlife Action Plan.

4.1.1. Action: Distribute information on SGCN conservation to the public.

4.1.2. Action: Support Citizen Science programs for SGCN monitor key SGCN

4.2. Objective: Consult with all levels of government and conservation partners to seek unity of effort in conserving SGCN.

4.2.1. Action: Share information regarding important SGCN, their habitats, and best- practices conservation actions with county and town governments to inform their land use decision making and conservation efforts. Measure: Number of local governments that adopt modified practices to conserve SGCN and their habitats.

4.3. Increase funding and support for local, state and federal conservation programs.

4.3.1. Action: Collaborate and support partners (NGO's, municipalities, state and federal agencies, etc.) to build capacity for conservation initiatives.

Literature Cited

- AFWA. 2012. Best Practices for State Wildlife Action Plans: Voluntary Guidance to States for Revision and Implementation. Teaming With Wildlife Committee's State Wildlife Action Plan Best Practices Working Group. November 2012. <http://www.fishwildlife.org/files/SWAPBestPractices.pdf>
- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity. Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT. www.vtfishandwildlife.com/library.cfm?libbase_=Reports_and_Documents
- Crisfield, E and the Northeast Fish and Wildlife Diversity Technical Committee. 2013. The Northeast Lexicon: Terminology Conventions and Data Framework for State Wildlife Action Plans in the Northeast Region. A report submitted to the Northeast Fish and Wildlife Diversity Technical Committee. Terwilliger Consulting, Inc., Locustville, VA.
- Duda, Mark D., Martin J., Beppler T., Bissell S.J., Center A., Criscione A., Doherty P., Hughes G.L., Kirkman T., Reilly C., Lanier A. 2015. Opinions on Fish, Wildlife, and Land Use among Vermont Residents, Hunters, and Anglers. Responsive Management National Office.
- U.S. Department of the Interior, U.S. Fish and Wildlife Service, and U.S. Department of Commerce, U.S. Census Bureau. 2011 National Survey of Fishing, Hunting, and Wildlife-Associated Recreation

Vermont's Wildlife Action Plan

Chapter 2: Vermont Overview

Draft • September, 2015

Vermont Overview.....	1
Vermont Cares about Wildlife Conservation	1
The Vermont Landscape—an Overview.....	1
Biophysical Regions of Vermont.....	3
Vermont's Landscape—an Historical Perspective.....	4
Vermont's Contemporary Land Use.....	4
Contemporary Problems and Threats to Vermont's Wildlife	5
Conservation Success! Keeping Common Species Common.....	10
The Importance of Education, Law Enforcement and Recreation to Wildlife Conservation.....	12
Conservation Education	12
Wildlife-Associated Recreation	13
Law Enforcement.....	14
All for one and one for all: Law enforcement, Education & Recreation	14
Literature Cited.....	15

Vermont Overview

Vermont Cares about Wildlife Conservation

It is no mystery why people enjoy living in and visiting Vermont. This state has what so many other once rural places have lost: a wealth of wildlife and scenic beauty, traditional working landscapes that support viable local economies, and desirable social and cultural attributes – low crime rate, helpful neighbors, and close-knit villages and towns.

Wildlife, scenic beauty, and the landscape that supports this way of life are not only vital parts of Vermont's rural character and identity, but are highly valued by Vermont residents. Based on 2011 public opinion survey results from the U.S. Fish and Wildlife Service, Vermont ranked first in the nation in percentage of residents that actively observed wildlife (53%). The results also show that resident and non-resident hunting, fishing, and wildlife viewing expenditures in Vermont totaled \$704 million. These statistics represent a significant contribution to the state's economy and underscore the strong connection Vermont residents and non-residents have to the land and wildlife.

Vermont's natural diversity, which include forests, clean waters, vibrant fisheries, healthy wildlife populations, rare species, significant natural communities, and a working landscape, provide people with the opportunity to, among other things, hunt, fish, trap, watch wildlife, hike and work the land.

The Vermont Landscape—an Overview

Vermont's landscape is a rich tapestry of mountains, valleys, woods and wetlands, with a fascinating [geological history](#). It is Vermont's natural landscape that enriches the lives of those who live here and draws so many visitors to the state. It is this same landscape that provides us with clean air, clean water, and habitat for thousands of species of plants and animals.

Understanding Vermont's natural heritage requires understanding the physical landscape. The configuration of mountains, valleys, wetlands, lakes, and rivers is crucial in determining the distribution of natural communities, habitats, and native species.

The following broad environmental factors influence the distribution of species, habitats and natural communities: climate, bedrock geology, surficial geology, topography, hydrology, and land use history. These factors that comprise and influence the Vermont landscape and subsequently the flora and fauna of the state are explained below.

Climate

Vermont's lowest land point is the shore of Lake Champlain, only 95 feet above sea level. Vermont's highest point is the Chin on Mount Mansfield, which juts to 4,393 feet. The distance between Lake Champlain and the summit of Mount Mansfield is only 20 miles, but in that short distance, the climate, topography, and vegetation change considerably. On warm limy sites in the Champlain Valley, where the growing season is 150 days, shagbark hickory trees grow. Apple orchards are common in this environment as well as dairy farming due to the influence of climate on growing season. On the summit of Mount Mansfield,

where the growing season is limited to 90 days, red spruce and balsam fir grows in stunted and contorted mats, bending to the direction of incessant winds.

Climate is major factor in determining the distribution of natural communities, habitats, plants and animals. Elevation provides a means for understanding the influence of climate on wildlife and habitats in Vermont because climate changes in relatively short distances with change in elevation. Thus climate's effect on biota can easily be observed.

Geology

Vermont's bedrock composition is varied and thus, Vermont's soils, hydrology, and subsequently plant distribution and abundance vary. The variations influence, in part, wildlife and plant distribution. The rocks comprising the Southern Green Mountains were formed more than 570 million years ago. The rocks of the Champlain Valley and the Northern Green Mountains date from a time 540 to 443 million years ago when Vermont was the edge of a warm, tropical sea. The remains of marine mammals that inhabited that sea can be found in the Champlain Valley's limestone rock. The youngest rocks in Vermont are the granites, including the stone that makes up the Barre granite quarries. These rocks were formed 200 to 400 million years ago as a result of deep underground magma welling up and hardening.

Whether the bedrock is limestone or granite—or some other rock type—is particularly important in the distribution of natural communities and plants because each rock type has its own particular physical and chemical composition. For instance, rich fens, a rare type of wetland with plants that require high levels of calcium, occur almost exclusively in areas where limestone or similar calcium-rich rock is found.

Vermont's surficial geology is defined by the sands, gravels, clays, peats, and other deposits found on top of the bedrock. They are a result of both glacial activity and post-glacial events (such as flooding) that continue today. Bedrock and surficial geology together have a profound influence on the soils in which Vermont's plants grow.

Topography

Topography describes the physical landscape and influences the distribution of plants, animals, and natural communities. The soil on the top of a mountain tends to be shallow and dry, whereas the soil at the base of a slope tends to be deep, moist, and rich in organic matter because of the downslope movement of plant litter and soil. Cliffs, for example, offer a specialized habitat for specialized groups of plants, as well as denning habitat for Bobcat and nesting sites for Peregrine Falcon. Certainly, topography influences the quality and distribution of winter habitat for White-tailed Deer in Vermont.

Hydrology

Water and its movement have a profound influence on animals, plants and natural communities, and ecosystem processes. Lakes, ponds, rivers, and streams provide habitat for a diversity of fish, aquatic plants, aquatic invertebrates, and other organisms. Wetlands form in waterlogged soils, either in low-lands where water collects by gravity, in uplands where impermeable soils create perched water tables, or at the highest elevations where fog and abundant rain provide a constant supply of water for wetland plants and animals.

Land Use History

Land use history has influenced the distribution of plants and animals across Vermont. For instance, the degree and type of forest cover have a great influence on the species that occur in an area. Vermont has more forest today (75%) than it had in the mid-1800s (25%), and the effect of this change on wildlife has been dramatic. Additionally, Vermont's agricultural activity also affected the soils and the plants that grow in them.

Biophysical Regions of Vermont

The five factors described previously combine to create nine distinct biophysical regions. It is important to consider Vermont's biophysical regions when assessing and planning for the conservation of wildlife (Fig 2.1). For example, what may be a common species in one biophysical region may be rare in another, thus, increasing the importance of conserving habitat for that species in the region in which it is rare. Vermont's biophysical regions are described below.

Northeast Highlands: Granite bedrock dominates this cool region, which is characterized by large wetlands, remote mountains, and lakes and ponds. Spruce and fir dominate the lowlands as well as the high elevations, whereas northern hardwoods cloak the mid-elevations. Forty-three percent of this region is conserved, the highest percentage of any of Vermont's biophysical regions.

Northern Vermont Piedmont: Calcium-rich soils combine with a cool climate to support mixed forests and Northern White Cedar Swamps, Fens and other interesting natural communities in this region. The uplands have fine agricultural soils, but a short growing season. Eight percent of the region is conserved.

Southern Vermont Piedmont: Calcium-rich soils and rolling hills make this a good place for agriculture. The climate is average for Vermont, except in the extreme southeast where it is quite warm. Northern hardwoods and red oak dominate the vegetation. Seven percent of the region is conserved.

Southern Green Mountains: A broad plateau is dotted with a few dominant peaks and several ski areas. Climate is cold and rainfall is relatively high. Northern hardwoods, spruce, and fir dominate, and there are a number of small lakes and ponds. Thirty-three percent of this region is conserved.

Northern Green Mountains: This area has a cool climate and high elevations and is mostly forested. Northern hardwoods dominate the side slopes, whereas high elevations




Fig 2.1: Vermont Biophysical Regions

Northeastern Highlands
Northern VT Piedmont
Southern VT Piedmont
Southern Green Mountains
Northern Green Mountains
Champlain Hills
Champlain Valley
Taconic Mountains
Vermont Valley

have spruce and fir as well as Alpine meadow habitat. Twenty-six percent of the region is conserved.

Champlain Hills: This region is transitional between the Champlain Valley and the Northern Green Mountains. Northern Hardwood Forests dominate on the low hills, but oak-pine forests extend up the major river valleys where there are warmer conditions.

Champlain Valley: This region of Vermont has a warm climate and abundant fertile farmland. The Champlain Valley contains both northern hardwood forest and also various species of oaks and hickory. It has some of the state's most significant natural diversity and also the state's most densely populated areas. Nine percent of the region is conserved.

Taconic Mountains: The slate belt of Vermont and New York is found in this region. The Taconics are dramatic wooded hills dominated by sugar maple, beech, and yellow birch forests. Dry oak and hickory forests are found on the lower elevation knolls, while spruce and fir occur at the highest elevations. Ten percent of the region is conserved.

Vermont Valley: The Marble Valley has marble and limestone with glacial deposits on the valley walls, abundant springs, and wetlands. About 10 percent of the region is conserved.

Vermont's Landscape—an Historical Perspective

Vermont's landscape has long been altered by people. Native cultures grew crops, harvested animals for food and clothing and lived in established settlements. During the 17th and 18th centuries, land was cleared for the development of agricultural economies. By the mid-1800s, 75% of Vermont's forests were cleared for agriculture, and in particular, sheep farming. These changes had an effect on the state's waters, forests, and wildlife. Even some wildlife species, such as beaver and deer, that had been common, nearly disappeared from the land. As other influences caused people to begin to move towards the western United States, lands were abandoned and forests began to regenerate.

With the return of the forest and the work of the Vermont Fish & Wildlife Department and partners the recolonization and reintroduction of animal species, Beaver, White-tailed Deer, Wild Turkey, Fisher, and others that had declined have now returned and are today abundant. These species and others stand as great testament to Vermont's commitment to wildlife conservation and the resiliency of the forests and wildlife. Many species of fauna and flora, however, have not recovered. The Passenger Pigeon, for instance, is now extinct, and some large predators such as wolves and mountain lions that once roamed the New England forests, are no longer present.

Vermont's Contemporary Land Use

Agriculture and forestry still support Vermont's economy in significant ways. These elements of Vermont's business and economic communities offer great opportunities for wildlife conservation because they allow private landowners to realize a financial return from their land while keeping the land in an undeveloped or natural condition. Many of these land-based business interests are excellent stewards of the land and wildlife.

Vermont non-industrial forestland owners have a long history of active engagement in the management of forest resources throughout the state. Since the advent of the [Vermont Use Value Appraisal Program](#) (a.k.a. Current Use Program) 16,000 landowners have brought more than 1.8 million acres of forestland under forest management. A 2008 update to the program now allows landowners to enroll lands as Ecologically Significant Treatment Areas rather than for timber production, if they support natural communities and wildlife habitats of statewide significance; rare, threatened, and endangered species; some riparian areas; vernal pools with amphibian breeding habitat; forested wetlands; and old forests. More than 2,800 acres are enrolled as Ecologically Significant Treatment Areas.

Many of Vermont forestland owners manage their lands for wildlife and forest resources and seek to enhance their management skill through their involvement in non-profit organizations advocating sustainable forest management such as [Vermont Coverts: Woodlands for Wildlife](#), [Vermont Woodlands Association](#), [Vermont Family Forests](#) and [Audubon Vermont's Forestry for the Birds](#) program. These stewards provide strong examples of Vermonters taking steps to conserve our wildlife resources.

Based on data from the U.S. Census Bureau's 2014 the population estimate, 626,562 people live in Vermont, an increase of 0.1% since 2010. While Vermont is considered the most rural of northeastern states, there is an increasing demand for residential and commercial development. The rate of development in Vermont is double the rate of population growth (Center for Rural Studies 2008) and this growth is occurring mostly in rural communities. Like other New England states, residential development is often dispersed in rural and suburban areas rather than in existing village and urban communities.

Parcelization describes the subdivision of land into smaller and smaller forested area and multiple ownerships--typically by roads, agriculture, utility corridors, subdivisions or other human development. This phenomenon drives habitat fragmentation that has been shown to reduce forest and habitat health, water quality and the sustainability of local forest product economies (VFPR 2015). With Vermont's population projected to increase by 85,000 in the next 15 years, the Vermont Forests, Parks & Recreation Department is leading efforts to balance population growth with Vermonters' interests in maintaining healthy forests and wildlife habitat, clean water and the state's traditional settlement pattern of village centers surrounded by fields, farms and working forests (VFPR 2015). Since 2006 the Vermont Natural Resources Council has hosted the [Vermont Forest Roundtable](#), a venue for information exchange among consulting foresters, professional planners, state agency officials (including VFWD and VFPR), landowners, sportsmen, forest products industry representatives, conservation groups, biomass energy organizations and academics. Addressing parcelization drivers has been a primary Roundtable focus.

Contemporary Problems and Threats to Vermont's Wildlife

The problems most frequently identified as impacting SGCN are loss of habitat (due to conversion, degradation, fragmentation and lack of needed successional stages), the impacts of roads and transportation systems, pollution and sedimentation, invasive species and climate change. Additionally, wildlife diseases and the decline of pollinators have been gaining greater attention in both the scientific community and among Vermonters in general. Each is summarized here:

Loss of Habitat: Due to Degradation, Conversion, Fragmentation or Lack of Needed Successional Stages

These four categories are not mutually exclusive and problems can often logically be placed into more than one category depending on the particular stress it causes for a species or habitat.

Habitat Conversion: The complete transformation or loss of a habitat by human action (examples include: filling a wetland to create a grassy field, converting a forest stand into a parking lot, or damming a stream to create a reservoir). Though many agencies and organizations work diligently to conserve important wildlife habitats, Vermont continues to lose approximately 480 acres of habitat each year to regulated development alone. According to the Vermont Environmental Board, regulated development in Vermont constitutes approximately one-third of the total development that occurs on an annual basis. Important habitats addressed by various statutes, largely Act 250, and include deer winter habitat, wetlands with significant wildlife functions, habitat for rare, threatened and endangered species and several types of habitat necessary for the survival of black bears. These habitats represent only a few of the many habitats that are affected by loss due to development.

Habitat Alteration/Degradation: A lessening of the quality of a habitat by human action stopping short of complete conversion (examples include: the reduction of mast (fruit and seed) production in a forest stand, riprapping a streambank, and significant land use changes adjacent to a habitat such as replacing a forest stand on the edge of a wetland with a housing development).

Habitat Fragmentation: The breaking up of habitats into smaller, non-contiguous patches as a result of habitat conversion (e.g., housing, commercial development, roads, utility lines). Fragmentation can: 1) render important habitats inaccessible (such as isolating a den site from a feeding site), 2) isolating populations (for example grassland butterflies, spotted salamander, and tiger beetles); and, 3) degrade remaining habitat patches through edge effects that favor edge-tolerant species such as raccoons and crows, as well as invasive exotic species that can out-compete native and rare species. The result of habitat fragmentation is often increased predation, increased mortality, reduced mobility and changes in habitat micro-climates. In the past decade several research into fragmentation and efforts to maintain and restore habitat connectivity have increased significantly in Vermont and elsewhere. It's also a problem for forestry and rural economies (VFPR 2015).

Inadequate Distribution of Successional Stages: The lack of either late, mid or early successional habitat in appropriate patch size and/or juxtaposition can be a problem for some SGCN especially as fragmentation makes it harder for species to move between forest patches (examples include ruffed grouse and woodcock which prefer early successional forest stands, American marten which prefers late-successional stands and Canada Lynx and Snowshoe Hare which depend on a mix of forest stages).

Climate Change

Long-term changes linked to global warming and other climate issues are expected to drive major changes in habitat availability (e.g., high elevation habitats, wintering areas and

migration stopovers) (Glick 2005), vegetative composition and location (e.g., the movement up in elevation or north in latitude, invasion by exotic pests), climate variability (e.g., change in snow depth, rainfall and/or natural disturbances). Many specific details as to how climate change is affecting Vermont's wildlife today remain unknown, but the pervasiveness and scale of the problem requires that we begin planning to address it now. Chapter three of this Wildlife Action Plan dives deep into the Climate Change threat.

Impacts of Roads and Transportation Systems

The number of vehicle miles traveled by Vermont residents is growing at seven times the rate of population growth, according to information from the Vermont Agency of Transportation (VTrans) (1999). Transportation systems can cause numerous problems for SGCN including: vehicle-wildlife collisions; reducing animal and fish passage, thus limiting habitat availability and isolating populations; vehicle emissions of pollutants such as ozone and greenhouse gases; and facilitating the spread of an exotic, invasive species into otherwise healthy areas. To address problems such as these, VFWD and VTrans established a joint Wildlife-Transportation Steering Committee in 2007 to guide and support interagency cooperation to make Vermont's transportation system safer for both people and wildlife. VTrans is now considered a leader nationally among transportation agencies in road ecology.

Pollution & Sedimentation

The introduction of exotic materials from point and non-point sources can significantly impact SGCN, particularly aquatic species. Pollutants and sediments include sands and silts, chemicals and toxins; excess nutrients from farm and municipal sewage plants; garbage and other solid waste; radioactive materials; road salt; excessive noise; excessive heat; and light pollution that disturbs animals and disrupts migration patterns. Sediments can be a problem for SGCN through their physical presence alone. For example, soils can wash into a stream from a construction site and smother fish eggs and other aquatic species living in the spaces between rocks and gravel streambed. The history of polluted runoff—rainwater and snowmelt draining from parking lots, roads and streets, logging roads, farm fields and croplands, and lawns—carrying sediments and nutrients, including phosphorus, into streams and rivers and directly into Lake Champlain is stimulating excessive growth of algae which is bad for lake species and people. The status of efforts to reduce the amount of pollutants getting into the lake can be found at [Restoring Lake Champlain](#).

Invasive Exotic Species

The introduction and spread of nuisance exotic and native species (plants and animals) may lead to the elimination of native wildlife populations, threaten long-term stability of habitats and even lead to extirpation by out-competing a native species, displacing its food source or altering a key process or function of a habitat. Invasive exotic species in Vermont include Eurasian watermilfoil, purple loosestrife, common buckthorn, Japanese knotweed, Morrow's honeysuckle, goutweed, black swallow-wort, hemlock wooly adelgid and zebra mussels.

Since the 2005 Wildlife Action Plan was released, the ecological and economic problems caused by invasive exotic species have gained much more attention in Vermont. A clearinghouse of information and resources to help combat the introduction and spread of these species is now available at [VermontInvasives.org](#). And, the state has an [Invasive Forest Pest Action Plan](#) ready for the eventual arrival of forest pests, including the Asian long-

horned beetle and emerald ash borer, in order to prevent the establishment and/or limit their spread within Vermont ([VFPR/VAAFM 2014](#)).

Diseases

Shortly after the first Wildlife Action Plans were submitted, White-Nose Syndrome (WNS) was discovered in New York and then Vermont in 2008 followed quickly by several other states. In a few short years WNS nearly wiped-out several bat populations in the Northeast. The Little Brown and Northern Long-eared Bats, decimated by the disease were added to Vermont's endangered species list and the Northern Long-eared bat was listed federally as threatened in 2015. The threat caused by WNS was added as an 'emerging issue' to the Action Plans of several states, including Vermont's. Subsequently, Snake Fungal Disease was discovered killing Rattlesnakes in Vermont and elsewhere, and *Heterosporis* appeared in Yellow Perch in Lake Champlain. Additionally, diseases not yet found in Vermont are potential threats. They include: Chronic Wasting Disease (deer), Viral Hemorrhagic Septicemia (fishes), *Batrachochytrium salamandrivorans* (salamanders) and Avian Influenza (birds). The Vermont Fish & Wildlife Department has steadily increased surveillance and is devoting additional resources to the threats posed by these and other diseases.

The Vermont Fish & Wildlife Department developed a fish health and biosecurity program to reduce the risk of disease importation and spread that includes strict regulations for fish importation, the prohibition of live transfer of fish from one body of water to another, strict baitfish use regulations, the use of water filtration and disinfection equipment at fish culture station water source(s), regular use of disinfectants at fish culture stations and health inspections of both naturally produced fish and fish produced at fish culture stations. The Department works in close coordination with its counterparts in the northeast states through the [Northeast Fish Health Committee](#) to develop and share procedures and protocols to maintain fish health and reduce the threat and movement of fish diseases.

VFWD also strictly controls access to bat hibernacula to prevent the spread of WNS and has regulations governing the movement of rabies vector species as well as rules to help prevent the introduction of Chronic Wasting Disease into the state. The VFWD, collaborates regionally through the [Northeast Wildlife Disease Cooperative](#), a consortium of states and veterinary diagnostic laboratories providing wildlife diagnostic services, expertise, training, and research support to state and federal wildlife agencies in the region.

Pollinators

While the 2005 Wildlife Action Plan included 33 species of butterflies and moths, many of them pollinators, their role as pollinators was not critical to their selection as SGCN. Since 2005, concerns over the status of pollinators in general (e.g. flies, wasps, moths, butterflies, beetles, bees, bats and hummingbirds), and native bees in particular, has become a worldwide concern. Pollination is defined as a mutually beneficial relationship between plants and pollinators wherein the plant provides pollen and/or nectar to the pollinator and the pollinator provides reproductive services for the plant (National Research Council, 2007). Roughly 75 percent of the 240,000 species of flowering plants world-wide rely on pollinators for flower reproduction (NRC, 2007). This includes many plant species that provide browse or forage for larger wildlife, as well as seeds and fruits to support birds and small mammals. These invertebrates also pollinate many commercial crops. In Vermont this includes blueberries, tomatoes, squash, apples, and other produce. The many drivers of

pollinator declines include habitat loss and degradation, intensive agricultural practices, use of certain pesticides, diseases and pathogens (Heinz Center, 2013). For this second Wildlife Action Plan nine bumble bee species and 31 species of butterflies and moths—including the Monarch butterfly—were selected as Species of Greatest Conservation Need.

Bad News & Good News

The preceding summary of threats is sobering to say the least. It challenges everyone concerned about wildlife and wildlife habitat to think smarter and work harder and more creatively. The good news here is that we can focus our limited conservation resources on the strategies that will provide the biggest bang for the buck, such as those outlined in this report that come up again and again including habitat restoration, encouraging wildlife-compatible resource use, providing education and technical assistance to landowners and managers, and providing economic incentives for conservation

But to do justice to this Action Plan, and to help the many SGCN and habitats in need, we need to add one problem to the list of major issues impacting Species of Greatest Conservation Need that our technical teams did not identify directly in their assessments, but it was often discussed during team meetings—the lack of sufficient funding for wildlife conservation. Without sufficient funding we will not be able to implement many of the conservation strategies identified in this report. The State Wildlife Grants program is a critical first step in funding SGCN conservation, but more is needed. And, to make the most of SWG funds, Vermont will have to develop the required matching stateside funds.

Conservation Success!

Keeping Common Species Common

In spite of the changes to the Vermont landscape, the fact is, Vermont remains a relatively rural state with an abundance of conserved land, private landowners who are excellent stewards of the environment, and many wildlife conservation success stories. The public opinion survey results (U.S. Fish & Wildlife Service 2001) speak volumes for the bright future of wildlife conservation in Vermont—that is, the public has a strong interest in and support for the conservation of Vermont’s natural heritage.

Moreover a review of past and ongoing wildlife conservation efforts provides proof of our collective ability to recover and conserve wildlife and the habitats required for their survival. It also identifies the key building blocks for successful conservation.

In 1724, when the first European settlement was established at Fort Dummer, near Brattleboro, the state was primarily forested and had abundant fish and wildlife populations including Passenger Pigeons, fisher, wolves, deer, black bear, beaver, and salmon. However, by 1865 many of these species would be present in far fewer numbers or on the cusp of extirpation because of unregulated harvests, habitat loss and habitat degradation.

Hunting and fishing license fees, soon after the turn of the 20th century, coupled with federal wildlife and sportfish restoration act dollars, enacted in the 1930’s and 1950’s respectively, established a financial framework in support of conservation. These monetary resources enabled Vermont, and the other states, to conduct inventories and research, acquire habitats, and provide conservation education to the public. Today, some of the species of low abundance 150 years ago are now once again common throughout the State. Consider, for example:

White-tailed Deer: Numbers were so low in the late 1800’s that no open season was offered and deer were transported from New York. Through extensive research, harvest management, and habitat protection, Vermont can now support in excess of 150,000 deer with 48 days of hunting opportunity, annually.

Wild Turkey: This bird was extirpated from the state in the 1800’s. Birds were reintroduced to the state in 1969. We now have more than 40,000 Turkey and both fall and spring hunting opportunities.

Fisher: This mid-sized carnivore was extirpated from the state. Animals were reintroduced to Vermont beginning in 1959 (to help control porcupine), and this predator now thrives on the Vermont landscape.

Anadromous fish on the Connecticut River: Migratory fish in the Connecticut River, including Atlantic Salmon, American Shad, Striped Bass and River Herring were reduced or eliminated in 1798 by a dam built in Turners Falls, Massachusetts. With the construction of fish passage at dams, and active restoration programs Shad, Stripers and Herring are now abundant in the lower river.

Trout and salmon in Lake Champlain: Landlocked Atlantic Salmon disappeared from Lake Champlain in the 1850's, and native Lake Trout were gone by 1929. A restoration program was begun in the 1970's in cooperation with the State of New York and the US Fish and Wildlife Service, and these fish are plentiful once again in Lake Champlain where they support a popular fishery that brings hundreds of millions of dollars into the regional economy each year.

Lake Sturgeon: A combination of dam construction, pollution and over-fishing reduced Lake Sturgeon populations in Lake Champlain in the early 1900's to the point that the commercial fishery was abandoned and all fishing for sturgeon was prohibited in 1967. Since this fishing closure Sturgeon conservation has benefited from water quality improvements, better water flows at the dams, and outreach to anglers to release any sturgeon they catch. Recent studies have documented successful natural reproduction of sturgeon in all four of their historic spawning rivers in Vermont.

Peregrine Falcon, Osprey, and Common Loon: These birds were gone or nearly gone from the state by the mid-1900's. Through focused management (e.g., the construction of artificial nesting platforms, water level management, banning use of the pesticide DDT and public education), each of these three species has recovered sufficiently that they've recently been removed from the state's endangered species list—a first for any species in Vermont.

Bald Eagle and Common Tern: Listed as endangered in Vermont, the known nesting population of Bald Eagles in Vermont has grown from zero in 2007 to 18 nesting pairs with 17 fledglings in 2014. Vermont's Common Tern population dropped from 300-400 breeding pairs in the 1980s to approximately 50 in 1988 when it was listed as endangered. With monitoring, management and the protection of nesting islands their numbers have increased since then. Breeding numbers have recently exceeded the levels recommended for down-listing to Threatened (but continuing low productivity delays down-listing).

These success stories suggest that new dollars coupled with patient and persistent efforts can produce new success stories for the future. In other words, the fish and wildlife profession has demonstrated the will and the competence to restore and manage wildlife. The Wildlife Action Plan, coupled with sustained funding and the dedicated participation of partners, will offer a template for advancing the success stories to a new suite of species.

The Importance of Education, Law Enforcement and Recreation to Wildlife Conservation

Through the State Wildlife Grants program (SWG) Congress provides every state with critically needed funds for wildlife conservation. Congress' intention is to support proactive and strategic efforts to prevent future Endangered Species Act listings—in other words, to keep common species common. To meet Congressional intent, states are compelled to employ all of their best conservation tools including education, wildlife-associated recreation and the creation and enforcement of wildlife protection laws and regulations. These are among the most proactive, strategic and time-tested tools in any conservation tool box.

The details of the SWG program legislation, however, currently preclude states from using SWG funds for law enforcement and recreation projects. A limited amount of SWG funds can be used for conservation education, but only in a supporting role in the implementation of a conservation strategy (e.g., signage explaining the purpose of a restoration project). This poses a dilemma for states trying to implement a truly comprehensive wildlife action plan because it restricts their use of three vital conservation tools. Moreover, it limits the participation of three significant conservation constituencies from participating in Wildlife Action Plan implementation—the law enforcement, education and outdoor recreation communities.

A limited number of education and law enforcement conservation strategies specific to particular species or habitat categories were addressed in the species and habitat conservation summaries of this report (Appendices A and B). We recognize that alternative funding sources are needed for their implementation. In this section of the Wildlife Action Plan report we present additional conservation strategies based on conservation education, wildlife-associated recreation and law enforcement. It is our hope that future renderings of the State Wildlife Grants program, along with other funding mechanisms, will more fully provide for the implementation of these strategies.

Conservation Education

Wildlife and human communities depend on healthy ecosystems and ecological processes. Their functions are essential for our quality of life and for the Vermont economy. Conservation strategies that follow a sound education model can foster healthy public behavior and attitudes toward land and wildlife conservation. Furthermore, strong educational programs that expand Vermonters' ecological literacy will enhance the credibility and effectiveness of other conservation efforts and build support for future efforts. Finally, the public plays a key role in influencing legislators, who in turn affect policy and funding decisions. Recommended strategies include:

- Foster and enhance educational partnerships to maximize efficiency (e.g., develop volunteers, outreach to teachers and youth group leaders to deliver programs)
- Ensure that sound messages, curricula, and best educational practices are followed to maximize our efforts (e.g., provide teacher training, curriculum support materials for teachers and students.
- Define a land stewardship message that promotes the conservation and ethical use of Vermont's fish, wildlife, and plants, and the habitats that sustain them.
- Focus outreach and education efforts to enable the public to make informed decisions on issues affecting ecosystems in Vermont such as: habitat degradation and fragmentation, threats to fish

and wildlife species and their habitats, the value of working rural landscapes and other rural lands, and the sustainable and ethical utilization of wildlife.

The connection of education to wildlife conservation is recognized nationwide. The Association of Fish and Wildlife Agencies developed the [North American Conservation Education Strategy](#) in 2010 to help make improve conservation education programs nationwide.

Wildlife-Associated Recreation

Hunting, fishing, trapping, and wildlife viewing have a long heritage in Vermont and Vermont leads the nation in wildlife viewing (U.S. Department of the Interior et al. 2011). By providing the means for more people to connect with wildlife, we can foster more and stronger relationships to the natural world. Applying the concept of stewardship through recreation Vermonters can become knowledgeable about and appreciate wildlife, natural communities, and conservation in ways that promote citizen interest in contributing to conservation. Recommended strategies include:

- Work with the broader community of recreation groups (e.g., outdoor guides, birders, sportsmen and women, hikers, paddlers, climbers, spelunkers, mountain bikers and snowmobile and ATV associations) to foster partnerships that build a stronger wildlife ethic among members.
- Expand educational programs on watchable wildlife, including such topics as birding, wildlife photography, animal track identification, and backyard habitat. Target population centers, with a focus on youths and families.
- Increase information available to the public on how and where to watch wildlife. Provide information to encourage watchable wildlife practices, such as viewing, photographing, and feeding, in a manner that is ethical, safe, and consistent with protecting the welfare wildlife resources.
- Foster a recreational ethic based on the concept of giving back to the natural world.
- Include an educational component in recreation activities making the connection between our actions and the impact on wildlife.
- Involve Vermonters in activities that will increase their understanding of wildlife and land stewardship and the influences of human activities on wildlife, in order to build public support for fish and wildlife conservation (e.g., citizen science projects such as the bird atlas, butterfly survey and other wildlife inventories, teacher training courses, streambank plantings, and field classrooms).
- Encourage responsible outdoor recreation through programs such as "Stop Aquatic Hitchhikers," "Leave No Trace," "Stay on the Trails," and "Be Bear Aware."

Statewide Comprehensive Outdoor Recreation Plan: In addition to the Wildlife Action Plan, states develop comprehensive plans for outdoor recreation as a requirement for receiving support from the federal Land and Water Conservation Fund (LWCF). The [Statewide Comprehensive Outdoor Recreation Plan](#) (2014-2018) developed by the Vermont Forests, Parks & Recreation Department helps guide efforts provide high quality outdoor recreation facilities, opportunities and experiences for Vermonters and visitors.

Law Enforcement

The creation and enforcement of fish and wildlife laws are among our society's oldest attempts to conserve wildlife. Vermont's first game wardens were appointed in 1779 to protect deer and were called "Deer Reeves." Law enforcement is an effective conservation tool and has been at the core of wildlife conservation ever since.

State game wardens prevent the illegal taking, trade, sale, collection and importation of wildlife by proactive enforcement of fish and wildlife laws. Game wardens also prevent and investigate the unlawful destruction of important habitat, trespass and disturbance of refuge areas and sensitive breeding grounds and enforce the regulations and permits that govern wildlife research, education and rehabilitation.

Law enforcement professionals strive to be proactive: Game wardens are an integral part of the Fish & Wildlife Department's outreach and education programs. Wardens teach conservation at schools, civic organizations and conservation camps and are often the first, and sometimes only, contact that the general public has with a conservation professional. Recommended strategies include:

- Maintain staffing of game wardens statewide sufficient to ensure the adherence of all laws pertaining to fish, wildlife and habitat conservation.
- Continue to conduct routine patrols, incidental to core duties, providing enforcement of boat, ATV and off road recreational vehicles to address the illegal operation and destruction of sensitive habitat and wildlife areas.
- Review, update, and enforce regulations controlling the importation and possession of exotic and potentially harmful fish and wildlife species and their pathogens.

Recent projects

As people interact more and more with wildlife, the number of wildlife-human conflicts increases. In 2014 Vermont game wardens responded to more than 500 calls from the public on issues such as rabies and damage to property. If not adequately addressed, members of the public might try to resolve the issues themselves in a manner unduly detrimental to wildlife. Many encounters require a physical response by a warden to prevent human injuries or disease exposure.

All for one and one for all: Law enforcement, Education & Recreation

It should be clear to a reader by this point that not only is each of these three tools critical to the long-term conservation of wildlife, but that they are all tightly intertwined. For example, our best opportunities to instill the message of conservation in the public are when they are out in nature recreating. And, state game wardens are often the ones to deliver the message. Furthermore, outdoor guides and other recreationalists often provide tips to wardens and compliance officers regarding habitat degradation or the illegal taking of wildlife, and by doing so they send a strong message to the general public that Vermonters care about wildlife.

Literature Cited

- Ecological Society of America's Committee on Land Use. 2000. Ecological Principles for Managing Land Use. Ecological Society of America, Washington, DC
- Forman, Richard T. 1995. Land Mosaics, The Ecology of Landscapes and Regions. Cambridge University Press. New York.
- Glick, P. The Waterfowlers' Guide to Global Warming. National Wildlife Federation. Washington, DC.
www.nwf.org/globalwarming/pdfs/waterfowlbook.pdf
- Hammond, F.M. 2002. The Effects of Resort and Residential Development on Black Bears in Vermont. Final Report. Fish & Wildlife Department, Agency of Natural Resources. Waterbury, Vermont Press.
- National Invasive Species Council 2005. National Agricultural Library, U.S. Department of Agriculture.
<http://www.invasivespecies.gov/>
- National Research Council. 2007. Status of Pollinators in North America. Washington, D.C.: The National Academies Press.
- The Heinz Center. 2013. Pollinators and the State Wildlife Action Plans: Voluntary Guidance for State Wildlife Agencies. Washington, DC.
- U.S. Department of the Interior, U.S. Fish and Wildlife Service, and U.S. Department of Commerce, U.S. Census Bureau. 2011 National Survey of Fishing, Hunting, and Wildlife-Associated Recreation
- Vermont Forests, Parks & Recreation Department. April, 2015. [Vermont Forest Fragmentation Report: Report to the Vermont Legislature](#), Submitted to House and Senate Committees on Natural Resources and Energy and the House Committee on Fish, Wildlife, and Water Resources.
- Vermont Forests, Parks & Recreation Department and Vermont Agency of Agriculture. 2014. [State of Vermont 2014 Invasive Forest Pest Action Plan](#).

Vermont's Wildlife Action Plan

Chapter 3 Climate Change & Conservation

Draft • September, 2015

Introduction	2
Historic Trends and Future Projections	3
Vermont's Changing Seasons.....	4
Impacts of Climate Change on Vermont's Ecology.....	5
Forests.....	5
Waterbodies.....	7
Wetlands	8
Habitats and Species	8
Conserving Wildlife in the Face of Climate Change.....	10
Regional Efforts to Conserve Wildlife in the Era of Climate Change	10
Efforts to Address Climate Change Impacts in the Northeast.....	11
Vermont Efforts to Understand and Address Climate Change Impacts to Species and Habitats	12
Recent Projects Helping Address Climate Impacts to Species and Habitats	17
Identifying Climate Adaptation Strategies for Vermont's Species and Habitats.....	20
Priority Actions to Address Climate Change Impacts to Species and Habitats	20
Managing Vermont's Wildlife and Habitats into the Future	21
Literature Cited	22

3. Climate Change & Conservation

Introduction

Ten years ago when states across the nation were completing their Wildlife Action Plans, wildlife managers were just beginning to think through basic questions such as: ‘How might climate change impact wildlife and wildlife habitat? ‘How can we systematically identify wildlife, plants and habitats that might be vulnerable to climate change? And, ‘What types of actions can be taken to help wildlife survive climate change?’

Ten years later, many states, including Vermont, have begun identifying how wildlife and habitats might be affected by climate change, which are most vulnerable, and how to best manage for the future. There are few clear or simple answers. Identifying vulnerable species and habitats requires teasing out myriad factors that climate change could influence, such as changes in precipitation rates, snow pack and soil moisture; changes in the number of freezing days; new diseases and invasive species; flooding, lack of flooding; movement of species and their habitats, and changes in predator-prey and plant-pollinator relationships. And, we don’t know the capacity of species to respond to a changing climate.

An honest response to these very complicated questions is ‘we don’t know how most species will fare in the coming decades.’ We can assume there will be winners and losers. Bicknell’s Thrush and other alpine plants and animals found only at the tops of Vermont’s highest peaks, for example, are likely to be losers here (though they may fare better north of here). Other species, such as the Virginia Opossum and Tufted Titmouse, which have been slowly moving north, might do better in Vermont in the coming decades.

What to do? We need to remain vigilant, expect to be surprised, foster adaptability and reduce other stresses on wildlife and wildlife habitat. In the short-term, riparian areas (the banks of rivers and streams) and adjacent land, including their floodplains, may be the best places to invest our conservation efforts because when healthy these areas are more resilient to flooding, reduce downstream flood impacts, keep waters cool and provide important habitat and connectivity to many wildlife species. Table 3.3.1 lists actions that are generally most important if we want to help wildlife in the coming decades.

Table 3.1. General adaptation goals to inform the identification of specific strategies that serve to increase the resiliency and/or adaptive capacity of wildlife and their habitats (Stain, B. et al. 2014).

Note: There is significant debate over the pros and cons of assisted migration (relocating organisms).

Adaptation Strategy	Definition
Reduce non-climate stresses	Minimize localized human stressors (e.g., pollution) that hinder the ability of species or ecosystems to withstand or adjust to climatic events
Protect key ecosystem features	Focus management on structural characteristics (e.g., geophysical stage), organisms, or areas (e.g., spawning sites) that represent important “underpinnings” or “keystones” of the current or future system of interest
Ensure connectivity	Protect, restore, and create landscape features (e.g., land corridors, stream connections) that facilitate movement of water, energy, nutrients, and organisms among resource patches
Restore structure and function	Rebuild, modify, or transform ecosystems that have been lost or compromised, in order to restore desired structures (e.g., habitat complexity) and functions (e.g., nutrient cycling)
Support evolutionary potential	Protect a variety of species, populations, and ecosystems in multiple places to bet-hedge against losses from climate disturbances, and where possible manage these systems to assist positive evolutionary change
Protect refugia	Protect areas less affected by climate change, as sources of “seed” for recovery (in the present) or as destinations for climate-sensitive migrants (in the future)
Relocate organisms	Engage in human-facilitated transplanting of organisms from one location to another in order to bypass a barrier (e.g., urban area)

The remainder of this chapter is a deeper exploration of the climate change threat. It includes a summary of historic climate trends and projections of future climate; a look at the ecological impacts of climate change on species, forests wetlands and aquatic habitats. It also delves deeper into efforts to conserve wildlife in the face of climate change both in Vermont and regionally. This chapter concludes with a list of conservation strategies to help Vermont's wildlife, plants and wildlife habitat.

Taken in a broader context, this chapter will help to illustrate the interconnection between climate and non-climate stressors and conclude with a list of conservation strategies to help Vermont's wildlife, plants, and habitat.

From the waters of our lakes and rivers to the forests and wetlands abutting our communities, Vermont's diverse ecosystems are essential to a healthy and sustainable future for wildlife and people.

Vermont's fish, wildlife and plants and their habitats are already responding to climate change. Plants are leafing out and blooming earlier; birds, butterflies, amphibians, and other wildlife are breeding, feeding, metamorphosing or migrating earlier; and many species that are able to migrate are shifting ranges northward and to higher elevations (Betts, A. K. 2011a, Stager, C., and Thill, M. 2010, U.S. Global Change Research Program 2009). Of particular concern is the potential for entire ecosystems to be disrupted. As diverse species and habitats in Vermont respond to climatic fluctuations in different ways, important inter-specific connections—such as between pollinators and the flowers they fertilize, or breeding birds and the insects on which they feed—will likely be disrupted. Further, the ecological impacts associated with climate change do not exist in isolation, but combine with and exacerbate other stresses on our natural systems. For instance, although climate pressures may be causing species ranges to shift, development and roadways have created a matrix of inhospitable habitat that may inhibit such movement. And while invasive species already have a major negative impact on many ecosystems in Vermont, many invasives may be favored under future climate conditions, making it even more difficult for native species to adjust and survive under new climatic regimes. These are just a few examples of the challenges that species and habitats in Vermont face under climate change.

Historic Trends and Future Projections

Since 1895, the average annual temperature in the U.S. has increased between 1.3°F and 1.9°F. Over the next 30 to 40 years, temperatures are expected to rise on average another 2°F to 4°F across most of the nation. By the end of the century, should carbon emissions continue to rise at the current rate (higher emission scenario), the U.S. can expect a rise in average annual temperature between 5°F to 10°F. On the other hand, if significant reductions in carbon emissions can be achieved (lower emission scenario), average annual temperature across the U.S. would rise approximately 3°F to 5°F by the end of the century (Melillo, J.M. et al. 2014) (Fig 1).

While the last decade has been the hottest on record, it is evident that there is significant regional variation - with northern latitudes experiencing greater warming. Therefore, the Northeast can expect a greater rise in average annual temperature. Depending on the different emission scenarios, temperatures in the Northeast are projected to increase by either 4.5°F to 10°F (under a higher emission scenario) or 3°F to 6°F (under a lower emissions scenario) by the 2080's (Horton, R. et al. 2014).

In Vermont, average annual temperatures have increased 1.6°F since 1960 and 0.9°F since 1990, with temperature increases even more significant in recent decades. For example, mountainous highland regions in Vermont have experienced average annual temperature increases of 1.8°F per decade from 1990-2012 and 2.5°F from 2000-2010. In addition, average annual winter temperatures are rising twice as fast as average summer temperatures. These trends are resulting in significant changes to Vermont's climate (Galford, G.L. et al. 2014).

Vermont's Changing Seasons

As temperatures continue to climb, the severity of Vermont's winters are decreasing. Compared to the 1940s-1960s, Vermont had a 20% decrease in the number of days below freezing in the last decade alone. As a result, lakes and ponds are experiencing an average of seven fewer freezing days per decade. Simultaneously, average precipitation rates have also increased in Vermont. For example, since 1960, decadal precipitation averages have increased by 0.9" in lowland areas and 2.3" in highland regions. The result has been increases in snowfall amounts at higher elevations. However, the projections suggest that while snowfall amounts will increase over the short term (20-25 years), as temperatures increase over the long-term winter precipitation will increasingly come in the form of rain (Galford, G.L. et al. 2014).

"Warmer winters will lead to reduced accumulation of snow during some years. Less snowpack may mean less runoff during the late winter/early spring thaw. This effect may be offset by increased rains falling on frozen ground, leading to greater runoff. However, if winter temperatures rise to levels that decrease the duration of frost conditions by late in the century, runoff may be moderated by increases in soil infiltration – as long as soils do not become saturated by a rain event (Galford, G.L. et al. 2014)."




Figure 1. Estimated average rise in global temperature (relative to the 1901-1960 average) for the higher emissions scenario (A2) and lower emissions scenario (B1). A higher emissions scenario assumes a continued increase in emissions throughout this century, whereas a lower emissions scenario assumes a significant reduction. Shading represents the range (5th to 95th percentile) of results from a suite of climate models. Average temperatures are predicted to increase in both scenarios, however the difference between lower and higher emissions pathways is significant (Melillo, J.M. et al. 2014) (Figure source: NOAA NCDC / CICS-NC).

Overall, projections suggest more winter and spring precipitation over the coming century. “Average monthly flows in January and March, as well as July, August, and October through December, have increased while average monthly flows in April and May have decreased (Galford, G.L. et al. 2014).”

Moreover, high intensity precipitation events greater than one inch increased an average of four days between 1960-1980 and an average of 7-10 days a year over the past two decades. One of the outcomes of more frequent high-intensity rain events is that stream high flows are larger, more frequent, and are projected to occur more frequently in winter. An increased number of high-intensity rain events in conjunction with warmer temperatures will result in more flooding along Vermont’s rivers and shorelines. At the same time, rising temperatures over the longer term and increased seasonal variability in rainfall suggest a potential increase in the number of short-term summer droughts resulting in periods of very low stream base flows and lake levels, and slow groundwater recharge rates by centuries end (Galford, G.L. et al. 2014).

Impacts of Climate Change on Vermont’s Ecology

Significant deviations in the variability around historic climatic norms, including increased variability in temperature, precipitation and extreme weather events, has direct implications on the vulnerability of species and the habitats upon which they depend. In this context, vulnerability is defined as “the susceptibility of a species, system or resource to be negatively affected by climate change” and interactions with other non-climate stressors such as habitat degradation and habitat loss (Staudinger, M.D. et al. 2015). Climate change already presents a variety of challenges for species and ecosystems across the Northeast. These include the reduction in the “quality and distribution of habitats, the availability of food, increases in the abundance of parasites and diseases, and the increased incidence of stress from heat and drought (Rustad, L. et al. 2012).” But how do the challenges driven by climate change affect the ecosystems in Vermont? To explore this question, we’ll take a look at how Vermont’s species and natural communities, including forests, wetlands, waterbodies and other habitats are currently responding to climate change and how such communities are likely to shift in abundance, composition, and range in the coming decades.

Forests

Vermont is predominately a forested landscape so climate-driven impacts to forests are of significant concern for Vermont wildlife as well as the forest products industry, rural communities and the Vermont way of life. Forest communities provide a number of important and complex ecosystem services including protecting water quality, reducing runoff, nutrient cycling, capturing air pollutants, providing wildlife habitat, and carbon sequestration among others (Galford, G.L. et al. 2014, Rustad, L. et al. 2012). However, predicting the specific future impacts of climate change on systems comprised in large part of long-lived species is difficult. Historic pollen and microfossil data tell us that in the past forest systems migrated, albeit slowly, to changes in climatic conditions (Rustad, L. et al. 2012). Over the past 12,000 years northern forests once dominated with spruce and jack pine have transitioned into larger ratios of white pine, oak, and eastern hemlock, and then to beech and maple dominance in the north, white pine, hickory and birch dominance in southern areas, and spruce and fir at higher elevations (Rustad, L. et al. 2012). While evidence suggest that forests have been able to migrate in the past, models of predicted

climates, suggest that past migration rates are too slow to keep up with current and future changes climatic conditions (Rustad, L. et al. 2012).

Current evidence indicates that forests will increase evapotranspiration and water use due to warmer seasonal temperatures and an increase in the number of growing days (Rustad, L. et al. 2012). This scenario may lower overall soil moisture, increasing the persistence of droughts, reduce forest productivity, and in turn increase the susceptibility of forests to insect and disease outbreaks (Rustad, L. et al. 2012).

As a result, several dominant tree species may undergo significant range shifts as forest communities adjust to new conditions over time (fig. 3. 2) (Rustad, L. et al. 2012).




Figure 3.2. Current and projected suitable habitat for major forest types in New England under low and high emissions scenarios. See Figure 5 for details of the scenarios. Under the low emissions scenario, the conditions will favor maple- birch-beech forests, while under the high emissions scenario suggest that conditions they will favor oak-hickory forests. Adapted from Iverson et al. 2008 (Rustad et al. 2012).

Models predicting the ways in which forests will respond to long-term climate trends show that species such as sugar maple and balsam fir are likely to lose suitable habitat in Vermont, while several species of oak are projected to expand in the extent of suitable habitat (Rustad, L. et al. 2012). Moreover, shorter winters, longer growing seasons, and increased concentrations of CO₂ should result in an overall increase in forest productivity (net forest growth rate) (Rustad, L. et al. 2012). However, many factors, including life spans, dispersal rates, invasive species, and changes in soil moisture, will make it difficult to predict when, and if, tree populations will be able to reach predicted, suitable habitat (Rustad, L. et al. 2012).

Forest managers can play a critical role addressing these uncertainties by promoting healthy forests and their successful adaptation to climate change. According to the Vermont Department of Forests, Parks and Recreation (VFPR) (2015), human activity is one of the primary drivers of change in Vermont's forested communities. Therefore, influencing change through forest management in ways that improve forest resiliency could increase future suitable habitat.

Table 3.2. Tree decline and associated climate factors. A review of decline episodes for five different tree species in the Northeast indicates that there have been important associations with changes in climate and weather-related conditions, which may be further exacerbated as climate changes in the future. Adapted from Mohan 2009 (Rustad et al. 2012).

Species/ Group	History	Role of Climate	Other Factors	References
Birch	Widespread declines since 1944	Maps of birch decline areas coincide with areas of experiencing extended winter thaw cycles	None	Balch 1944 Bourque et al. 2005 Braathe 1995
Sugar maple	26 widespread decline episodes between 1912 and 1986	Prolonged thaw-freeze events and associated fine root damage have been implicated in sugar maple decline	Insects, disease, loss of soil nutrients	Millers et al 1989 Bertrand et al. 1994 Decker et al. 2003 Fitzhugh et al. 2003
Oak	Large areas of oak mortality recorded in New England and the Appalachian Mountains in the early 1900s	Drought stresses have been reported as important initiating factors in oak decline.	Insects, secondary pathogens	Millers et al. 1989
Ash	Widespread dieback in the Northeast since 1920	Drought and freezing damage have been identified as inciting factors, with drought playing a particularly important role	Phytoplasmal disease, Asian beetle, emerald ash borer	Millers et al. 1989 Poland and McCullough 2006
Red spruce	Widespread decline through the Northeast after 1960, increasing over the last few decades	Reduced cold tolerance leads to winter injury which is intensified rapid rates of thaw and subsequent exposure to refreezing	Acid deposition, anomalous weather	Friedland et al. 1984 Johnson 1992 Schaberg and DeHayes 2000 Bourque et al. 2005

Waterbodies

In Vermont and across the Northeast, hydrological cycles and processes are being affected by climate change. Increased average annual rainfall in conjunction with a larger fraction of precipitation falling as rain is increasing average annual streamflows resulting in more frequent and greater magnitude high flows (Galford, G.L. et al. 2014). Moreover, these high flows are occurring more frequently in winter months as a result of early snowpack thaw dates driven by warming temperatures (Galford, G.L. et al. 2014). Despite the fact that recent decades have seen higher summer base flows, projections suggest that warming temperatures in conjunction with greater variability in seasonal rainfall may increase the likelihood of short term summer droughts (Galford, G.L. et al. 2014). Together these changes in hydrological processes have serious implications for northeastern ecosystems.

One concern is the likely increase in frequency and amount of stormwater runoff. Nutrient and sediment loading into rivers exacerbates siltation and algae production and can limit the capacity of the waterway to support macroinvertebrates, fish, freshwater mussels, and other aquatic organisms. “In lakes, warmer temperatures and greater nutrient loading can result in more frequent blue-green algae blooms which are known to be detrimental to both animals

and humans”(Galford, G.L. et al. 2014). Another concern is the expected increase in the a “wider range of lake level fluctuations over the course of the year (Galford, G.L. et al. 2014, Stager, J. & Thill, M. 2010).

Changes in river hydrology, with larger and more frequent peak discharges, may trigger significant changes in river channel formation (Allan, 1995) which can result in greater channel instability and channel movement. The response to such instability in the past has been to channelize and harden riverbanks to prevent channel movement. This practice often have deleterious and homogenizing effects on in-stream and riparian habitat features (Poff and Zimmerman 2009). Moreover, catastrophic results can be expected when hardened and channelized banks fail during very high flow situations—as was seen many times over during Tropical Storm Irene-driven flooding.

Wetlands

Vermont has more than 290,000 acres of wetlands that provide critical habitat for many species of fish, wildlife and plants. Wetlands play an important role in the attenuation of storm water, sediment transport, and naturally improve water quality (Galford, G.L. et al. 2014). Similar to waterbodies in Vermont, increases in variability and overall seasonal extent of precipitation are a significant challenge to wetland ecosystems, with potential for greater frequency of flood events and short to mid-term droughts. Increased flooding will bring about changes to wetland shorelines, may facilitate the spread of invasive aquatic species (e.g., Japanese Knotweed), and cause an influx of runoff sediment. Reduced water clarity due to flooding will, reduce light penetration and greatly affect the productivity of aquatic organisms (Galford, G.L. et al. 2014). In addition, projected increases in summer temperatures and earlier snowmelt present a threat to wetlands as water levels may intermittently decrease or completely dry up as a result of drought in conjunction with increased evapotranspiration rates. If this were to occur, the breeding cycles and survival of amphibians, such as the Spotted Salamander, would be adversely impacted (Galford, G.L. et al. 2014). The potential for increased dry spells and low water conditions will have water quality implications too. With shallower water depths and warm temperatures toxins and nutrients will be more concentrated and algae blooms will be more frequent (Galford, G.L. et al. 2014). Furthermore, non-climate stressors, such as habitat fragmentation and pollution, also threaten wetland ecosystems in Vermont. In assessing the vulnerability of wetland habitats, the Vermont Fish and Wildlife Department (VFWD), National Wildlife Federation (NWF) and others found Basin Swamps & Wetlands and Wet Shores to be particularly vulnerable to climate-driven impacts.

Habitats and Species

The interactions between climate-driven stressors including changes in temperature, precipitation, and storm events and non-climate stressors such as development-driven habitat loss and habitat degradation is currently impacting Vermont’s habitats and the species that depend on them. Moreover, synergistic impacts are likely to increase over the coming decades.

Projections suggest that warming temperatures will expand the extent of suitable habitat for tree species such as oak, hickory, and red maple, while significantly decreasing the extent of suitable habitat for more cold-tolerant species such as balsam fir currently found in Vermont’s higher elevations (Galford, G.L. et al. 2014). With earlier leaf out and flowering periods because of this warming, many species may be more susceptible to pests and

pathogens (Galford, G.L. et al. 2014). Changes in habitat suitability for many species can prompt a reshuffling and unraveling of currently recognized species assemblages and natural communities. Invasive species, which generally exhibit a competitive edge under warmer conditions, could further spread. Species that are ecological specialists are particularly vulnerable as their ranges are restricted, they are limited to a specific habitat, or they are geographically isolated (Rustad, L. et al. 2012). For example, ecological specialists [such as the Fowlers Toad, found in sandy outwash areas along Connecticut River], or those with populations already in decline [such as the Rusty Blackbird, a boreal wetland species], may be particularly vulnerable under changing habitat conditions (Rustad, L. et al. 2012). Bicknell's thrush is limited to high-elevation spruce-fir forests and is a good example of a species particularly vulnerable to climate-driven habitat change. Moreover, data suggests that many migratory bird species are arriving and breeding earlier in Vermont, and that the ranges of numerous migratory bird species are changing in response to climate change (Rustad, L. et al. 2012).

Climate change is also affecting the phenology, or timing of life-history events of many plant and animal species. Due to increasing spring air temperatures, many plants are leafing out and blooming earlier and numerous wildlife species are breeding or migrating earlier than they did during the previous century (Betts, 2011; Stein, B.A. et al. 2014). Current evidence, including shifts in entire ecoregions in some locations, suggests that over time these changes could exceed the ability of many species to adjust, leading to predictions of species declines and higher extinction rates globally (Loarie, S.R. et al. 2009). However, predictions ought to be species-specific as different species respond to climate-driven impacts in different ways and at different rates (Stein, B.A. et al. 2014). Evidence suggests that there are significant differences in response capacity between “short-lived species with high dispersal capacity (such as birds) and long-lived species with limited dispersal capacity (such as many trees)” (Stein, B.A. et al. 2014). Those species that have a greater capacity to adapt to new conditions, or have greater climatic tolerances, will have a competitive advantage and may expand their ranges while those with narrower climatic tolerances may experience range contraction.

In addition, climate-driven impacts from extreme weather events, direct thermal stress, changes in habitat availability, and increases in parasites and diseases will effect native wildlife “at all levels of organization from the physiology of individual animals to changes at the population level” (Rustad, L. et al. 2012). Two species that provide examples of the extent to which different suites of climate-driven stressors can increase vulnerability are the Little Brown Bat and Moose. The Little Brown Bat relies largely on insects with aquatic larval stages as a food resource (Rustad, L. et al. 2012). Changing precipitation patterns, alterations to stream flow and reduced soil moisture can significantly affect the availability of these insects (Rustad, L. et al. 2012). Apart from the vulnerability of Little Brown Bats to altered hydrology, there have been [significant declines in the population](#) as a result of [White Nose Syndrome](#). Altered precipitation and disease may interact synergistically to exacerbate stress on this species in Vermont, though more water and shorter hibernation periods in a warmer climate could potentially be a direct benefit.

Moose appears to be vulnerable to climate change in Vermont. As a cold-adapted species, Moose begin to reduce food intake in response to high summer temperatures (Renecker and Hudson 1986). As annual summer high temperatures increase under climate change, the habitat range of Moose may shift northward. Warming winter temperatures may reduce the

area of permanent winter snowpack. The reduction in snowpack may increase contact between Moose and White-tailed Deer, which are carriers of a brain parasite (*Parelaphostrongylus tenuis*) that is potentially lethal to Moose (Whitlaw and Lankester 1994). In addition, Winter Ticks (*Dermacentor albipictus*) are becoming an increasing problem for Moose with the early loss of snowpack. When ticks fall off Moose in the spring, they have a greater likelihood of survival if they fall onto soil as opposed to snow.

Amphibians are particularly vulnerable to the interactions between altered hydrology and increasing temperature. With the majority of amphibians breeding in water, the hydroperiod (the time period that there is standing water) of ephemeral ponds is a critical component of their life cycle (Rustad, L. et al. 2012). Altered precipitation and increased seasonal warming which drive evaporation and the frequency and severity of droughts are of particular concern (Rustad, L. et al. 2012). “A reduced hydroperiod can increase competition, decrease size at metamorphosis, and kill larvae as ponds dry out (Rustad, L. et al. 2012).”

Conserving Wildlife in the Face of Climate Change

What makes the issue of climate change particularly intractable is the breadth of interconnections between these climate-driven impacts and numerous non-climate stressors such as habitat fragmentation, habitat degradation, and pollution – which have been the focus of conservation efforts for some time. Given the magnitude of climate impacts, rates of increasing variability, and the synergy between climate and non-climate stressors on wildlife and the habitats upon which they depend, there is an urgent need to prepare for and respond to these impacts (Stein, B. et al. 2014).

In planning for this new future, we face questions such as: (a) which species and habitats are likely to be more vulnerable; (b) which may benefit or be unaffected by climate change; (c) how will non-climate stressors contribute to vulnerability; (d) how might species adapt; (e) which strategies will be most effective and where should we apply our efforts; and, (f) how can we feasibly monitor species and habitats to inform our management actions? Significant efforts have been made at the Federal, regional, State and local levels to address these questions and identify strategies that benefit wildlife and their habitats under climate change.

Regional Efforts to Conserve Wildlife in the Era of Climate Change

The Northeast is well-known for collaborative conservation efforts between states, municipalities, conservation organizations, and federal entities. A prime example is the Northeast Association of Fish and Wildlife Agencies (NEAFWA), which is a collaboration between State Fish and Wildlife Departments across the thirteen Northeastern States. Since 2007, the NEAFWA sponsored [Regional Conservation Needs Grant Program](#) has funded a large number of projects focused on identifying both the threats facing wildlife and their habitats under climate change as well as identifying strategies to benefit those species in a warming world.

Recognizing the need to conserve and restore habitat connectivity in the Northeast, the [Staying Connected Initiative](#) (SCI) was developed among a partnership of twenty-four private and public entities across New York, Vermont, New Hampshire, Maine, and Canada. The mission of this collaboration is to sustain connectivity for wildlife by protecting against habitat fragmentation and climate change. In doing so, SCI brings its partners together in an interdisciplinary approach that utilizes tools of conservation science, land protection,

community outreach, land use planning, transportation, and policy to ensure that connectivity across the landscape is healthy for both wildlife and human communities.

In September 2012, the National Wildlife Federation (NWF) in partnership with Manomet Center for Conservation Sciences (MCCS) completed the Northeast's first regional vulnerability assessment (RVA) (MCCS & NWF 2012). The assessment utilized facilitated expert elicitation from a panel of 27 natural resource practitioners including staff from state fish and wildlife agencies in all of the Northeastern states, as well as other state and federal habitat professionals. The goal of the assessment was to (a) quantify the vulnerabilities to climate change of fish and wildlife habitats, and how these vulnerabilities vary spatially across the region, (b) project how the status and distributions of these habitats and species may be affected by climate change, and (c) to work with states to increase their institutional knowledge and capabilities to respond to climate change.

Another important regional collaboration working to “to address increasing land use pressures and widespread resource threats and uncertainties amplified by a rapidly changing climate” is the [North Atlantic Landscape Conservation Cooperative](#) (NALCC). “The partners and partnerships in the cooperative address these regional threats and uncertainties by agreeing on common goals for land, water, fish, wildlife, plant and cultural resources and jointly developing the scientific information and tools needed to prioritize and guide more effective conservation actions by partners toward those goals (NALCC).”

Efforts to Address Climate Change Impacts in the Northeast

State-level land conservation efforts vary considerably from state to state across the Northeast. They include efforts by state agencies, departments, and programs such as Open Space Programs and Land Conservation Programs across a number of different state environmental agencies. All the northeastern States are developing Action Plans that address climate change as a significant stressor for wildlife and their habitats. Several of them have specifically conducted climate change vulnerability assessments on priority SGCN and associated habitats.

In 2011 New York conducted a species vulnerability assessment on 119 SGCN using NatureServe's Climate Change Vulnerability Index (Schlesinger, M.D. et al. 2011). The assessment found that nearly all of the species identified as Highly or Extremely Vulnerable were associated with aquatic or seasonally wet habitats, with mussels rated as particularly vulnerable due to limited mobility (Schlesinger, M.D. et al. 2011). Another interesting finding was that “vulnerability was only weakly associated with conservation status (Schlesinger, M.D. et al. 2011).” In addition, a vulnerability assessment of key habitats, conducted by NWF in partnership with the New York Department of Environmental Conservation's Division of Fish, Wildlife and Marine Resources, found Acadian- Appalachian Montane Spruce-Fir and Coastal Plain Basin Peat Swamp as Vulnerable to climate change, and Boreal Bog and Alpine Tundra habitats as Highly Vulnerable (Hilke, C. & Galbraith, H. 2013).

New Hampshire Fish and Game Department (NHFGD) (2013) conducted a habitat vulnerability assessment in 2014 on 25 key habitats across the state. Due to climate impacts on the hydrology of freshwater habitats, findings suggested that species that are “more tolerant of a wide range of hydrologic conditions will be favored, and the total species richness may decrease (NHFGD 2013).” For terrestrial habitats, high-elevation spruce fir forest was identified as particularly vulnerable to climate change (NHFGD 2013). Moreover,

findings suggested that “hardwood-pine forests will move northwards and up slope, while Appalachian oak-pine forests are likely to increase in extent” as a result of long-term warming trends (NHFGD 2013). Other findings of interest identified Pine Barrens as less vulnerable to climate change given the predisposition of species within those habitats to warmer and drier conditions, and more generally predicted an increase in early successional habitats as a result of increased disturbance (NHFGD 2013).

In 2010, Massachusetts conducted the first climate change habitat vulnerability assessment in the Northeast. The assessment was directed towards answering: “(a) how do the targeted fish and wildlife habitats rank in terms of their likely comparative vulnerabilities to climate change; (b) how will the representation of these habitats in Massachusetts be altered by a changing climate; (c) which vertebrate Species in Greatest Need of Conservation [SGCN] are likely to be most vulnerable to climate change; and (d) what degree of confidence can be assigned to the above predictions” (MCCS & MA Department of Fisheries and Wildlife 2010)? The assessment was conducted under two emission scenarios, a doubling and a tripling of atmospheric CO₂. Spruce-fir forests, smaller coldwater lakes and ponds, spruce-fir boreal swamp, brackish marsh, and intertidal mudflats and sandflats were identified as highly vulnerable to climate change under both emission scenarios (MCCS & MA Department of Fisheries and Wildlife 2010). The highly vulnerable representative SGNC associated with those habitats include the Northern Leopard Frog, Green Heron, American Eel, Blackpoll Warbler, Moose and Bobcat (MCCS & MA Department of Fisheries and Wildlife 2010).

Table 3.3. Numbers and percentages of vertebrate SGNC [SGCN] most at risk from doubling (2X) and tripling (3X) of atmospheric CO₂ concentration (Manomet & MA DFW 2010).

	Amphibians (7)	Reptiles (19)	Fish (28)	Birds (63)	Mammals (20)
2X CO₂	2 (28%)	1 (5%)	2 (7%)	26 (41%)	3 (15%)
3X CO₂	4 (57%)	4 (21%)	14 (50%)	36 (57%)	7 (35%)

Vermont Efforts to Understand and Address Climate Change Impacts to Species and Habitats

Vermont has been actively involved in addressing the impacts of climate change for some time. In 2011, Governor Peter Shumlin established the Vermont Climate Cabinet which was charged with coordinating climate change efforts specific to greenhouse gas emission reduction, reliance of fossil fuels, as well as the implementation of climate adaptation efforts across State agencies. While initial efforts to address climate change focused largely on mitigation efforts, the Vermont Agency of Natural Resources developed a series of white papers addressing climate adaptation across its Programs and Divisions. In 2012 and 2013 the Vermont Agency of Natural Resources developed the Climate Change Adaptation Framework to gather information about climate change in Vermont as it relates to natural resources and to propose a strategic framework for continued climate change vulnerability assessment and action planning (Tetra Tech Inc. 2013).

Building on this assessment the Fish & Wildlife Department and National Wildlife Federation developed the Species & Habitat Climate Vulnerability Assessment for 18 key species, 20 upland habitats, 11 wetland habitats, and 13 freshwater habitats as part of our Action Plan revision (Table 3.4 and Appendix D). Species included SGNC and important “surrogate” species that are widely considered representative of particular habitat types.

Species assessments culminated in an overall vulnerability rating for climate-specific and non-climate stressors and an associated confidence score.

The most important lesson taken from this exercise is that species responses to climate change will not be uniform. For some, climate change may not be a significant threat, however if that species is already subjected to other stresses, climate change impacts may push that species over the edge. This is an important consideration to take into account.

Vulnerability rating criteria were standardized and applied across all assessments. Criteria were selected from similar assessments conducted by other states and from current literature. Vulnerability scores were designated for each species, habitat and system. Ratings account for the extent of vulnerability (extremely vulnerable, highly vulnerable etc.) based upon percentage changes in abundance or extent expected by 2050 and include an associated confidence rating of high, medium or low based upon the percentage certainty of the vulnerability score (Low = Not very confident, 0-30% certainty in vulnerability score). See Vermont Climate Vulnerability Assessment Rating Key Appendix E for details.

Non-climate Stressors: Non-climate stressors included acidity & pollution, habitat alteration & altered hydrology, invasive species, channel erosion & sedimentation, encroachments, land erosion, nutrient loading, thermal stress, toxic substances/pollution, and habitat fragmentation.

Sensitivity Factors: assessments factored in how likely a species, habitat, or ecosystem is to be affected by or respond to climate change given (1) habitat specificity, (2) edge of range, (3) environmental or physiological tolerance, (4) interspecific or phenological dependence, (5) mobility, and (6) exotic pathogens or invasive species.

Exposure & Key Climate Change Factors: The four categories of climate change and exposure factors are temperature, hydrology, extreme events, and phenology. Within each category, several factors were selected along with an associated trend and specific projections for each trend (see Appendix D. Vermont Vulnerability Assessment Rating Key for details).

Temperature: Annual temperature, seasonal temperature, number of hot days, number of cold days, and variability. Each factor has an associated trend and specific projections. As an example, annual temperature (code=A) had an increasing trend with projections suggesting a 3.7 to 5.8°F increase by 2050, and a 5.0 to 9.5°F increase by 2100.

Hydrology: Ten hydrology factors were selected, including annual precipitation, seasonal precipitation, heavy rainfall events, soil moisture, snow, spring flows, summer low flows, ice dynamics, fluctuating lake levels, and lake stratification.

Extreme Events: Extreme weather events include flood events, number of short-term droughts, storms, and fire.

Phenology: Phenological factors include length of growing season, onset of spring, onset of fall, and biological interactions.

Of the 18 species assessed, eight were identified as highly vulnerable to climate-driven impacts. Of those, five are SGCN: Jefferson Salamander, Canada Lynx, Brook Trout, Eastern Pearlshell Mussel and Bicknell's Thrush.

Climate change vulnerability assessments were similarly conducted for Vermont's upland and wetland natural communities. For efficiency, the 95 natural community types were grouped into categories based on the environmental factors that drive their development and that could affect their susceptibility to climate change. Some natural community types were assessed individually.

Of the upland habitats, Subalpine Krummholz, Alpine Meadow, Cold-Air Talus Woodland, and Dry Oak Woodland were identified as highly vulnerable to climate-driven impacts, and Upland Shores and Hemlock Forests were identified as particularly vulnerable to non-climate stressors. Of the 11 wetland habitats, Basin Swamps and Wetlands, Floodplains, Floodplain Forests, and Wet Shores were identified as highly vulnerable to climate change. As with the species analyses, habitat assessments culminated in an overall vulnerability rating for climate-specific and non-climate stressors and an associated confidence score. The assessments also detailed key climate change/exposure factors and non-climate stressors that contributed to the overall vulnerability rating.

The effort also included an assessment of 13 freshwater habitats including six river types and seven lake types. River types were delineated primarily by stream order and pH, and include; high gradient, cold-water acidic, 1-2 order, high gradient, cold-water, not acidic, 1-2 order, medium-sized rivers (4-6 order) etc. Lake types were delineated by trophic type and extent of stratification and included; Oligotrophic lake (stratified), Mesotrophic-Eutrophic lake (stratified), Mesotrophic-Eutrophic Lake (unstratified) etc. For the 13 freshwater habitats, High gradient, cold water, not acidic (1-2 order), and High gradient, cold water acidic (1-2 order) lakes were also identified as highly vulnerable to climate change.

Table 3.4: Vermont Species & Habitat Climate Vulnerability Assessment (SGCN in bold)**Vulnerability Rating Key** (Abundance and/or range extent in Vermont change by 2050)**E** = Extremely Vulnerable: Extremely likely to substantially decrease (>75% loss) or disappear**H** = Highly Vulnerable: Likely to decrease significantly (25-75% loss)**M** = Moderately Vulnerable: Likely to decrease (10-25% loss)**L** = Slightly Vulnerable: Available evidence does not suggest change (decrease, 5-10% loss)**N** = Not Vulnerable, No Effect: Likely to increase or decrease by less than 5%**B** = Increase Possible or Likely: Likely to increase (>15% increase)**U** = Unknown/Uncertain: Available evidence not available or not conclusive at this time.**Confidence Ratings****H** = High: Very confident (>60% certainty in vulnerability score)**M** = Moderate: Somewhat confident (30-60% certainty in vulnerability score)**L** = Low: Not very confident (0-30% certainty in vulnerability score)

Species and Habitats	Climate Vulnerability Rating	Confidence Score	Non-climate stressors Vulnerability
Key Species			
Jefferson salamander	H	H	H
Northern white cedar	M	M	H
Fingernail clam	L	H	M
Beaver	N	H	M
Bobcat	L	M	M
Lynx	H	H	M
Brook Trout	H	H	H
Wood Turtle	M	H	H
Pearlshell mussel	H	M	M
Fallfish	L	M	L
Smelt	M	H	M
Lake trout	H	H	H
Bald Eagle	L	M	L
Bicknell's Thrush	H	H	H
Common Loon	M	L	L
Red Oak	N	M	N
Sugar Maple	H	M	L
West Virginia White (butterfly)	H	H	M
Wetland Habitats			
Cattail Marsh	L	M	L
Shallow Emergent Marsh	M	M	M
Marsh and Sedge Meadow (Formation)	M	M	M
Alluvial Shrub Swamp	M	M	L
Basin swamps & wetlands	H	M	M
Floodplains	H	M	M
Ground water seepage & Flood swamp	L	M	L
Open peatlands (precipitation-dependent)	M	M	L
Open peatlands (ground-fed)	L	M	M

Species and Habitats	Climate Vulnerability Rating	Confidence Score	Non-climate stressors Vulnerability
Floodplain Forests	H	M	M
Wet Shores	H	M	M
Upland Habitats			
Alpine Meadow	E	M	M
Spruce-Fir-Northern Hardwood	M	M	M
Northern Hardwood Forest	L	M	L
Oak-Pine–Dry Mesic Forests & Woodlands w/deeper soils	L	M	M
Oak-Pine Northern Rocky–Northern Dry Rocky Forests and Woodlands	L	M	L
Oak-Pine Southern Rocky–Southern Dry Rocky Forests and Woodlands	M	L	L
Outcrops and upland meadows	N	L	L
Cliffs and Talus	L	M	L
Upland shores	M	M	H
Subalpine Krummholz	E	M	
Montane Spruce-fir	M	M	
Red Spruce-Heath Rocky Ridge	M	M	
Montane Yellow Birch-Red Spruce Forest	M	M	
Red Spruce-Northern Hardwood	L	M	
Lowland Spruce-Fir Forest	M	M	
Boreal Talus Woodland	M	M	
Cold-Air Talus Woodland	H	M	
Limestone Bluff Cedar-Pine Forest	L	M	
Transition Hardwood Talus Woodland	L	M	
Dry Oak Woodland	H	L	
Freshwater Habitats			
Medium-sized river (4-6 order)	M	M	
Large river (7+ stream order)	L	M	
High gradient, cold water acidic, 1-2 order	H	H	
High gradient, cold water, not acidic, 1-2 order	H	H	
Low gradient marsh	M	M	
Lake Champlain valley	M	M	
High Elevation Lake	M	L	L
Dystrophic Lake	M	L	L
Lake–Oligotrophic, Stratified	L	H	L
Mesotrophic-Eutrophic Lake (stratified)	M	M	M
Mesotrophic-Eutrophic Lake (unstratified)	H	M	M
Unstratified lakes	M	H	H
Stratified Lakes	M	H	H

Recent Projects Helping Address Climate Impacts to Species and Habitats

Vermont has developed a number of tools that provide a strong foundation for addressing the impacts of climate change on wildlife and their habitats. Examples include VFWD's 2014 [BioFinder](#) project, which identifies Vermont's lands and waters supporting high priority ecosystems, natural communities, habitats, and species. BioFinder's 21 datasets represent biological, ecological, and natural heritage data stacked together for a co-occurrence analysis which identified the locations of greatest overlap (concentration) for priority ranking at the statewide scale. The Habitat Block project, another spatial analysis tool, prioritized 4,055 habitat blocks and identifies likely wildlife road crossing locations for all of Vermont's roads. Maintaining a connected network of unfragmented habitat blocks is recognized as a primary strategy for conserving biological diversity in the face of a rapidly changing climate, and is one of the most widely used of VFWD's datasets in conservation planning technical assistance provided to towns and Regional Planning Commissions.

Vermont's [iMap Invasives](#) database aids in assessing the spread of invasive species anticipated under climate change. The database allows ANR, TNC, and other land management organizations to document and track the spread of invasive species throughout the state, provides an early warning of the arrival of new invasives into the state or into new areas of the state, and allows ANR to follow management actions taken to control any given population. In 2015, a new ANR Invasive Plant Coordinator position was created to help further invasive species management.

Vermont is engaged in a number of wildlife species monitoring and management efforts to further our understanding of climate impacts, which may help us develop management strategies. VFWD is monitoring Moose populations through indices of ticks, hair loss, and mortality; assessing the impacts of climate change on the interactions between Fisher and American Marten; surveying for recently documented Canada Lynx; and is actively engaged in management activities to raise the elevation of targeted turtle nesting beaches in response to higher Lake Champlain water levels. Similar activities include monitoring deer populations and the implications of deer browsing under climate change. Projects such as the [Vermont Breeding Bird Atlas](#) may also have significant value for monitoring changes over time that may be related to climate change.

VFWD has recently developed a sampling regime that includes annual trout population monitoring concurrent with stream temperature data. The resulting datasets will allow biologists to monitor changes in fish abundances and stream temperatures at specific locations. Stream temperature monitoring also takes place in various streams across the state for specific fisheries management projects.

Changes to state lands management planning: 1. Beginning in 2010 the ANR began addressing climate change in new Long Range Management Plans (LRMPs) for state lands. 2. ANR is currently developing new LRMP for Victory State Forest and Wildlife Management Area with climate change as a key concept to drive management decisions. Key factors include boreal habitat conditions and the presence of rare and edge-of-range species (e.g., Canada Lynx, Spruce Grouse, American Marten, and Black-backed Woodpecker). 3. In an effort to address the uncertainty of assisted migration as a climate change strategy, the ANR Lands Team is reviewing options, impacts and management guidelines for state lands.

ANR has a new state lands management policy and guidelines for management and protection of riparian buffers. Good management of riparian areas is often referred to as a “No Regrets” climate strategy because it makes such good sense to maintain vegetated habitats along our rivers and streams for a variety of benefits. While the buffer policy is a work in progress, a draft is complete and is based on all currently available science on the values and protection strategies for riparian areas. The policy is not based solely on climate and flood resilience, but it intentionally addresses these important considerations.

In 2015 VFPR presented a report to the legislature on forest fragmentation. Strategies to address forest fragmentation complement habitat conservation efforts needed for climate change. The Vermont Forest Roundtable is focused on implementing solutions to address forest fragmentation that will be essential to wildlife species habitat protection goals.

Other important State efforts to address the impacts of climate change on wildlife species and habitats include:

- The ANR Stewardship Team funded a pilot project on four areas of state lands where the Department of Environmental Conservation (VDEC), in coordination with the Forests, Parks & Recreation Department (VFPR) and VFWD, is investigating flood resilience conditions and related management opportunities. This was intentionally developed to better understand state lands management decisions relative to climate change and flood resilience.
- VFWD and VDEC Rivers Program have been evaluating culverts for fish and aquatic organism passage and stream geomorphic compatibility since 2005. The objectives are to gain a greater understanding of the scope of fish and aquatic organism passage (AOP) barriers and undersized culverts that may be having a negative impact on physical stability and quality of stream habitat in Vermont and work toward addressing these issues where appropriate; and improve the understanding and communication between VFWD, VDEC, state and local road managers and state and federal regulators in addressing AOP and geomorphic compatibility issues at stream-road crossings.
- The Vermont Dam Task Force is dedicated to restoring rivers through the assessment, prioritization, and facilitation of dam removal or modification. This work is particularly important in light of climate change because dams can significantly degrade a river’s water quality (e.g., temperature and dissolved oxygen), aquatic habitat, the movement of aquatic organisms and the transport of sediments downstream. The Dames Task Force is a statewide cooperative effort among federal and state agencies (including VFWD and VDEC), private organizations, and individuals allows participants to share information and dam removal advice; coordinate efforts to accomplish the removal of dams whose negative effects exceed their benefits; and, to reach out to the public.
- The Narrows Wildlife Management Area in West Haven was selected for a pilot project by VFPR for assessing and monitoring timber management and its attendant consequences on climate change factors including the presence and movement of invasive species, and regeneration success. Additional demonstration areas are in development as part of the Mount Philo State Park, the Putnam State Forest, and the Okemo State Forests.

- VFWD recently updated the Conserving Vermont's Natural Heritage document to include a section that discusses climate change, how it may impact habitats, natural communities, plants and animals, and how to consider those effects in the context of local and regional land use planning.
- VFPR is working with private landowners to conserve and manage one of five priority habitat linkages identified by Two Countries, One Forest as critical to maintaining the integrity of the entire Northern Forest. This landowner cooperative involves over 13 properties known as Cold Hollow to Canada and employs climate smart forest management strategies.
- Forest changes are monitored by FPR through a number of forest health monitoring plot networks to track current and future habitat suitability and tree species distributions.
- VFWD's Community Wildlife Program (CWP) provides technical assistance to municipalities and Regional Planning Commissions emphasizes the importance of integrating climate change effects on land, fish and wildlife conservation into planning. VFWD intentionally shifted to a greater focus on landscape and habitat connectivity through the efforts of this program for a number of reasons, including our understanding that creating a well-connected landscape is important for landscape resilience relative to climate change.
- Since Tropical Storm Irene in 2011, VFWD and VDEC has worked together with Vermont Agency of Transportation (VTrans) to change how we address road-stream crossings and other river/road conflicts. The net result is larger stream crossing structures, a more rigorous application of Aquatic Organism Passages (AOP), and a more holistic approach to river management practices driven in part by flood concerns related to climate-driven impacts. VFWD, VDEC and VTrans created a rivers and roads training program targeted at state and municipal employees and contractors that work on roads to increase the knowledge base of river/road [best management practices](#), especially as they pertain to post-flood scenarios when the likelihood is greatest for in-channel work that can harm channel stability and in-stream habitat.
- Currently, VFWD staff members are developing a guidance document that sets out a vision for future land acquisition efforts. This guidance will take into consideration important factors related to climate change and landscape resilience, in particular landscape and habitat connectivity.
- VFWD staff and ANR's Office of Planning and Legal Affairs collaborated on revisions to Act 250 criterion 8 to better address necessary wildlife habitat, rare, threatened and endangered species and their habitats, rare and irreplaceable natural areas, forest blocks, and landscape connectivity. Riparian corridors are identified as especially important for wildlife movement, biological diversity, and river stability. The primary impetus for this work was to improve how Act 250 addresses habitat fragmentation and landscape connectivity, two factors recognized as being critical to climate change adaptation.

- And in an effort to maintain the reciprocal flow of information between the State and Federal programs, Governor Shumlin serves on the Presidential Task Force on Climate Preparedness and Resilience. In this capacity, the Governor works to ensure that lessons learned from Vermont’s climate-related activities inform Federal climate change considerations.

Identifying Climate Adaptation Strategies for Vermont’s Species and Habitats

Before selecting actions to help wildlife species and their habitats respond to climate change it is important to bear in mind the ways in which climate change may impact the evaluation of management alternatives (Stein, B. et al. 2014).

Performance: Changing climate conditions that could affect the outcome of some conservation strategies. For example, shifts in the intensity of peak flows or extent of low stream flow may affect the performance of some of fish passages structures.

New constraints: Climate change may add new constraints, limiting what is technologically, ecologically, or culturally achievable. Changing conditions may make local persistence of some species or habitats impossible, or climate-related shifts in land uses may create new obstacles to species movements.

Relative weight: Because climate considerations could increase costs of some actions significantly these costs should be considered up front.

Perceived value: Climate change may affect the perceived value of various resources. For example, as floods become more frequent and severe in some places, marshes and wetlands may become increasingly valued for their ability to mitigate flood risk.

Priority Actions to Address Climate Change Impacts to Species and Habitats

1. Protect large habitat blocks, riparian habitats and climate refugia, and promote landscape integrity and connectivity to facilitate the movement of species across habitats based on the VFWD report “Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape” (Sorenson et al. 2015), the Aquatic Organism Passage program, River Corridor Planning and other conservation plans.
2. Increase riverine, floodplain, and riparian connectivity based on Aquatic Organism Passage program and River Corridor Planning recommendations.
3. Restore rivers through the assessment, prioritization, and facilitation of dam removal or modification.
4. Support the [Staying Connected Initiative](#), [Cold Hollow to Canada](#), and similar programs focused on maintaining and enhancing landscape integrity and connectivity.
5. Complete and implement a state river corridor protection plan, riparian buffer policy and vernal pool management guidelines.
6. Expand river corridor and floodplain protections similar to the Lakeshore Protection Act.
7. Support and help implement the recommendations of VFPR’s Creating and Maintaining Resilient Forests in Vermont: Adapting Forests to Climate Change

7. Support and help implement the recommendations of VFPR's Creating and Maintaining Resilient Forests in Vermont: Adapting Forests to Climate Change
8. Support and help implement the recommendations of VFPR's Forest Fragmentation Plan.
9. Implement Vermont's Clean Water Initiative to improve water quality statewide particularly in Lake Champlain.
10. Protect and expand riparian habitats to increase habitat connectivity, increase cold water habitats, reduce the spread of invasive species, accommodate river channel dynamics and mitigate the impacts of flooding (Hilke, C. & Galbraith, H. 2013).
11. Protect ecosystem health and stability by preventing new introductions of invasive species and pests and controlling infestations of existing species via integrated pest management programs.
12. Minimize climate change impacts by employing management strategies that sustain fundamental ecological functions, promote habitat resiliency and adaptive capacity, and restore habitats with future conditions in mind.
13. Invest in research and monitoring programs that can inform species and habitat management with future conditions in mind such as: monitoring indicators of changes in species distribution and abundance; stream temperatures, cold water refugia and cold groundwater inputs to streams, early detection of invasive species, pests and pathogens; identification of genetically adapted species.
14. Develop incentive, education and technical assistance programs to help landowners, land managers, municipalities and others adopt and implement climate smart conservation programs, such as: culvert replacement, streamside shading, stream bank stabilization, sediment control river corridor easements and other types of conservation easements.

Managing Vermont's Wildlife and Habitats into the Future

As states across the country grapple with climate change impacts to our natural heritage, the need to set management priorities based on a sound understanding of projected impacts is becoming increasingly apparent. Climate change typically amplifies existing ecological stressors including rates of change, disturbance, habitat degradation and fragmentation. As such, isolating specific climate change impacts is difficult because it interacts with and compounds a host of non-climate stressors. To that end, this chapter provides important baseline information about how to protect Vermont's species and habitats under climate change including, (a) the climate-driven threats facing Vermont's wildlife and habitats, the vulnerability of key habitats and species to climate change, and (c) potential strategies for increasing the resiliency and/or adaptive capacity of select habitats and species. Climate change is not the singularly most detrimental stressor to Vermont's ecological systems, but it cannot be brushed aside with the thinking that what we are already doing is sufficient. Complex problems often require holistic problem solving. Vermont's Wildlife Action Plan can serve as an important role in an integrated approach to protecting our wildlife and the habitats upon which they depend.

Literature Cited

- Allan, J. D., 1995. Stream Ecology: Structure and Function of Running Waters, Kluwer Academic Publishers, Boston, MA.
- Anderson, M. C. (2012). Resilient sites for terrestrial conservation in the northeast and mid-atlantic region. The Nature Conservancy, Eastern Conservation Science, 168 p.
- Association of Fish and Wildlife Agencies (AFWA), State Wildlife Action Plan (SWAP) webpage, Accessed 01/2015, <http://www.wildlifeactionplan.org>.
- Betts, A.K. 2011. Vermont Climate Change Indicators. Weather, Climate and Society. Vol. 3. Pp 106-115. DOI: 10.1175/2011WCAS1096.1
- Chen, I.-C., J.K. Hill, R. Ohlemuller, et al. 2011. Rapid shifts of species associated with high levels of climate warming. Science. 333: 1024–1026.
- Dukes, J. P. (2009). Responses of insect pests, pathogens, and invasive plant species to climate change in the forests of northeastern North America: What can we predict? *Journal of Canadian Forest Research* 39: 231-248.
- Galford, G.L. et al. 2014. Considering Vermont's Future in a Changing Climate: The First Vermont Climate Assessment. Gund Institute for Ecological Economics, p.20
- Hilke, C. and Galbraith, H. 2013. Assessing the Vulnerability of Key Habitats in New York: A Foundation for Climate Adaptation Planning. National Wildlife Federation, Northeast Regional Center. Montpelier, VT.
- Horton, R. et al. 2014. Ch. 16: Northeast. Climate Change Impacts in the United States: The Third National Climate Assessment, J.M. Melillo (T.C.) Richmond, and G.W. Yohe, Eds., U.S. Global Change Research Program, 16-1.
- Jacobson, G. (2009). *Past and future forest responses to climate change in New England: The urgency of maintaining healthy forested landscapes*. Report of the blue ribbon commission on land conservation: New England Governor's Conference, Inc. 2009. Staying Connected Initiative, Two Countries-One Forest.
- Kelly, A.E., and M.L. Goulden. 2008. Rapid shifts in plant distribution with recent climate change. Proceedings of the National Academy of Sciences 105: 11823–11826.
- Loarie, S.R., P.B. Duffy, H. Hamilton, et al. 2009. The velocity of climate change. Nature 462: 1052–1055.
- Manomet Center for Conservation Sciences and National Wildlife Federation. 2012. The Vulnerabilities of Fish and Wildlife Habitats in the Northeast to Climate Change. A report to the Northeastern Association of Fish and Wildlife Agencies and the North Atlantic Landscape Conservation Cooperative. Manomet, MA.
- Manomet Center for Conservation Sciences & Massachusetts Division of Fisheries and Wildlife. 2010. Massachusetts Department of Fish and Game. A report to the Commonwealth of Massachusetts. P. 1-59
- Melillo, Jerry M. et al. 2014. Climate Change Impacts in the United States: The Third National Climate Assessment. U.S. Global Change Research Program, 841 pp. doi:10.7930/J0Z31WJ2
- New Hampshire Fish and Game Department. 2013. Ecosystems and Wildlife Climate Change Adaptation Plan. Amendment to the New Hampshire Wildlife Action Plan. Pp. 1-133
- North Atlantic Landscape Conservation Cooperative. <http://northatlanticlcc.org/the-cooperative>. Website accessed 01/2015
- Poff, N. L and J.K.L Zimmerman. 2009. Ecological responses to altered flow regimes: a literature review to inform the science and management of environmental flows. *Freshwater Biology* 55: 194-205
- Renecker, L. A. and R. J. Hudson. 1986. Seasonal energy expenditures and thermoregulatory responses of moose. *Canadian Journal of Zoology* 64: 322-327.

- Rustad, L. et al. 2012. Changing Climate, Changing Forests: The Impacts of Climate Change on Forests of the Northeastern United States and Eastern Canada. United States Department of Agriculture. United States Forest Service; Northern Research Station. General Technical Report NRS-99, pp. 1-56
- Schlesinger, M.D., J.D. Corser, K.A. Perkins, and E.L. White. 2011. Vulnerability of at-risk species to climate change in New York. New York Natural Heritage Program, Albany, NY.
- Stager, J. Curt and Mary Thill, 2010, Climate Change in the Champlain Basin: What natural resource managers can expect and do, The Nature Conservancy, available at:
www.nature.org/cs/groups/webcontent/@web/@vermont/documents/docume nt/prd_002936.pdf
- Staudinger, M. D., T. L. Morelli, and A. M. Bryan. 2015. Integrating Climate Change into Northeast and Midwest State Wildlife Action Plans. DOI Northeast Climate Science Center Report, Amherst, MA.
- Stein, B.A., P. Glick, N. Edelson, and A. Staudt (eds.). 2014. Climate-Smart Conservation: Putting Adaptation Principles into Practice. National Wildlife Federation, Washington, D.C.
- Tetra Tech Inc. 2013. Vermont Agency of Natural Resources Climate Change Adaptation Framework. A report for VT ANR. p.1-157
- The Nature Conservancy. (2012). Best management practices for the prevention and treatment of terrestrial invasive plants in Vermont woodlands: Field guide for land managers, forestry professionals and landowners. Montpelier, VT.
- U.S. Global Change Research Program (USGCRP). 2009. Global Climate Change Impacts in the United States. A report of the USGCRP. USGCRP, Washington, D.C. <http://www.globalchange.gov/usimpacts/> (accessed August 7, 2010).
- Vermont Department of Forests, Parks and Recreation. 2015. Creating and Maintaining Resilient Forests in Vermont: Adapting Forests to Climate Change.
http://fpr.vermont.gov/sites/fpr/files/Forest_and_Forestry/The_Forest_Ecosystem/Library/Climate%20change%20report_final_v6-18-15a.pdf.
- Vermont Department of Forests, Parks and Recreation. 2015. Vermont Forest Fragmentation Report. Submitted to House and Senate Committees on Natural Resources and Energy and the House Committee on Fish, Wildlife, and Water Resources. Pp 62. <http://fpr.vermont.gov/node/1300>
- Whitlaw, H.A. and M. W. Lankester. 1994. The co-occurrence of moose, white-tailed deer and *Paralaphostrongylus tenuis* in Ontario. Can. J. Zool. 72:1-7.

Vermont's Wildlife Action Plan

Chapter 4 Conservation at Multiple Scales

Draft • September, 2015

Introduction	2
Selection of Classification Systems.....	3
Habitat Succession, Species of Greatest Conservation Need & the Action Plan	4
Literature Cited.....	7

4. Conservation at Multiple Scales

Introduction

This chapter explains how conservation is organized in this Wildlife Action Plan.

Vermont's list of Species of Greatest Conservation Need (SGCN) comprises 133 vertebrate species 200 invertebrate species (such as the Tawny Emperor Butterfly, Cobblestone Tiger Beetle, and Giant Floater mussel) and 813 plants (vascular and bryophytes). Developing individual conservation plans for each of these species would have been exhausting and impractical. Moreover, implementing so many individual plans would be impossible due to insufficient staffing, resources and funds. In short, it would be monumentally inefficient.

Fortunately an efficient approach exists. It consists of designing and implementing conservation at multiple scales. This is commonly referred to as the “coarse filter-fine filter” approach and is widely accepted by scientists, wildlife managers and planners. The underlying concept is that if examples of all coarse-filter features are conserved at the scale at which they naturally occur, many of the species they contain—from the largest trees and mammals to the smallest insects—may also be conserved (Hunter 1991; NCASI 2004; Schulte et al. 2006). The coarse-filter approach is well documented in the scientific literature (Jenkins 1985; Noss 1987; Hunter et al. 1988; Hunter 1991; Noss and Cooperrider 1994; Haufler et al. 1996; Jenkins 1996; Poiani et al. 2000; USDA 2004). Habitat management historically practiced by Fish and Wildlife agencies to create young forests and shrublands that benefit dozens of “shrub and early-successional species” including Moose, American Woodcock and Ruffed Grouse is an example of a ‘habitat-scale’ coarse filter.

To most efficiently conserve all our SGCN, this Wildlife Action Plan focuses on three scales of conservation:

1. **Landscapes:** Include the features that contribute to ecological function at the state and regional levels, including a network of large, connected habitat blocks and another of aquatic habitats and riparian areas. Species requiring large habitat block, mixes of forest, wetlands and waters and connections between them will benefit most from landscape-level conservation but most other SGCN can also benefit.
2. **Habitats and Natural Communities:** Include the range of naturally occurring and anthropogenic habitats (such as young forest and grasslands). Terrestrial natural communities follow the classification system developed by Sorenson and Thompson (2005) which ties in with the ecological systems classification developed for the Northeast Association of Fish & Wildlife Agencies (Gawler 2008) for the 13 northeastern states. Aquatic communities follow the classification developed by Langdon et.al. (1998).
3. **Species and Groups of Species:** these are the SGCN for which we have identified specific conservation needs that would not be covered sufficiently by conservation efforts at the other two scales.

Not all species, however, are best conserved by coarse-filters alone. For example, species dependent on multiple habitats at different times during their life cycles, those that occur in small geographic areas, those with highly specialized needs or unique threats, those that travel across large geographic areas and those that are particularly rare often require focused attention. To ensure that the needs of these species are also addressed, fine filter conservation strategies are also needed. Species-specific conservation summaries can be found in Appendix A.

Efficiency in conservation effort can be realized by first identifying landscape conservation priorities that will effectively capture many natural communities, habitats, and species found within them. Natural community and habitat level conservation can effectively capture many of the remaining species. And finally, species-specific conservation action will be required for those species that are not captured at landscape or habitat/natural community scales. Typically these are species that are very rare, are declining across their range, aggregate for breeding, and/or require large home ranges.

Given the species focus of the congressional requirements for Wildlife Action Plan development, we began at the species level by assessing SGCN individually (Appendix A). Then SGCN were organized by taxonomic group and by the habitats they use. This resulted in conservation strategies at the three levels listed above (and in table 4.1).

Table 4.1 Organization of Conservation Information in this Report

Level	Organization	Location in this Action Plan
1-Species	6 group summaries (amphibians & reptiles, birds, fishes, invertebrates, mammals and plants)	Chapter 4
	133 individual species and 15 invertebrates group summaries	Appendix A
2-Habitats & Natural Communities	125 communities & cultural habitats grouped into 24 summaries	Appendix B
3-Landscapes	Statewide and regional conservation strategies	Chapter 1
	Landscapes Landscape Report	Chapter 5 Appendix F

Selection of Classification Systems

We delineated landscapes based on the following elements: Interior Forest Blocks, Connectivity Blocks, Surface Waters and Riparian Areas, Riparian Areas for Connectivity, Physical Landscape Diversity Blocks, and Wildlife Road Crossings. Landscape conservation is discussed in chapter 5 and Appendix F of this Wildlife Action Plan.

Though great strides have been made in developing vegetation classification systems that function at the site, landscape, region and national scales (Barnes 1979, Allen and Starr 1982, Forman and Godron 1986, Cleland et. al 1997, Grossman et. al 1998), they are incomplete. In particular, no system satisfactorily integrates aquatic and terrestrial communities and cultural habitats¹ used by wildlife nationwide.

In lieu of a unified habitat classification system, Vermont's Action Plan technical teams selected the best features of five peer-reviewed vegetation classification systems that can be crosswalked with those used in other states to support broader scale conservation efforts—regionally, nationally, and internationally. Forest Cover Types (Eyre 1980) and U.S Forest Service Forest Inventory & Analysis Types (USDA 2003) were used for early successional stage forests. Natural Communities (Thompson and Sorenson 2000) were the basis for most terrestrial vegetation. "A Classification of the Aquatic Communities of Vermont" by Langdon et al. (1998) was adapted for aquatic community designations and cultural habitats¹ were adapted from Reschke (1990). Landscape scale communities were adapted from Poiani et.al. (2000).

One hundred twenty-five aquatic and natural community types, cultural habitats and land cover types, capturing most of the habitat required by SGCN were selected from the five systems (table

¹ Cultural habitats are communities and sites that are either created and/or maintained by human activities or are modified by human influence to such a degree that the physical condition is substantially different from what existed prior to human influence (adapted from Reschke 1990).

4.2). Each was assigned to one of 22 categories. Because Lake Champlain and the Connecticut River harbor most of the fish diversity in Vermont, these two waterbodies were broken out from the taxonomy to provide for a more targeted assessment. Technical teams then developed assessment summaries for each that include descriptions and general locations; current conditions; desired conditions based on the needs of associated SGCN; priority problems; conservation strategies to address problems (along with the identification of potential conservation partners and funding sources); and a listing of relevant plans and planning processes pertinent to a habitat type.

Our terresitral classification is designed to roll up to the Northeast Terrestrial Habitat Classification System (Gawler 2008) with standardized terminology and compatible habitat classifications. It allows the Action Plan to describe the aspects of conservation which are particular to Vermont, while facilitating conservation at a broader regional level. A Companion to the Terrestrial and Aquatic Maps has been published by The Nature Conservancy (Anderson et al. 2013). It includes profiles of each habitat type in the Northeast, distribution maps, state acreage figures, SGCN identification concern, and an assessment of overall conditions in the region. Habitat conservation summaries can be found in Appendix B.

Habitat Succession, Species of Greatest Conservation Need & the Action Plan

Plant succession produces cumulative change in the types of plant species occupying a given area through time. Succession is complicated by factors such as disturbance (large and small), local conditions, seed banks and soil legacies (Oliver 1981). A highly simplified timeline begins when land is cleared. Pioneer species typically return first followed by other species generally better adapted to the new and changing conditions created by the previous suite of species. Given sufficient time and appropriate conditions the area moves roughly through early, middle, and late successional stages—often referred to as mature or old growth. A disturbance, if sufficiently large, can re-set the clock anytime and succession begins again. The best known examples are forest succession but it occurs in virtually all vegetated areas. For example, lichen communities on granite mountaintops experience successional changes (Wessels 2002).

Succession can significantly impact habitat for Species of Greatest Conservation Need and other wildlife as in the edge habitat example noted earlier. Generally as succession moves from early (young forests) to late stages some wildlife will lose out (e.g., Spruce Grouse, American Woodcock, Cottontail Rabbit) and others will benefit (e.g., American Marten, Northern Goshawk). Others still prefer a mix of successional stages in appropriate configurations (e.g., Black Bear, Canada Lynx).

Over the past two centuries the mix of successional stages available to Vermont's wildlife has changed dramatically in both distribution and abundance. Though precise estimates (current and historic) are unavailable, prior to 1800 a significant percentage of Vermont's forests were in late-successional stages (>150-300 years and older) and forest stands provided greater structural diversity. One-hundred years later young forests (early-successional stages of 1-15 years) dominated the state and today mid-successional forests (60-100 years) are most abundant. Wildlife populations have responded in turn. Vermont's SGCN list contains relatively few species requiring mid-successional forests and more that thrive in early and late-successional representations.

Because the loss of late-successional forests in the eastern U.S. occurred prior to the advent of modern wildlife biology and the current scarcity of later-successional stages (particularly northern hardwood forest types) our understanding how wildlife utilized these stages is not as advanced as our knowledge of wildlife in early successional stages. Historic records and research in late-successional areas elsewhere indicate that the distribution and abundance of some wildlife species was much greater when late-successional forests were in greater abundance—even if these species

can survive without them. Given the lack of this condition on the landscape it is advisable to increase its availability to wildlife.

The habitat, community and landscape summaries that follow here and in Chapter 6 and Appendix B address the habitat the needs of Species of Greatest Conservation Need that use that vegetation types in one or more successional stages. Conservation strategies address these particular successional stage needs as well as those of species that prefer a mosaic of successional stages.

Table 4.2: Landscape, Community, Habitat & Cover Type Categories

* Categories marked with an asterisk "*" are considered major categories for the purposes of organizing this report (24 in all). Conservation summaries were developed addressing characteristics and location, current and desired condition, SGCN using this habitat category, priority problems impacting this category, conservation strategies to address the problems and a list of other plans and planning entities with significant interest in this area.

***Landscapes**

Interior Forest Blocks
Connectivity Blocks
Surface Waters and Riparian Areas
Riparian Areas for Connectivity
Physical Landscape Diversity Blocks
Wildlife Road Crossing

Aquatic Communities

***Riparian Areas**

***Riverine** (Langdon et.al. 1998)

Brook trout community
Brook trout-slimy sculpin community
Blacknose dace-slimy sculpin community
Blacknose dace-bluntnose minnow community
Blacknose dace-creek chub community
Tessellated darter-fallfish community
Blacknose dace-slimy sculpin community
White sucker-tessellated darter community

***Lower Connecticut River**

(Atlantic salmon-American shad community)

***Lower Lake Champlain Tributaries**

(Redhorse-lake sturgeon community)

Cultural Habitats

(Reschle 1990)

***Building & structures**

***Mine & Gravel Pits**

***Grassland & Hedgerows**

Grasslands
Hedgerow
Old field/shrub
Orchard

***Lakes & Ponds**

Dystrophic lakes
Meso-eutrophic lakes
Oligotrophic lakes
High elevation acidic lakes

***Lake Champlain**

***Young Forests**

(Successional Stages, Forest Cover Types, Eyre 1980, US Dept of Agriculture 2003)

Stages: Seedling/Sapling Sapling/Pole Timber, Pole Timber

Cover types

Boreal Conifers

Balsam fir
Black spruce
White spruce

Boreal Hardwoods

Aspen
Pin cherry
Paper birch

Spruce-Fir

Red spruce
Red spruce-balsam fir
Paper birch-red spruce-balsam fir

Pine and Hemlock

Eastern white pine

Table 4.2 continued: Terrestrial Natural Communities (Thompson & Sorenson 2005)

Open or Shrub Wetlands

***Open Peatlands**

Alpine peatland
Dwarf shrub bog
Black spruce woodland bog
Pitch pine woodland bog
Poor fen
Rich fen
Intermediate fen

***Marshes & Sedge Meadows**

Deep bulrush marsh
Deep broadleaf marsh
Shallow emergent marsh
Sedge meadow
Cattail marsh
Wild rice marsh

***Wet Shores**

Calcareous riverside seep
River cobble shore
Lakeshore grassland
Riverside sand or gravel shore
Outwash plain pondshore
River mud shore
Rivershore grassland

***Shrub Swamps**

Buttonbush basin swamp
Alder swamp
Alluvial shrub swamp
Sweet gale shoreline swamp
Buttonbush swamp

Forested Wetlands

***Floodplain Forests**

Silver maple-ostrich fern riverine floodplain forest
Lakeside floodplain forest
Silver maple-sensitive fern riverine floodplain forest
Sugar maple-ostrich fern riverine floodplain forest

***Hardwood Swamps**

Red maple-black ash swamp
Red maple-northern white cedar swamp
Calcareous red maple-tamarack swamp
Red or silver maple-green ash swamp
Red maple-black gum swamp
Red maple-white pine-huckleberry swamp

***Softwood Swamps**

Northern white cedar swamp
Spruce-fir-tamarack swamp
Black spruce swamp
Hemlock swamp

***Seeps & Vernal Pools**

Vernal pools
Seeps

Open Upland Communities

***Upland shores**

Riverside outcrop
Lake sand beach
Lake shale or cobble beach
Erosional river bluff
Sand dune

***Outcrops & Upland Meadows**

Alpine meadow
Boreal outcrop
Serpentine outcrop
Temperate acidic outcrop
Temperate calcareous outcrop

***Cliffs & Talus**

Boreal acidic cliff
Boreal calcareous cliff
Temperate acidic cliff
Temperate calcareous cliff
Open talus

Upland Forests & Woodlands

***Spruce-Fir Northern Hardwood Forest**

Subalpine krummholz
Montane spruce-fir forest
Lowland spruce-fir forest
Montane yellow birch-red spruce forest
Boreal talus woodland
Cold-air talus woodland
Red spruce-northern hardwood forest
Red Spruce-Heath Rocky Ridge Forest

***Northern Hardwood Forest**

Northern hardwood forest
Rich northern hardwood forest
Mesic red oak-northern hardwood forest
Hemlock forest
Hemlock-northern hardwood forest
Northern hardwood talus woodland

***Oak-Pine-Northern Hardwood Forest**

Limestone bluff cedar-pine forest
Mesic maple-ash-hickory-oak forest
Mesic Clayplain Forest
White pine-red oak-black oak forest
Dry oak forest
Dry Red Oak-White Pine Forest
Pine-oak-heath sandplain forest
Dry oak-hickory-hophornbeam forest
Red cedar woodland
Red pine forest or woodland
Pitch pine-oak-heath rocky summit
Dry oak woodland
Sand-Over-Clay Forest
Temperate Hemlock Forest
Temperate Hemlock-Hardwood Forest
Transition Hardwoods Limestone Forest

Literature Cited

- Allen, T., F.H.; Starr, 1982. *Hierarchy: perspectives for ecological complexity*. Chicago: The University of Chicago Press.
- Anderson, M.G., M. Clark, C.E. Ferree, A. Jospe, A. Olivero Sheldon, and K.J. Weaver. 2013. Northeast Habitat Guides: A companion to the terrestrial and aquatic habitat maps. Submitted to the Regional Conservation Needs Grants Program of the Northeast Association of Fish and Wildlife Agencies. The Nature Conservancy, Eastern Conservation Science, Eastern Regional Office. Boston, MA.
<http://rcngrants.org/content/guide-terrestrial-habitat-map>.
- Association of Fish and Wildlife Agencies (AFWA), Teaming With Wildlife Committee, State Wildlife Action Plan Best Practices Working Group. 2012. *Best Practices for State Wildlife Action Plans—Voluntary Guidance to States for Revision and Implementation*. Washington, DC: Association of Fish and Wildlife Agencies. www.fishwildlife.org/files/SWAPBestPractices.pdf.
- Barnes, R.B. 1979. Wildlife habitat from a forest resource inventory: is it possible? *Transactions of the Northeast Section of the Wildlife Society* 36:151-159.
- Cleland D.T, P.E. Avers, W.H McNab, M.E. Jensen, R.G. Bailey, T. King, and W.E. Russell 1997. *National Hierarchical Framework of Ecological Units. Ecosystem Management: Applications for Sustainable Forest and Wildlife Resources.*, Mark S. Boyce and Alan Haney, editors. Yale University Press, New Haven.
- Eyre, F. H. ed. 1980. *Forest Cover Types of the United States*. Society of American Foresters. Washington, DC.
- Forman, R.T.T.; Godron, M. 1986. *Landscape Ecology*. New York: Wiley and Sons
- Gawler, S. C. 2008. *Northeastern Terrestrial Wildlife Habitat Classification*. Report to the Virginia Department of Game and Inland Fisheries on behalf of the Northeast Association of Fish and Wildlife Agencies and the National Fish and Wildlife Foundation. NatureServe, Boston, Massachusetts.
- Grossman, D.H., D. Faber-Langendoen, A.W. Weakley, M. Anderson, P. Bourgeron, R. Crawford, K. Goodin, S. Landaal, K. Metzler, K.D. Patterson, M. Pyne, M. Reid and L. Sneddon. 1998. *International Classification of Ecological Communities: Terrestrial Vegetation of the United States. Volume I: The National Vegetation Classification Standard*. The Nature Conservancy.
- Langdon, R., J. Andrews, K. Cox, S. Fiske, N. Kamman, and S. Warren. 1998. *A classification of the aquatic communities of Vermont*. The Nature Conservancy and the Vermont Biodiversity Project, Montpelier, VT.
- Oliver, C.D. 1981. Forest Development in North America Following Major Disturbances. *Forest Ecology and Management* 3:153-168
- Poiani, K.P., B.D. Richter, M.G. Anderson, and H.E. Richter. 2000. Conservation at multiple scales: functional sites, landscapes, and networks. *BioScience* 50:133-146
- Smith, R.L. 1980 *Ecology and Field Biology*, 3th edition. Harper & Row. New York.
- Reschke, Carol. 1990. *Ecological Communities of New York State*. Latham, NY: New York Natural Heritage Program, NYSDEC, www.dec.state.ny.us/website/dfwmr/heritage/EcolComm.htm
- Simberloff D. 1998. Flagships, umbrellas, and keystones: Is single-species management passe in the landscape era? *Biological Conservation* 83:247–257.
- Thompson, E. H., and E. R. Sorenson. 2005. *Wetland, Woodland, Wildland—A guide to the natural communities of Vermont*. University Press of New England, Hanover.
- U.S. Department of Agriculture, Forest Service. 2003. *Forest inventory and analysis national core field guide, volume 1*. U.S. Department of Agriculture, Forest Service, Forest Inventory and Analysis, 201 14th St., Washington, D.C., 20250.
- Wessels, T. 2002. *The Granite Landscape: A Natural History of America's Mountain Domes, From Acadia to Yosemite*. The Countryman Press. Woodstock, VT

Vermont's Wildlife Action Plan

Chapter 5

Species of Greatest Conservation Need

Draft • September, 2015

Vermont's Species of Greatest Conservation Need	1
SGCN Lists.....	4
Conserving Vermont's Amphibians & Reptiles.....	14
Conserving Vermont's Birds.....	20
Conserving Vermont's Fishes	28
Conserving Vermont's Invertebrates.....	35
Conserving Vermont's Mammals	46
Conserving Vermont's Plants	55

5. Vermont's Species of Greatest Conservation Need

Vermonters love their wildlife. And wildlife love Vermont. During the past century many wildlife species once rare or missing from the state have returned in larger numbers. The resurgence of Vermont's forests is a significant reason. From a low of 40% forest cover in the 1840s the state is now nearly 80% forested. However, more trees are not the whole story. Restoring wildlife to the state also required the hard work and dedication of scientists, wildlife and habitat managers, sportsmen, and other conservationists. Signature species such as deer, moose, beaver, fisher, osprey, peregrine falcon, and loon, all missing or in perilously low numbers just decades ago are now faring well.

Keeping wildlife populations healthy offers a host of benefits: healthier ecosystems upon which we all depend; more wildlife to enjoy; and, fewer species on the brink of extirpation means fewer regulatory mandates.

Our work, however, is not complete. A significant number of wildlife species need help because of threats such as habitat loss, fragmentation and degradation; invasive exotic species; diseases; and climate change.

The State Wildlife Grants program helps Vermont meet these challenges. It provides federal funds for conservation to prevent fish and wildlife populations from becoming endangered. Per Congressional requirements, the Wildlife Action Plan is centered on the identification and conservation of Species of Greatest Conservation Need (SGCN).

Selecting Species of Greatest Conservation Need

Vermont's Species of Greatest Conservation Need include 133 amphibians and reptiles, birds, fishes, invertebrates, mammals (vertebrates) of 470 in the state and 200 invertebrate species from out of an estimated 21,000. For plants, 645 species of an estimated 2000 vascular and non-vascular species were selected.

Some of these SGCN have official protection status (e.g., threatened, endangered) whereas others may be in decline but are not currently listed as part of either the Federal or State Endangered Species program. One guiding principle of the Wildlife Action Plan is to direct conservation attention to species and habitats *before* they become imperiled and recovery becomes more difficult and costly. Some of the species on the list may be relatively common including some game species. It is our goal to keep them common.

In Vermont, six Species Teams, with expertise in amphibians and reptiles, birds, fish, invertebrates, mammals and plants assessed the status of Vermont's native species. They applied assessment criteria such as the degree of species rarity, species designated as at-risk, population trends, species whose habitat are vulnerable to loss, habitat fragmentation, habitat conversion or succession changes and species threatened by exotic plants or animals. Changes to the SGCN list are summarized in table 4.1 and details of SGCN selection process can be found in Appendix C—Developing Vermont's Wildlife Action Plan.

Table 5.1 Changes to SGCN Lists 2005:2015

Taxon	2005 SGCN	2015 SGCN	Change Notes
Amphibians & Reptiles	19	19	No changes
Birds	58	51	Long-eared Owl, Henslow's Sparrow, Osprey, Cooper's Hawk, Barn Owl, Veery, Blue-winged Teal removed
Fishes	33	27	Arctic Char, Atlantic Salmon (anadromous), Brassy Minnow, Muskellunge, Quillback and the CT River Sea Lamprey removed
Invertebrates	191	200	19 species removed from the list; 28 added, including 9 bumble bee species
Mammals	33	34	Mink removed. Moose and Snowshoe Hare added
Plants	577	645	

Teams used the best information available at the time from local, regional, and national sources. However, while a wealth of information is available for some species, others (especially invertebrates, fish, small mammals and some reptiles and amphibians) are poorly known. Species were ranked with a conservation priority of high, medium or low. Those ranked medium and high constitute Vermont's Species of Greatest Conservation Need. Those ranked low priority are considered reasonably secure. It is expected that low priority species will benefit from conservation efforts directed toward species ranked medium and high as well as from other ongoing wildlife management programs (e.g., federal aid to sportfish and wildlife).

Ongoing wildlife monitoring will help track species status and progress toward greater security. Regular Action Plan review and revision will provide opportunities to add additional species to the list as warranted and to remove those species deemed secure.

Use of and Changes to this List

The list of Species of Greatest Conservation Need will help prioritize the allocation of State Wildlife Grant and other conservation funds. The list will also provide a quick measure of our success conserving Vermont's wildlife. It also raises the profile of a species to a wider audience of conservation partners and can encourage others to initiate projects that may benefit the species. It is important to note that presence on this list does not necessarily mean that conservation resources *will* be directed towards the animal or plant, but that conservation actions for the animal species are *eligible* for State Wildlife Grants funding, and may be more competitive for other grant programs.

The Species of Greatest Conservation Need list can be amended if and when important information becomes available about a species' status. For example, there are a number of current and pending inventory and assessment projects funded by State Wildlife Grants that could significantly increase our understanding of a species' status.

Big Game: White-Tailed Deer & Wild Turkey

Nearly 20 game and sportfish species are listed on the following pages as Species of Greatest Conservation Need (SGCN) due to concerns about population declines and loss of habitat. White-tailed deer and wild turkey, however, were not selected as SGCN. Though absent or

nearly extirpated from the state by 1865, their populations are now sufficiently large and stable. And, relative to SGCN, our knowledge of deer and turkey biology and management is great.

White-tailed deer and wild turkey rank high among Vermont's greatest wildlife restoration successes. Still their management remains of utmost concern because of the great importance they have to Vermonters and because of the significant roles they play in their ecosystems. Fortunately, management plans (developed with significant public involvement), harvest regulations and monitoring protocols have long been in place for these species and dependable implementation funds come through license fees and the Federal Aid to Wildlife Restoration Act.

Vermont Fish & Wildlife Department's [Big Game Management Plan \(2010-2020\)](#) is incorporated into the Wildlife Action Plan as Appendix I.

Amphibian & Reptile Species of Greatest Conservation Need

¹ Regional Species of Greatest Conservation Need in the Northeastern United States (Terwilliger, 2013)

High Priority

Jefferson Salamander (*Ambystoma jeffersonianum*- and hybrids)¹
Mudpuppy (*Necturus maculosus*)¹
Fowler's Toad (*Bufo fowleri*)¹
Boreal Chorus Frog (*Pseudacris maculata*)
Spotted Turtle (*Clemmys guttata*)¹
Wood Turtle (*Glyptemys insculpta*)¹
Spiny Softshell Turtle (*Apalone spinifera*)¹
Five-lined Skink (*Plestiodon fasciatus*)
North American Racer (*Coluber constrictor*)¹
Eastern Ratsnake (*Pantherophis alleghaniensis*)
Timber Rattlesnake (*Crotalus horridus*)¹

Medium Priority

Blue-spotted Salamander (*Ambystoma laterale*)¹
Spotted Salamander (*Ambystoma maculatum*)
Four-toed Salamander (*Hemidactylium scutatum*)
Common Musk Turtle (*Sternotherus odoratus*)
Northern Water Snake (*Nerodia sipedon*)
DeKay's Brownsnake (*Storeria dekayi*)¹
Eastern Ribbonsnake (*Thamnophis sauritus*)¹
Smooth Greensnake (*Liochlorophis vernalis*)¹

Bird Species of Greatest Conservation Need

High Priority

Pied-billed Grebe (*Podilymbus podiceps*)¹
Least Bittern (*Ixobrychus exilis*)¹
American Black Duck (*Anas rubripes*)¹
Bald Eagle (*Haliaeetus leucocephalus*)
Northern Harrier (*Circus cyaneus*)¹
Red-shouldered Hawk (*Buteo lineatus*)¹
American Kestrel (*Falco sparverius*)¹
Spruce Grouse (*Falcapennis canadensis*)¹
Upland Sandpiper (*Bartramia longicauda*)¹
Common Tern (*Sterna hirundo*)¹
Black Tern (*Chlidonias niger*)¹
Common Nighthawk (*Chordeiles minor*)¹
Whip-poor-will (*Caprimulgus vociferus*)¹
Chimney Swift (*Chaetura pelagica*)¹
Olive-sided Flycatcher (*Contopus cooperi*)¹
Gray Jay (*Perisoreus canadensis*)
Sedge Wren (*Cistothorus platensis*)¹
Bicknell's Thrush (*Catharus bicknelli*)¹
Wood Thrush (*Hylocichla mustelina*)¹
Brown Thrasher (*Toxostoma rufum*)¹
Blue-winged Warbler (*Vermivora pinus*)¹
Golden-winged Warbler (*Vermivora chrysoptera*)¹
Blackpoll Warbler (*Dendroica striata*)
Canada Warbler (*Wilsonia canadensis*)¹
Eastern Towhee (*Pipilo erythrophthalmus*)¹
Vesper Sparrow (*Pooecetes gramineus*)¹
Grasshopper Sparrow (*Ammodramus savannarum*)¹
Eastern Meadowlark (*Sturnella magna*)¹
Rusty Blackbird (*Euphagus carolinus*)¹

Medium Priority

Common Loon (*Gavia immer*)¹
American Bittern (*Botaurus lentiginosus*)¹
Great Blue Heron (*Ardea herodias*)
Black-crowned Night-heron (*Nycticorax nycticorax*)¹
Northern Goshawk (*Accipiter gentilis*)¹
Peregrine Falcon (*Falco peregrinus*)¹
Ruffed Grouse (*Bonasa umbellus*)¹
Sora (*Porzana carolina*)¹
Lesser Yellowlegs (*Tringa flavipes*)
American Woodcock (*Scolopax minor*)¹
Black-billed Cuckoo (*Coccyzus erythrophthalmus*)¹
Barn Owl (*Tyto alba*)¹
Short-eared Owl (*Asio flammeus*)¹
Black-backed Woodpecker (*Picoides arcticus*)
Purple Martin (*Progne subis*)
Chestnut-sided Warbler (*Dendroica pensylvanica*)
Black-throated Blue Warbler (*Dendroica caerulescens*)¹
Prairie Warbler (*Dendroica discolor*)¹
Bay-breasted Warbler (*Dendroica castanea*)¹
Cerulean Warbler (*Dendroica cerulea*)¹
Field Sparrow (*Spizella pusilla*)¹
Bobolink (*Dolichonyx oryzivorus*)¹

Fish Species of Greatest Conservation Need

¹Regional Species of Greatest Conservation Need in the Northeastern United States (Terwilliger, 2013)

High Priority

Northern Brook Lamprey (*Ichthyomyzon fossor*)¹
Silver Lamprey (*Ichthyomyzon unicuspis*)¹
American Brook Lamprey (Lethenteron appendix)¹
Lake Sturgeon (*Acipenser fulvescens*)¹
Round Whitefish (*Prosopium cylindraceum*)¹
Bridle Shiner (*Notropis bifrenatus*)¹
Blackchin Shiner (*Notropis heterodon*)
Blacknose Shiner (*Notropis heterolepis*)
Greater Redhorse (*Moxostoma valenciennesi*)
Stonecat (*Noturus flavus*)¹
Eastern Sand Darter (*Ammocrypta pellucida*)¹
Channel Darter (*Percina copelandi*)¹
Sauger (*Sander canadense*)¹

Medium Priority

Blueback Herring (CT River population) (*Alosa aestivalis*)¹
Atlantic Salmon (naturally reproducing populations-Lake Champlain & Memphremagog basins) (*Salmo salar*)¹
Mottled Sculpin (*Cottus bairdi*)
American Eel (Lk Champlain pop) (*Anguilla rostrata*)¹
American Eel (CT River pop) (*Anguilla rostrata*)¹
American Shad (*Alosa sapidissima*)¹
Mooneye (*Hiodon tergisus*)¹
Cisco or Lake Herring (*Coregonus artedii*)
Lake Whitefish (*Coregonus clupeaformis*)¹
Brook Trout (naturally reproducing populations) (*Salvelinus fontinalis*)¹
Lake Trout (naturally reproducing populations) (*Salvelinus namaycush*)¹
Redfin Pickerel (*Esox americanus*)
Silver Redhorse (*Moxostoma anisurum*)
Shorthead Redhorse (*Moxostoma macrolepidotum*)
Redbreast Sunfish (*Lepomis auritus*)¹

Invertebrate Species of Greatest Conservation Need

MP=Medium Priority SGCN; HP=High Priority SGCN.

¹Regional Species of Greatest Conservation Need in the Northeastern United States (Terwilliger, 2013)

²Regional responsibility species identified in (White et al. 2014).

Ant Group (2)

An ant (*Myrmica lobifrons*) MP
A Slave-making Ant (*Temnothorax pilagens*) HP

Bumble Bee Group (9)

Rusty-patched Bumble Bee (*Bombus affinis*) HP
Ashton Cuckoo Bumble Bee (*Bombus ashtoni*) HP
Lemon Cuckoo Bumble Bee (*Bombus citrinus*) HP
Fernald's Cuckoo Bumble Bee (*Bombus fernaldae*) HP
Yellow Bumble Bee (*Bombus fervidus*) HP
American Bumble Bee (*Bombus pensylvanicus*) HP
Confusing Bumble Bee (*Bombus perplexus*) HP
Red-belted Bumble Bee (*Bombus rufocinctus*) MP
Yellow-banded Bumble Bee (*Bombus terricola*) HP

Beetles-Tiger Beetle Group (7)

Boulder-beach Tiger Beetle (*Cicindela ancocisconensis*)¹ HP
Hairy-necked Tiger Beetle (*Cicindela hirticollis hirticollis*) HP
Boreal Long-lipped Tiger Beetle (*Cicindela longilabris*) HP
Cobblestone Tiger Beetle (*Cicindela marginipennis*)¹ HP
Northern Barrens Tiger Beetle (*Cicindela patruela*)¹ HP
Puritan Tiger Beetle (*Cicindela puritana*)¹ HP
Eastern Red-bellied Tiger Beetle (*Cicindela rufiventris*) HP

Butterflies-Grassland Group (4)

Dusted Skipper (*Atrytonopsis hianna*) MP
Cobweb Skipper (*Hesperia metea*) MP
Monarch (*Danaus plexippus*) HP
Regal Fritillary (*Speyeria idalia*) MP

Butterflies-Hardwood Forest Group (4)

Early Hairstreak (*Erora laeta*) HP
Hackberry Emperor (*Asterocampa celtis*) MP
Tawny Emperor (*Asterocampa clyton*) MP
West Virginia White (*Pieris virginiensis*) HP

Butterflies-Wetland Group (6)

Two-spotted Skipper (*Euphyes bimacula*) HP
Black Dash (*Euphyes conspicua*) HP
Dion Skipper (*Euphyes dion*) MP
Mulberry Wing (*Poanes massasoit*) HP
Bog Copper (*Lycaena epixanthe*) HP
Jutta Arctic (*Oeneis jutta*) HP

Ground Beetles-Carabid Group (67)

Agonum crenistriatum HP
Agonum darlingtoni HP
Agonum moerens MP
Agonum picicornoides HP
Agonum punctiforme MP
Agonum superioris MP
Amara erraticus HP
Amara laevipennis MP
Apristus latens HP
Atraneus pubescens MP
Bembidion affine MP
Bembidion cordatum MP
Bembidion grapii HP
Bembidion mutatum HP
Bembidion quadratulum HP
Bembidion robusticollis MP
Bembidion rolandi MP
Bembidion rufotinctum HP
Blethisa hudsonica MP
Blethisa julii HP
Blethisa quadricollis HP
Carabus goryi MP
Carabus maeander MP
Dicaelus dilatatus dilatatus HP
Dicaelus teter HP
Dicheirotichus cognatus HP
Diplocheila impressicollis MP
Diplocheila striatopunctata HP
Dyschirius brevispinus MP
Dyschirius erythrocerus MP
Dyschirius politus politus HP
Elaphropus dolosus MP
Elaphropus levipes MP
Elaphrus fuliginosus HP
Geopinus incrassatus HP
Harpalus fulvilabris HP
Harpalus indigens MP
Harpalus providens MP
Lophoglossus scrutator HP
Nebria suturalis HP
Notiobia sayi MP
Notiophilus aquaticus MP
Notiophilus borealis HP
Notiophilus nemoralis HP
Notiophilus novemstriatus MP
Olisthopus micans HP
Patrobus foveocollis HP
Pentagonica picticornis MP
Pericompsus ephippiatus MP
Philodes alternans HP
Philodes rectangulus MP
Platynus cincticollis MP
Platypatrobus lacustris MP
Pseudamara arenaria MP
Pterostichus brevicornis brevicornis HP
Pterostichus castor MP
Pterostichus pinguedineus HP
Pterostichus punctatissimus HP
Scaphinotus bilobus MP
Schizogenius ferrugineus MP

Sericoda obsoleta MP
Sericoda quadripunctata MP
Sphaeroderus nitidicollis HP
Tachys oblitus MP
Tachys rhodeanus HP
Tetragonoderus fasciatus MP
Tetraleucus picticornis MP

Moths Group

A Noctuid Moth (*Zale submediana*) HP
Pine Barrens Zanclognatha (*Zanclognatha martha*) HP
Currant Spanworm (*Speranza ribearia*) HP
A Ghost Moth (*Sthenopsis thule*) MP
A Noctuid Moth (*Lasionycta taigata*) MP
A Noctuid Moth (*Lemmeria digitalis*) MP
Franclemont's Lithophane (*Lithophane franclemonti*) HP
An Autumnal Noctuid Moth (*Pachypolia atricornis*) HP
Ostrich Fern Borer (*Papaipema* sp. 2 nr. *Pterisii*) HP
Barrens Moth (*Properigea costa*) MP
A Noctuid Moth (*Xestia fabulosa*) MP
A Noctuid Moth (*Xestia homogenea*) HP
Pine Imperial Moth (*Eacles imperialis pini*) HP
New England Buckmoth (*Hemileuca lucina*) MP
Plum Sphinx (*Sphinx drupiferarum*) HP
Clemens' Sphinx (*Sphinx luscitiosa*) HP
A tortricid moth (*Eana georgiella*) MP

Mayflies/Stoneflies/Caddisflies Group (14)

A Caddisfly (*Ceraclea submacula*) HP
A Caddisfly (*Polycentropus glacialis*) HP
A Caddisfly (*Polycentropus iculus*) HP
A Caddisfly (*Rhyacophila amicus*) HP
A Caddisfly (*Rhyacophila brunnea*) HP
A Mayfly (*Ameletus browni*) HP
A Mayfly (*Baetisca rubescens*) HP
A Mayfly (*Eurylophella bicoloroides*) HP
Roaring Brook Mayfly (*Epeorus frisoni*) HP
Tomah Mayfly (*Siphonisca aerodromia*) HP
A Mayfly (*Siphonurus demaryi*) HP
Lawrence Sallfly (*Alloperla voinae*) HP
Appalachian Stonefly (*Hansonoperla appalachia*) HP
Spiny Salmonfly (*Pteronarcys comstocki*) HP

Odonates-Bog/Fen/Swamp/Marshy Pond Group (15)

Mottled Darner (*Aeshna clepsydra*) HP
Zigzag Darner (*Aeshna sitchensis*)² HP
Subarctic Darner (*Aeshna subarctica*)² HP
Comet Darner (*Anax longipes*) HP
Swamp Darner (*Epiaeschna heros*) HP
Cyrano Darner (*Nasiaeschna pentacantha*) HP
Spatterdock Darner (*Rhionaeschna mutata*) HP
Subarctic Bluet (*Coenagrion interrogatum*) HP
Petite Emerald (*Dorocordulia lepida*) HP
Ski-tailed Emerald (*Somatochlora elongata*)² HP
Forcinate Emerald (*Somatochlora forcipata*)² HP
Delicate Emerald (*Somatochlora franklini*)² HP
Kennedy's Emerald (*Somatochlora kennedyi*)² HP
Ebony Boghaunter (*Williamsonia fletcheri*) HP
Black Meadowhawk (*Sympetrum danae*)² HP

Odonates-Lakes/Ponds Group (7)

New England Bluet (*Enallagma laterale*)² HP
Slender Bluet (*Enallagma traviatum*) HP
Lilypad Forktail (*Ischnura kellicotti*) HP
Ringed Emerald (*Somatochlora albicincta*)² HP
Lake Emerald (*Somatochlora cingulata*) HP
Banded Pennant (*Celithemis fasciata*)² HP
Carolina Saddlebags (*Tramea carolina*) HP

Odonates-River/Stream Group (17)

American Rubyspot (*Hetaerina americana*) HP
Blue-fronted Dancer (*Argia apicalis*) HP
River Bluet (*Enallagma anna*)² HP
Rainbow Bluet (*Enallagma antennatum*)² HP
Big Bluet (*Enallagma durum*) HP
Stygian Shadowdragon (*Neurocordulia yamaskanensis*) HP
Spine-crowned Clubtail (*Gomphus abbreviatus*) HP
Midland Clubtail (*Gomphus fraternus*) HP
Rapids Clubtail (*Gomphus quadricolor*) HP
Cobra Clubtail (*Gomphus vastus*) HP
Skillet Clubtail (*Gomphus ventricosus*)² HP
Brook Snaketail (*Ophiogomphus aspersus*) HP
Riffle Snaketail (*Ophiogomphus carolus*) HP
Maine Snaketail (*Ophiogomphus mainensis*) HP
Rusty Snaketail (*Ophiogomphus rupinsulensis*) HP
Riverine Clubtail (*Stylurus amnicola*)² HP
Zebra Clubtail (*Stylurus scudder*) HP

Crustaceans Group (3)

Appalachian Brook Crayfish (*Cambarus bartonii*) HP
Taconic Cave Amphipod (*Stygobromus borealis*) HP
An Amphipod (*Diporeia hoyi*) HP

Freshwater Mussels Group (13)

Eastern Pearlshell (*Margaritifera margaritifera*)¹ MP
Dwarf Wedgemussel (*Alasmodonta heterodon*)¹ HP
Elktoe (*Alasmodonta marginata*)¹ HP
Brook Floater (*Alasmodonta varicosa*)¹ HP
Alewife Floater (*Anodonta implicata*)¹ MP
Cylindrical Papershell (*Anodontoides ferussacianus*)¹ MP
Pocketbook (*Lampsilis ovata*)¹ HP
Creek Heelsplitter (*Lasmigona compressa*)¹ HP
Fluted-shell (*Lasmigona costata*) HP
Fragile Papershell (*Leptodea fragilis*)¹ HP
Black Sandshell (*Ligumia recta*)⁴ HP
Pink Heelsplitter (*Potamilus alatus*) HP
Giant Floater (*Pyganodon grandis*) MP

Freshwater Snails Group (15)

Buffalo Pebblesnail (*Gillia altilis*) HP
Squat Duskysnail (*Lyogyrus granum*) HP
Pupa Duskysnail (*Lyogyrus pupoideus*) HP
Canadian Duskysnail (*Lyogyrus walkeri*) HP
Boreal Marstonia (*Marstonia lustrica*) HP
Spindle Lymnaea (*Acella haldemani*) HP
Mammoth Lymnaea (*Bulimnaea megasoma*) HP
Country Fossaria (*Fossaria rustica*) HP
Star Gyro (*Gyraulus crista*) MP
Dusky Ancyloid (*Laevapex fuscus*) MP
Thicklip Rams-horn (*Planorbula armigera*) MP
Liver Elimia (*Goniobasis livescens*) HP
Sharp Hornsnail (*Pleurocera acuta*) MP
Fringed Valvata (*Valvata lewisi*) HP
Mossy Valvata (*Valvata sincera*) HP

Mammal Species of Greatest Conservation Need

High Priority

Water Shrew (*Sorex palustris*)¹
Long-tailed or Rock Shrew (*Sorex dispar*)¹
Pygmy Shrew (*Sorex hoyi*)
Little Brown Bat (*Myotis lucifugus*)¹
Indiana Bat (*Myotis sodalis*)¹
Small-footed Bat (*Myotis leibii*)¹
Northern Long-eared Bat (*Myotis septentrionalis*)¹
Silver-haired Bat (*Lasionycteris noctivagans*)¹
Tri-colored Bat (*Perimyotis subflavus*)¹
Eastern Red Bat (*Lasiurus borealis*)¹
Hoary Bat (*Lasiurus cinereus*)¹
New England Cottontail (*Sylvilagus transitionalis*)¹
Rock Vole (*Microtus chrotorrhinus*)
Woodland Vole (*Microtus pinetorum*)
Northern Bog Lemming (*Synaptomys borealis*)
American Marten (*Martes americana*)¹
Canada Lynx (*Lynx canadensis*)

Medium Priority

Cinereus or Masked Shrew (*Sorex cinereus*)
Smoky Shrew (*Sorex fumeus*)¹
Hairy-tailed Mole (*Parascalops breweri*)¹
Big Brown Bat (*Eptesicus fuscus*)¹
Snowshoe Hare (*Lepus americanus*)
Southern Flying Squirrel (*Glaucomys volans*)
Northern Flying Squirrel (*Glaucomys sabrinus*)
Muskrat (*Ondatra zibethicus*)
Southern Bog Lemming (*Synaptomys cooperi*)¹
Wolf (*Canis sp?*)
Gray Fox (*Urocyon cinereoargenteus*)
Black Bear (*Ursus americanus*)
Long-tailed Weasel (*Mustela frenata*)
Mink (*Mustela vison*)
Northern River Otter (*Lontra canadensis*)
Bobcat (*Lynx rufus*)¹
Eastern Mountain Lion (*Puma concolor cougar*)
Moose (*Alces alces*)

Plant Species of Greatest Conservation Need

MP=Medium Priority SGCN; HP=High Priority SGCN

Conifers and relatives

Creeping Juniper (*Juniperus horizontalis*) HP

Ferns and relatives

Aleutian Maidenhair-fern (*Adiantum aleuticum*) HP

Green Mountain Maidenhair-fern (*Adiantum viridimontanum*) HP

Mountain Spleenwort (*Asplenium montanum*) HP

Green Spleenwort (*Asplenium viride*) HP

Upswept Moonwort (*Botrychium ascendens*) HP

Prairie Moonwort (*Botrychium campestre*) HP

Common Moonwort (*Botrychium lunaria*) HP

Mingan Moonwort (*Botrychium minganense*) HP

Blunt-lobed Grapefern (*Botrychium oneidense*) HP

Rugulose Grape-fern (*Botrychium rugulosum*) HP

Spatulate Moonwort (*Botrychium spathulatum*) MP

Shade Moonwort (*Botrychium tenebrosum*) HP

Weft Fern (*Crepidomanes intricatum*) HP

Laurentian Bladder Fern (*Cystopteris laurentiana*) HP

Northern Ground-cedar (*Diphasiastrum complanatum*) MP

Ground-fir (*Diphasiastrum sabinifolium*) MP

Male Fern (*Dryopteris filix-mas*) HP

Fragrant Fern (*Dryopteris fragrans*) MP

Marsh Horsetail (*Equisetum palustre*) MP

Mountain Fir Clubmoss (*Huperzia appressa*) HP

Fir Clubmoss (*Huperzia selago*) HP

Engelmann's Quillwort (*Isoetes engelmannii*) MP

Lake Quillwort (*Isoetes lacustris*) MP

River-bank Quillwort (*Isoetes riparia*) MP

Tuckerman's Quillwort (*Isoetes tuckermanii*) MP

Green Mountain Quillwort (*Isoetes viridimontana*) MP

Northern Adder's-tongue (*Ophioglossum pusillum*) HP

Massachusetts Fern (*Parathelypteris simulata*) MP

Stiff Clubmoss (*Spinulum canadense*) MP

Alpine Woodsia (*Woodsia alpina*) HP

Smooth Woodsia (*Woodsia glabella*) HP

Virginia Chain-fern (*Woodwardia virginica*) MP

Flowering Plants

Slender Copperleaf (*Acalypha gracilens*) MP

Yellow Giant Hyssop (*Agastache nepetoides*) HP

Purple Giant Hyssop (*Agastache scrophulariifolia*) HP

Boreal Bentgrass (*Agrostis mertensii*) MP

Wild Garlic (*Allium canadense* var. *canadense*) MP

Siberian Chives (*Allium schoenoprasum*) MP

Water Hemp (*Amaranthus tuberculatus*) HP

Small Round-leaved Orchis (*Amerorchis rotundifolia*) HP

Champlain Beach Grass (*Ammophila breviligulata* ssp. *champlainensis*) HP

Long-headed Thimbleweed (*Anemone cylindrica*) MP

Early Thimbleweed (*Anemone multifida* var. *multifida*) HP

Alpine Sweet-grass (*Anthoxanthum monticola* ssp. *monticola*) HP

White Camas (*Anticlea glauca*) HP

Putty-root (*Aplectrum hyemale*) HP

Lyre-leaved Rock-cress (*Arabidopsis lyrata*) MP

Dwarf Mistletoe (*Arceuthobium pusillum*) MP

Arethusa (*Arethusa bulbosa*) HP

Green Dragon (*Arisaema dracontium*) MP

Spiked Grass (*Aristida longespica* var. *geniculata*) MP

Boreal Wormwood (*Artemisia campestris* ssp. *canadensis*) HP

Beach Wormwood (*Artemisia campestris* ssp. *caudata*) HP

Blunt-leaved Milkweed (*Asclepias amplexicaulis*) MP

Whorled Milkweed (*Asclepias verticillata*) MP

Canada Milk-vetch (*Astragalus canadensis* var. *canadensis*) HP

Jesup's Milk-vetch (*Astragalus robbinsii* var. *jesupii*) HP

Blake's Milk-vetch (*Astragalus robbinsii* var. *minor*) HP

Smooth False-foxglove (*Aureolaria flava* var. *flava*) MP

Feverweed (*Aureolaria pedicularia*) MP

Downy False-foxglove (*Aureolaria virginica*) MP

Yellow Bartonian (*Bartonia virginica*) MP

Dwarf Birch (*Betula minor*) HP

Small Bidens (*Bidens discoidea*) MP

Downy Wood-mint (*Blephilia ciliata*) HP

Smooth Wood-mint (*Blephilia hirsuta* var. *glabrata*) HP

Hairy Wood-mint (*Blephilia hirsuta* var. *hirsuta*) HP

Strawberry Blite (*Blitum capitatum*) MP

Drummond's Rock-cress (*Boechera stricta*) MP

Green Rock-cress (*Borodinia missouriensis*) HP

Northern Rock-cress (*Braya humilis*) HP

Wild Chess (*Bromus kalmii*) MP

Langsdorf's Bluejoint (*Calamagrostis canadensis* var. *langsdorfii*) HP

Short-flower Bluejoint (*Calamagrostis canadensis* var. *macouniana*) HP

Pickering's Reed-grass (*Calamagrostis pickeringii*) MP

Bentgrass (*Calamagrostis stricta* ssp. *inexpansa*) HP

Northern Water-starwort (*Callitriche hermaphrodita*) HP

Large Water-starwort (*Callitriche heterophylla*) MP

Fairy Slipper (*Calypso bulbosa* var. *americana*) HP

Twin-flower Hedge Bindweed (*Calystegia silvatica* ssp. *fraterniflora*) HP

Low Bindweed (*Calystegia spithamea* ssp. *spithamea*) HP

Spring Cress (*Cardamine bulbosa*) MP

Cuckoo Flower (*Cardamine dentata*) HP

Small-flower bittercress (*Cardamine parviflora*) MP

Emmon's Sedge (*Carex albicans* var. *emmonsii*) MP

Foxtail Sedge (*Carex alopecoidea*) HP

Contracted Sedge (*Carex arcta*) MP

Awed Sedge (*Carex atherodes*) HP

Atlantic Sedge (*Carex atlantica* var. *atlantica*) MP

Howe's Sedge (*Carex atlantica* var. *capillacea*) MP

Blackish Sedge (*Carex atratiformis*) HP

Bicknell's Sedge (*Carex bicknellii*) HP

Bigelow's Sedge (*Carex bigelowii* ssp. *bigelowii*) HP

Bush's Sedge (*Carex bushii*) HP

Buxbaum's Sedge (*Carex buxbaumii*) MP

Capillary Sedge (*Carex capillaris* ssp. *capillaris*) HP

Creeping Sedge (*Carex chordorrhiza*) HP

Clustered Sedge (*Carex cumulata*) MP

Davis' Sedge (*Carex davisii*) HP
 Urchin Sedge (*Carex echinodes*) MP
 Bog Sedge (*Carex exilis*) MP
 Bronze Sedge (*Carex foenea*) MP
 Garber's Sedge (*Carex garberi*) HP
 Flaccid Sedge (*Carex glaucoidea*) HP
 Slender Sedge (*Carex gracilescens*) HP
 Pale Sedge (*Carex livida*) HP
 False Hop Sedge (*Carex lupuliformis*) MP
 Fernald's Sedge (*Carex merriitt-fernaldii*) MP
 Michaux Sedge (*Carex michauxiana*) MP
 Troublesome Sedge (*Carex molesta*) HP
 Nerveless Muehlenberg Sedge (*Carex muehlenbergii* var. *enervis*) HP
 Muehlenberg's Sedge (*Carex muehlenbergii* var. *muehlenbergii*) MP
 Few-fruited Sedge (*Carex oligocarpa*) HP
 Richardson's Sedge (*Carex richardsonii*) HP
 Schweinitz's Sedge (*Carex schweinitzii*) HP
 Scirpus-like Sedge (*Carex scirpoidea* ssp. *scirpoidea*) HP
 Hay Sedge (*Carex siccata*) MP
 Dioecious Sedge (*Carex sterilis*) HP
 Thin-flowered Sedge (*Carex tenuiflora*) HP
 Sheathed Sedge (*Carex vaginata*) HP
 Wiegand's Sedge (*Carex wiegandii*) MP
 Willdenow's Sedge (*Carex willdenowii*) HP
 Pignut Hickory (*Carya glabra*) MP
 Pale Painted-cup (*Castilleja septentrionalis*) HP
 Prairie Redroot (*Ceanothus herbaceus*) HP
 Nodding Chickweed (*Cerastium nutans* ssp. *nutans*) MP
 Wild Sensitive Plant (*Chamaecrista nictitans* var. *nictitans*) MP
 Bush's Goosefoot (*Chenopodium berlandieri* var. *bushianum*) HP
 Fogg's Goosefoot (*Chenopodium foggii*) HP
 Field Thistle (*Cirsium discolor*) MP
 Virginia Spring Beauty (*Claytonia virginica*) HP
 Small-flowered Collinsia (*Collinsia parviflora*) HP
 Canada Horse-balm (*Collinsonia canadensis*) MP
 Autumn Coral-root (*Corallorhiza odontorhiza* var. *odontorhiza*) MP
 Flowering Dogwood (*Cornus florida*) MP
 Golden Corydalis (*Corydalis aurea*) HP
 Pygmyweed (*Crassula aquatica*) MP
 Biltmore Hawthorn (*Crataegus biltmoreana*) MP
 Stinking Hawthorn (*Crataegus boyntonii*) HP
 Precocious Hawthorn (*Crataegus chrysocarpa* var. *praecox*) HP
 Dodge's Hawthorn (*Crataegus dodgei*) HP
 Zigzag Hawthorn (*Crataegus irrasa* var. *irrasa*) MP
 Kennedy's Hawthorn (*Crataegus kennedyi*) HP
 Oakes' Hawthorn (*Crataegus oakesiana*) HP
 Pea Hawthorn (*Crataegus pisifera*) HP
 Poplar Hawthorn (*Crataegus populnea*) HP
 Fleshy Hawthorn (*Crataegus succulenta* var. *succulenta*) HP
 Plains Frostweed (*Crocianthemum bicknellii*) MP
 Rattlebox (*Crotalaria sagittalis*) MP
 Buttonbush Dodder (*Cuscuta cephalanthi*) MP
 Northern Wild Comfrey (*Cynoglossum virginianum* var. *boreale*) HP
 Low Cyperus (*Cyperus diandrus*) MP
 Houghton's Cyperus (*Cyperus houghtonii*) HP
 Ram's Head Lady's-slipper (*Cypripedium arietinum*) HP
 Makasin's Yellow Lady's-slipper (*Cypripedium parviflorum* var. *makasin*) HP
 Tansy Mustard (*Descurainia pinnata* var. *brachycarpa*) HP
 Large-bracted Tick-trefoil (*Desmodium cuspidatum*) HP
 Perplexed Tick-trefoil (*Desmodium perplexum*) MP
 Prostrate Tick-trefoil (*Desmodium rotundifolium*) MP
 Diapensia (*Diapensia lapponica* ssp. *lapponica*) HP
 Few-flowered Panc-grass (*Dichanthelium oligosanthos* ssp. *scribnerianum*) MP
 Spherical Panic-grass (*Dichanthelium sphaerocarpon*) MP
 Rock Draba (*Draba arabisans*) HP
 Hoary Draba (*Draba cana*) HP
 Smooth Draba (*Draba glabella*) HP
 American Dragonhead (*Dracocephalum parviflorum*) HP
 American Waterwort (*Elatine americana*) HP
 Small Waterwort (*Elatine minima*) MP
 Tidal Spikerush (*Eleocharis aestuum*) HP
 Flat-stem Spikerush (*Eleocharis compressa* var. *compressa*) HP
 Wright's Spikerush (*Eleocharis diandra*) HP
 Olive Spikerush (*Eleocharis flavescens* var. *olivacea*) MP
 Slender Spikerush (*Eleocharis nitida*) HP
 Few-flowered Spikerush (*Eleocharis quinqueflora*) MP
 Robbins Spikerush (*Eleocharis robbinsii*) MP
 MacGregor's Wild Rye (*Elymus macgregorii*) HP
 Southern Wild-rye (*Elymus villosus* var. *arkansanus*) HP
 Hairy Wild-rye (*Elymus villosus* var. *villosus*) MP
 Black Crowberry (*Empetrum nigrum*) HP
 Marsh Willow-herb (*Epilobium palustre*) MP
 Hyssop-leaved Fleabane (*Erigeron hyssopifolius*) HP
 Provancher's Dwarf Fleabane (*Erigeron philadelphicus* var. *provancheri*) HP
 Slender Cotton-grass (*Eriophorum gracile*) MP
 Rough Cotton-grass (*Eriophorum tenellum*) MP
 Sessile-leaved Boneset (*Eupatorium sessilifolium*) MP
 Nodding Spurge (*Euphorbia nutans*) HP
 Rough-leaved Aster (*Eurybia radula*) MP
 Shortleaf Fescue (*Festuca brachyphylla* ssp. *brachyphylla*) HP
 Autumn Fimbristylis (*Fimbristylis autumnalis*) MP
 False Mermaid-weed (*Floerkea proserpinacoides*) HP
 Limestone Swamp Bedstraw (*Galium brevipes*) HP
 Bog Bedstraw (*Galium labradoricum*) HP
 Hairy Bedstraw (*Galium pilosum*) MP
 Fringe-top Closed Gentian (*Gentiana andrewsii*) HP
 Felwort (*Gentianella amarella*) HP
 Stiff Gentian (*Gentianella quinquefolia*) MP
 Spring Avens (*Geum vernum*) HP
 Sharp Manna-grass (*Glyceria acutiflora*) MP
 Eastern Manna-grass (*Glyceria septentrionalis*) MP
 Giant Rattlesnake-plantain (*Goodyera oblongifolia*) HP
 Nodding Stickseed (*Hackelia deflexa* ssp. *americana*) HP
 Spurred Gentian (*Halenia deflexa*) MP
 Alpine Sweet-broom (*Hedysarum alpinum*) MP
 Sneezeweed (*Helenium autumnale*) MP
 Mare's-tail (*Hippuris vulgaris*) MP
 Longleaf Bluet (*Houstonia longifolia*) MP

Beach Heather (*Hudsonia tomentosa*) HP
 Green Violet (*Hybanthus concolor*) HP
 Golden-seal (*Hydrastis canadensis*) HP
 Broad-leaved Waterleaf (*Hydrophyllum canadense*) HP
 Great St. John's-wort (*Hypericum ascyron*) MP
 Orange-grass St. John's-wort (*Hypericum gentianoides*) MP
 Smooth Holly (*Ilex laevigata*) MP
 Large Whorled Pogonia (*Isotria verticillata*) MP
 Tapering Rush (*Juncus acuminatus*) MP
 Alpine Rush (*Juncus alpinoarticulatus*) MP
 Greater Poverty Rush (*Juncus anthelatus*) MP
 Greene's Rush (*Juncus greenii*) MP
 Secund Rush (*Juncus secundus*) MP
 Woodland Rush (*Juncus subcaudatus*) MP
 Torrey's Rush (*Juncus torreyi*) HP
 Highland Rush (*Juncus trifidus*) HP
 Vasey Rush (*Juncus vaseyi*) HP
 Hairy Lettuce (*Lactuca hirsuta*) HP
 Beach Pea (*Lathyrus japonicus* var. *maritimus*) MP
 Pale Vetchling (*Lathyrus ochroleucus*) HP
 Marsh Vetchling (*Lathyrus palustris*) MP
 Lesser Pinweed (*Lechea minor*) HP
 Hairy Pinweed (*Lechea mucronata*) MP
 Minute Duckweed (*Lemna perpusilla*) HP
 Turion Duckweed (*Lemna turionifera*) HP
 Violet Bush-clover (*Lespedeza frutescens*) MP
 Hairy Bush-clover (*Lespedeza hirta* ssp. *hirta*) MP
 Trailing Bush-clover (*Lespedeza procumbens*) MP
 Large White-flowered Ground-cherry (*Leucophysalis grandiflora*) HP
 Stiff Yellow Flax (*Linum medium*) HP
 Grooved Yellowflax (*Linum sulcatum* var. *sulcatum*) HP
 Lily-leaved Twayblade (*Liparis liliifolia*) HP
 Tulip Tree (*Liriodendron tulipifera*) MP
 American Shore-grass (*Littorella americana*) MP
 Great Blue Lobelia (*Lobelia siphilitica* var. *siphilitica*) HP
 Hairy Spike Lobelia (*Lobelia spicata* var. *hirtella*) HP
 Hairy Honeysuckle (*Lonicera hirsuta*) HP
 Swamp Fly-honeysuckle (*Lonicera oblongifolia*) MP
 Many-fruited False-loosestrife (*Ludwigia polycarpa*) HP
 Wild Lupine (*Lupinus perennis*) HP
 Spiked Wood-rush (*Luzula spicata*) HP
 Virginia Bugleweed (*Lycopus virginicus*) MP
 Lance-leaved Loosestrife (*Lysimachia hybrida*) MP
 Winged-loosestrife (*Lythrum alatum* ssp. *alatum*) HP
 White Adder's-mouth (*Malaxis monophyllos* var. *brachypoda*) HP
 Green Adder's-mouth (*Malaxis unifolia*) MP
 Mountain Sandwort (*Minuartia groenlandica*) HP
 Marcescent Sandwort (*Minuartia marcescens*) HP
 Marble Sandwort (*Minuartia rubella*) HP
 Large-leaved Sandwort (*Moehringia macrophylla*) HP
 Dotted Horsemint (*Monarda punctata*) HP
 Red Mulberry (*Morus rubra*) HP
 Schreber's Muhly (*Muhlenbergia schreberi*) MP
 Sprout Muhly (*Muhlenbergia sobolifera*) MP
 Woodland Muhly (*Muhlenbergia sylvatica*) MP

Smaller Forget-me-not (*Myosotis laxa*) MP
 Spring Forget-me-not (*Myosotis verna*) MP
 Low Water-milfoil (*Myriophyllum humile*) MP
 Boott's Rattlesnake-root (*Nabalus boottii*) HP
 Glaucous Rattlesnake-root (*Nabalus recemosus*) MP
 Slender Naiad (*Najas gracillima*) MP
 Guadalupe Naiad (*Najas guadalupensis*) MP
 Auricled Twayblade (*Neottia auriculata*) HP
 Southern Twayblade (*Neottia bifolia*) HP
 Dwarf Water-lily (*Nymphaea leibergii*) HP
 Bog Aster (*Oclemena nemoralis*) MP
 Nodding Evening-primrose (*Oenothera nutans*) HP
 Woodland Cudweed (*Omalothea sylvatica*) MP
 Blunt-fruited Sweet-cicely (*Osmorhiza depauperata*) HP
 Violet Wood-sorrel (*Oxalis violacea*) HP
 Stiff Witch-grass (*Panicum flexile*) HP
 Philadelphia Panic-grass (*Panicum philadelphicum* var. *philadelphicum*) MP
 Smooth Forked Chickweed (*Paronychia canadensis*) HP
 Slender Paspalum (*Paspalum setaceum* var. *muhlenbergii*) MP
 Pale Beardtongue (*Penstemon pallidus*) MP
 Carey's Smartweed (*Persicaria careyi*) MP
 Sweet Coltsfoot (*Petasites frigidus* var. *palmatus*) MP
 American Reed (*Phragmites australis* ssp. *americanus*) HP
 Strawberry-tomato (*Physalis grisea*) MP
 Obedient Plant (*Physostegia virginiana*) MP
 Black-seeded Clearweed (*Pilea fontana*) HP
 Butterwort (*Pinguicula vulgaris*) HP
 Slender Mountain-rice (*Piptatheropsis pungens*) MP
 White-fringed Orchid (*Platanthera blephariglottis* var. *blephariglottis*) MP
 Tubercled Orchid (*Platanthera flava* var. *herbiola*) MP
 Hooker's Orchid (*Platanthera hookeri*) MP
 Large Roundleaf Orchid (*Platanthera macrophylla*) MP
 Roundleaf Orchid (*Platanthera orbiculata*) MP
 Glaucous Bluegrass (*Poa glauca* ssp. *glauca*) HP
 Inland Bluegrass (*Poa interior*) HP
 Wavy Bluegrass (*Poa laxa* ssp. *fernaldiana*) HP
 Lax Bluegrass (*Poa saltuensis* ssp. *languida*) HP
 May-apple (*Podophyllum peltatum*) HP
 Riverweed (*Podostemum ceratophyllum*) MP
 Eastern Jacob's Ladder (*Polemonium vanbruntiae*) HP
 Racemed Milkwort (*Polygala polygama*) MP
 Ambiguous Milkwort (*Polygala verticillata* var. *ambigua*) HP
 Whorled Milkwort (*Polygala verticillata* var. *verticillata*) MP
 Common Solomon's-seal (*Polygonatum biflorum*) MP
 Douglas' Knotweed (*Polygonum douglasii*) MP
 Erect Knotweed (*Polygonum erectum*) HP
 Slender Knotweed (*Polygonum tenue*) MP
 White-flowered Leafcup (*Polymnia canadensis*) HP
 Snail-seed Pondweed (*Potamogeton bicupulatus*) MP
 Tuckerman's Pondweed (*Potamogeton confervoides*) MP
 Vasey's Pondweed (*Potamogeton vaseyi*) MP
 Ogden's Pondweed (*Potamogeton x ogdenii*) HP
 Bird's-eye Primrose (*Primula mistassinica*) HP
 Low Sand Cherry (*Prunus pumila* var. *depressa*) MP

Susquehanna Sand Cherry (*Prunus susquehanae*) MP
 Pinedrops (*Pterospora andromedea*) HP
 Hoary Mountain-mint (*Pycnanthemum incanum*) MP
 Blunt Mountain-mint (*Pycnanthemum muticum*) MP
 Bog Wintergreen (*Pyrola asarifolia* ssp. *asarifolia*) MP
 Lesser Pyrola (*Pyrola minor*) HP
 Scarlet Oak (*Quercus coccinea*) MP
 Scrub Oak (*Quercus ilicifolia*) MP
 Dwarf Chinquapin Oak (*Quercus prinoides*) MP
 Allegheny Crowfoot (*Ranunculus allegheniensis*) MP
 Bristly Buttercup (*Ranunculus hispidus* var. *hispidus*) HP
 Virginia Meadow-beauty (*Rhexia virginica*) MP
 Roseroot (*Rhodiola rosea*) HP
 Great Laurel (*Rhododendron maximum*) HP
 Pinxter-flower (*Rhododendron periclymenoides*) MP
 Capillary Beak-rush (*Rhynchospora capillacea*) HP
 Lake-cress (*Rorippa aquatica*) HP
 Needle-spine Rose (*Rosa acicularis* ssp. *sayi*) HP
 Shining Rose (*Rosa nitida*) MP
 Black-eyed Susan (*Rudbeckia hirta* var. *hirta*) HP
 Western Dock (*Rumex occidentalis*) HP
 Small Pearlwort (*Sagina decumbens*) HP
 Peach-leaf Willow (*Salix amygdaloides*) HP
 Bog Willow (*Salix pedicellaris*) MP
 Satiny Willow (*Salix pellita*) MP
 Tea-leaved Willow (*Salix planifolia*) HP
 Bearberry Willow (*Salix uva-ursi*) HP
 Water Pimpernel (*Samolus parviflorus*) MP
 Canada Burnet (*Sanguisorba canadensis*) MP
 Long-styled Snakeroot (*Sanicula canadensis* var. *grandis*) HP
 Yellow Mountain Saxifrage (*Saxifraga aizoides*) HP
 Purple Mountain Saxifrage (*Saxifraga oppositifolia*
oppositifolia) HP
 White Mountain Saxifrage (*Saxifraga paniculata*) HP
 Pod-grass (*Scheuchzeria palustris*) MP
 Smith's Bulrush (*Schoenoplectiella smithii* var. *smithii*) MP
 Slender Bulrush (*Schoenoplectus heterochaetus*) HP
 Torrey's Bulrush (*Schoenoplectus torreyi*) MP
 Barbed-bristle Bulrush (*Scirpus ancistrochaetus*) HP
 Georgia Bulrush (*Scirpus georgianus*) HP
 Shale Barren Skullcap (*Scutellaria parvula* var. *missouriensis*) HP
 Small Skullcap (*Scutellaria parvula* var. *parvula*) HP
 Wild Senna (*Senna hebecarpa*) HP
 Starry Catchfly (*Silene stellata*) HP
 Eastern Blue-eyed-grass (*Sisyrinchium atlanticum*) MP
 Cutler's Goldenrod (*Solidago leiocarpa*) HP
 Snowy Aster (*Solidago ptarmicoides*) HP
 River-ledge Goldenrod (*Solidago racemosa*) HP
 Elm-leaved Goldenrod (*Solidago ulmifolia*) MP
 Branching Bur-reed (*Sparganium angrocladum*) HP
 Shiny Wedgegrass (*Sphenopholis nitida*) HP
 Blunt Sphenopholis (*Sphenopholis obtusata*) HP
 Case's Ladies'-tresses (*Spiranthes casei* var. *casei*) HP
 Rough Dropseed (*Sporobolus compositus*) MP
 Small Dropseed (*Sporobolus neglectus*) HP
 Marsh Woundwort (*Stachys pilosa* var. *pilosa*) MP
 Trailing Stitchwort (*Stellaria alsine*) MP
 Slender Pondweed (*Stuckenia filiformis*) HP
 Hybrid Thread-leaved Pondweed (*Stuckenia x fennica*) MP
 Ontario Aster (*Symphotrichum ontarionis*) HP
 Crooked-stem Aster (*Symphotrichum prenanthoides*) HP
 Small White Aster (*Symphotrichum racemosum*) MP
 Tradescant Aster (*Symphotrichum tradescantii*) MP
 White-arrow Aster (*Symphotrichum urophyllum*) HP
 Yellow Pimpernel (*Taenidia integerrima*) HP
 Rue-anemone (*Thalictrum thalictroides*) MP
 Border Meadow-rue (*Thalictrum venulosum*) HP
 Sticky False-asphodel (*Triantha glutinosa*) MP
 Deer-hair Sedge (*Trichophorum cespitosum*) MP
 Bashful Bulrush (*Trichophorum planifolium*) MP
 False Pennyroyal (*Trichostema brachiatum*) HP
 Common Arrow-grass (*Triglochin maritima*) HP
 Three-bird Orchid (*Triphora trianthophora*) MP
 Hairy-glumed False Oats (*Trisetum spicatum* var. *pilosiglume*) MP
 Cork Elm (*Ulmus thomasii*) HP
 Inflated Bladderwort (*Utricularia radiata*) MP
 Northeastern Bladderwort (*Utricularia resupinata*) MP
 Perfoliate Bellwort (*Uvularia perfoliata*) MP
 Boreal Blueberry (*Vaccinium boreale*) MP
 Dwarf Bilberry (*Vaccinium caespitosum*) MP
 Deerberry (*Vaccinium stamineum*) MP
 Alpine Bilberry (*Vaccinium uliginosum*) HP
 Mountain Cranberry (*Vaccinium vitis-idaea*) MP
 Marsh Valerian (*Valeriana uliginosa*) HP
 Narrow-leaved Vervain (*Verbena simplex*) HP
 Water-speedwell (*Veronica catenata*) HP
 Culver's-root (*Veronicastrum virginicum*) HP
 Squashberry (*Viburnum edule*) MP
 Lance-leaved Violet (*Viola lanceolata* ssp. *lanceolata*) MP
 Early Blue Violet (*Viola palmata*) MP
 Lobed Violet (*Viola subsinuata*) HP
 Eight-flowered Fescue (*Vulpia octoflora*) HP
 Northern Yellow-eyed-grass (*Xyris montana*) MP
 Horned Pondweed (*Zannichellia palustris*) MP
Hornworts
 A Hornwort (*Anthoceros agrestis*) MP
Liverworts
 A Liverwort (*Anastrophyllum helleranum*) MP
 A Liverwort (*Anastrophyllum michauxii*) MP
 A Liverwort (*Anastrophyllum saxicola*) MP
 A Liverwort (*Aneura maxima*) MP
 A Liverwort (*Athalamia hyalina*) MP
 A Liverwort (*Barbilophozia hatcheri*) MP
 A Liverwort (*Barbilophozia kunzeana*) MP
 A Liverwort (*Calypogeia suecica*) MP
 A Liverwort (*Cephalozia connivens*) MP
 A Liverwort (*Cephaloziella arctica*) MP
 A Liverwort (*Cephaloziella elachista*) MP
 A Liverwort (*Cephaloziella massalongii*) MP
 A Liverwort (*Cephaloziella rubella* var. *elegans*) MP
 A Liverwort (*Cephaloziella rubella* var. *rubella*) MP

A Liverwort (*Cephaloziella stellulifera*) MP
 A Liverwort (*Chandonanthus setiformis*) MP
 A Liverwort (*Chiloscyphus pallescens* var. *fragilis*) MP
 A Liverwort (*Fossombronina foveolata*) MP
 A Liverwort (*Frullania brittoniae*) MP
 A Liverwort (*Frullania inflata*) MP
 A Liverwort (*Frullania oakesiana*) MP
 A Liverwort (*Frullania plana*) MP
 A Liverwort (*Frullania selwyniana*) MP
 A Liverwort (*Gymnocolea inflata*) MP
 A Liverwort (*Gymnomitrium concinnum*) MP
 A Liverwort (*Harpanthus drummondii*) MP
 A Liverwort (*Harpanthus scutatus*) MP
 A Liverwort (*Jubula pennsylvanica*) MP
 A Liverwort (*Jungermannia caespiticia*) MP
 A Liverwort (*Jungermannia evansii*) MP
 A Liverwort (*Jungermannia pumila*) MP
 A Liverwort (*Jungermannia sphaerocarpa*) MP
 A Liverwort (*Kurzia pauciflora*) MP
 A Liverwort (*Lejeunea lamacerina* ssp. *gemminata*) MP
 A Liverwort (*Lophocolea cuspidata* var. *alata*) MP
 A Liverwort (*Lophocolea minor*) MP
 A Liverwort (*Lophozia alpestris*) MP
 A Liverwort (*Lophozia ascendens*) MP
 A Liverwort (*Lophozia badensis* var. *badensis*) MP
 A Liverwort (*Lophozia collaris*) MP
 A Liverwort (*Lophozia excisa*) MP
 A Liverwort (*Lophozia heterocolpos* var. *heterocolpos*) MP
 A Liverwort (*Lophozia laxa*) MP
 A Liverwort (*Lophozia sudetica*) MP
 A Liverwort (*Marchantia alpestris*) MP
 A Liverwort (*Marchantia aquatica*) MP
 A Liverwort (*Marsupella sphacelata*) MP
 A Liverwort (*Metzgeria crassipilis*) MP
 A Liverwort (*Mylia taylorii*) MP
 A Liverwort (*Nardia scalaris* ssp. *scalaris*) MP
 A Liverwort (*Pellia megaspora*) MP
 A Liverwort (*Plagiochila austinii*) MP
 A Liverwort (*Radula obconica*) MP
 A Liverwort (*Riccia huebeneriana* ssp. *sullivantii*) MP
 A Liverwort (*Scapania cuspiduligera* var. *cuspiduligera*) MP
 A Liverwort (*Scapania gymnostomophila*) MP
 A Liverwort (*Scapania irrigua* ssp. *irrigua*) MP
 A Liverwort (*Scapania lingulata* var. *lingulata*) MP
 A Liverwort (*Scapania mucronata* ssp. *mucronata*) MP
 A Liverwort (*Scapania paludicola* var. *paludicola*) MP
 A Liverwort (*Scapania umbrosa*) MP
 A Liverwort (*Tritomaria exsectiformis* ssp. *exsectiformis*) MP
 A Liverwort (*Tritomaria quinqueidentata* var. *quinqueidentata*) MP

Mosses

A Moss (*Amphidium lapponicum*) MP
 A Moss (*Amphidium mougeotii*) MP
 Knothole Moss (*Anacamptodon splachnoides*) MP
 A Moss (*Andreaea rothii*) MP
 A Moss (*Anomobryum filiforme*) MP
 A Moss (*Aphanorhegma serratum*) MP

Arctoa Moss (*Arctoa fulvella*) MP
 A Moss (*Atrichum tenellum*) MP
 A Moss (*Aulacomnium androgynum*) MP
 A Moss (*Barbula indica* var. *indica*) MP
 A Moss (*Brachythecium acutum*) MP
 A Moss (*Brachythecium campestre*) MP
 A Moss (*Brachythecium digastrum*) MP
 A Moss (*Brachythecium erythrorrhizon*) MP
 Falcate Feather Moss (*Brachythecium falcatum*) MP
 A Moss (*Brachythecium turgidum*) MP
 A Moss (*Bryoandersonia illecebra*) MP
 A Moss (*Bryohaplodadium microphyllum*) MP
 A Moss (*Bryum pallens*) MP
 A Moss (*Bryum pallescens*) MP
 A Moss (*Bryum weigeli*) MP
 A Moss (*Bucklandiella microcarpa*) MP
 Elf Cap Moss (*Buxbaumia aphylla*) MP
 Hump-backed Elves (*Buxbaumia minakatae*) MP
 A Moss (*Calliergon obtusifolium*) MP
 A Moss (*Calliergon trifarium*) MP
 A Moss (*Campylium polygamum*) MP
 A Moss (*Campylium radicale*) MP
 A Moss (*Cinclidium stygium*) MP
 A Moss (*Cirriphyllum piliferum*) MP
 A Tree Moss (*Climacium kindbergii*) MP
 A Moss (*Codriophorus aduncoides*) MP
 A Moss (*Codriophorus fascicularis*) MP
 A Moss (*Conardia compacta*) MP
 A Moss (*Cynodontium alpestre*) MP
 A Moss (*Cynodontium strumiferum*) MP
 A Moss (*Cynodontium tenellum*) MP
 A Moss (*Cyrt-hypnum pygmaeum*) MP
 A Moss (*Cyrtomnium hymenophylloides*) MP
 A Moss (*Dichelyma capillaceum*) MP
 A Moss (*Dichelyma falcatum*) MP
 A Moss (*Dichelyma pallescens*) MP
 A Moss (*Dicranella cerviculata*) MP
 A Moss (*Dicranella schreberiana*) MP
 A Moss (*Dicranodontium denudatum*) MP
 A Moss (*Dicranum muehlenbeckii*) MP
 Ontario Dicranum Moss (*Dicranum ontariense*) MP
 A Moss (*Didymodon fallax*) MP
 A Moss (*Didymodon ferrugineus*) MP
 A Moss (*Didymodon rigidulus* var. *rigidulus*) MP
 A Moss (*Didymodon tophaceus*) MP
 A Moss (*Distichium capillaceum*) MP
 A Moss (*Ditrichum flexicaule*) MP
 A Moss (*Ditrichum tortuloides*) MP
 A Moss (*Drepanocladus longifolius*) MP
 A Moss (*Drummondia prorepens*) MP
 A Moss (*Entodon brevisetus*) MP
 An Emerald Dewdrops Moss (*Ephemerum cohaerens*) MP
 An Emerald Dewdrops Moss (*Ephemerum spinulosum*) MP
 A Moss (*Eurhynchium hians*) MP
 Small Pocket Moss (*Fissidens exilis*) MP
 A Moss (*Fissidens subbasilaris*) MP

A Moss (*Fontinalis hypnoides* var. *duriaei*) MP
 A Moss (*Forsstroemia trichomitria*) MP
 A Moss (*Grimmia hartmanii*) MP
 A Moss (*Grimmia longirostris*) MP
 A Moss (*Grimmia muehlenbeckii*) MP
 A Moss (*Grimmia pilifera*) MP
 A Moss (*Grimmia trichophylla*) MP
 A Moss (*Grimmia unicolor*) MP
 A Moss (*Hamatocaulis vernicosus*) MP
 Rock Thread Moss (*Haplohymenium triste*) MP
 Blandow's Helodium Moss (*Helodium blandowii* var. *elodioides*) MP
 A Moss (*Helodium paludosum*) MP
 A Moss (*Heterocladium dimorphum*) MP
 Closter's Brook-hypnum (*Hygrohypnum closteri*) MP
 A Moss (*Hygrohypnum duriusculum*) MP
 A Moss (*Hygrohypnum luridum*) MP
 A Moss (*Hygrohypnum micans*) MP
 A Moss (*Hygrohypnum molle*) MP
 A Moss (*Hygrohypnum montanum*) MP
 A Moss (*Hygrohypnum subeugyrium*) MP
 A Moss (*Hylocomiastrum pyrenaicum*) MP
 A Moss (*Hyophila involuta*) MP
 A Moss (*Hypnum fauriei*) MP
 A Moss (*Hypnum plicatulum*) MP
 A Moss (*Hypnum recurvatum*) MP
 A Moss (*Isopterygiopsis pulchella*) MP
 A Moss (*Leptodictyum humile*) MP
 A Moss (*Leskea gracilescens*) MP
 A Moss (*Leskea obscura*) MP
 A Moss (*Leucodon brachypus* var. *brachypus*) MP
 A Moss (*Limprichtia cossonii*) MP
 A Moss (*Limprichtia revolvens*) MP
 Lindberg's Maple-moss (*Lindbergia brachyptera*) MP
 A Moss (*Loeskeobryum brevirostre*) MP
 Triangular Swan Moss (*Meesia triquetra*) MP
 A Moss (*Microbryum davallianum*) MP
 Micromitrium Moss (*Micromitrium tenerum*) MP
 A Moss (*Mnium thomsonii*) MP
 A Moss (*Myurella julacea*) MP
 A Neckera Moss (*Neckera bessi*) MP
 A Moss (*Neckera complanata*) MP
 A Moss (*Niphotrichum canescens* ssp. *canescens*) MP
 A Moss (*Orthotrichum ohioense*) MP
 A Moss (*Orthotrichum pumilum*) MP
 A Moss (*Paludella squarrosa*) MP
 A Moss (*Palustriella commutata*) MP
 A Moss (*Philonotis marchica*) MP
 A Moss (*Philonotis muehlenbergii*) MP
 A Moss (*Physcomitrium immersum*) MP
 A Moss (*Plagiobryum zieri*) MP
 A Moss (*Plagiomnium drummondii*) MP
 A Moss (*Plagiomnium rostratum*) MP
 A Moss (*Platydictya jungermannioides*) MP
 A Moss (*Platydictya subtilis*) MP
 A Moss (*Pogonatum dentatum*) MP
 Andalusian Pohlia Moss (*Pohlia andalusica*) MP
 A Moss (*Pohlia annotina*) MP
 A Moss (*Pohlia bulbifera*) MP
 A Moss (*Pohlia drummondii*) MP
 A Moss (*Pohlia elongata* var. *elongata*) MP
 A Moss (*Pohlia prolifera*) MP
 A moss (*Pohlia sphagnicola*) MP
 A Moss (*Polytrichastrum formosum*) MP
 A Moss (*Polytrichastrum longisetum*) MP
 A Moss (*Ptychomitrium incurvum*) MP
 Felt Round Moss (*Rhizomnium pseudopunctatum*) MP
 Blue Dew (*Saelania glaucescens*) MP
 A Moss (*Schistidium liliputanum*) MP
 A Moss (*Schistidium papillosum*) MP
 A Moss (*Schistidium viride*) MP
 Luminous Moss (*Schistostega pennata*) MP
 A Moss (*Schwetschkeopsis fabronia*) MP
 A Moss (*Scorpidium scorpioides*) MP
 A Moss (*Sematophyllum adnatum*) MP
 A Moss (*Sematophyllum demissum*) MP
 A Moss (*Sematophyllum marylandicum*) MP
 A Moss (*Sphagnum andersonianum*) MP
 A Moss (*Sphagnum angermanicum*) MP
 A Moss (*Sphagnum atlanticum*) MP
 A Moss (*Sphagnum austinii*) MP
 Bartlett's Peatmoss (*Sphagnum bartlettianum*) MP
 A Moss (*Sphagnum brevifolium*) MP
 Low Peatmoss (*Sphagnum compactum*) MP
 A Moss (*Sphagnum contortum*) MP
 Henry's Peatmoss (*Sphagnum henryense*) MP
 A Moss (*Sphagnum inundatum*) MP
 A Moss (*Sphagnum isoviitae*) MP
 Lindberg's Sphagnum (*Sphagnum lindbergii*) MP
 A Moss (*Sphagnum mcqueenii*) MP
 A Moss (*Sphagnum platyphyllum*) MP
 Beautiful Peatmoss (*Sphagnum pulchrum*) MP
 Five-ranked Bogmoss (*Sphagnum quinquefarium*) MP
 Recurved Peatmoss (*Sphagnum recurvum*) MP
 A Moss (*Sphagnum riparium*) MP
 A Peatmoss (*Sphagnum subfulvum*) MP
 Delicate Peatmoss (*Sphagnum tenellum*) MP
 Giant Peatmoss (*Sphagnum torreyanum*) MP
 A Moss (*Sphagnum viride*) MP
 A Moss (*Syntrichia ruralis*) MP
 A Moss (*Thelia asprella*) MP
 A Moss (*Timmia megapolitana* ssp. *megapolitana*) MP
 A Moss (*Tomenthypnum falcifolium*) MP
 A Moss (*Tortella fragilis*) MP
 A Moss (*Tortella inclinata* var. *densa*) MP
 A Moss (*Tortella inclinata* var. *inclinata*) MP
 A Moss (*Tortula mucronifolia*) MP
 A Moss (*Tortula obtusifolia*) MP
 A Moss (*Trichostomum crispulum*) MP
 A Moss (*Weissia muhlenbergiana*) MP

Conserving Vermont's Amphibians & Reptiles

Reptile & Amphibian Team

Doug Blodgett (team leader) Vermont Fish & Wildlife Department
Steve Parren, Vermont Fish & Wildlife Department
Jim Andrews, Middlebury College
Steve Faccio, Vermont Institute of Natural Science
Chris Slesar, Vermont Agency of Transportation

Team Charge

The Reptile and Amphibian Team was charged with identifying Species of Greatest Conservation Need (SGCN); describing the distribution and habitat usage for each SGCN; evaluating threats impacting SGCN and their habitat; identifying priority research needs to improve our ability to conserve these species; and, developing conservation strategies to address priority threats.

Introduction

For much of the year Vermont's 40 species of amphibians and reptiles, collectively known as herps or herptiles (from the Greek *Herpeton*), are secretive creatures shunning the fuss made over our more charismatic mega-fauna. But stand beside a Vermont wetland, pond or vernal pool on an early spring evening and the cacophony of calls from wood frogs, spring peepers, green frogs, and others and these enigmatic micro-fauna will make themselves noticed.

Vermont's reptiles and amphibians certainly deserve notice. As if their penchant for feasting on black flies, mosquitoes, garden slugs, rodents and other pests isn't reason enough to conserve them (some frogs are reported to eat as many as 3,000 insects a year), many also play critical roles in ecosystems, and serve as excellent indicators of the health of natural systems due to their sensitivity to toxic chemicals and habitat change.

Amphibians and reptiles face many conservation challenges in today's world, be it crossing high-traffic roads or the loss of habitat and connections between habitat patches. It could be argued that all 21 amphibians and 19 reptiles known to be extant in Vermont deserve Species of Greatest Conservation Need (SGCN) designation. The Action Plan Reptile and Amphibian Team took a conservative approach to selecting SGCN in order to highlight those species thought to be most in need of conservation assistance so that scarce resources can be directed toward their conservation.

Implementing the 2005 Wildlife Action Plan

A substantial amount of work, primarily through SWG funded projects, has been accomplished since the 2005 Action Plan to advance our knowledge of specific Vermont herps. Our on-going Rare Snakes of Vermont project, established in 2010 with SWG funding, focuses specifically on rare snake SGCN. It's significantly added to our knowledgebase of rattlesnake and ratsnake home ranges, their movements, mapping of critical SGCN snake habitats, population demographics, genetic assessment and other life history information. This investigation also revealed the presence of the lethal Snake Fungal Disease (SFD) in Vermont's rattlesnake population. A newly emerging skin fungus afflicting myriad snake species, SFD, is now an issue of serious regional concern and alarm in the eastern U.S. and has recently been detected in a dozen eastern states. The Rare Snakes project

produced significant new findings and completion of a report on Vermont’s two-year rattlesnake research study (Spear et. al.). Two Vermont Recovery Plans were also completed since 2005; one for the threatened spiny softshell turtle and the second for the endangered timber rattlesnake.

A wealth of reliable documentation has been added to The Vermont Reptile & Amphibian Atlas throughout the state resulting from hundreds of additional logged sightings/locations of reptiles and amphibians. A survey of vernal pools (also SWG-funded) has catalogued some 2500 vernal pool amphibian breeding sites statewide. Turtle species have also received research attention, including documenting/protecting nesting sites for the endangered spotted turtle, the threatened spiny softshell, and wood turtles – a species of special concern. Additional genetic research/assessment has been directed to the Mudpuppy revealing two distinct populations in Vermont (one in the Lake Champlain basin and the second, in the Connecticut River basin). In 2015, the Fowler’s Toad was listed as endangered in Vermont.

Northeast Partners in Reptile and Amphibian Conservation ([NEPARC](#)) and its national affiliate [PARC](#) have been particularly active recently in attempting to stem the tide of the ubiquitous international amphibian import trade to help protect US amphibians from imported, exotic diseases such as the newly identified salamander fungus *Batrachochytrium salamandrivorans* (Martel, A. et al.).

Selecting Amphibian & Reptile SGCN

The Herp Taxa Team deliberated extensively on SGCN selection criteria, and ultimately made no changes to the 2005 SGCN list of herp species, nor its priority rankings. No species were added nor deleted. Scientific nomenclature for several species was updated. Selection criteria included knowledge about current listing as endangered and threatened, population declines, rarity, vulnerability of habitat, life history traits, vulnerability to collection or take, other impacts from humans, and dispersal capability. Each species was examined across all criteria and the five-person team utilized a high, medium, and low conservation need ranking to attempt to separate species with greater need from those that may be more secure, at least in the short term. We assigned numerical rankings that assisted our assignment to high, medium, and low priority categories. This approach resulted in the selection of the same 12 species of high conservation need and seven of medium conservation need (table 1)

Table 1. Amphibian & Reptile Species of Greatest Conservation Need

High Priority

Jefferson Salamander (*Ambystoma jeffersonianum*- and hybrids)¹
Mudpuppy (*Necturus maculosus*)¹
Fowler’s Toad (*Bufo fowleri*)¹
Boreal Chorus Frog (*Pseudacris maculata*)
Spotted Turtle (*Clemmys guttata*)¹
Wood Turtle (*Glyptemys insculpta*)¹
Spiny Softshell Turtle (*Apalone spinifer*)¹
Five-lined Skink (*Plestiodon fasciatus*)
North American Racer (*Coluber constrictor*)¹
Eastern Ratsnake (*Pantherophis alleghaniensis*)
Timber Rattlesnake (*Crotalus horridus*)¹

Medium Priority

Blue-spotted Salamander (*Ambystoma laterale*)¹
Spotted Salamander (*Ambystoma maculatum*)
Four-toed Salamander (*Hemidactylium scutatum*)
Common Musk Turtle (*Sternotherus odoratus*)
Northern Water Snake (*Nerodia sipedon*)
DeKay’s Brownsnake (*Storeria dekayi*)¹
Eastern Ribbonsnake (*Thamnophis sauritus*)¹
Smooth Greensnake (*Liochlorophis vernalis*)¹

¹ Regional Species of Greatest Conservation Need in the Northeastern United States (Terwilliger, 2013)

Though some of Vermont’s SGCN are at the periphery of their range (e.g., Boreal Chorus Frog, Mudpuppy, Fowlers Toad, and Mink Frog, a finding that challenges conventional wisdom is that

species populations have been documented to be more at risk of loss at the core of their range than at the periphery (Channel & Lomolino 2000, Lomolino 1995). This argues for us giving serious consideration to SGCN that may be peripheral in Vermont.

Full reports on each Species of Greatest Conservation Need are in Appendix A of this document. The following is a summary of those reports.

Habitat Needs

Since many reptiles and amphibians use a variety of habitats annually and over the course of their lives, maintaining healthy populations entails maintaining connectivity between habitats. Connectivity also enables individuals to find alternative cover, food sources, breeding, or over-wintering sites when natural disasters occur. Furthermore, connectivity between populations ensures vital genetic exchange and allows for the re-colonization of areas where populations have been eliminated due to drought, winterkill, disease, or anthropogenic forces. This can only occur if the landscape is permeable to these animals—that is, development proceeds in a way that allows amphibians and reptiles to move freely across the landscape. To conserve our native amphibians and reptiles, especially those considered SGCN, it will be essential to maintain a network of interconnected sites where natural processes are allowed to occur.

Discussion of Threats to Vermont's Amphibians & Reptiles

The threats identified most frequently for Vermont's reptile and amphibian populations are all closely related to habitat degradation: trampling and direct impacts, road and transportation system impacts, habitat fragmentation, habitat alteration, and habitat conversion.

We do not understand all the ramifications, but the pattern seen elsewhere in the US and the world is that increased human population density, higher consumption of land and other resources, and lack of awareness of the impacts to other species can lead to devastating losses of native biota (TWS 2004). Vermont is not immune from these sorts of impacts and our landscape is continuing to be developed (DeVillars 1999). Habitat alteration and loss is a near universal challenge to native amphibians and reptiles.

In an attempt to address this threat, Vermont's Agency of Natural Resources remains vigilant in its efforts to minimize loss of critical habitats under its jurisdiction thru formal regulatory authority and mechanisms provided under Vermont's Act 250 development law, Act 248 and wetlands protection permit processes. The Vermont Fish and Wildlife Dept. has developed some powerful new mapping (Natural Resources Atlas) and habitat/natural community analysis tools (Biofinder) to assist in identifying the states' most diverse, valuable and vulnerable lands as targets for eventual conservation.

A significant, newly identified threat is a lethal Snake Fungal Disease (SFD) which causes significant and debilitating skin lesions on snakes. Widespread detection of SFD in the eastern US has prompted an extensive, on-going, regional research investigation into this novel fungus to assess the causes and conservation significance of this extremely serious threat to free-ranging snakes. Concern has also recently arisen over a novel, salamander fungus *Batrachochytrium salamandrivorans* (Martel, A. et al.) identified on the Asian and European continents, and the threat of spread to North America via a ubiquitous international pet trade. The national conservation organization Partners in

Amphibian and Reptile Conservation (PARC) is now attempting to promulgate US/international herptile trade regulations to protect against this threat.

Crossing roads is a real problem for both amphibians and reptiles in Vermont. Vernal migrations of salamanders and frogs to breeding pools result in many dead and wounded animals when a busy road must be crossed. At some sites in Vermont, thousands of amphibians are killed during a single night, which may overwhelm the reproductive capacity to sustain the populations and, according to the Vermont Agency of Transportation, constitutes a public safety issue (C. Slesar, VTrans, pers. comm). Female turtles seeking nest sites are more at risk of being killed on roads than more sedentary males, resulting in a sex bias in some populations and raises questions about population persistence (Sheen & Gibbs. 2004, Marchand & Litvaitis 2004). The still abundant, but believed to be declining, wood turtle often encounters roads in Vermont during its annual movements along riparian corridors. Snakes emerging from hibernation often bask on warm pavement, increasing their risk of being struck by vehicles.

An emerging awareness of herps' need for improved connectivity and safer wildlife crossings is taking hold in Vermont and the construction of several highway culverts and underpasses at strategic wildlife crossings are being planned and/or contemplated. Some of these structures are relatively expensive and require a good deal of up-front planning, but collaborative efforts are increasingly embraced and accepted in the transportation and wildlife management communities. VTrans, which recognizes that better crossings not only protect wildlife but also protect motorists, is a leader among state transportation agencies in transportation ecology.

Other factors that may negatively impact amphibians and reptiles now and in the foreseeable future include pollution, changes in hydrology, sedimentation, and global climate change.

Research & Monitoring Needs and Conservation Strategies

The Reptile and Amphibian Team identified priority research and monitoring projects to improve our ability to conserve Vermont's reptile and amphibian Species of Greatest Conservation Need. The Team also developed conservation strategies to address threats impacting each SGCN. Those recommended most frequently include:

Research & Monitoring Needs

1. Better determine habitat needs, identify significant breeding sites, vernal pools and habitat connections.
2. Better determine the distribution and relative abundance of populations in Vermont.
3. Better identify and evaluate threats.
4. Monitor trends in population size, distribution and habitat.

Conservation Strategies

5. Help people better value reptiles and amphibians and to understand the essential needs of all life stages, especially upland habitat in proximity to breeding pools.
6. Encourage reports of road-killed specimens, road crossings, and road basking areas to VFWD, VTrans, and the Vermont Reptile and Amphibian Atlas Project. Develop safer crossings at significant sites when roads are being upgraded.

7. Maintain habitat through appropriate management, direct habitat disturbance and site roadways away from sensitive sites such as breeding pools.
8. Continue to work cooperatively with landowners, habitat management agencies, towns and communities to protect habitat and maintain connectivity. Develop management guidelines for owners and managers of appropriate habitat.
9. Conserve known critical habitat through fee simple purchase, acquisition of development rights or easements, management agreements and education of private landowners and managers.
10. If loss of important sites is likely due to development, consider creating or enhancing other pools that might allow some adults to transfer to the new site if they encounter it or develop a new breeding population from dispersal of colonizers.
11. Protect turtle nests and adults by predator trapping.
12. Work with biologists to minimize impacts to SGCN populations and habitats during and following management activities for sport fish and game wildlife.
13. Participate in existing regulatory processes (e.g., Act 250, stream alteration permits) to protect and restore critical habitats.
14. Maintain continued vigilance, monitoring and research efforts as regards novel and lethal emerging herp disease issues including snake fungal disease, *Batrachochytrium salamandrivorans*, chytrid fungus, etc.
15. Begin implementation of vernal pool management guidelines as described by VFWD staff in “Conserving Pool-Specialist Amphibian Habitat”. (Mark Ferguson, VFWD White Paper, 2015)

Conclusion

Vermont's reptiles and amphibians are fortunate for a number of reasons. We have a much less developed and rural landscape than many states. For example, even the eastern newt, a very abundant species in Vermont, is declining in Rhode Island due to development and roads (C.J. Raithel, RI Dept of Environmental Management pers comm). We have an engaged Scientific Advisory Group on Reptiles and Amphibians that provides advice to the Vermont Endangered Species Committee. We also have a well-developed Reptile and Amphibian Atlas Project (<http://cat.middlebury.edu/herpatlas/>) that, mostly through volunteer efforts, has collected, and continues to collect valuable information on the distribution of reptile and amphibian species in Vermont and raises awareness of conservation need in Vermont. Interest is increasing with schools and groups in Vermont that host ‘salamander nights’, helping small amphibians cross roads safely and raise awareness about the impacts of traffic. The Vermont Fish and Wildlife Department continues to work on conservation projects that raise awareness and benefit reptiles and amphibians, including species listed here as SGCN, and we are working collaboratively with other agencies including the Vermont Transportation Agency, as well as an expanding number of conservation partners.. More needs to be done, but with the foundation we already have in place and the awareness and strengthening of partnerships promoted by Action Plan, we expect more conservation actions in our shared future.

See Appendix A for full reports on each reptile and amphibian Species of Greatest Conservation Need.

Literature Cited

- Channel, R. and M.V. Lomolino. 2000. Dynamic biogeography and conservation of endangered species. *Nature* 403:84-86.
- (DEC) Vermont Department of Environmental Conservation. 6-2004. Biomonitoring and Aquatic Studies. <http://www.anr.state.vt.us/dec/waterq/bass.htm>
- DeVillars, J.P. 1999. An Open Letter to the People of Vermont. Vermont Keeping In Touch. U.S. Environmental Protection Agency Vermont State Unit. September 1999.
- Galbraith, D.A., R.J. Brooks, and G.P. Brown. 1997. Can management intervention achieve sustainable exploitation of turtles? Proceedings: Conservation, Restoration, and Management of Tortoises and Turtles-An International Conference, pp. 186-194).
- Garber, S.D. and J. Burger. 1995. A 20-yr study documenting the relationship between turtle decline and human recreation. *Ecological Applications*. 5(4): 1151-1162).
- Lomolino, M.V. and R. Channel. 1995. Splendid isolation: patterns of geographic range collapse in endangered mammals. *Journal of Mammalogy* 76(2): 335-347.)
- Marchand, M.N. and J.A. Litvaitis. 2004. Effects of habitat features and landscape composition on the population structure of a common aquatic turtle in a region undergoing rapid development. *Conservation Biology* 18(3): 758-767).
- Martel, A. et al. (2014) Recent introduction of a chytrid fungus endangers Western Palearctic salamanders. *Science* 346, 630; DOI: 10.1126/science.1258268
- Sheen, D.A. and J.P. Gibbs. 2004. Effects of roads on the structure of freshwater turtle populations. *Conservation Biology* 18(4):1143-1149.
- Spear, S., J. Bauder, D. Blodgett, C. Jenkins, K. Briggs. 2013. The Ecology of Timber Rattlesnakes (*Crotalus horridus*) in Vermont. Vermont Fish and Wildlife Department, Rutland, Vermont, USA.
- Terwilliger Consulting, Inc. and the Northeast Fish and Wildlife Diversity Technical Committee. 2013. [Taking Action Together: Northeast Regional Synthesis for State Wildlife Action Plans](#). A report submitted to the Northeast Fish and Wildlife Diversity Committee. Locustville, VA.
- The Wildlife Society (TWS). 2004. Position on Economic Growth. Adopted by the TWS Council September 2004.

Conserving Vermont's Birds

Birds Team

John Buck, Wildlife Biologist, Vermont Fish & Wildlife Department. (team leader)

Margaret Fowle, Conservation Biologist, Audubon Vermont

John Gobeille, Wildlife Biologist, Vermont Fish & Wildlife Dept.

Mark Labarr, Conservation Biologist, Audubon Vermont

Dr. Allan Strong, University of Vermont

Dr. Rosalind Renfrew, Vermont Center for Ecostudies

Erin Talmadge, Director, Birds of Vermont Museum

Chip Darmstadt, Director, North Branch Nature Center

Dr. William Barnard, Norwich University

Sally Laughlin, Editor, First Vermont Bird Atlas

David Sausville, Wildlife Biologist, Vermont Fish and Wildlife Department

Team Charge

The Bird was charged with identifying Species of Greatest Conservation Need (SGCN); describing the distribution and habitat usage for each SGCN; evaluating threats impacting SGCN and their habitat; identifying priority research needs to improve our ability to conserve these species; and, developing conservation strategies to address priority threats.

Introduction

Vermont serves as host to 268 bird species for some, if not all, of their annual life cycle. These species occupy a wide variety of habitats ranging from Lake Champlain wetlands at about 100 feet above sea level to montane spruce-fir forests at elevations greater than 4000 feet above sea level. Included among these highs and lows are nine distinct biophysical regions all compressed into the 9600 square miles that constitute the small state of Vermont. Naturally following this diverse range of habitats are the diverse guilds of bird species that occupy them. Open water and marsh birds, grassland and shrub birds, deciduous, coniferous, boreal, and montane forest birds are all found within the small state of Vermont. Of the 268 species, 12 of them are listed as endangered or threatened. Another 39 are considered Species of Greatest Conservation Need.


The Action Plan Bird Team convened in October of 2014 to assess the status of SGCN species identified in the 2005 Action Plan species and determine the appropriate category ranking for each of them in the 2015 plan. Each team member reviewed a suite of species and updated the existing narratives with current information about population trends, management actions, and recent research. The result of this effort was identifying 22 changes in prioritization. No new species were added to the priority list except Sandhill Crane as an emerging issue species. Four species were revised from High to Medium, 11 were revised from Medium to High, and seven species were revised from Medium to Low. The remaining 36 species retained their priority status from the 2005 assessment. This assessment resulted in modifying the Species of Greatest Conservation Need list from 58 to 51 species.

The issue of lead in the environment continues to present itself most visibly in the Common Loon. This species continues to experience lead-based morbidity and mortality from residual lead fishing tackle. The loss of shrubland and grassland habitat to natural plant succession and human

development pose a threat to the habitat specialists of these two habitats. These species include, Eastern Meadowlark, Grasshopper Sparrow, Bobolink, Golden-winged Warbler, Blue-winged Warbler, and Eastern Towhee.

Perhaps the single most significant emerging issue impacting birds in Vermont during the last 10 years has been the conversion of forest and grassland habitat to utility-scale wind and solar energy generation. Although descriptors such as ‘renewable’, ‘sustainable’, and ‘environmentally friendly’ create an image of energy development that is preferable to fossil fuel, however, wind and solar energy development still involves habitat loss and impairment. In some cases the habitat involved is rare and sensitive (i.e. montane forest) and supports species of the same status (e.g. Bicknell’s Thrush). Furthermore, collision deaths of resident and migratory birds directly related to wind turbines presents a population loss factor that is difficult to assess due to the relatively small amount of comparative mortality data and the cumulative mortality impact as birds migrate north and south along the Atlantic Flyway. The seemingly benign impact of alternative energy development to the public will create a challenge to VFWD biologists when developing equitable habitat mitigation. A sound depiction of the problem and reasonable solutions will have to be narrated by the Department to achieve bird conservation success.

We are particularly pleased to note that since 2005 Peregrine Falcon, Common Loon, and Osprey were all de-listed. Thanks to 30 years of effort by the VFWD and many partners, populations of these species have recovered sufficiently to where they no longer require heightened protection as Threatened or Endangered species (2005). Looking forward, possibilities for additional delistings in coming years are positive as Common Tern appears to be reaching its down-listing goal and nesting results for Bald Eagle suggest a steady increase in its population since the first Action Plan. Additional population data has provided the background for revising the status of Coopers Hawk, and Veery from ‘Medium’ to ‘Low’.



Implementing the 2005 Wildlife Action Plan

Conservation efforts expanded beyond the restoration of Osprey, Peregrine Falcon, and Common Loon during the 10-years since the first Wildlife Action Plan include more focused attention on Common Tern, Black Tern, Grasshopper Sparrow, Spruce Grouse, and Bald Eagle. Each of the latter species benefit from intensive survey efforts to better determine their respective population status and corresponding listing status. Closer working relations with state airport managers was cultivated during the last action plan period resulting in greater awareness of bird species needs and their uses of grassland habitat at airports. This led to a more diversified response to aircraft safety concerns. One profound outcome during this period is the completion and publication of Vermont's Second Breeding Bird Atlas. Very much a collaborative effort, the Atlas was developed with more than 350 citizen scientists contributing more than 50,000 hours over five years to document every bird species breeding across the state. The effort was led by the Vermont Center for Ecostudies, the Fish & Wildlife Department and others, with funding, in part, from the SWG program.

Selecting Bird SGCN

In contrast to lesser-known taxa, the Bird Team benefited from the relative wealth of available data on bird distribution and abundance. Data from Vermont's original (1982) and current (2013) Breeding Bird Atlases and the USFWS Breeding Bird Surveys helped immensely in selecting the 51 SGCN. In addition to this hard data source, ongoing bird conservation programs, including the Vermont Endangered Species Scientific Advisory Group on Birds (SAG-Birds), Partners-In-Flight, North American Bird Conservation Initiative, National Audubon Society's Watch List, and the American Bird Conservancy's Green List all contributed to our understanding of which species belonged on Vermont's SGCN list.

Selection criteria included current listing as endangered or threatened, population declines, rarity, vulnerability of habitat, life history traits, impacts from humans, and recent range expansion or contraction. Each species was examined across all criteria and the team developed a high, medium, and low conservation need ranking to attempt to separate species with greater need from those that may be more secure, at least in the short term (table 1).

Of the 58 species identified by the bird team in 2005, 37 (64%) retained their priority ranking from that list. Four were moved from 'High' to 'Medium'. These were Peregrine Falcon, Common Loon, Purple Martin, and American Bittern. Moving from 'Medium' to 'Low', were Long-eared Owl, Henslow's Sparrow, Osprey, Cooper's Hawk, Barn Owl, Veery, and Blue-winged Teal. These seven were moved for different reasons that include better population information (Veery, and Cooper's Hawk), unlikely conservation opportunities (Long-eared Owl, Blue-winged Teal, and Henslow's Sparrow) and significant increases in state-wide population (Osprey). A total of 11 species were moved from 'Medium' to 'High'. By and large these changes were made based on a combination of long-term population data (i.e. national Breeding Bird Survey) and the recently completed second Breeding Bird Atlas. Of the 11 species newly identified as high priority, only Rusty Blackbird was listed (Endangered) during the last 10 years. The other 10 species, while not meeting listing criteria, were raised to high priority based on the combination of habitat loss and long-term population decline and the likelihood that these two factors will continue during the next 10 years. The 10 High Priority species can be grouped by loss of their habitats. Wood Thrush (Deciduous Forest), American Kestrel, Red-shouldered Hawk, Eastern Meadowlark (Grassland), Brown Thrasher, Blue-

winged Warbler (Shrubland), Olive-sided Flycatcher, Gray Jay, and Black-poll Warbler (Boreal Forest).

Table 1. Bird Species of Greatest Conservation Need

High Priority

Pied-billed Grebe (*Podilymbus podiceps*)¹
 Least Bittern (*Ixobrychus exilis*)¹
 American Black Duck (*Anas rubripes*)¹
 Bald Eagle (*Haliaeetus leucocephalus*)
 Northern Harrier (*Circus cyaneus*)¹
 Red-shouldered Hawk (*Buteo lineatus*)¹
 American Kestrel (*Falco sparverius*)¹
 Spruce Grouse (*Falcapennis canadensis*)¹
 Upland Sandpiper (*Bartramia longicauda*)¹
 Common Tern (*Sterna hirundo*)¹
 Black Tern (*Chlidonias niger*)¹
 Common Nighthawk (*Chordeiles minor*)¹
 Whip-poor-will (*Caprimulgus vociferus*)¹
 Chimney Swift (*Chaetura pelagica*)¹
 Olive-sided Flycatcher (*Contopus cooperi*)¹
 Gray Jay (*Perisoreus canadensis*)
 Sedge Wren (*Cistothorus platensis*)¹
 Bicknell's Thrush (*Catharus bicknelli*)¹
 Wood Thrush (*Hylocichla mustelina*)¹
 Brown Thrasher (*Toxostoma rufum*)¹
 Blue-winged Warbler (*Vermivora pinus*)¹
 Golden-winged Warbler (*Vermivora chrysoptera*)¹
 Blackpoll Warbler (*Dendroica striata*)
 Canada Warbler (*Wilsonia canadensis*)¹
 Eastern Towhee (*Pipilo erythrophthalmus*)¹
 Vesper Sparrow (*Pooecetes gramineus*)¹
 Grasshopper Sparrow (*Ammodramus savannarum*)¹
 Eastern Meadowlark (*Sturnella magna*)¹
 Rusty Blackbird (*Euphagus carolinus*)¹

Medium Priority

Common Loon (*Gavia immer*)¹
 American Bittern (*Botaurus lentiginosus*)¹
 Great Blue Heron (*Ardea herodias*)
 Black-crowned Night-heron (*Nycticorax nycticorax*)¹
 Northern Goshawk (*Accipiter gentilis*)¹
 Peregrine Falcon (*Falco peregrinus*)¹
 Ruffed Grouse (*Bonasa umbellus*)¹
 Sora (*Porzana carolina*)¹
 Lesser Yellowlegs (*Tringa flavipes*)
 American Woodcock (*Scolopax minor*)¹
 Black-billed Cuckoo (*Coccyzus erythrophthalmus*)¹
 Barn Owl (*Tyto alba*)¹
 Short-eared Owl (*Asio flammeus*)¹
 Black-backed Woodpecker (*Picoides arcticus*)
 Purple Martin (*Progne subis*)
 Chestnut-sided Warbler (*Dendroica pensylvanica*)
 Black-throated Blue Warbler (*Dendroica caerulescens*)¹
 Prairie Warbler (*Dendroica discolor*)¹
 Bay-breasted Warbler (*Dendroica castanea*)¹
 Cerulean Warbler (*Dendroica cerulea*)¹
 Field Sparrow (*Spizella pusilla*)¹
 Bobolink (*Dolichonyx oryzivorus*)¹

¹ Regional Species of Greatest Conservation Need in the Northeastern United States (Terwilliger, 2013)

Bird species rare in Vermont did not always make the SGCN listing. Species that have expanded their range in recent decades due to a proliferation of winter bird feeders, such as Tufted Titmouse, were excluded, as we did not consider Vermont to be a geographic area of responsibility for that species. Other species for which Vermont is on the extreme periphery of their breeding range, and for which confirmed breeding records are very infrequent, such as the Three-toed Woodpecker, were also not selected.

Full reports on each Species of Greatest Conservation Need are in Appendix A of this document. The following is a summary of those reports.

Birds and Their Habitat Needs

Vermont's bird SGCN utilize a variety of habitats from open and shrub-dominated wetlands, mature hardwood or coniferous forests, young regenerating forests, old fields, grasslands, and other cultural habitats such as buildings and structures. As birds are generally more mobile relative to most species from other taxa, they are usually better able to exploit smaller, more widely distributed habitat patches. However, most species benefit from the larger assemblages of similar habitat types, such as a contiguous forest area or large, agricultural (grassland) complex.

The Bird Team organized most birds into one of several habitat guilds, for which a particular conservation strategy would often be appropriate for all species in the guild. These guilds match the major habitat categories used in this report:

- Northern hardwood forest & Oak-pine-northern hardwood forest
- Spruce-fir northern hardwood forest
- Sub-alpine krummholz & Montane spruce-fir forests (high elevation areas)
- Early successional forest stages
- Riparian
- Lacustrine (lakes and ponds)
- Wetlands-(open, shrub and forested wetlands)
- Cliff & Talus
- Grassland
- Grassland/Edge
- Urban

Discussion of Threats Impacting Bird SGCN

New strains of avian influenza have been identified around the world, including western and central North America. To date these new viruses have had the greatest impact on commercial poultry farms and to some degree on the human population in Southeast Asia. Effects on wild bird populations have not been significant to date. However, the potential for viruses to adapt and thrive in a concentrated wintering ground could produce devastating effects. The impact of neonicotinoid pesticides on the continent's insect populations and ultimately on avian insectivores, remains poorly understood. More research is needed to fully understand any far reaching consequences of this new breed of pesticide. It took more than thirty years and untold amount of money and effort to recover populations of Eagle, Osprey, and other raptors after the 1972 ban of the pesticide DDT. It is hoped we will learn from this history lesson and not put bird species in a similar situation in the future. The impact of wind and solar electric generation facilities is still in its infancy. While the detrimental impact of habitat loss from the facilities is readily apparent, the cumulative impacts to populations created by local and cumulative collision mortality are not well understood yet. The rate of growth of these two industries far outpaces the rate of knowledge growth about their relative impacts to continental bird populations. Finally, the habitats estimated to be at risk to climate change are too wrought with dynamic system drivers to permit resource managers to take a direct and meaningful response to the threats. However, it is generally agreed that large blocks of contiguous habitat are in the best are positions to resist or more gradually shift due to climate change. A discussion of any of these emerging issues cannot be complete in the absence of habitat loss.

The loss of habitat is the primary source of bird population declines. As the land area available for birds to exist continues to shrink, ever declining bird populations are increasingly concentrated into the remaining, sometimes marginal quality space, and places species at greater risk of nest failure, predation, and disease. The threats to Vermont's bird populations most frequently identified for are all related to changes in habitat. In the case of the 51 SGCN, conversion of habitat was identified 82% of the time (42 SGCN). Habitat alteration (67%), habitat fragmentation (47%), and distribution of successional stages (35%) were all identified in descending order of frequency. This is the same pattern that emerged in the first Wildlife Action Plan. Many bird species find optimum habitat in young regenerating forests, which have declined statewide in recent decades. Similarly, grassland-dependent species, which are declining throughout the northeast, are finding less and less suitable habitat in Vermont as farms are managed more intensively or sold and either developed or revert to forestland. The increased roads, housing units, utility corridors and other attendant disturbances associated with human development further fragment habitat into smaller and smaller habitat units.

In these smaller units nesting birds are more vulnerable to habitat generalist predators such as raccoons, skunks, and crows. Greater access is also available to the nest parasitizing Brown-headed Cowbird. Finally, the increasing numbers of houses in bird habitat often come with house pets, namely dogs and cats. These familiar companion animals, not in need of shelter or food, add an additional layer of disturbance and predation to nesting birds. Pet cats in particular have been attributed to over 250 million bird deaths nationally each year. See appendix A for full reports on each SGCN.

Research & Monitoring Needs and Conservation Strategies

The research and monitoring needs and conservation strategies most frequently identified by the Bird Team and those best applied for multiple bird SGCN are as follows:

Research & Monitoring Needs

1. Better determine habitat requirements and habitat availability.
2. Better determine the distribution and relative abundance of populations in Vermont.
3. Better identify and evaluate threats including the impacts of wind and solar energy infrastructure and emerging diseases.
4. Obtain better knowledge of basic life history traits.

Conservation Strategies

1. Habitat Restoration efforts on public lands and conservation payments or other financial incentives, fee simple purchase, easements, management guidelines, and cooperative agreements with user groups and private landowners. Existing technical assistance/cost-share programs (EQIP, CRP) were frequently identified as potential funding sources to implement conservation on private lands. Important Bird Area designations can aid in the development of needed funds. Common habitat restoration themes include incentives and planning to slow the rate of fragmentation and development and maintain blocks of contiguous forest, grasslands, early and late-successional habitats.
2. Species Restoration projects, which may involve active translocation of individuals or eggs from a source population into suitable Vermont habitats, and/or may involve efforts to

provide suitable nesting sites and reduce predation or human disturbances around nesting sites.

3. Raising awareness within the general public to build support and opportunities for conservation techniques. Important Bird Area designations can help focus public attention on opportunity areas.
4. Developing and evaluating forestry practices that can enhance habitat suitability such as maintaining or increasing aspen stands or the retention of coarse woody debris and snags. Provide technical assistance to landowners and communities about best management practices (e.g., Vermont Fish and Wildlife Department's Forest Landowner Assistance Program and Audubon's [Forest Bird Initiative](#)).
5. Support and participate in an international effort to maintain large blocks of undeveloped forests linked together by habitat corridors in order to provide a network of interconnected habitats throughout northeastern New England and southeastern Canada.
6. Identify, prioritize and conserve existing contiguous forest blocks and associated linkages that allow for upward and northward movement in response to climate change.
7. Participate in existing regulatory processes (e.g., Act 250) to protect and restore critical habitats.

Conclusion

We are fortunate in that Vermont bird species may be in a better place than those in other parts of their distribution. Still, 51 of the 268 (19%) species remain on the list of Species of Greatest Conservation Need and 12 of those are listed as threatened or endangered. The integration of the state's rural landscape and an economy that is largely fueled by agricultural and forest commodities and tourism along with a human population that values their natural resources to the point they are willing to commit public tax dollars to their state natural resource agency and to land-use legislation. We also have recent conservation successes in the form of peregrine, loon, and osprey de-listing to emulate. However, these are but 3 species out of 51 (5.8%) listed as SGCN and their restoration required funding and human resource levels that would be unimaginable to apply to the remaining species of greatest conservation need. Good collaborative work with our partners (Audubon Vermont, Vermont Center for Ecostudies, University of Vermont, and U.S. Fish and Wildlife Service) is underway toward the restoration of Common Tern and Bald Eagle as well as important monitoring efforts for Black Tern, Grasshopper Sparrow, and Least Bittern. Future conservation opportunities exist with the expansion of grassland habitat conservation through cooperative landowner agreements and land development regulation. Continued work with the state airport managers will also add to the security of this limited habitat. Broader scrutiny of mountain top developments is required if the rare montane forest habitat is to be conserved as significant development pressure is placed on these fragile locations. For Bicknell's Thrush and Blackpoll Warblers to coincide with human use of their habitat a better understanding of the limits of human disturbance is needed. Continued work with private forest landowners and state and local governments is needed to retain the habitat values associated with stable, long-term ownership of contiguous forestland. The Vermont Department of Forests, Parks, and Recreation, Audubon Vermont, the Vermont Woodlands Association and Vermont Coverts will be valuable partners in this effort. Finally, more intensive and purposeful inventory of the secretive marsh bird guild is necessary to accurately assess the status of these hard-to-sample species.

See Appendix A for full reports on each bird Species of Greatest Conservation Need.

Literature Cited

Sauer, J. R., J. E. Hines, J. E. Fallon, K. L. Pardieck, D. J. Ziolkowski, Jr., and W. A. Link. 2014. The North American Breeding Bird Survey, Results and Analysis 1966 - 2012. Version 02.19.2014 USGS Patuxent Wildlife Research Center, Laurel, MD.

The Atlas of Breeding Birds of Vermont. 1985. Laughlin and Kibbe, eds. U. Press of New England. Hanover, NH.

The Second Atlas of Breeding Birds of Vermont, 2013. R. Renfrew, ed. University Press of New England, Hanover and London.

Terwilliger Consulting, Inc. and the Northeast Fish and Wildlife Diversity Technical Committee. 2013. Taking Action Together: Northeast Regional Synthesis for State Wildlife Action Plans. A report submitted to the Northeast Fish and Wildlife Diversity Committee. Locustville, VA.

Conserving Vermont's Fishes

Fish Team

Kenneth Cox, Vermont Fish & Wildlife Department (team leader)
Dr. William Barnard, Norwich University
Dr. Douglas Facey, Saint Michael's College
Mark Ferguson, Vermont Fish & Wildlife Department
Eric Howe, Lake Champlain Basin Program
Richard Langdon, Vermont Department of Environmental Conservation

Team Charge

The Fish Team was charged with identifying Species of Greatest Conservation Need (SGCN); describing the distribution and habitat usage for each SGCN; evaluating problems impacting SGCN and their habitat; identifying priority research needs to improve our ability to conserve these species; and, developing conservation strategies to address priority problems. Details of the Species Teams and other Action Plan team and committee charges can be found in Appendix D of this document.

Introduction

Vermont with its estimated 7,100 miles of rivers and streams and 809 lakes and ponds supports populations of 92 fish species (Langdon et al. 2006). Eighty of these are recognized as being native to the state. A native species is one that was present in the state prior to early European colonization. The remaining 12 species are non-indigenous to Vermont. These fishes were either purposely introduced, legally and illegally, to waters of the state, such as for sport fish enhancement (e.g., Brown and Rainbow trout), or gained access inadvertently to the state via interstate waterways, such as canals (e.g., Gizzard Shad). Lake Champlain has the most diverse fish community of any Vermont water with about 71 species documented to exist there.

Vermonters are probably aware of the existence of about one-third of the fish species occurring in the state. Our familiarity with most of these fishes is rooted in sport fishing; that is, their recognized value as game fish and to a lesser degree their use as bait fish. As for the remaining two-thirds of Vermont species, many exist here largely out-of-sight of the general public and others are viewed more or less with ambivalence. Nonetheless, the diversity of Vermont's ichthyofauna contributes significantly to the functional ecological complexity of our aquatic systems. Many species are excellent indicators of the health of our environment, such as their sensitivity to toxic chemicals (e.g., mercury and PCBs) and habitat change. Additionally, sport fisheries, utilized and valued by the public, are dependent directly and indirectly on healthy communities and ecosystems.

Native fishes face many conservation challenges. The threats of habitat alteration, loss and fragmentation are pervasive in Vermont's rapidly changing landscape. The introduction of non-indigenous fishes, including associated aquatic pathogens and parasites, also pose risks to aquatic ecosystem health and native species conservation. Just within the past 20 years, seven non-native fishes have shown up in state and interstate waters. Whirling disease, caused by the parasite *Myxobolus cerebralis*, first appeared in native Brook Trout inhabiting Vermont sections of the Batten Kill as recently as 2002. Two viral diseases have also recently appeared in Vermont waters. Largemouth Bass virus was first detected in Lake Champlain in 2002 and a year later in Lake St.

Catherine; and esocid lymphosarcoma infecting Lake Champlain Northern Pike in 2002 (http://www.vtfishandwildlife.com/fisheries_health.cfm). Viral hemorrhagic septicemia (VHS), which was first documented in 2005 infecting fishes in Lake Ontario has spread rapidly to the other Great Lakes and nearby inland waters and has been responsible for large and small scale fish mortalities. Of 37 species of fish known to be susceptible to the VHS virus, 26 are found in the Great Lakes and 24 in Vermont. Unregulated or illegal transportation of fishes from out-of-state sources and between in-state waters is likely cause for the increasing incidences of disease-causing organisms appearing in fish populations in Vermont and other states. To date VHS infections have not been identified in Vermont fish populations; nonetheless, the Vermont Fish & Wildlife Department has taken proactive action to reduce the risks that the virus will find its way here, such as adopting more stringent bait fish and fish transportation regulations.

Implementing the 2005 Wildlife Action Plan

Since adoption of the 2005 Wildlife Action Plan, State Wildlife Grant funds have been used to increase our understanding and conservation of fish SGCN in the state. Some of the projects conducted under the 2005 Wildlife Action Plan include:

- Assessment of Lake Sturgeon in Large Vermont Tributaries to Lake Champlain;
- Development of Metrics to Assess the Quality of Riverine Habitat for Coldwater Fish Based on Stream Temperature;
- Aquatic Invasive Species – Hazard Analysis & Critical Control Point Training for Commercial Baitfish Dealers;
- Development of Guidelines for the Design of Stream Road Crossings for Passage of Aquatic Organisms in Vermont and the Vermont Culvert Aquatic Organism Passage Screening Tool;
- Aquatic Organism Passage at Stream & Road Crossings and Aquatic Organism Passage Barrier Assessments;
- Muskellunge Management & Conservation Planning in the Vermont Lake Champlain Basin;
- Development of In-House Capability to Detect Fish Disease Organisms by Polymerase Chain Reaction;
- Survey Design and Standard Operating Procedures for Forage Fish Assessment in Lake Champlain;
- Landlocked Atlantic Salmon Fry Stocking Evaluation;
- Development of Triploid Brook Trout Production Capability;
- Genetics of Lake Trout Populations in Northeastern Vermont Lakes
- Survey & Inventory of Round Whitefish Populations in Northeastern Vermont Lakes
- Survey of Fishes in the Large Lake Champlain Tributaries in Vermont
- Lacustrine Shoreline Planning
- Anadromous Atlantic Salmon Genetics
- Assessment of Stonecat Populations in the LaPlatte and Missisquoi Rivers of Vermont

- Genetic Examination of Lake Whitefish Population Sub-Structuring in Lake Champlain
- Development and implementation of Disease Spread Prevention Systems to Minimize the Inadvertent Spread of Fish Diseases from Fisheries Management Activities, including the Rearing and Stocking of Cultured Fish, to Waters Supporting Fish SGCN.
- Fish SGCN Selection

Selection criteria included 27 categories reflecting our knowledge about current listing as endangered and threatened; species rarity; population declines; vulnerability of habitats; life history traits; vulnerability to collection, harvest or other taking; other impacts from humans; and dispersal capability. Only native species were considered. Each species was examined across all criteria by the team. Based on this evaluation process the team assigned a high, medium and low rank to attempt to separate species with greater conservation needs from those with more secure status, at least in the short term. See Appendix C of this report, “Developing Vermont’s Action Plan,” for details on selection criteria and process. This approach resulted in 27 species making either the rank of high conservation need or medium conservation need (table 1). Six species listed as SGCN in 2005 were removed from the list (Artic Char, Atlantic salmon-anadromous, Brassy Minnow, Muskellunge, Quillback, and the Connecticut River Sea Lamprey population). No species were added though the American Eel is listed as two populations (Lake Champlain and Connecticut River) because their conservation needs differ.

Table 1. Fish Species of Greatest Conservation Need

High Priority

Northern Brook Lamprey (*Ichthyomyzon fossor*)¹
 Silver Lamprey (*Ichthyomyzon unicuspis*)¹
 American Brook Lamprey (*Lethenteron appendix*)¹
 Lake Sturgeon (*Acipenser fulvescens*)¹
 Round Whitefish (*Prosopium cylindraceum*)¹
 Bridle Shiner (*Notropis bifrenatus*)¹
 Blackchin Shiner (*Notropis heterodon*)
 Blacknose Shiner (*Notropis heterolepis*)
 Greater Redhorse (*Moxostoma valenciennesi*)
 Stonecat (*Noturus flavus*)¹
 Eastern Sand Darter (*Ammocrypta pellucida*)¹
 Channel Darter (*Percina copelandi*)¹
 Sauger (*Sander canadense*)¹

Medium Priority

Blueback Herring (CT River pop) (*Alosa aestivalis*)¹
 Atlantic Salmon (naturally reproducing populations-Lake Champlain & Memphremagog basins) (*Salmo salar*)¹
 Mottled Sculpin (*Cottus bairdi*)
 American Eel (Lake Champlain pop) (*Anguilla rostrata*)¹
 American Eel (CT River population) (*Anguilla rostrata*)¹
 American Shad (*Alosa sapidissima*)¹
 Mooneye (*Hiodon tergisus*)¹
 Cisco or Lake Herring (*Coregonus artedii*)
 Lake Whitefish (*Coregonus clupeaformis*)¹
 Brook Trout (naturally reproducing populations) (*Salvelinus fontinalis*)¹
 Lake Trout (naturally reproducing populations) (*Salvelinus namaycush*)¹
 Redfin Pickerel (*Esox americanus*)
 Silver Redhorse (*Moxostoma anisurum*)
 Shorthead Redhorse (*Moxostoma macrolepidotum*)
 Redbreast Sunfish (*Lepomis auritus*)¹

¹ Regional Species of Greatest Conservation Need in the Northeastern United States (Terwilliger, 2013)

Species of Greatest Conservation Need status for Blueback Herring is limited to the population residing in the Connecticut River. Similarly, Atlantic Salmon, Lake Trout and Brook Trout are defined with limitations. Two populations of American eel—Connecticut River and Lake

Champlain—are recognized individually as the conservation needs of these two runs differ significantly.

Although a disproportionate number of Vermont's SGCN are at the periphery of their range, this should not diminish the importance of these species to the state's biodiversity or in terms of their ecological significance. To illustrate this, of the 80 native Vermont fish species, nearly half of these are here on the eastern edge of each of the species' natural North American range.

Full reports on each Species of Greatest Conservation Need are in Appendix A of this document. The following is a summary of those reports.

Habitat Needs

Vermont's fish species use a variety of habitats: small ponds, large lakes, rivers, streams, and wetlands. Some habitats are used year round and others are occupied seasonally, such as for spawning. Within water bodies, SGCN have specific habitat needs for example, riffles or pools in streams or deep, cold areas of lakes. Loss or degradation of any one critical habitat component can threaten the survival of the species in that particular water.

While most of our fishes are completely freshwater dependent, others spend portions of their lives in both freshwater and marine environments. Three SGCN (American Eel, Blueback Herring, and American shad. are dependent on both. Herring and shad have anadromous life cycles, that is spawning and at least a portion of the juvenile life occurs in freshwater; to attain maturity the fish must go to sea for a period of years. In contrast, eel are catadromous. Maturity is attained in freshwater and reproduction occurs in the ocean. Consequently, whether anadromous or catadromous, these species not only face problems at the Vermont landscape level but also those at the regional and international scopes. To conserve our native fishes, and in particular SGCN, it is essential that we protect, enhance and restore habitat not only within Vermont but also, where appropriate, beyond our borders.

Discussion Threats Impacting Fish SGCN

Factors affecting the security of SGCN are classified as either habitat or non-habitat problems. The most frequently identified habitat related problems impacting aquatic systems are habitat alteration, habitat fragmentation, habitat conversion, invasive non-indigenous species, and climate change. Habitat alteration includes activities, which diminish the quality and/or quantity of habitat features critical to the survival and maintenance of fish populations and other biota on which SGCN are dependent, including stream flows and lake water levels, water temperature regimes, and habitat diversity. Sedimentation is a form of habitat alteration by which the composition of the stream or lake bottoms are altered by greater than normal deposition of fine materials (e.g., silt, sand, organic matter) changing the composition and suitability of substrates to the detriment of their spawning, cover and food production values. Habitat conversion results in the total or near complete loss of function as a result of extreme habitat alteration. Examples of habitat conversion are loss of active floodplains, wetland draining and stream impoundments. Habitat fragmentation occurs when artificial structures, such as dams, impassable bridge structures, and dewatered stream channels, interfere with the movements of fish preventing their access to critical spawning areas or seasonal refugia. Habitat fragmentation also interferes with the natural dispersal of fish and genetic flow within and between populations. Climate change threatens several SGCN at the regional scale by altering (warming) their required thermal regimes. Invasive species, such as nonnative aquatic plants

and zebra mussels, can impact aquatic habitats in a variety of ways. Exotic plants represent a “double edged sword” with respect to the conservation of certain fish species requiring abundant aquatic vegetation. Invasive plant species, such as Eurasian milfoil, may displace native plant communities on which fish are dependent for refugia, food production, and spawning. And, on the other hand, invasive vegetation control programs may eliminate these functions before native plants are restored to desired levels.

While virtually all fishes identified as SGCN are impacted by one or more problems to their habitats, non-habitat related problems are generally more variable from species to species. In some cases, non-habitat problems are a consequence of impacts on habitat. Those affecting SGCN include competition from other species, predation, loss of prey base, water pollution, disease and parasites, and over-harvest. The Sea Lamprey problem in Lake Champlain poses a challenging dilemma. Sea Lamprey has been identified as a known or potential parasite/predator on several SGCN. On the other hand, other SGCN may be threatened by certain control methods needed to control Sea Lamprey abundance and parasitism rates in the lake. Further research and monitoring is required to ensure that successful control measures minimize harm to SGCN.

Research & Monitoring Needs and Conservation Strategies

Priority research and monitoring projects and needs to improve our ability to conserve Vermont’s fish SGCN are identified. Conservation strategies to address problems impacting each SGCN were developed. Those cited most frequently and those most effectively applied for multiple fish SGCN include:

Research & Monitoring Needs

1. Better determine the distribution and relative abundance of populations in Vermont.
2. Acquire better information on species’ life histories, biology and habitat requirements.
3. Monitor and assess populations and habitats for current condition and future changes.
4. Identify and monitor problems for species and their habitats.
5. Establish a centralized fish database within the Agency of Natural Resources to manage fish and other aquatic data, track permits and management projects that impact aquatic species.

Conservation Strategies

1. Protect and restore aquatic and riparian habitats through improved water quality; flow, water level and temperature regimes; sediment reduction; establishment of streamside buffers (see ANR buffer guidance document); and suitable aquatic habitat structure, diversity and complexity.
2. Maintain and restore aquatic organism passage and habitat connectivity at barriers (e.g., dams, culverts) to provide access to critical habitats and maintain ecological connectivity.
3. Assess, monitor and manage as appropriate potential negative and beneficial effects of the Lake Champlain Sea Lamprey control program on SGCN and other non-target fishes.
4. Participate in existing regulatory processes (e.g., Act 250, stream alteration permits) to protect and restore critical habitats.

5. Implement measures and programs to prevent the introduction and expansion of non-indigenous species to Vermont waters; develop and execute appropriate invasive species control programs.
6. Support and cooperate with inter-agency programs for the restoration of anadromous and catadromous fishes to the Connecticut River basin.
7. Support efforts to curb global climate change and its negative impacts on SGCN.
8. Support state and regional efforts to require reduction in emissions from coal burning power plants and other sources of acid precipitation.

See Appendix A for full reports on each fish Species of Greatest Conservation Need.

Conclusion

Over the past decade, since implementation of the 2005 Vermont Wildlife Action Plan, meaningful progress has been made on several fronts to increase our understanding of SGCN and undertake efforts to enhance their conservation in the state. Adult Lake Sturgeon have been documented ascending the Missisquoi, Lamoille and Winooski rivers to spawn and reproduction has been confirmed in these rivers (MacKenzie 2015). A sturgeon restoration plan is now under development.

Technical guidance was developed for the identification of stream crossing structures (e.g., culverts, bridges) where aquatic organism passage problems occur and to assist with the design of passable road-stream crossings. Even though structural improvements emphasize fish passage, other SGCN taxa are also beneficiaries, including numerous amphibian, reptile, mammal and invertebrate species.

Recent surveys of four northeastern Vermont lakes, where there had been historic occurrences of Round Whitefish populations, now appear to be limited to a single population in Lake Willoughby. With this knowledge, actions to conserve this population from potential threats are a critical need and deserve appropriate measures moving forward to secure the species presence in the state.

Other Fish SGCN that are currently receiving attention are Stonecat and Lake Whitefish. Additionally, the ANR continues to adopt and implement programs to minimize the introduction of invasive fish species and diseases to our native fish fauna. Since adoption of the 2005 Wildlife Action Plan four rules have been adopted into statute: (1) a list of prohibited, restricted and unrestricted fish species that may be imported and/or possessed in Vermont; (2) restrictions on the commercial harvest and sale of baitfish, as well as the personal harvest, use and movement of baitfish to minimize the risk of transporting aquatic invasive species, unwanted fish species, and fish diseases and pathogens; (3) a prohibition on the transport of fish in a manner which attempts to keep them alive and represents a risk of introducing unwanted organisms to new waters; and (4) procedures for a Rapid Response General Permit authorizing the ANR to quickly respond to and control an invasive nonnative species. All of these accomplishments establish a good base from which future conservation initiatives will emerge and continued progress will be made on those already on the ground.

Lastly, climate change represents an emerging threat to several fish species indigenous to the Vermont as well as to the integrity and function of aquatic communities and systems. This will require considerable commitments of agency and stakeholder resources to stem the threats that this global problem poses to our fauna and the ecological and economic values and uses that are so important to the citizens of Vermont.

Literature Cited

- (ANR) VT Agency of Natural Resources. 1/20/2005. Riparian Buffer Guidance, and Riparian Buffers and Corridors Technical Papers www.anr.state.vt.us/dec/dec.htm
- Langdon, R.W., M.T. Ferguson, and K.M. Cox. 2006. Fishes of Vermont. Vermont Department of Fish and Wildlife, Waterbury, Vermont.
- MacKenzie, C. 2015. Vermont restoration plan for lake Sturgeon in Lake Champlain. Draft 3. Vermont Department of Fish & Wildlife, Montpelier.
- Terwilliger Consulting, Inc. and the Northeast Fish and Wildlife Diversity Technical Committee. 2013. [Taking Action Together: Northeast Regional Synthesis for State Wildlife Action Plans](#). A report submitted to the Northeast Fish and Wildlife Diversity Committee. Locustville, VA.

Conserving Vermont's Invertebrates

Invertebrate Team

Mark Ferguson, Vermont Fish & Wildlife Department (team leader)
Steve Fiske, Vermont Dept of Environmental Conservation
Trish Hanson, Vermont Forest Parks & Recreation Department
Kent McFarland, Vermont Center for Ecostudies
Bryan Pfeiffer, Consulting Entomologist

Team Charge

The Invertebrate Team was charged with identifying Species of Greatest Conservation Need (SGCN); describing the distribution and habitat usage for each SGCN; evaluating problems impacting SGCN and their habitat; identifying priority research needs to improve our ability to conserve these species; and, developing conservation strategies to address priority problems.

Introduction

The role of invertebrates in our world is mostly unrecognized by humans. But once we get beyond the buzz of mosquitoes and our annoyance with blackflies, our reliance upon these tiny animals slowly unfolds. Within cool forest streams, stonefly and mayfly nymphs consume leaves that fall from forest trees and provide a food source for brook trout and other fishes. In the gardener's corner, bees, flies, wasps, and butterflies pollinate the flowers that will later yield the anticipated fruits and vegetables. Spiders wait to ambush flies in our homes. Dragonflies patrol the stream shores for their insect prey. Beetles, flies, and other invertebrates consume the wastes produced by the human world, leaving fertile soil in exchange. Mostly unnoticed and even avoided these smallest of creatures provide an amazing array of functions that we depend upon in our everyday life. The diversity of species we are so fortunate to have is, itself, something to marvel.

Of the thousands of species that occur in Vermont, several are rare or threatened enough to be at risk of disappearing from the state in the future. The causes that lead to their predicament vary among species. One of the greatest obstacles in taking action to help conserve these "at risk" invertebrates has been the scarcity of information that exists on their distribution, abundance, habitat requirements, life history characteristics, population trends, and threats. It is necessary to assess the status and needs of each species to adequately conserve populations and track the success of these actions. The invertebrate conservation outlined in our first Wildlife Action Plan (2005) therefore focused on obtaining this baseline information.

In the past decade, however, great strides were made in augmenting our knowledge of hundreds of species including bumble bee, butterflies, giant silkmoths, odonates (dragonflies and damselflies), carabid beetles and mussels. While only a fraction of the more than 21,400 invertebrate species believed to be in Vermont, it's a significant advancement for invertebrate conservation and the 2015 Wildlife Action Plan includes specific conservation actions for several species and species groups that were not possible just a decade ago.

Pollinators: While the 2005 Wildlife Action Plan included 33 species of butterflies and moths, many of them pollinators, their role as pollinators was not critical to their selection as SGCN. Since 2005,

concerns over the status of pollinators in general (e.g. flies, wasps, moths, butterflies, beetles, bees, and hummingbirds), and native bees in particular, has become a worldwide concern. Pollination is defined as a mutually beneficial relationship between plants and pollinators wherein the plant provides pollen and/or nectar to the pollinator and the pollinator provides reproductive services for the plant (National Research Council, 2007). Roughly 75 percent of the 240,000 species of flowering plants world-wide rely on pollinators for flower reproduction (NRC, 2007). This includes many plant species that provide browse or forage for larger wildlife, as well as seeds and fruits to support birds and small mammals. These invertebrates also pollinate many commercial crops. In Vermont this includes blueberries, tomatoes, squash, apples, and other produce. The many drivers of pollinator declines include habitat loss and degradation, intensive agricultural practices, use of certain pesticides, diseases and pathogens (Heinz Center, 2013). For this second Wildlife Action Plan nine bumble bee species and 31 species of butterflies and moths—including the Monarch butterfly—were selected as Species of Greatest Conservation Need.

Implementing the 2005 Wildlife Action Plan

Since the adoption of Vermont's first Wildlife Action Plan in 2005 several significant advances were made in the realm of invertebrate research and conservation, including the following—most of which were funded at least in part by the State Wildlife Grants program:

Vermont Butterfly Atlas: The Vermont Butterfly Atlas (2002-2007) surveyed the entire state and analyzed historic records and collections to document the distribution of 103 butterfly species, including 12 species new to Vermont, and giant silkworm moths (Saturnids) too. The project was a collaboration between VFWD, the Vermont Center for Ecostudies, and the Vermont Institute of Natural Sciences with more than 125 citizen scientists participating. Assessing the conservation status of each species and establishing a baseline for understanding future changes was the principle goal of the Atlas. Fifteen species were listed as Species of Greatest Conservation Need representing three ecological groups—wetlands, grasslands, and hardwood forests. The vulnerability to climate change was calculated for 14 butterfly SGCN currently found in Vermont. Three species were found to be extremely vulnerable to climate change, five were highly vulnerable, one moderately vulnerable, three presumed stable, and two could likely increase in numbers in Vermont (Table 1). We expect to see the Butterfly Atlas repeated in ~2027 as a comparison to this baseline.

Peatland and Large River Dragonfly and Damselfly Survey: The first statewide assessment of odonate populations in Vermont, focused on peatlands and large river habitat, was completed in 2009 and provides vital species distribution and occurrence information which has broadened our understanding of rare habitat-specialist dragonfly and damselfly SGCN. Habitat data collected as part of the study provides a comparative baseline for future population trend monitoring. Among the investigation's results were: new encounters with *Gomphus abbreviatus* (S1S2) on two rivers and a moderate gain in the knowledge of *Ophiogomphus* spp. (four species), particularly on the White River; discovery of at least two previously unknown populations of *Neurocordulia yamaskinensis* (S3), a species that had been rarely encountered in Vermont; an expanded knowledge of *Somatochlora* spp. (seven species) distribution in and around peatlands, including *S. albicincta* (S1), *S. cingulata* (S1S2) and *S. franklini* (S1S2), revealing that Silvio Conte National Wildlife Refuge and West Mountain Wildlife Management Area have some of the highest *Somatochlora* diversity in New England; and two new peatland sites for *Williamsonia fletcheri* (S1S2). Future efforts toward odonate SGCN conservation will continue to rely on the information resulting from this and future field studies. For more information see <http://vtecostudies.org/blog/atlas-of-vermont-dragonflies-updated/>.

Table 1. Results of a climate change vulnerability assessment of butterfly SGCN in Vermont (from McFarland, and Zahendra. 2010).

Common Name	Climate Change Vulnerability Index	Confidence	GRank	SRank
West Virginia White	Extremely Vulnerable	Very High	G4	S3S4
Bog Copper	Extremely Vulnerable	Very High	G4	S2
Edwards' Hairstreak	Extremely Vulnerable	Very High	G4	SU
Early Hairstreak	Highly Vulnerable	Very High	G4	S2S3
Hackberry Emperor	Increase Likely	Very High	G5	S2
Tawny Emperor	Increase Likely	Very High	G5	S2
Jutta Arctic	Moderately Vulnerable	Very High	G5	S1
Cobweb Skipper	Presumed Stable	Low	G4	S1
Mulberry Wing	Highly Vulnerable	Very High	G4	S2
Broad-winged Skipper	Presumed Stable	Very High	G5	S2
Black Dash	Highly Vulnerable	Very High	G4	S1S2
Dion Skipper	Highly Vulnerable	Very High	G4	S2
Two-spotted Skipper	Highly Vulnerable	Very High	G4	S2
Dusted Skipper	Presumed Stable	Very High	G4	S1

Vermont Bumble Bee Survey: Growing concerns about the decline of pollinators nationwide and locally prompted this survey. From 2012-2013 biologists and trained citizen scientists led by the Vermont Center for Ecostudies searched more than 1,500 locations across the state and recorded more than 10,000 individual bumble bee encounters. Survey data was then compared with historic data gleaned from public and private collection. The results provided sobering news about the status of Vermont's 15 bumble bee species: more than one-quarter of these species have either vanished or are in serious decline. Harmful parasites accidentally imported from Europe and a class of pesticides toxic to bees are believed to account for North American bumble bee declines (Hatfield et. al. 2012).

Endangered Species Protection for Three Bumble Bee Species: Based on the results of the bumble bee survey, three species—the Rusty-patched Bumble Bee, Yellow-banded Bumble Bee, and Ashton Cuckoo Bumble Bee—were given protection under Vermont's Endangered Species law in 2015. The Rusty-patched Bumble Bee, for example, was fairly common in Vermont until the 1990s. But not a single specimen was found during the 2012-2013 statewide survey. Vermont's bumble bee species appear to be in decline due to parasites imported from Europe and possibly the widespread use of a group of systemic insecticides referred to as 'neonicotinoids' (Hatfield et. al. 2012).

Freshwater Mussel Surveys: Freshwater mussels are recognized as the most endangered group of aquatic organisms in North America, with over two thirds of species considered extinct, endangered, or in need of special protection. In Vermont, ten of the native eighteen species, or 55%, are listed under the state endangered species law, and several others are considered rare. One species, the dwarf wedgemussel, is federally endangered. Over the past decade surveys were conducted to determine the current status of Vermont's freshwater mussel populations in order to determine habitat needs, fish hosts and to establish appropriate species population goals and conservation strategies with a particular

focus on the Brook Floater (*Alasmidonta varicosa*), Fluted-shell (*Lasmigona costata*), and Creek Heelsplitter (*L. compressa*)—all SGCN.

Tiger Beetle Surveys: Two state-threatened tiger beetle species, the Cobblestone Tiger Beetle the Hairy-necked Tiger Beetle were the focus of dedicated surveys from 2005-2010 in Vermont. Data gathered in this study was combined with existing information to help inform the drafting of recovery plans for each species.

Development of the Vermont Invertebrate Database: The paucity of basic information such as species presence, geographic distribution, habitat associations, and life history has limited our ability to direct conservation actions for most groups of invertebrates. However, a substantial amount of information exists that could be gleaned from past invertebrate collecting and research in Vermont, although it is scattered among various collections, government offices, research facilities, published works, gray literature, and other sources. In 2006 the Vermont Invertebrate Database project began the task of compiling and organizing information into a format that can easily be accessed so that it could provide a baseline from existing sources for a broad spectrum of invertebrate taxa for future invertebrate conservation and research planning efforts. This ongoing project is accessible to biologists, researchers, conservationists, land-use planners, educators, and other interested parties.

Compilation of the Ross Bell Carabid Beetle Collection: For more than six decades, Dr. Ross Bell of UVM and dean of Vermont's Entomologists collected carabid beetles in Vermont and across the world. It took a concerted effort by many of the entomologists that followed in his footsteps to catalog and map the thousands of beetles he collected over the years.

Restoration of the Carl Parsons Insect Collection: UVM's Zoological Museum includes more than half a million pinned and identified insects emphasizing Vermont species. This wealth of data covering invertebrate diversity over the past two hundred years had fallen into disrepair from neglect and lack of funding. A renewed interest in conserving this collection, led by VCE, over the past decade has led to its near complete restoration.

Selecting Invertebrate SGCN

The task of assessing the conservation needs of Vermont's invertebrates is daunting. The number of species that occur within the state is not known; however, current estimates hover around 21,000 different species. In addition, many of our invertebrates have not yet been scientifically described. Life history, distribution, and abundance information is available for a small minority of Vermont's invertebrates that would be considered as conservation targets, such as freshwater mussels and some tiger beetles. Thus the Invertebrate Team had to determine how best to assess conservation needs with limited information to draw upon. State and regional experts, as well as entomological hobbyists, have compiled a valuable knowledge base for selected groups of invertebrates over the last century. Although distributional information is often limited, an understanding of the natural history of many of these species enabled the team to move forward. It was the team's decision that identification of SGCN would focus on species and species groups for which adequate information was available.

SGCN selection criteria included: current listing as endangered and threatened; population declines; rarity; vulnerability of habitat; life history traits; vulnerability to collection or take; population limitations; regional status; historic occurrence; disjunct populations; habitat specialization; impacts by exotics; and dispersal capability. A review using these criteria resulted in a SGCN list of 200

species. 141 of these species are ranked High Priority SGCN and 59 are ranked Medium Priority. These species were organized into 15 taxonomic groups to aid in the development of conservation reports (table 2).

Table 2. Invertebrate Groups VT Action Plan 2015	# Species/ Group
Ant Group	2
Bumble Bee Group	9
Beetles-Carabid Group	67
Beetles-Tiger Beetle Group	7
Butterflies-Grassland Group	4
Butterflies-Hardwood Forest Group	4
Butterflies-Wetland Group	6
Moths Group	17
Mayflies/Stoneflies/Caddisflies Group	14
Odonates-Bog/Fen/Swamp/Marshy Pond Group	15
Odonates-Lakes/Ponds Group	7
Odonates-River/Stream Group	17
Crustaceans Group	3
Freshwater Mussels Group	13
Freshwater Snails Group	15
Total	200

The list of species within each of these groups can be found in table 3.

Table 3. Invertebrate Species of Greatest Conservation Need

MP=Medium Priority SGCN; HP=High Priority SGCN.

¹Regional Species of Greatest Conservation Need in the Northeastern United States (Terwilliger, 2013)

²Regional responsibility species identified in (White et al. 2014).

Ant Group (2)

An ant (*Myrmica lobifrons*) MP

A Slave-making Ant (*Temnothorax pilagens*) HP

Bumble Bee Group (9)

Rusty-patched Bumble Bee (*Bombus affinis*) HP

Ashton Cuckoo Bumble Bee (*Bombus ashtoni*) HP

Lemon Cuckoo Bumble Bee (*Bombus citrinus*) HP

Fernald's Cuckoo Bumble Bee (*Bombus fernaldae*) HP

Yellow Bumble Bee (*Bombus fervidus*) HP

American Bumble Bee (*Bombus pensylvanicus*) HP

Confusing Bumble Bee (*Bombus perplexus*) HP

Red-belted Bumble Bee (*Bombus rufocinctus*) MP

Yellow-banded Bumble Bee (*Bombus terricola*) HP

Beetles-Tiger Beetle Group (7)

Boulder-beach Tiger Beetle (*Cicindela ancocisconensis*) ¹ HP

Hairy-necked Tiger Beetle (*Cicindela hirticollis hirticollis*) HP

Boreal Long-lipped Tiger Beetle (*Cicindela longilabris*) HP

Cobblestone Tiger Beetle (*Cicindela marginipennis*) ¹ HP

Northern Barrens Tiger Beetle (*Cicindela patruela*) ¹ HP

Puritan Tiger Beetle (*Cicindela puritana*) ¹ HP

Eastern Red-bellied Tiger Beetle (*Cicindela rufiventris*) HP

Butterflies-Grassland Group (4)

Dusted Skipper (*Atrytonopsis hianna*) MP

Cobweb Skipper (*Hesperia metea*) MP

Monarch (*Danaus plexippus*) HP

Regal Fritillary (*Speyeria idalia*) MP

Butterflies-Hardwood Forest Group (4)

Early Hairstreak (*Erora laeta*) HP

Hackberry Emperor (*Asterocampa celtis*) MP

Tawny Emperor (*Asterocampa clyton*) MP

West Virginia White (*Pieris virginiensis*) HP

Butterflies-Wetland Group (6)

Two-spotted Skipper (*Euphyes bimacula*) HP

Black Dash (*Euphyes conspicua*) HP

Dion Skipper (*Euphyes dion*) MP

Mulberry Wing (*Poanes massasoit*) HP

Bog Copper (*Lycaena epixanthe*) HP

Jutta Arctic (*Oeneis jutta*) HP

Ground Beetles-Carabid Group (67)

Agonum crenistriatum HP

Agonum darlingtoni HP

Agonum moerens MP

Agonum picicornoides HP

Agonum punctiforme MP

Agonum superioris MP

Amara erraticus HP

Amara laevipennis MP

Apristus latens HP

Atranus pubescens MP

Bembidion affine MP

Bembidion cordatum MP

Bembidion gratii HP

Bembidion mutatum HP

Bembidion quadratum HP

Bembidion robusticollis MP

Bembidion rolandi MP

Bembidion rufotinctum HP

Blethisa hudsonica MP

Blethisa julii HP

Blethisa quadricollis HP

Carabus goryi MP

Carabus maeander MP

Dicaelus dilatatus dilatatus HP

Dicaelus teter HP

Dicheirotrichus cognatus HP

Diplocheila impressicollis MP

Diplocheila striatopunctata HP

Dyschirius brevispinus MP

Dyschirius erythrocerus MP

Dyschirius politus politus HP

Elaphropus dolosus MP

Elaphropus levipes MP

Elaphrus fuliginosus HP

Geopinus incrassatus HP

Harpalus fulvilabris HP

Harpalus indigens MP

Harpalus providens MP

Lophoglossus scrutator HP

Nebria suturalis HP

Notiobia sayi MP

Notiophilus aquaticus MP

Notiophilus borealis HP

Notiophilus nemoralis HP

Notiophilus novemstriatus MP

Olisthopus micans HP

Patrobus foveocollis HP

Pentagonica picticornis MP

Pericompsus ephippiatus MP

Philodes alternans HP

Philodes rectangulus MP

Platynus cincticollis MP

Platypatrobus lacustris MP

Pseudamara arenaria MP

Pterostichus brevicornis brevicornis HP

Pterostichus castor MP

Pterostichus pinguedineus HP

Pterostichus punctatissimus HP

Scaphinotus bilobus MP

Schizogenius ferrugineus MP

Sericoda obsoleta MP

Sericoda quadripunctata MP

Sphaeroderus nitidicollis HP

Tachys oblitus MP

Tachys rhodanus HP

Tetragonoderus fasciatus MP

Tetraleucus picticornis MP

Moths Group

A Noctuid Moth (*Zale submediana*) HP
Pine Barrens Zanclognatha (*Zanclognatha martha*) HP
Currant Spanworm (*Speranza ribearia*) HP
A Ghost Moth (*Sthenopis thule*) MP
A Noctuid Moth (*Lasionycta taigata*) MP
A Noctuid Moth (*Lemmeria digitalis*) MP
Franclemont's Lithophane (*Lithophane franclemonti*) HP
An Autumnal Noctuid Moth (*Pachypolia atricornis*) HP
Ostrich Fern Borer (*Papaipema* sp. 2 nr. *Pterisii*) HP
Barrens Moth (*Properigea costa*) MP
A Noctuid Moth (*Xestia fabulosa*) MP
A Noctuid Moth (*Xestia homogena*) HP
Pine Imperial Moth (*Eacles imperialis pini*) HP
New England Buckmoth (*Hemileuca lucina*) MP
Plum Sphinx (*Sphinx drupiferarum*) HP
Clemens' Sphinx (*Sphinx luscitiosa*) HP
A tortricid moth (*Eana georgiella*) MP

Mayflies/Stoneflies/Caddisflies Group (14)

A Caddisfly (*Ceraclea submacula*) HP
A Caddisfly (*Polycentropus glacialis*) HP
A Caddisfly (*Polycentropus iculus*) HP
A Caddisfly (*Rhyacophila amicis*) HP
A Caddisfly (*Rhyacophila brunnea*) HP
A Mayfly (*Ameletus browni*) HP
A Mayfly (*Baetisca rubescens*) HP
A Mayfly (*Eurylophella bicoloroides*) HP
Roaring Brook Mayfly (*Epeorus frisoni*) HP
Tomah Mayfly (*Siphonisca aerodromia*) HP
A Mayfly (*Siphonurus demaryi*) HP
Lawrence Sallfly (*Alloperla voianae*) HP
Appalachian Stonefly (*Hansonoperla appalachia*) HP
Spiny Salmonfly (*Pteronarcys comstocki*) HP

Odonates-Bog/Fen/Swamp/Marshy Pond Group (15)

Mottled Darner (*Aeshna clepsydra*) HP
Zigzag Darner (*Aeshna sitchensis*)² HP
Subarctic Darner (*Aeshna subarctica*)² HP
Comet Darner (*Anax longipes*) HP
Swamp Darner (*Epiaeschna heros*) HP
Cyrano Darner (*Nasiaeschna pentacantha*) HP
Spatterdock Darner (*Rhionaeschna mutata*) HP
Subarctic Bluet (*Coenagrion interrogatum*) HP
Petite Emerald (*Dorocordulia lepida*) HP
Ski-tailed Emerald (*Somatochlora elongata*)² HP
Forcipate Emerald (*Somatochlora forcipata*)² HP
Delicate Emerald (*Somatochlora franklini*)² HP
Kennedy's Emerald (*Somatochlora kennedyi*)² HP
Ebony Boghaunter (*Williamsonia fletcheri*) HP
Black Meadowhawk (*Sympetrum danae*)² HP

Odonates-Lakes/Ponds Group

New England Bluet (*Enallagma laterale*)² HP
Slender Bluet (*Enallagma traviatum*) HP
Lilypad Forktail (*Ischnura kellicotti*) HP
Ringed Emerald (*Somatochlora albicincta*)² HP
Lake Emerald (*Somatochlora cingulata*) HP

Banded Pennant (*Celithemis fasciata*)² HP
Carolina Saddlebags (*Tramea carolina*) HP

Odonates-River/Stream Group (17)

American Rubyspot (*Hetaerina americana*) HP
Blue-fronted Dancer (*Argia apicalis*) HP
River Bluet (*Enallagma anna*)² HP
Rainbow Bluet (*Enallagma antennatum*)² HP
Big Bluet (*Enallagma durum*) HP
Stygian Shadowdragon (*Neurocordulia yamaskanensis*) HP
Spine-crowned Clubtail (*Gomphus abbreviatus*) HP
Midland Clubtail (*Gomphus fraternus*) HP
Rapids Clubtail (*Gomphus quadricolor*) HP
Cobra Clubtail (*Gomphus vastus*) HP
Skillet Clubtail (*Gomphus ventricosus*)² HP
Brook Snaketail (*Ophiogomphus aspersus*) HP
Riffle Snaketail (*Ophiogomphus carolus*) HP
Maine Snaketail (*Ophiogomphus mainensis*) HP
Rusty Snaketail (*Ophiogomphus rupinsulensis*) HP
Riverine Clubtail (*Stylurus amnicola*)² HP
Zebra Clubtail (*Stylurus scudleri*) HP

Crustaceans Group (3)

Appalachian Brook Crayfish (*Cambarus bartonii*) HP
Taconic Cave Amphipod (*Stygobromus borealis*) HP
An Amphipod (*Diporeia hoyi*) HP

Freshwater Mussels Group (13)

Eastern Pearlshell (*Margaritifera margaritifera*)¹ MP
Dwarf Wedgemussel (*Alasmodonta heterodon*)¹ HP
Elktoe (*Alasmodonta marginata*)¹ HP
Brook Floater (*Alasmodonta varicosa*)¹ HP
Alewife Floater (*Anodonta implicata*)¹ MP
Cylindrical Papershell (*Anodontoides ferussacianus*)¹ MP
Pocketbook (*Lampsilis ovata*)¹ HP
Creek Heelsplitter (*Lasmigona compressa*)¹ HP
Fluted-shell (*Lasmigona costata*) HP
Fragile Papershell (*Leptodea fragilis*)¹ HP
Black Sandshell (*Ligumia recta*)¹ HP
Pink Heelsplitter (*Potamilus alatus*) HP
Giant Floater (*Pyganodon grandis*) MP

Freshwater Snails Group (15)

Buffalo Pebblesnail (*Gillia altilis*) HP
Squat Duskysnail (*Lyogyrus granum*) HP
Pupa Duskysnail (*Lyogyrus pupoideus*) HP
Canadian Duskysnail (*Lyogyrus walkeri*) HP
Boreal Marstonia (*Marstonia lustrica*) HP
Spindle Lymnaea (*Acella haldemani*) HP
Mammoth Lymnaea (*Bulimnaea megasoma*) HP
Country Fossaria (*Fossaria rustica*) HP
Star Gyro (*Gyraulus crista*) MP
Dusky Ancylicid (*Laevapex fuscus*) MP
Thicklip Rams-horn (*Planorbula armigera*) MP
Liver Elimia (*Goniobasis livescens*) HP
Sharp Hornsnail (*Pleurocera acuta*) MP
Fringed Valvata (*Valvata lewisi*) HP
Mossy Valvata (*Valvata sincera*) HP

Reports on each invertebrate species group are in Appendix A of this document.

Habitat Needs

As invertebrates are the most diverse of Vermont's animals, the breadth of habitats they occupy is great. From deep lakes and slow rivers to the alpine peaks of our highest mountains, from the leaf litter of lowland floodplain forests to treetops in upland beech stands, there are invertebrates utilizing an amazing array of niches in every corner of Vermont. Many of these species have fairly general habitat requirements, or live in natural communities that are common and secure within the state. A number of these are so abundant that they are treated as forest and agricultural pests. Such species do not normally require special conservation attention.

In contrast, habitat specialization is also a common strategy among invertebrates. Examples of habitats that host specialized invertebrates include fens, black spruce bogs, river cobble shores, large rivers, and alpine meadows. Certain herbivorous invertebrates feed only on specific plant hosts, exhibiting another form of specialization. While such specialization is often advantageous when the required habitat or plant host is plentiful, it creates a risk to these invertebrates when the habitat or host is rare, widely scattered, or also at risk (e.g., the Monarch butterfly is suffering from limited winter habitat and loss of milkweed—its host plant for egg laying). In such cases, conservation attention is sometimes needed to ensure that these specialized invertebrates remain a part of Vermont's fauna.

Certain habitats in Vermont support highly diverse wildlife assemblages, including SGCN invertebrates. Good examples include Lake Champlain and its lower tributaries, where many of our freshwater mussel SGCN are located and peatlands for dragonfly and damselfly SGCN. These species-rich areas provide us the opportunity to help conserve many SGCN simultaneously.

Discussion of Threats Impacting Invertebrate SGCN

The greatest problems faced by SGCN invertebrates in Vermont relate to the loss, degradation, and fragmentation of their habitats. Poorly planned construction is ever-increasing on the landscape, often whittling away the wetland and upland habitats available to these creatures when these areas are not protected. As small habitat units disappear from the landscape, those remaining become more distant from one another; this presents an obstacle to those invertebrates that are limited to short-distance movement.

Declines in pollinators from bumble bees to the Monarch butterfly has been noted nationwide and Vermont is no exception. The drivers of these declines likely vary from region to region and from species to species. In most cases it is a combination of threats that are probably responsible. One commonly cited threat is of a group of systemic insecticides referred to as 'neonicotinoids'. These pesticides are used on agricultural crops, and are also used in concentrated doses on home gardens, lawns, and ornamental trees. Several types of neonicotinoids are highly toxic to bees, in addition to making them more susceptible to parasites and pathogens.

Surface runoff from developed and agricultural lands can carry pollutant and sediment loads that find their way to rivers and streams, particularly during heavy rain events. The buildup of sediments on river bottoms embeds the natural substrate and can smother the invertebrates that reside there. Other pollutants entering streams and rivers can be detrimental to sensitive aquatic species.

Exotic species and diseases are negatively impacting several invertebrate SGCN, and will likely present increased challenges to conservation in the future as new foreign species invade our lands and waters. Parasites accidentally imported from Europe have ravaged Vermont's bumble bee

populations and are particularly deadly to those bees weakened by exposure to the neonicotinoid pesticides mentioned previously. Native freshwater mussels have been eliminated from several large areas of Lake Champlain by the ongoing Zebra Mussel invasion. A small exotic fly (*Tachinid spp*) originally introduced to control gypsy moths instead preys upon many native woodland moth species, including some of our giant silk moths. This may prompt the need for future inclusion as SGCN such species as the Luna, Polyphemus, and Cecropia silkmoths.

Some of the challenges faced by invertebrate SGCN stem from their dwindling numbers and their life history characteristics. Low natural recruitment of offspring into the adult populations can hinder population recovery when numbers are low, such as with freshwater mussels. Other factors shared by several invertebrate SGCN groups that limit or impact populations include trampling/direct impacts, limited localized populations, and the requirement of specialized habitats.

Research & Monitoring Needs and Conservation Strategies

Over the past decade significant efforts have been made to address the lack of knowledge about many invertebrate SGCN. Data has been gathered through both field surveys and through the analysis of existing collections both public and private. Despite these efforts, however, our greatly expanded invertebrate datasets still pale in comparison those of other taxonomic groups and additional research is still needed to better guide conservation efforts. The Invertebrate Team therefore identified priority research and monitoring projects to improve our ability to conserve Vermont's invertebrate SGCN. The Team also developed conservation strategies to address problems impacting each SGCN. Those used most frequently and those best applied to multiple invertebrate SGCN include:

Research & Monitoring Needs

1. Define habitat requirements of SGCN within Vermont, utilizing current knowledge of researchers and field investigations as well as important life history characteristics when such information is lacking.
2. Obtain baseline SGCN distributional and abundance data by conducting surveys throughout the state particularly for additional invertebrate groups, such as spiders, moths, land snails and Orthoptera (grasshoppers, crickets).
3. Freshwater mussels:
 - A. Centralize freshwater mussel data currently managed by multiple entities.
 - B. Resurvey sites known to support rare mussels in the past that have not been surveyed in at least the past decade.
 - C. Determine lampricide impacts on juvenile mussels and the long-term effects on adults.
4. Determine associations between invertebrate SGCN and targeted habitat types and/or natural communities (e.g., wetlands and wetland butterflies) to determine hotspots for conservation planning. Begin with habitat specialists and uncommon/threatened habitats.
5. Develop a threat analysis for odonates and coordinate with regional threat analyses.
6. Assess potential and existing impacts of threats to SGCN populations and their habitats.
7. Monitor trends in SGCN population size and structure, and in habitat.
8. Monitor current and potential threats to SGCN species.

Priority Conservation Strategies

1. Conserve high priority SGCN sites through acquisition, easements, technical assistance, and other cooperative means.
2. Protect and restore aquatic habitats on which SGCN are dependent through pollution abatement, riparian buffers (ANR 2005), flow regulation, easements, and other means.
3. Develop mowing plans for state lands (e.g., parks, roadsides (FWHA 2007, rest stops, old fields and rights-of-way) to benefit SGCN pollinators and to limit the spread of invasive plants. Develop similar mowing BMPs for use by partners, municipalities and VTTrans.
4. Continue developing recovery plans for listed species including freshwater mussels and tiger beetles plans. Develop Best Management Practices (BMPs) and management guidelines where appropriate (e.g., for bumble bees).
5. Work with farmers and other landowners to promote the growth and retention of milkweed and other wildflowers in old fields and pastures to benefit bumble bees, Monarch butterflies and other pollinators.
6. Work with foresters to avoid impacts to SGCN populations and habitats during forest management activities.
7. Work with biologists to minimize impacts and maximize benefits to SGCN invertebrate populations and habitats during and following management activities for sport fish and game wildlife.
8. Participate in existing regulatory processes (e.g., Act 250, stream alteration permits) to protect and restore critical habitats.

Conclusion

The work to conserve our invertebrate SGCN is well underway. We now have statewide data for butterflies and bumble bees, and targeted data for rare dragonflies, damselflies, and freshwater mussels. Efforts to gather and organize invertebrate data from private individuals, museums, and universities are in progress. This expanded base of knowledge can help everyone interested in invertebrate conservation focus on the species, habitats and threats that are most in need of attention and launch new initiatives directed at invertebrate conservation.

See Appendix A for reports on each Invertebrate Group of Greatest Conservation Need.

Literature Cited

- (ANR) VT Agency of Natural Resources. 1/20/2005. Riparian Buffer Guidance, and Riparian Buffers and Corridors Technical Papers www.anr.state.vt.us/dec/dec.htm
- (FWHA) Federal Highway Administration. 2007. A Manager's Guide to Roadside Revegetation Using Native Plants. FHWA-WFL/TD-07-006.
- Hatfield, R., S. Jepsen, E. Mader, S. H. Black, and M. Shepherd. 2012. Conserving Bumble Bees. Guidelines for Creating and Managing Habitat for America's Declining Pollinators. The Xerces Society for Invertebrate Conservation. Portland, OR.
- McFarland, K., and S. Zahendra. 2010. Vermont Butterfly Survey 2002-2007; final report to the Natural Heritage Program of the Vermont Department of Fish and Wildlife. Vermont Center for Ecostudies.

- National Research Council. 2007. Status of Pollinators in North America. Washington, D.C.: The National Academies Press.
- Pfeiffer, Bryan (Wings Environmental). 2009 An Investigation of Odonata in Vermont Rivers and Peatlands. Submitted to the Vermont Fish & Wildlife Department 3/11/2010.
- The Heinz Center. 2013. Pollinators and the State Wildlife Action Plans: Voluntary Guidance for State Wildlife Agencies. Washington, DC.
- White, Erin L., Pamela D. Hunt, Matthew D. Schlesinger, Jeffrey D. Corser, and Phillip G. deMaynadier. 2014. [Δ conservation status assessment of Odonata for the northeastern United States](#). New York Natural Heritage Program, Albany, NY.

Conserving Vermont's Mammals

Mammal Team Members

Alyssa Bennett, Vermont Fish & Wildlife Department
Chris Bernier, Vermont Fish & Wildlife Department (team leader)
Dr. William Kilpatrick, University of Vermont
Dr. James Murdoch, University of Vermont
Dr. Peter Smith, Green Mountain College
Christopher Spatz, Cougar Rewilding Foundation/Northeast Wolf Coalition

Species-specific contributions from:

Moose: Cedric Alexander (Vermont Fish & Wildlife Department),
New England Cottontail: John Gobeille (Vermont Fish & Wildlife Department),
Wolf: John Benson, PhD, Diane Bentivegna, Maggie Howell and Adam Katrick
(Northeast Wolf Coalition),
Wolf and Cougar: John Laundre, PhD. (Cougar Rewilding Foundation and Northeast
Wolf Coalition).

Team Charge

The Mammal Team was charged with identifying mammals of greatest conservation need (SGCN), describing the distribution and habitat usage for each SGCN; evaluating threats impacting SGCN and their habitat; identifying priority research needs to improve our ability to conserve these species; and, developing conservation strategies to address priority threats.

Introduction

For a relatively small state, Vermont is characterized by an impressive diversity of habitat types. This diversity is of course a function of the state's variable climate, geological past and rich human history. The nine distinct biophysical regions that comprise Vermont range in character from that of the low, warm and comparatively dry Champlain Valley where farmers make good use of its productive agricultural soils to that of the cold and largely forested Northeastern Highlands where the underlying granite may be the only thing harder than the animals that call this remote part of the state home. This landscape diversity provides the underpinnings for a similarly diverse suite of mammals ranging from boreal species such as the Canada Lynx and Northern Bog Lemming to those that are better known inhabitants of the south such as Gray Fox and Southern Flying Squirrel. In total, sixty-one mammal species presently exist in Vermont or were here just prior to European settlement. While many of these species are abundant and readily recognizable on the landscape such as deer and woodchuck, other once common species such as wolf and Mountain Lion are now believed to be extirpated and are clearly in need of a concerted conservation strategy if they are to be restored. In between these extremes, however, exist a host of mammals that are either poorly understood, occur in low numbers and/or in specific habitats, are known to be in decline, or are susceptible to any number of identified threats. The updating of Vermont's Wildlife Action Plan was necessary not only to reevaluate the status of these more vulnerable mammals but also to reassess our current knowledge and understanding of the challenges

that lie ahead for all of Vermont's wildlife. It's also an opportunity to measure the progress made to date in addressing these challenges.

While all of the threats identified in the 2005 Wildlife Action Plan (e.g., habitat conversion alteration and fragmentation, competition, pollution, loss of prey base, impacts of roads) continue to be of concern today, the knowledge and experience gained over the past ten years has positioned us to better understand the implications of these threats and the actions we must take to address them. Since 2005, for example, the decimation of bat populations throughout the region as a result of White-nose Syndrome (WNS) *provides a stark demonstration of the vulnerability of Vermont's wildlife to the spread of exotic diseases and highlights the importance of not only implementing measures to avoid the introduction and spread of such pathogens but the value of a solid foundation of baseline population data for these species as well. Similarly, mounting evidence of resident Canada Lynx in the northeast corner of Vermont since 2005 provides clear indication of the critically important role a connected landscape plays in terms of maintaining wildlife diversity as well as of the importance of conserving and managing the unique habitats upon which such specialized species depend.* Although many of Vermont's mammals are extremely adaptable and resilient such as the ubiquitous raccoon, red fox, and skunk, others are sensitive to any number of threats and will require continued vigilance in our efforts to better understand these threats and to implement appropriate conservation strategies. While some threats may be relatively simple, readily identifiable and/or reasonably preventable, others will continue to challenge us through the future and will require comprehensive, multifaceted solutions.

Some of the mammal SGCN presently appear to be secure such as moose and bobcat but could be at risk in the foreseeable future due to loss of critical habitats or to population declines resulting from a variety of environmental threats such as climate change, interspecific competition and disease. A number of species are facing immediate known threats such as with several of the bat species (disease) and American Marten (climate change) and could easily exist only as a memory on the Vermont landscape in the absence of appropriate and timely action. Others are listed primarily because little is known about their population status and/or distribution in Vermont such as with several of the smaller, more secretive species like the voles and shrews. Despite the specific challenges facing these SGCN, the Mammal Team interpreted the selection criteria for listing fairly broadly in the hopes of preventing further declines in any of Vermont's native mammals.

Implementing the 2005 Wildlife Action Plan

Since its adoption in 2005, Vermont's Wildlife Action Plan has guided the implementation of numerous mammal related initiatives aimed at filling critical knowledge gaps and addressing the challenges they face. While most of this work has been focused primarily in Vermont, the research and monitoring needs identified in the plan, as well as the conservation strategies, have been used to justify the state's participation in a number of regional and even national initiatives.

Perhaps the best example of such is Vermont's response to White-nose Syndrome. Prior to the availability of State Wildlife Grant (SWG) funds in 2003, Vermont's efforts to monitor bat populations were limited to periodic hibernacula surveys. But since then, the Vermont Fish & Wildlife Department (VFWD) developed and implemented what has become one of

the more significant state bat conservation initiatives in the region. Major elements of this initiative are:

- Collection of a broad array of statewide bat population data on both summer range and winter hibernacula to determine species composition and relative abundance across the state;
- A study of Indiana Bat maternity colony distribution, size, and habitat use throughout the state;
- Detailed risk assessments to determine wind energy facility impacts to Vermont's bat populations; and,
- A technical assistance and outreach program for land managers and conservation organizations to develop and support the management and protection of important Indiana bat habitat.

Then, in 2008, when White-nose Syndrome was identified in the state, the VFWD was thrown into triage mode to conduct disease surveillance, collect diagnostic samples and coordinate at the state, regional and national levels. However, the experience and knowledge gained by VFWD staff from the original SWG-funded work proved instrumental in positioning the state as a leader in the nation's response to this unprecedented challenge. In the past few years our bat conservation efforts have focused on ongoing surveillance, protection of hibernacula and maternity colony sites, and research into the development of alternative hibernaculum.

Similarly, the work conducted here in Vermont on Canada Lynx has also contributed to the conservation/restoration of this species in the region. Guided by the strategies outlined in the 2005 Wildlife Action Plan, the VFWD continues to partner with other states and organizations in the region, for example, to monitor for its presence (2012 to present – Vermont Trappers Association, US Fish and Wildlife Service, US Forest Service, New Hampshire Fish and Game) in recognition of the critically important role such information plays in the implementation of appropriate conservation strategies. Also, the VFWD remains engaged with its regional partners to identify and conserve critical connective corridors facilitating the continued existence of lynx and many other species across the northern Appalachians (*Staying Connected Initiative* -<http://stayingconnectedinitiative.org/>). Since 2009, Staying Connected Initiative partners have permanently conserved more than 300,000 forested and wetland acres that include wildlife corridors and road crossings essential to healthy wildlife populations across the region.

Since 2005, the VFWD has also undertaken initiatives to study American Marten and Black Bear. The bear research aims to determine the level of impact that wind power facilities have on bear use of adjacent beech stands, but study's findings have significance for the region in terms of critical habitat protection for the species, the permitting requirements of future energy developments and a better understanding of how bears utilize the landscape and maintain genetic diversity across potential anthropogenic barriers.

Ongoing Marten research and monitoring was spurred by the discovery of individual Marten in southern Vermont in the vicinity of a previously deemed failed reintroduction attempt

(1989-1991). Although this work is largely focused on mapping the current distribution and abundance of the species in Vermont, several aspects of this work have regional utility; particularly the testing of a marten occurrence model in collaboration with the New York Department of Environmental Conservation, an assessment of the genetic structure of Marten populations across New England, and the evaluation of various measures to minimize the incidental take of marten in traps set for other species. Working in accordance with Vermont's Wildlife Action Plan, the effort to evaluate the status of mammals here in the state and to assess the challenges they face is clearly of regional, and even national, significance.

As noted above, we knew very little about the status and needs of many of our small mammal SGCN, so in 2007 Vermont initiated its Small Mammal Atlas to determine the distribution, relative abundance, and habitat requirements for all small mammal species. Field surveys yielded the capture of 2,844 small mammals representing 20 different species and distribution maps based on historical and current records were constructed for all small mammal species in Vermont.

In addition to the projects highlighted above, the 2005 Wildlife Action Plan was used to justify and develop a number of key SWG funded research projects designed to fill critical knowledge gaps and address specific threats, including:

- An evaluation of Bobcat habitat use and movements to identify Bobcat home range requirements and key habitat and resource needs;
- A Beaver Wetland Conservation Technical Assistance program to help landowners and land managers resolve conflicts with beaver on their properties while allowing beaver to continue maintaining the wetlands they create for the benefit of beaver and the many SGCN that rely on these incredibly productive habitats; and
- A detailed GIS analysis and prioritization of more than 4,000 forest blocks, the corridors connecting these blocks, and the locations across the state where wildlife crosses roads in significant numbers.

Selecting Mammal Species of Greatest Conservation Need

Of the sixty-one mammal species native to Vermont, the Mammal Team opted to list 34 as Species of Greatest Conservation Need. Those species that were the most vulnerable (faced with immediate threats to survival or showing a significant population decline) were ranked as high. In addition, species that were extirpated locally but were known to exist in adjacent states were included on the high list. In all, 17 species were designated as having high conservation priority (table 1). Eighteen additional species were ranked as medium priority.

The Mammal Team was influenced by the Congressional intent of the State Wildlife Grants program of “keeping common species common” so some of the species in the medium category are those that might be well-distributed and even locally abundant at the present time, but that team members felt were at risk in the foreseeable future due to the increasing potential for mortality, habitat loss/fragmentation or other identifiable threat. Mammals may have been included in the medium category either because little was known about their

population status, distribution, and/or trends in Vermont or they have been considered extirpated in the region (table 1).

Table 1. Mammal Species of Greatest Conservation Need

High Priority

Water Shrew (*Sorex palustris*)¹
 Long-tailed or Rock Shrew (*Sorex dispar*)¹
 Pygmy Shrew (*Sorex hoyi*)
 Little Brown Bat (*Myotis lucifugus*)¹
 Indiana Bat (*Myotis sodalis*)¹
 Small-footed Bat (*Myotis leibii*)¹
 Northern Long-eared Bat (*Myotis septentrionalis*)¹
 Silver-haired Bat (*Lasionycteris noctivagans*)¹
 Tri-colored Bat (*Perimyotis subflavus*)¹
 Eastern Red Bat (*Lasiurus borealis*)¹
 Hoary Bat (*Lasiurus cinereus*)¹
 New England Cottontail (*Sylvilagus transitionalis*)¹
 Rock Vole (*Microtus chrotorrhinus*)
 Woodland Vole (*Microtus pinetorum*)
 Northern Bog Lemming (*Synaptomys borealis*)
 American Marten (*Martes americana*)¹
 Canada Lynx (*Lynx canadensis*)

Medium Priority

Cinereus or Masked Shrew (*Sorex cinereus*)
 Smoky Shrew (*Sorex fumeus*)¹
 Hairy-tailed Mole (*Parascalops breweri*)¹
 Big Brown Bat (*Eptesicus fuscus*)¹
 Snowshoe Hare (*Lepus americanus*)
 Southern Flying Squirrel (*Glaucomys volans*)
 Northern Flying Squirrel (*Glaucomys sabrinus*)
 Muskrat (*Ondatra zibethicus*)
 Southern Bog Lemming (*Synaptomys cooperi*)¹
 Wolf (*Canis sp?*)
 Gray Fox (*Urocyon cinereoargenteus*)
 Black Bear (*Ursus americanus*)
 Long-tailed Weasel (*Mustela frenata*)
 Northern River Otter (*Lontra canadensis*)
 Bobcat (*Lynx rufus*)¹
 Eastern Mountain Lion (*Puma concolor cougar*)
 Moose (*Alces alces*)

¹ Regional Species of Greatest Conservation Need in the Northeastern United States (Terwilliger, 2013)

Of the 33 species identified as SGCN in Vermont's 2005 Wildlife Action Plan, only Mink was delisted during this revision process. The removal of Mink from the list (from medium in 2005 to low in 2015) was precipitated by the fact that very little evidence could be found in the scientific literature supporting the primary concern regarding their vulnerability to environmental toxins. Therefore, in consideration of this and of the existing framework for monitoring the species (trapper derived harvest, catch per unit effort and pelt sales data), the widespread and abundant nature of the current population, and the continued listing of other indicator species such as the Northern River Otter, team members concluded the species was secure for the foreseeable future and that processes were well established for detecting and reacting to changes in the species vulnerability to environmental toxins.

In its 2015 revision process, the Mammal Team identified specific threats to two additional species and assigned them both as medium priority SGCN. Since 2005, several threats for Moose have emerged having the potential to impact populations at a regional scale. The most significant is the recently documented, unprecedented mortality resulting from acute winter tick infestations across parts of its range. Much work is currently underway to assess the effects of this tick related mortality as well as to evaluate other potential stressors influencing the population health of this species along the southern edge of its range.

Similar to Moose, Snowshoe Hare also exist in Vermont at the southern periphery of its range which alone makes it vulnerable to certain threats such as a changing climate but, unlike Moose, the species is a habitat specialist reliant upon early successional northern forest habitat types. Although Hare populations appear secure at present, forest management trends in recent decades have led to declines in early successional habitat throughout the state creating concern for the species' long-term persistence. To further justify the SGCN status for Hare, team members also acknowledged that the species serves as primary prey for

an array of furbearers including other SGCN species (e.g. Canada Lynx and American Marten) and that its security in the state could be jeopardized by shifting carnivore communities responding to climate change.

In addition to removing and adding species to the SGCN list, the Mammal Team also reassigned two previously identified SGCN from medium to high (Little Brown Bat and Northern Long-eared Bat) and two from high to medium (Southern Bog Lemming and Woodland Vole). The former changes were based entirely on the emergence of WNS in 2008 that resulted in the dramatic decimation of these previously healthy bat populations and the latter on the findings of the SWG-funded Small Mammal Atlas which indicated populations of these species to be more secure than was previously believed.

Full reports on each of the Species of Greatest Conservation Need are in Appendix A of this document.

Habitat Needs

The habitat requirements of the mammals listed as SGCN are as diverse as the species themselves and are reflective of Vermont's varied landscape. While some species are habitat generalists (e.g., Moose, Black Bear) simply requiring undeveloped open space, others are specialists dependent upon very specific habitat conditions to fulfill their life cycles (e.g., Indiana Bat, Northern Bog Lemming, and American Marten). Collectively, the habitat needs of the SGCN encompass nearly every identifiable habitat type in the state from the most common and ubiquitous northern hardwood forests to the more scarce and unique alpine meadows. Depending on the species in question, even some of the cultural habitat types can play an important role in the conservation of these species. Thus, in general, maintaining healthy populations of Vermont's native mammals requires the conservation of critical habitats, both specific and broad in nature, and the important connecting corridors linking key habitats across the state and region. It also means conserving large blocks of contiguous forestland with corridors, such as riparian buffers, to provide a network of interconnected habitat blocks suitable for the wide-ranging species such as Lynx, and Black Bear as well as for the numerous, less travelled species that make use of the many niches such a conserved landscape provides.

Discussion of Threats to Mammal Species of Greatest Conservation Need

The threats most frequently identified to the 34 SGCN mammals were: Conversion of Habitat (28), Habitat Alteration (28), Loss of Prey Base (16), Competition (14), Disease (12), Genetics (12) and Climate Change (11).

We do not understand all the ramifications, but the pattern seen elsewhere in the US and the world is that increased human population density, higher consumption of land and other resources, and a lack of awareness of impacts to other species can lead to devastating losses of native biota (TWS 2004). Vermont is not immune from these sorts of impacts and our landscape is continuing to be developed (DeVillars 1999). For example, Vermont lost an average of 4,800 acres of wildlife habitat each year to development between 1997 and 2007 (Plumb). Habitat alteration and loss is a near universal challenge to many native mammal SGCN.

Occupying only those limited portions of the state where suitable habitat conditions prevail, some species are found in low numbers and/or in isolated patches making their long term persistence susceptible to direct habitat impacts and reliant upon functioning connective corridors. Similarly, other species, while abundant to either the north or south of Vermont, exist in the state at the furthest extent of their ranges making them vulnerable to a changing climate and the resulting shifts in biotic communities. Despite our successes at conserving large tracts of land in the state in response to these threats, current trends in the forest products industry and applied forest management practices complicate our ability to manage lands for some species particularly those that rely on early successional habitats. Adequately protecting and managing the landscape to meet the needs of Vermont's SGCN through the future is undoubtedly amongst the greatest challenge we face.

In recent years, the emergence of several pathogens such as WNS, chronic wasting disease (CWD), and winter ticks provide more than ample evidence of the severe consequences such agents can inflict upon whole populations of mammals. In a matter of a few years, for example, we witnessed once thriving populations of bats dwindle to alarming numbers leaving at least one species, the Northern Long-eared Bat, subject to the protections of the federal Endangered Species Act and another, the Little Brown Bat, subject to Vermont's endangered species law. In other states, we watched as biologists scrambled to prevent the further spread of CWD and to minimize the disease's impact on local deer herds. These experiences not only demonstrate the potentially grave consequences such diseases and pathogens have for Vermont's wildlife, but also illustrate the importance of stemming the flow of such agents into the state and implementing sound response protocols should new diseases be discovered within our borders.

Pollution was also identified as a potential threat to several species including bats and Otter. Industrial pollutants and heavy metals such as PCBs and mercury can build up in the bodies of animals exposed to these toxins (Novak, 1987). In Vermont, for example, trace amounts of mercury were readily detected in the tissue of several Otters sampled during annual necropsy work. Although the ramifications are not clear, it is likely that the biomagnification of these toxins negatively affects reproduction and survival. Bats are particularly susceptible to pesticides and other environmental poisons because they store some lipophilic (fat soluble) pesticides in brown adipose fat tissue. These stores are released as bats use their fat reserves during hibernation. Bats can, therefore, be exposed to both chronic and acute poisoning which can result in death. In addition, broad spectrum insecticides can deplete insect diversity and limit the food sources available for a number of the smaller, insectivorous mammals such as the bats and shrews.

Perhaps the biggest challenge for some species such as bats, wolf, and mountain lion is the public's acceptance of and desire to conserve them. Sensational and often inaccurate presentations of public health issues, property damage and potential risk factors involving these species have created an exaggerated fear of these ecologically important animals. The resulting unwarranted negative public perception presents an especially serious threat to the recovery and conservation of these species. It is clear that for some species recovery efforts must begin with a public outreach and education effort.

Research and Monitoring Needs and Conservation Strategies

The Mammal Team developed research, monitoring, and conservation strategies for each individual SGCN species. Below is a compilation of the strategies that arose most frequently:

Research and Monitoring

1. Determine the distribution and relative abundance of populations in Vermont.
2. Identify, evaluate and monitor threats.
3. Determine critical habitat needs and connectivity requirements.
4. Determine life history requirements.

Conservation Strategies

1. Develop outreach and education programs that promote the conservation of SGCN and the habitats that they depend on, and increase awareness of the importance of maintaining or restoring these species.
2. Identify the habitat requirements of SGCN and develop strategies for conservation and protection through fee simple purchase, easements, management guidelines, and cooperative agreements with user groups and landowners, etc. (i.e. bat hibernaculums and maternity roost trees, bobcat denning sites, reverting field habitat for New England cottontail, bear-scarred beech stands, connective corridors, etc.).
3. Initiate an international effort to maintain large blocks of undeveloped forests linked together by habitat corridors in order to provide a network of interconnected habitats throughout northeastern New England and southeastern Canada.
4. Maintain riparian buffers along streams (see ANR 2005).
5. Maintain and restore habitat connectivity and minimize fragmentation of forest blocks. Identify and prioritize wildlife road crossing locations. Work with the Agency of Transportation and adjacent landowners to reduce wildlife mortality and increase the potential for movement from one side of the road to the other.
6. Work to eliminate pollution that causes acid rain, the deposition of heavy metals, and global climate change.
7. Continue to work cooperatively with landowners, towns, and communities to protect critical habitats and maintain connectivity. Provide *Conserving Vermont's Natural Heritage* to municipal and regional planners (Austin et.al. 2004)
8. Participate in existing regulatory processes (e.g., Act 250, stream alteration permits) to protect and restore critical habitats.

See Appendix A for full reports on each mammal Species of Greatest Conservation Need.

Conclusion

Vermont is at a crossroad. Due primarily to conscious choices made by her citizens in the last 100 years (restoration of deer, beaver, turkey, fisher populations, enactment of Act 250 legislation and wetland regulations, etc.), as well as, economic forces that essentially allowed the state to bypass the Industrial Revolution (Bryan, pers com), Vermont has remained predominantly rural throughout the 20th century. Many mammal species, therefore, are at population levels that are likely higher than they were prior to European settlement (fisher, red fox, white-tailed deer, raccoon, bobcat). Today, however, with Vermont's population growing, development pressures increasing and increased roads and traffic the potential for significant habitat destruction in the next ten years is high. In addition, global climate change is already influencing the potential residency of some native mammal populations in Vermont (Royar, pers com). The decisions made by Vermonters today will chart the course for the future and influence the long-term viability of our native wildlife populations.

Literature Cited

- (ANR 2005) VT Agency of Natural Resources. 1/20/2005. Riparian Buffer Guidance, and Riparian Buffers and Corridors Technical Papers www.anr.state.vt.us/dec/dec.htm
- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity. Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
www.vtfishandwildlife.com/library.cfm?libbase_=Reports_and_Documents
- Bryan, F. Political Science Department, University of Vermont, Burlington, VT
- Cool, K. L. 1997. Michigan Gray Wolf Recovery and Management Plan for the Michigan Department of Natural Resources, Lansing, Michigan. Prepared by the Michigan Gray Wolf Recovery Team.
- North American Bat Conservation Partnership Strategic Plan, www.batcon.org/nabcp/newsite/index.html
- Novak, M., J.A. Baker, M.E. Obbard, and B. Malloch. Wild Furbearer Management and Conservation in North America. Ontario Trappers Assoc. 1150pp.
- Northeast Endangered Species and Wildlife Diversity Committee. 1999. Wildlife Species of Regional Conservation Concern in the Northeastern United States. Northeast Wildlife V.54.
- Plumb. 2011 [Environmental Trends Report: The Population Connection](#). Vermonters for Sustainable Population.
- Royar, Kim. Vermont Fish & Wildlife Department.
- Sanders, Chris. 2004. Vermont Species Bat Conservation and Recovery Plan. 55 pp.
- Terwilliger Consulting, Inc. and the Northeast Fish and Wildlife Diversity Technical Committee. 2013. [Taking Action Together: Northeast Regional Synthesis for State Wildlife Action Plans](#). A report submitted to the Northeast Fish and Wildlife Diversity Committee. Locustville, VA.
- The Wildlife Society (TWS). 2004. Position on Economic Growth. Adopted by the TWS Council September 2004.
- Wildlife Species of Regional Conservation Concern in the Northeastern United States. Northeast Wildlife Vol. 54, 1999.
- Wydevan, Adrian P., T.K. Fuller, W. Weber, K. MacDonald. 1998. The potential for wolf recovery in the northeastern United States via dispersal from southeastern Canada. Wildlife Society Bulletin, 26 (4): 776-784.

Conserving Vermont's Plants

Plant Team

Bob Popp, Botanist, Vermont Fish & Wildlife Department
Aaron Marcus, Botanical Information Manager, Vermont Fish & Wildlife Department
Everett Marshall, Information Manager, Vermont Fish & Wildlife Department
Eric Sorenson, Ecologist, Vermont Fish & Wildlife Department
Bob Zaino, Ecologist, Vermont Fish & Wildlife Department
Charlie Hohn, Natural Community Information Manager, Vermont Fish & Wildlife Dept.

Partners

Flora Advisory Group to the Endangered Species Committee (FLAG)
Vermont Flora Task Force
Vermont Forest Parks and Recreation
Vermont Department of Environmental Conservation
US Fish & Wildlife Service
US Forest Service
Natural Resource Conservation Service
Vermont Land Trust
St. Michaels College
University of Vermont
Green Mountain College
Vermont Botanical and Bird Club
New England Wildflower Society
Vermont Center for Ecostudies
NatureServe
Network of Natural Heritage Programs and Conservation Data Centers

Team Charge

The Plant Team identified Species of Greatest Conservation Need (SGCN); described the habitats and natural community types they occur in; evaluated impacts to SGCN and their habitat; identified priority research needs to improve our ability to conserve these species; and developed conservation strategies to address priority problems.

Introduction

Vermont is home to approximately 2,000 species of native plants. This includes 1,200 native vascular plants (seed and flowering plants, ferns and fern allies) and 800 non-vascular plants also known as bryophytes (mosses, liverworts, and hornworts). Many species are quite common (e.g., sugar maple, jewelweed) while others are exceedingly rare (e.g., the Green Mountain Quillwort which is found only in Vermont). For a small, northern state such diversity is remarkable.

Plant diversity is an important part of Vermont's biodiversity and they provide food, habitat and shelter for many animal species. Most plant species occur widely enough on the landscape that human activities do not put them at risk. Rare plants, however, often require

specialized habitats and occur in relatively few locations. Some species are rare because availability of their habitats has always been limited or they are at the edge of their range in Vermont. Others have recently become rare as land uses have affected their traditional habitat. Rare species may require management or protection to ensure their survival in a working or natural landscape.

Vermont's plant diversity is driven, in part, by the different biomes that inhabit the state.

While the majority of the state is dominated by northern hardwood forest, there are also extensive areas of boreal forest in the higher elevations and the northern part of the state, and oak-hickory forests in the Champlain and Connecticut River Valleys. There are even remnant alpine tundra and coastal beach species.

Plant distribution and diversity is also determined by the following factors: the type of the bedrock; surficial deposits (gravels, sands, silts, and clays) that were laid down during and after the last glaciation; soil chemistry; climate, elevation, topography; and past land use history. Vermont has extensive areas of calcareous (limy) bedrock that is conducive to high plant diversity. While acidic soils or bedrock areas have distinctly less plant species diversity, they still contribute to the overall diversity in the state in that certain species are adapted to these conditions.

Vermont Fish and Wildlife Department's botanists and ecologists met to develop SGCN selection criteria and to cross-walk rare plants with natural community types (Thompson and Sorenson 2005). The Team also benefited by work by field botanists in the last 35 years and the rich documentation in locally and regional herbaria for information going back to the 18th century.

Vascular plants identified as SGCN include gymnosperms (includes conifers), angiosperms (flowering plants), and ferns and allies (seedless vascular plants that disperse by spores). Bryophytes collectively are mosses, hornworts, and liverworts. The emphasis of this report is on vascular plants as there is much greater knowledge of them, in part because they are easier to identify. However, bryophytes can be an important component of many habitats and natural community types and even dominate certain types, such as dwarf shrub bog.

Selecting Plant SGCN

The team selected plants as Species of Greatest Conservation Need from the Vermont Natural Heritage Inventory's (VNHI) list of rare plants. Of the approximately 1,200 vascular and 800 bryophytes native to Vermont, 813 were selected as Species of Greatest Conservation Need, 493 vascular plants and 320 bryophytes.

The 439 vascular plants were further prioritized by means of the New England Plant Conservation Program's [Flora Conservanda](#), which identifies those vascular plants that are rare globally, regionally, and locally. Those species identified as rare globally and regionally (222) are ranked High Priority and those considered locally rare, 271 species, are ranked Medium Priority. A full list can be found beginning on page 8 of this chapter.

The list of rare VNHI plants include those species that the rarest and often have threats to some or all the populations. These ranks, explained in a following paragraph are based on

knowledge of experts, field research over the last 35 years, and more than 125 years of historical records from the literature and specimens documented in regional herbaria. The plant ranks are based on the number of presumed occurrences and the threats to these populations.

A comprehensive review and update of the ranks of vascular plants was completed in the fall of 2014 by the VFWD working with members of the Flora Advisory Group to the Endangered Species Committee and others with knowledge of Vermont's flora. The taxonomy was updated to match the New Flora of Vermont (Gilman, 2015) and Flora Novae Angliae (Haines, 2011).

Dorothy Allard, a member of the Flora Advisory Group and a professional bryologist, ranked all the Bryophyte species in the state and maintains a list in conjunction with VNHI that was last updated on April 2, 2011.

The VNHI rarity ranks are defined as: S1– very rare, thought to have 5 or fewer populations with some degree of threat; S2 – rare, thought to have 20 or fewer populations with some degree of threat; and SH – historical, documented from the state, but no currently known populations. VNHI also tracks uncommon species that contain a state rank of S3. These species are generally thought to have 21 to 100 populations. They can be considered watch list species and if their numbers decline substantially they may be considered rare in the future.

Plants and Their Habitat Needs

Vermont plant SGCN occupy a broad and diverse array of natural habitats and community types found in the state. In some cases, these species take advantage of anthropogenic dominated areas that mimic natural habitats, such as wet agricultural meadows, transmission line rights-of-way, or sandy opening in developed areas.

The New Flora of Vermont (Gilman, 2015) generally describes the habitats for all vascular plants, including SGCN. The habitat for SGCN bryophytes can be found in the literature listed in references.

Vascular plant SGCN were cross-walked to the natural community types where they are commonly found (table X). The crosswalk was not completed for bryophytes. The natural community approach of conserving rare species conserves representative examples of each natural community type in different physiographic regions of the state. This approach is a more cost effective way to protect a multitude SGCN, than trying to protect individual species occurrences.

Conserving Vermont's Plant SGCN

The following are highlights of plant conservation work in the state since 2005.

- Monitored hundreds of the rare plant populations state-wide through efforts of VFWD staff, Vermont (plant) Task Force members, New England Plant Conservation volunteers, and our partners.

- Discovered hundreds of new rare plant locations over the past 25 years while conducting natural community inventories, such as the bog-fen and oak-pine, and state lands.
- Documented plant SGCN in the Natural Heritage Inventory database by entering field data for thousands of new and updated records of rare and uncommon species occurrences.
- In 2014 revised the Natural Heritage Inventory rare plant ranking list to include new species and varieties that are now recognized due to taxonomic revisions and nomenclatural changes. This is the first major revision of the list since 2009. The project entailed several hundred taxonomic and rank changes and standardized the ranking of subspecies, varieties, and exotic sub-taxa.
- Collected seeds from numerous high and medium priority SGCN for storage at New England Wildflower Society seed banking facility.
- Developed a list of plant species that are threatened by collection within the state and regionally to improve data sharing while protecting rare species. This collective solution was completed because herbarium images are now available online through the Consortium of Northeast Herbaria [portal](#) and the locations of certain rare species are suppressed.
- Helped protect and mitigate impacts to threatened and endangered plant species from development and facilitated scientific research of these species through the state's endangered species permitting program.
- Discovered a plant species new to science in 2013 a quillwort (*Isoetes viridimontana*), currently known only from one pond site in Vermont.
- Managed invasive plant species that were impacting rare plant populations at multiple sites across the state.
- Supported the 2015 publication of the New Vermont Flora by Arthur Gilman.

Discussion of Problems Impacting Plant SGCN

The most significant near-term threats to plant SGCN across the state is conversion, alteration, and fragmentation of natural habitats and invasive plants and animals. Other sometime less obvious threats to plant SGCN include pollinator declines; plant diseases; suppression of natural processes; an overabundance of certain animals; air pollution, including acid deposition; and how natural and anthropogenic plant habitats are managed. Long-term threats are from increasing human population and footprint; and the many issues related to climate change. We can expect that there will be dramatic shifts in plant communities and diversity in the coming decades and centuries from a warming climate. This inevitability is one that we should start planning for, as there is no turning back from much carbon dioxide we have put into the atmosphere.

For decades state plant conservation efforts in Vermont, and nationwide, have been at a significant disadvantage compared with fish and wildlife conservation, due to a dearth of

federal funding. Since 1937 states have received federal Wildlife Restoration Program funds (Pittman Robertson) for wildlife conservation (birds and mammals) and since 1950 Sportfish Restoration Program funds (Dingell Johnson). While the State Wildlife Grants program provides funds for fish and wildlife that don't directly benefit from these other programs, plant conservation is not eligible. The short-lived federal Landowner Incentive Program (2002-2005) provided funding for plant conservation but was cancelled by Congress in 2006. The only federal funds currently available for plant conservation are through the Endangered Species Act for federally endangered plants. In Vermont only three plant species are eligible, the Jesup's Milk-vetch, Barbed-bristle Bulrush and Small Whorled Pogonia.

Research & Monitoring Needs and Conservation Strategies

The research and monitoring needs most frequently identified by the Plant Team and those would benefit multiple plant SGCN are as follows:

Research & Monitoring Needs

1. Document the distribution, abundance, and viability of populations
2. Monitor species
3. Assess condition and viability of associated habitat or natural community
4. Determine life history, such as seedling establishment, pollination needs, threats from invasives, and seed dispersal strategies.
5. Refine the plant/natural community crosswalk, to facilitate using natural communities in conservation planning to protect assemblages of rare plant species.
6. Follow species distribution changes over time, to document changes related to climate shifts.

Conservation Strategies

1. Implement the Vermont Flora Task Force's annual priority Action List of species to inventory and stewardship priorities in coordination with the New England Plant Conservation Program (NEPCoP). Conduct site visits with VFWD staff, NEPCoP trained volunteers and Task Force members.
2. Offer landowners and land managers technical assistance through inventory and stewardship to support private lands stewardship for plant SGCN. Inform landowners how they can manage their forests for rare plant populations when enrolled in the state's Use Value Appraisal Program (Current Use).
3. Work with landowners and partners to develop conservation easements on lands containing plant SGCN. The Vermont Land Trust and The Nature Conservancy in consultation with the Natural Heritage Inventory provide opportunities for protection of rare plants through inclusion in easements and with monitoring of populations.
4. Manage information on plant SGCN in the Natural Heritage Inventory database. Update the rarity ranks of plants based on new information. Include documentation of rare plant abundance, location, viability, habitat description, threats, and

- landowner contact information and permission. Incorporate rare plant data from VFWD field inventories and various partners, researchers, and citizen science.
5. Support species restoration through habitat and natural community management and restoration to restore or mimic natural processes. Carry on restoration work at the Vermont Army National Guard's Camp Johnson with planned burns of the Pine-Oak-Heath Sandplain community, in part for the benefit of rare sandplain plants. Where necessary, remove competing vegetation to maintain a necessary seral stage. Partner with utility companies and develop and implement annual work plans for habitats with plant SGCN on utility properties and rights-of-way.
 6. Manage rare plant populations that are threatened by invasive species. Update the Vermont Flora Task Force Action list with species that have management or stewardship needs. Coordinate with partners who provide stewardship services or manage the land.
 7. Share the data with partners. Also, provide generalized information to public through the Agency's Natural Resource Atlas and the Vermont Center for Geographic information to be used in project and conservation planning.
 8. Facilitate species restoration through the banking of live plant material by collecting seeds or cuttings that may be used to restore or enhance existing populations threatened with extirpation. NEPCoP in conjunction with the Vermont Task Force prioritizes which plants need to have live material collected. The collection effort is to be coordinated with annual species inventory and stewardship priorities. Plan for species restoration through assisted migration and provide guidance and advice to independent efforts to restore or introduce new populations of SGCN.
 9. Educate the public through outreach about Vermont's plants, particularly SGCN. Tell natural history stories about the habitat in which they occur and include their interaction with animal species.
 10. Develop conservation strategies for suites of plant species at the natural community formation level or similar higher level grouping by generalized habitat type.
 11. Determine the pollination needs of plant SGCN. Conserve and manage habitat of pollinators that are important to plant SGCN.
 12. Develop a long-term funding plan for the VFWD's Plant Program to support conservation of our state's plant diversity. Currently the VFWD's funding for plant inventory, monitoring and stewardship is almost solely from the Department's state funds. Explore funding options from a variety of state, federal and private sources. Create a plant funding committee.
 13. Coordinate with state agencies on management, monitoring, and information sharing. Incorporate plant SGCN into the long range management of Agency of Natural Resources lands. Review annual work plans for potential conflict and where necessary conduct site visits to avoid or mitigate the impacts. Coordinate and share information with ANR's Lakes and Ponds' aquatic plant survey. Coordinate with Vermont Agency of Transportation and Department of Buildings and General Services on SGCN that occur in lands that they manage or own.

14. Share species data with NatureServe, which serves as an umbrella organization for the international network of natural heritage programs. NatureServe create global datasets with range-wide maps for each species with a corresponding species rarity rank. They also reconciliation in taxonomic differences between different jurisdictions.
15. Work with our federal, municipal and NGO partners to prioritize management practices and stewardship needs and enhance information sharing. The U.S. Forest Service currently monitors numerous populations of rare plants on their lands. The U.S. Fish and Wildlife Service monitors rare species on its Wildlife Refuges. Also, VFWD monitors federally listed species through the U.S. Fish and Wildlife Service's Cooperative Endangered Species Conservation Fund (Section 6 of the Endangered Species Act).
16. Develop management plans that include rare plants for landowners with Natural Resource Conservation Service. Coordinate with the Vermont Land Trust who monitors populations of rare plants that are incorporated into their easements. Work with municipal governments who provide protection to rare species and monitor populations.
17. Coordinate and prioritize research with universities and colleges to encourage a focus on high priority conservation needs.
18. Identify, prioritize and maintain existing contiguous forest blocks and associated linkages that allow for movement in response to climate change with consideration to the physical landscape, especially with the makeup of the bedrock. That is, certain plants require a calcareous (limy) bedrock type while others thrive on more acidic conditions.
19. Participate in regulatory processes, such as Acts 250 and 248, the Vermont Wetlands Rule, and the endangered species law to protect rare threatened and endangered plants. Some of these laws only afford protection to those legally listed as threatened or endangered in Vermont. There has been an effort to add protection of habitat to the endangered species law, which would increase the viability of populations in or near developments.
20. List additional species, as warranted, as threatened or endangered to provide additional protection for vulnerable species.

Literature Cited

- Vermont Fish and Wildlife. No date. Vermont Natural Heritage Inventory Database.
- Vermont Dept. Forest, Parks and Recreation. 2105. Voluntary Harvesting Guidelines for Landowners in Vermont
- Natureserve. 2015. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.0. NatureServe, Arlington, VA. U.S.A. Available <http://explorer.natureserve.org>
- Thompson E.H. and E.R. Sorenson. 2005. Wetland, Woodland, Wildland: A Guide to the Natural Communities of Vermont Fish and Wildlife and The Nature Conservancy.

Vascular plants

- Brumback, W.E. and J. Gerke. 2013. Flora Conservanda: New England 2012. The New England Plant Conservation Program (NEPCoP) List of Plants in Need of Conservation. Rhodora: October 2013, Vol. 115, No. 964, pp. 313-408.
- New England Wild Flower Society. 2015. State of the Plants: Challenges and Opportunities for Conserving. New England's Native Flora.
- Gilman, A.V. 2015. New Flora of Vermont. The New York Botanical Garden Press.
- Haines, A. Flora Novae Angliae. 2011. A Manual for the Identification of Native and Naturalized Higher Vascular Plants of New England. New England Wildflower Society.
- Seymour, F.C. 1969. The flora of Vermont; a manual for the identification of ferns and flowering plants growing without cultivation in Vermont. Vermont Agricultural Experiment Station Bulletin.
- Flora of North America Editorial Committee, eds. 1993+. Flora of North America North of Mexico. 16+ vols. New York and Oxford. Vol. 1, 1993; vol. 2, 1993; vol. 3, 1997; vol. 4, 2003; vol. 5, 2005; vol. 7, 2010; vol. 8, 2009; vol. 19, 2006; vol. 20, 2006; vol. 21, 2006; vol. 22, 2000; vol. 23, 2002; vol. 24, 2007; vol. 25, 2003; vol. 26, 2002; vol. 9, 2014
- Bryophytes**
- Flora of North America Editorial Committee, eds. 2007 and 2014. Flora of North America North of Mexico. New York and Oxford. vol. 27, 2007; vol. 28, 2014
- Flora of North America Editorial Committee, eds., Vol. 29. In prep. Flora of North America Editorial Committee, Oxford University Press.
- Anderson, Lewis E., Howard A. Crum and William R. Buck. 1990. List of the Mosses of North America North of Mexico. The Bryologist. Vol. 93, No. 4 pp. 448-499
- Stotler, R. and B. Crandall-Stotler. 1977. A checklist of the liverworts and hornworts of North America. The Bryologist 80:405–428.
- Stotler, R. E. and B. Crandall-Stotler. 2005b. A revised classification of the Anthocerotophyta and a checklist of the hornworts of North America, North of Mexico. The Bryologist 108:16–26.

Vermont's Wildlife Action Plan

Chapter 6

Landscape Conservation

Draft • September, 2015

Introduction	2
Landscape Condition	2
Implementing the 2005 Wildlife Action Plan.....	4
Species of Greatest Conservation Need Benefitting from Landscape Conservation	6
Landscape Characteristics	6
The Ecologically Functional Landscape	9
Threats & Information Needs	11
Priority Conservation Strategies	11
Literature Cited.....	13

6. Landscapes and the Conservation of Vermont's Species of Greatest Conservation Need

Introduction

Maintaining and enhancing landscape integrity and ecological function across Vermont is fundamental to conserving our natural heritage and Species of Greatest Conservation Need. Large, connected landscapes are particularly important for wide-ranging species (e.g., Northern Goshawk, Bobcat, Black Bear, Canada Lynx, and American Marten). And because landscape conservation is the most efficient strategy for ensuring the persistence of the many smaller-scale habitats found within a landscape, it is critical to the conservation of many not-so-wide-ranging SGCN as well. Moreover, healthy, intact landscapes enhance the capacity of species and communities to shift and adapt to the changing climate. For these reasons landscape-scale conservation is a fundamental strategy of this Wildlife Action Plan.

Wide-ranging species require large areas encompassing a variety of habitats in order to find sufficient food, shelter and mates. The home range requirements of our wide-ranging SGCN vary greatly from species to species, as do requirements of habitat quality and the number of individuals needed to sustain a population. For example, some area-sensitive birds may require a minimum forest block size of 7,500 acres (Robbins et. al. 1989). Bobcat populations of 250 breeding females require approximately 2,000 square miles, and maintaining Vermont's black bear population may require as much as 6,000 square miles of habitat (Vermont Fish & Wildlife Black Bear Management Plan 1999). Canada Lynx, Wolf, and American Marten range so widely that Vermont alone can meet only a portion of their populations' current or potential habitat needs. Therefore our landscape conservation efforts cannot stop at the state's borders.

This chapter describes the condition of Vermont's landscapes (historic, current and desired), provides a framework for identifying and prioritizing landscapes important to SGCN and natural heritage conservation based on six key landscape components (Interior Forest Blocks, Connectivity Blocks, Surface Waters and Riparian Areas, Riparian Areas for Connectivity, Physical Landscape Diversity Blocks, and Wildlife Road Crossings), identifies SGCN benefitting from landscape conservation, and identifies significant threats and priority conservation strategies. Additional details and maps of our landscape conservation approach can be found in the Landscape Conservation Report (Appendix F).

Landscape Condition

Historical condition: Forests have dominated the Vermont landscape for most of the last 4,500 years—predominantly Northern Hardwood, Spruce-Fir Northern Hardwood and Oak-Pine Northern Hardwood Forests. The forests were continuous, covering mountains and valleys, with intact riparian zones, except in those areas with significant, long-term Native American settlement. Wildlife and plants moved freely, streams and rivers meandered across natural floodplains and natural processes were intact. It has been estimated that 95% of Vermont was forested when Europeans first arrived in the early 1600s. The population of Native Americans in the Champlain Valley and Connecticut River valley in the early 1600s was only 8,000 and only a small amount of forestland was cleared for agriculture, primarily in the river valleys (Klyza and Trombulak 1999). Significant forest clearing began with the arrival of European settlers, however, primarily for lumber, fuelwood, potash, and agriculture. It has been roughly estimated that the percent of forest cover in

Vermont was reduced to 82% by 1790, 47% by 1850, and reached a low of 37% by 1880, after which the area of forest began to increase as farms were abandoned (various sources in Klyza and Trombulak 1999). According to Harper (1918), by 1850 more than 60% of the land in New England had been cleared for agriculture.

The impact to Vermont's landscape was not limited to these cleared areas. Forests in the region that were not cleared were typically on steep slopes, stony ground, or poorly drained soils. Many of these were heavily harvested for timber and or used as woodland pastures, with the result that virtually all of our forests have been altered by human activity (Whitney 1994). In general, our forests today are much younger than the presettlement forests. The composition of presettlement forests was also different from our present-day forests, as has been described in several studies of early land survey records that documented witness and boundary line trees (Siccama 1971, Cogbill 1998, Cogbill 2000, Cogbill et al. 2002). These studies indicate that beech was much more abundant in presettlement forests, whereas sugar maple and white pine were less abundant. Red spruce was more abundant in mid-elevation presettlement forests, whereas red maple, white birch, and poplars – species now associated with younger forests and human activity – were much less abundant in the presettlement forests (Cogbill 2000).

Aquatic habitat degradation was another result of the extensive land clearing for forestry and agriculture given that aggressive stream clearing of boulders and coarse woody debris was engaged in for stream log driving and flood control, and by dam construction and railroad and road building. Such activities have resulted in the relocation and straightening of stream and river channels throughout Vermont, resulting in an overall decrease in available riverine habitat. For example, a recent assessment of the upper White River watershed between Granville and Stockbridge shows that 93% (17.8 of 19.1 miles) of the length of the mainstem White River has been channelized in the past, 13 miles of which are still in channelized form (Vermont Department of Environmental Conservation 2004). In addition, the extensive removal of natural substrates, such as boulders and coarse woody debris, has reduced overall stream habitat complexity throughout the Northeast (Verry et al. 2000). The hard armoring of channels combined with the construction of flood control dams means that many of Vermont's river channels have not regained their historic sinuosity. Furthermore, the slow regrowth of the Northeast's forests means that large woody debris contribution to stream and river channels has yet to reach historic levels (Verry 2000).

Prior to European settlement in the northeastern United States, natural disturbance (including wind, fire, and flooding) were the primary forces affecting the region's forests. In Vermont, wind has been the primary source of natural disturbance in upland forests, ranging from frequent local blowdowns of individual trees to infrequent hurricane events that can affect thousands of acres. A recent study, based on the review of many sources of information, provides figures on the expected percentage of the presettlement regional landscape occupied by different age classes (Lorimer and White 2003). For northern hardwood forest, the expected percentage occupied by uneven aged forest over 150 years ranges from 70 to 89 percent, depending on the assumptions and models used. In these forests, from 1.1 to 3.0 percent was occupied by early successional forests (1-15 year age class). For spruce-northern hardwood forest, the expected percentage occupied by uneven aged forest over 150 years ranges from 35 to 78 percent, depending on the assumptions and models used. In these forests, from 2.4 to 7.1 percent was occupied by early successional forests (1-15 year age class).

Current Condition: Currently, Vermont is approximately 8 percent water, 71 percent forest and 21 percent non-forest (including open, agricultural and developed land). These statistics do not, however,

express the degree of intactness of these terrestrial and aquatic habitats, nor the barriers in our forest and stream systems.

VFWD conducted an analysis of unfragmented forest blocks in Vermont (Sorenson and Osborne 2014). Each of 4,055 forest blocks was analyzed and ranked for biological and physical diversity factors. Vermont's largest block is 153,000 acres. The average block size statewide is 1,000 acres. But block size is not evenly distributed across the landscape. As seen in figure 1, the largest habitat blocks occur along the spine of the Green Mountains and in the northeastern portion of the State. In the Northeastern Highlands biophysical region, for example, the average block size is 6,810 acres and 2,694 acres in the Green Mountains. In the Piedmont average block size is 830 acres and in the Champlain Valley it is only 413 acres.


Desired Condition (SGCN Needs): To maintain the full complement of Vermont's Species of Greatest Conservation Need and particularly of wide-ranging species such as Black Bear, American Marten, Canada Lynx and Wolf, Vermont needs landscapes of large, connected habitat blocks with interior forests, surface waters and riparian areas where ecological processes and native species are most likely to persist and adapt to climate change. These areas should represent all natural communities (in all successional stages), habitats and physical landscape diversity. Distributed across all Vermont biophysical regions, these landscapes should be connected locally and regionally, now and in the future as land use and climate change, by way of smaller blocks, riparian areas and rivers to allow for plant and animal movement and migration. Structural and functional connectivity should be maintained and enhanced across and under roads and other transportation structures.

Several wide-ranging wildlife species will not persist or re-establish without linkages to other states and Canada. Therefore, regional connectivity (i.e., linkages to New York, New Hampshire, and Canada) must be maintained. Linkages along riparian habitats will also provide connectivity for both semi-aquatic and upland species.

Implementing the 2005 Wildlife Action Plan

Actions by the Vermont Fish & Wildlife Department and partners to implement landscape conservation recommendations of the Wildlife Action Plan since 2005 include:

Contiguous forest/habitat blocks and associated linkages were identified and prioritized as part of the "habitat block project" conducted from 2007 to 2014. Using GIS analysis of existing data, this project identified 4,055 unfragmented forest blocks in Vermont and ranked each block for its biological and physical landscape diversity values. The project also identified a modeling tool for identifying likely wildlife corridors in Vermont. Partners included Vermont Land Trust, the Forests, Parks & Recreation Department, The Nature Conservancy, Audubon Vermont, and Green Mountain National Forest. The project results are now used extensively in VFWD technical



assistance to towns. The project report is “Vermont Habitat Blocks and Habitat Connectivity: An Analysis using Geographic Information Systems.”

VFWD has acquired in fee and through conservation easements many high priority sites that further landscape conservation and provide critical landscape connectivity. From 2005-2013, the Department acquired 41 separate parcels (excluding fishing access areas) in fee totaling more than 4,100 acres to be added to WMAs or to create new WMAs. VFWD also acquired more than 2,300 acres under conservation easement during the same period. All of these projects either directly or indirectly benefit SGCN. Partner organizations including the Forests, Parks & Recreation Department, The Nature Conservancy, The Trust for Public Land, Vermont Land Trust and many local land trusts acquired and managed lands similarly benefitting SGCN.

VFWD provided technical assistance to every Vermont Regional Planning Commission and nearly every town on a variety of wildlife and land planning related issues, including SGCN conservation, habitat blocks, and wildlife corridors. [Conserving Vermont's Natural Heritage](#) (Austin et.al. 2004) was reprinted and distribution of this planning document continues.

The [Vermont Forest Roundtable](#) first convened in 2006 as a venue for information exchange on keeping Vermont’s forests as forests. Organized by the Vermont Natural Resources Council, the Roundtable regularly hosts consulting foresters, professional planners, state agency officials (including VFWD and VFPR), landowners, sportsmen, forest products industry representatives, conservation groups, biomass energy organizations and academics. The Roundtable formed with an initial focus on parcelization and forest fragmentation issues. It’s since facilitated discussions on trends in Vermont’s real estate market and rising forestland values, property tax policy, land use and conservation planning, estate planning, landowner incentive programs such as the Current Use Program, and the long-term sustainability of the forest products industry.

VFWD and the Vermont Agency of Transportation (VTrans) established a joint Wildlife-Transportation Steering Committee in 2007 to guide and support interagency cooperation to make Vermont’s transportation system safer for both people and wildlife. VTrans published its [Vermont Transportation & Habitat Connectivity Guidance Document](#) in 2012. Together they currently support three wildlife camera and road tracking projects to advance our understanding of wildlife’s use of transportation infrastructure. These studies are providing VTrans with improved infrastructure design criteria and VFWD with an enhanced understanding of wildlife movement at key locations in the state.

The [Staying Connected Initiative](#) was established in 2008 to maintain and improve landscape connectivity across the Northern Appalachian/Acadian region of the eastern U.S. and Canada (NY, VT, NH, ME, MA and the eastern provinces) through research, land use planning, land management, land protection and road barrier mitigation. The comprehensive approach of the partnership allows the targeting of specific wildlife movement pinch points and coordinated action and affords some assurance that expensive state investment in wildlife-friendly transportation infrastructure is not undone by conflicting land uses in the near vicinity beyond the transportation right-of-way. Partners include VFWD, TNC, VNRC, VTrans, NWF, Wildlife Conservation Society, and the fish and wildlife and transportation agencies of partner states). VFWD has also worked closely with the North Atlantic Landscape Conservation Cooperative on a pilot conservation design for the Connecticut River watershed.

In 2014-2015 VFWD and partners including Vermont Land Trust, Vermont Forests, Parks & Recreation, The Nature Conservancy, and the Northwoods Stewardship Center produced “Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape” (Sorenson et al. 2015). This report identifies coarse-filter conservation targets for landscape scale features including forest blocks, riparian areas, wildlife and landscape connectivity, and physical landscape diversity that are necessary to effectively conserve many finer scale conservation elements in the face of climate change and habitat loss, including natural communities, rare species, and SGCN.

Species of Greatest Conservation Need Benefitting from Landscape Conservation

Without landscape-scale conservation, some species are unlikely to remain on our landscape. These are wide-ranging species, including the American Marten, Canada Lynx, Black Bear, Bobcat, Northern River Otter, Bald Eagle, Red-shouldered Hawk and Northern Goshawk. Wolf and Eastern Mountain Lion likely could not return without secure landscapes. Landscape conservation, however, is also expected to benefit most of Vermont’s other Species of Greatest Conservation Need as well.

Landscape Characteristics

As part of the Wildlife Action Plan revision, Vermont conducted a broad-based assessment of landscape-level biological and ecological data to identify lands and waters that are of highest priority and value for maintaining Vermont’s ecological integrity. The resulting report, Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape (Sorenson et al. 2015) identified six landscape elements as most effective at capturing the needs of many Species of Greatest Conservation Need and their habitats. They are Interior Forest Blocks, Connectivity Blocks, Surface Waters and Riparian Areas, Riparian Areas for Connectivity, Physical Landscape Diversity Blocks, and Wildlife Road Crossings and are described below.

The assessment identified the blocks, riparian areas and road crossings that are a parsimonious solution to conserving a functional landscape. High priority areas for each element were mapped and will be made available to conservation practitioners and others via the [BioFinder website](#). The report and maps can also be found in Appendix F.

The goal is to maintain the ecological functions provided by each landscape element. For example, the goal for Interior Forest Blocks is to maintain the unfragmented, interior forest of these areas that provides critical habitat for many species of plants and animals. There is considerable leeway on what can happen within a forest block and still maintain interior forest function. For example, most responsible forest management activities are compatible with maintaining the long-term interior forest functions for these blocks. Each section below provides guidelines on what is needed to maintain ecological functions for that element.

While each landscape element is important on its own, it cannot function in isolation. Maintaining or enhancing an ecologically functional landscape in Vermont depends on both the specific function of the element and the ability of landscape elements to function together. Interactions between elements are what support Vermont’s environment and are essential for long-term conservation of Vermont’s biological diversity and natural heritage.

By ‘conservation’ we mean a wide range of activities, from private land stewardship to public ownership and other activities that help maintain ecological function. Many tools can be used to achieve the overall goal. With approximately 80% of Vermont’s land privately owned, management

and stewardship of private lands will be essential to achieving these goals. Other tools include local planning and zoning, state regulations, conservation easements, and ownership by a state or federal agency or a private conservation organization. This document and these maps do not provide detail as to which of these tools are best suited to specific places, but there are recommendations for further prioritization filters that users can apply to make these decisions.

Interior Forest Blocks: Areas of contiguous forest and other natural communities and habitats (such as wetlands, ponds, and cliffs) that are unfragmented by roads, development, or agriculture.

Forest blocks were identified, mapped, and ranked by Sorenson and Osborne (2014). These forest blocks provide many ecological and biological functions critical for protecting SGCN and the integrity of natural systems (Austin et al. 2004), including:

- Supporting natural ecological processes such as predator-prey interactions and natural disturbance regimes;
- Helping to maintain air and water quality and flood resilience;
- Supporting the biological requirements of many plant and animal species, especially those that require interior forest habitat or require large areas to survive;
- Supporting viable populations of wide-ranging animals by allowing access to important feeding habitat, reproduction, and genetic exchange; and
- Serving as habitat for source populations of dispersing animals for recolonization of nearby habitats that may have lost their original populations of those species.

In addition, large, topographically diverse forest blocks will allow many species of plants and animals to shift to suitable habitat within a forest block in response to climate change within the next century without having to cross developed areas to other forest blocks. (Beier 2012)

Connectivity Blocks: The network of forest blocks that together provide terrestrial connectivity at the regional scale (across Vermont and to adjacent states and Québec) and connectivity between all Vermont biophysical regions.

Landscape connectivity refers to the degree to which blocks of suitable habitat are connected to each other (Noss and Cooperrider 1994). There is a high level of connectivity within individual forest blocks. The proximity of one forest block to another and the characteristics of the intervening roads, agricultural lands, or development determine the effectiveness of the network of Connectivity Blocks in a particular area.

A network of Connectivity Blocks allows wide-ranging animals to move across their range, allows animals to find suitable habitat for their daily and annual life needs, allows young animals to disperse, allows plant and animal species to colonize new and appropriate habitat as climate and land uses change, and contributes to ecological processes, especially genetic exchange between populations (Austin et al. 2004). There is general agreement among conservation biologists that landscape connectivity and wildlife corridors can mitigate some of the adverse effects of habitat fragmentation on wildlife populations and biological diversity (Beier and Noss 1998; Noss and Cooperrider 1994; Haddad et al. 2003; Damschen et al. 2006). Specifically, climate change adaptation is enhanced if the long distance movements of plants and animals is supported by a combination of

short movements within large, topographically diverse forest blocks and short corridor movements between forest blocks (Beier 2012).

Surface Waters and Riparian Areas: The network of all lakes, ponds, rivers, and streams, their associated riparian zones and valley bottoms in which geophysical processes occur, and their connections to groundwater.

Vermont's rivers, streams, lakes, and ponds provide vital habitat for a rich assemblage of aquatic species, including fish, amphibians, reptiles, invertebrates (e.g., insects, mussels, snails, worms, freshwater sponges), and plants. This represents an enormous contribution to Vermont's biological diversity. The ecological integrity of an aquatic system is dependent on the condition of the watershed in which it occurs, but is also critically tied to the condition of the riparian area adjacent to the stream or pond. For stability, rivers and streams must have access to their floodplains and freedom to meander within their valley bottoms or river corridors. Naturally vegetated riparian areas provide many critical ecological functions, including stabilizing shorelines against erosion, storage of flood waters, filtration and assimilation of sediments and nutrients, shading of adjacent surface waters to help moderate water temperatures, and direct contribution of organic matter to the surface water as food and habitat structure. Riparian areas are also critical habitat for many species of wildlife that are closely associated with the terrestrial and aquatic interface, including mink, otter, beaver, kingfisher, spotted sandpiper, and wood turtle. Furthermore, the shorelines and riparian areas of rivers and lakes support floodplain forests, several other rare and uncommon natural communities, and many species of rare plants and animals. In addition to these ecological functions that are tied to aquatic systems, the linear network of riparian areas provides a critical element of landscape connectivity for plant and animal movement in response to climate change (Beier 2012). Although many riparian areas and river corridors are highly altered by agriculture, roads, and urbanization, the risk of flooding serves as a natural deterrent for future development. Riparian areas also respond rapidly to restoration efforts (Beier 2012).

Riparian Areas for Connectivity: The connected network of riparian areas in which natural vegetation occurs, providing natural cover for wildlife movement and plant migration.

In addition to supporting the integrity of the lakes, ponds, rivers, and streams that they border, naturally vegetated riparian areas are especially important for providing cover for wildlife movement and other important wildlife habitat, such as nesting habitat for birds. Many wildlife species use riparian corridors for travel to find suitable habitat to meet their life requisites, but certain species are almost entirely restricted to riparian areas, including mink, otter, beaver, and wood turtle. The linear nature of riparian areas contributes to their function as movement corridors for wildlife. Roads, development, and agricultural lands fragment the Vermont landscape. The combination of Riparian Areas for Connectivity and Connectivity Blocks provide the best available paths for connectivity across the landscape.

Physical Landscape Diversity Blocks: Blocks that include physical landscape diversity features that are either rare in Vermont or under-represented in the land and water areas identified as highest priority for Interior Forest Blocks, Connectivity Blocks, and Surface Waters and Riparian Areas.

The Physical Landscape Diversity Blocks complement the other block types and riparian areas in order to more fully represent the full spectrum of physical landscape diversity that is important for an ecologically functional landscape. Physical landscape diversity is represented in this conservation

design by rare Land Type Associations (Ferree and Thompson 2008) and Ecological Land Units stratified by elevation, adapted from Ferree and Anderson (2008).


Physical landscapes (often referred to as enduring features) are the parts of the landscape that resist change. They are the hills and valleys, the underlying bedrock, and the deposits left behind by glaciers. They remain constant even when changes in land cover and wildlife occur, as plants and animals move, and even as the climate changes. However, these physical landscapes cannot continue to drive ecological processes or support plants, animals, or natural communities if they are developed or otherwise significantly altered by human activities.

Wildlife Road Crossings: Sections of road that cross a wildlife corridor where the adjacent landscape quality and permeability are high, usually because the road is adjacent to a forest block, and the road is the primary impediment to animal movement. Likely wildlife road crossings are identified statewide in VFWD's habitat block project (Sorenson and Osborne 2014).

Wildlife corridors (also referred to as wildlife connecting habitats) are lands and waters that connect larger patches of habitat together within a landscape and allow the movement, migration, and dispersal of animals and plants (Austin et al. 2004). Roads represent a barrier to wildlife movement and dispersal of many other species, including some plants. Sections of roads that have suitable habitat on both sides are more likely to allow wildlife movement and dispersal of other species and, therefore, these sections of roads are critical components of maintaining or enhancing an interconnected, ecologically functional landscape. Wildlife road crossings that provide connectivity over or under roads are critically important between adjacent forest blocks and along linear riparian area networks.

The Ecologically Functional Landscape

Maintaining and enhancing an ecologically functional landscape in Vermont depends on conservation of the five landscape level elements described above. It is the specific functions of each alone and their complementarity functioning together that are critical for long-term conservation of much of Vermont's biological diversity and natural heritage. The following map shows the ecologically functional landscape conservation design developed for this Wildlife Action Plan revision. It excludes the Surface Waters and Riparian Areas elements which are difficult to display at this scale.



Map 1. The Highest Priority and Priority portions of the Ecologically Functional Landscape, including Interior Forest Blocks, Connectivity Blocks, and Physical Landscape Diversity Blocks (excluding the Surface Waters and Riparian Areas which are difficult to interpret at this scale).

Threats & Information Needs

Problem/ Information Need Category	Problem/ Information Need Detail	Rank
Determine SGCN Habitat Requirements	Some SGCN and RTE species need to be more confidently captured by landscape and natural community/habitat conservation.	High
Habitat Conversion	Permanent conversion of large blocks of forest to housing development, commercial development, and roads	High
Habitat Fragmentation	Fragmentation of large forest blocks, riparian corridors, and migration paths disrupts animal movement and ecological processes.	High
Impacts of Roads	Roads and road usage disrupt animal movements, alter water quality and stream migration and provide pathways for introduction of invasive species.	High
Climate Change	Species will need to shift their distributions in response to climate-driven habitat and environmental changes. This will be more difficult in a fragmented landscape.	High
Invasive Exotic Species	Invasive species can degrade some landscape scale processes such as species movement and migration. For example, riparian corridors dominated by Japanese knotweed are less suitable for native plant and animal movement.	High
Habitat Conversion	River channel straightening and bank hardening contributes to loss of floodplain connectivity, habitat loss, and downstream erosion and flooding.	High
Habitat Fragmentation	Interruption of movement corridors to and from breeding, feeding, and seasonal habitats via conversion, degradation, and road mortality (herps).	High

Priority Conservation Strategies

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Develop a system to track the loss and conversion of habitat.		ANR, VTrans, ACCD, UVM, VNRC	SWG
Complete phase 2 of the Conservation Design Project: Identifying habitat and species-level conservation goals for SGCN species that are confidently captured by landscape and natural community/habitat conservation		FPR, TNC,	SWG, PR
Refine models of habitat connectivity (e.g., BioFinder's network of Connected land, Local Road Crossings, Staying Connected Linkage Models, Structural pathways)	Number of suitable habitat patches available, miles of riparian corridors & linkages conserved.	TNC, USFWS, USFS	SWG, VHCB, FPR, TNC
Support conservation through fee simple purchase or easements on high priority sites	Number of acres conserved	ANR, VLT, TNC, VHCB, and other land trusts	VHCB, VLT, USFS, USFWS, LWCF,

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
			Forest Legacy
Protect from inappropriate development the highest priority areas identified in the Landscape Conservation Design Report	Number of acres protected	ANR, VLT, TNC, TPL, VHCB, Towns, RPCs, and other land trusts	VHCB, VLT, USFS, USFWS, LWCF, Forest Legacy
Provide Technical assistance to private landowners, user groups and forest managers to reduce habitat fragmentation and degradation and to restore and enhance degraded habitats.	Number of landowners managing for species of greatest conservation need	NRCS, TNC, VFWD, FPR, Coverts, SAF VWA, NWF	SWG
Restore riparian areas to enhance riparian connectivity at sites identified in Landscape Conservation report.	Increase in number of acres of riparian habitat restored and/or conserved	ANR, Agency of Agric., VTrans, Rivers Conservancy Municipal Road Managers	
Financial incentives for private landowners to reduce problems and fragmentation to habitats for wide ranging species and to restore and enhance degraded habitats	Number of acres affected/restored	VFWD, NRCS	EQIP
Technical assistance to town and Regional Planning Commissions. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns incorporating wide-ranging species into planning	VFWD, VNRC, RPCs, VFS, AVCC, SAF, VWA, Coverts, Keeping Track	VFWD
Technical assistance to state and federal land management agencies	Number of state and federal land management plans providing for Lynx and Marten habitat	ANR, USFWS, USFS	ANR
Increase cooperation/coordination between adjacent states and provinces to support and encourage trans-jurisdictional actions to address issues such as global climate change, acid rain and connectivity.	Implementation of trans-jurisdictional actions.	USFWS, USFS, ANR, other states, VTrans, USDOT, TNC, Quebec,	USFWS, AFWA
Monitor, protect and restore water quality from excessive nutrient sediment loading, other pollutants.	Miles of SGCN habitat meeting water quality standards.	ANR, USFWS, NRCS, USFS, Lake & Watershed Associations	
Support efforts to reduce the long range transport of acid rain pollutants to Vermont.	Reduction in acidity levels in monitored high elevation waterbodies	ANR, USFS, AG office, Legislature, Congress.	
Restore aquatic connectivity based on Aquatic Organism			

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Passage recommendations			
Support efforts to manage flow regulation projects to minimize impacts on SGCN	Decrease in number of river miles with altered flow regimes	ANR, ACOE, VT Dam Task Force, USFWS, watershed orgs	LBCP, USFWS, ACOE, SWG
Provide technical assistance to VTrans, to identify and maintain (or restore) terrestrial & riparian habitat connectivity and improve aquatic organism passage	Number of functional linkages across highways/roads Increase in % or number of road crossings that do not impede aquatic organism movement	ANR, VTrans, Better Back Roads, USFWS, USFS, AVCC	SWG, USFWS, LCBP, VTrans
Fund and support a natural resource planner in each RPC	Number of Regional Planning Commissions with natural resource planner. Number of Regional Planning Commissions requesting technical assistance	ANR, USFWS, USFS, EPA, RPC, VFWD, VNRC, Staying Connected	ANR, USFWS, USFS, EPA,
Develop road management BMPs for habitat connectivity		VTrans, VFWD, Staying Connected	FHWA
Increase the number of road structures meeting fish and wildlife passage guidelines	Number of improved/upgraded structures	VTrans, VFWD, Staying Connected, USFWS	FHWA,
Provide more fish, wildlife & natural resource oriented technical assistance to constituent towns for town plan rewrites and bylaw changes	Number of town plans and bylaws with improved language	RPC, VFWD	
Provide natural resource planning support in each RPC in a similar manner to the current transportation planning model.	Number of Regional Planning Commissions with natural resource planners	ANR, USFWS, USFS, EPA	ANR, USFWS, USFS, EPA, General Fund
Conduct municipal-scale natural resource inventories	Number of towns with completed inventories of their natural resources.	VFWD, Enviro Consultants	

Literature Cited

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity. Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT. www.vtfishandwildlife.com/common/pages/DisplayFile.aspx?itemId=111191.
- Beier, P. 2012. Conceptualizing and designing corridors for climate change. *Ecological Restoration* 30(4): 312-319.
- Beier, P. and R. F. Noss. 1998. Do habitat corridors provide connectivity? *Conservation Biology* 12:1241-1252.
- Carleton, Frank H. 1899-1900. Why forest preservation should interest fisherman. 15th Biennial report of the Commissioners of Fisheries and Game of the State of Vermont. p. 98-110

- Cogbill, Charles V. 1998. Report on the Vermont presettlement forest database. Unpublished report for the Vermont Biodiversity Project.
- Cogbill, Charles V. 2000. Vegetation of the presettlement forests of northern New England and New York. *Rhodora*, Volume 102, No. 911, 250-276.
- Cogbill, Charles V., John Burk, and G. Motzkin. 2002. The forests of presettlement New England, USA: spatial and compositional patterns based on town proprietor surveys. *Journal of Biogeography*, Volume 29, 1279-1304.
- Damschen, E. I., N. M. Haddad, J. L. Orrock, J. J. Tewksbury, and D. J. Levey. 2006. Corridors increase plant species richness at large scales. *Science* 313:1284-1286.
- Ferree, C. and E. Thompson. 2008. Land Type Associations Descriptions for Vermont. Vermont Department of Forests, Parks, and Recreation.
- Ferree, C. and M.A. Anderson. 2008. Ecological Land Units. Version 11/2008. The Nature Conservancy Eastern Conservation Science Office, Boston, MA.
- Haddad, N. M., D. R. Bowne, A. Cunningham, B. J. Danielson, D. J. Levey, S. Sargent, and T. Spira. 2003. Corridor use by diverse taxa. *Ecology* 84:609-615.
- Harper, R.M. 1918. Changes in the forest area of New England in three centuries. *Journal of Forestry*, Volume 16, 442-452.
- Klyza, Christopher M. and Steven T. Trombulak. 1999. *The Story of Vermont: A Natural and Cultural History*. Middlebury College Press and University Press of New England. 240 pp.
- Lorimer, Graig G., and Alan S. White. 2003. Scale and frequency of natural disturbances in the northeastern US: implications for early successional forest habitats and regional age distributions. *Forest Ecology and Management*, Volume 186, 41-64.
- Noss, R.F. and A.Y. Cooperrider. 1994. *Saving nature's legacy*. Defenders of Wildlife. Island Press. Washington, D.C.
- Robbins, C.S., D.K. Dawson, and B.A. Dowell. 1989. Habitat area requirements of breeding forest birds of the Middle Atlantic States. *Wildlife Monographs* 103. The Wildlife Society, Bethesda, MD.
- Siccama, Thomas G. 1971. Presettlement and present forest vegetation in northern Vermont with special reference to Chittenden County. *American Midland Naturalist*. Volume 85, No. 1, 153-172.
- Sorenson, E. and J. Osborne. 2014. *Vermont Habitat Blocks and Habitat Connectivity: An Analysis using Geographic Information Systems*. Vermont Fish and Wildlife Department, Montpelier, Vermont.
- Sorenson, E., R. Zaino, J. Hilke and E. Thompson. 2015. *Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape*. Vermont Fish & Wildlife Department and Vermont Land Trust. Montpelier, VT.
- Vermont Agency of Transportation. 2012. [Vermont Transportation & Habitat Connectivity Guidance Document](#). Montpelier, VT
- Vermont Department of Environmental Conservation. 2004. Stream geomorphic assessment data available from the DEC River Management Program, Water Quality Division, Waterbury, VT.
- Vermont Fish & Wildlife Department. 1999. *Black Bear Management Plan*. Montpelier, VT.
- Verry, E. S., J. W. Hornbeck, and C. A. Dolloff (eds). 2000. *Riparian management in forests of the continental Eastern United States*. Lewis Publishers, Boca Raton, FL.
- Whitney, Gordon G. 1994. *From Coastal Wilderness to Fruited Plain: A History of Environmental Change in Temperate North America from 1500 to Present*. Cambridge University Press.

Vermont's Wildlife Action Plan

Chapter 7

Monitoring, Implementation & Review

Draft • September 2015

Monitoring & Adaptive Management.....	1
Current Survey and Monitoring of Vermont's Wildlife and Habitats	2
Meeting the Congressional Requirements for Monitoring and Adaptive Management	4
Adapting Conservation Actions and the Wildlife Action Plan in Response to New Information or Changing Conditions	8
Action Plan: Implementation	9
Implementation and Participation	9
Coordination and Collaboration	10
Prioritizing Conservation Need.....	10
Action Plan Review	11
Literature Cited.....	12

Monitoring & Adaptive Management

“However beautiful the strategy, you should occasionally look at the results.”

—Winston Churchill

Elements five and six of the Eight Required Elements for Wildlife Action Plans outline Congressional expectations for monitoring and plan review:

5. Proposed plans for monitoring species [of Greatest Conservation Need] and their habitats, the effectiveness of the conservation actions proposed in the 4th element [strategies], and for adapting these conservation actions to respond appropriately to new information or changing conditions.
6. Descriptions of procedures to review the strategy at intervals not to exceed ten years.

Just as a doctor checks a patient's blood pressure at every visit, wildlife monitoring allows biologists to identify changes in the health of wildlife (e.g., population changes, the spread of disease, changes to the landscape). Biologists can also monitor the impact of strategies to determine effectiveness just as doctors assess the efficacy of treatments and compare competing medical practices. The goal is not simply to cure one patient but improve the standard of care for all patients.

Taken together elements five and six speak to the need for a program to track changes in wildlife populations and their habitats, and to hone the effectiveness of actions. Adaptive management is a formalized method for learning from experience where design, management, and monitoring are integrated to test assumptions in order to adapt, learn and improve (Salafsky et. al. 2001). Instead of relying on a fixed conservation goal and an inflexible plan for achieving the goal, adaptive management provides a framework of planning, acting and monitoring for midcourse corrections (Figure 5.1).




Fig 7.1 Basic steps in an adaptive management process (adapted from Noss & Cooperider 1994)

In the initial **planning** phase for this Wildlife Action Plan teams identified threats and problems limiting SGCN and their habitats and then developed of conservation strategies and research recommendations that the could be implemented during an **action** phase. Measuring the effectiveness and success of the action would occur in the **monitoring** phase. The cycle begins anew with the fine-tuning of goals and objectives before action is renewed.

In addition to monitoring the status of species and threats impacting their populations, implementation, effectiveness and validation monitoring (Derr et. al 2005) can help ensure that goals and objectives are achieved and SWG funds are spent wisely.

- **Implementation Monitoring:** Assessing the degree to which a conservation strategy was implemented (e.g., were trees planted in a riparian area?).
- **Effectiveness Monitoring:** Measuring the impact or effect of a conservation strategy (e.g., did planting trees in the riparian area stabilize the streambank?—the strategy’s objective).
- **Validation Monitoring:** Checking the assumptions upon which the conservation strategy was based (e.g., did stabilizing the streambank actually reduce sedimentation of spawning beds downstream, producing more brook trout fry? —the project’s goal). Validation monitoring can help answer questions such as: Is the conservation strategy worth repeating or might another strategy produce results faster, more economically, or meet with better social acceptance?

Current Survey and Monitoring of Vermont’s Wildlife and Habitats

The list of current survey and monitoring programs providing relevant data for the conservation and management of SGCN is remarkably long. The Fish & Wildlife Department, sister departments at the Agency of Natural Resources the Department of Environmental Conservation (DEC), the Department of Forests, Parks & Recreation (FPR), and other conservation partners monitor the status of several threatened and endangered species, as well as some rare species, some habitats and some uncommon natural communities. The following is a cursory review of survey and monitoring efforts in that may benefit SGCN conservation and management. It is not meant to be comprehensive.

Species Surveys and Monitoring

Amphibians & Reptiles: The Vermont Reptile and Amphibian Atlas (Middlebury College) is an ongoing citizen science research and monitoring project begun in 1995 that tracks the distribution of reptiles and amphibians in Vermont. VFWD monitors threatened and endangered reptiles and amphibians including the Rattlesnake and Spiny Softshell Turtle, and a SWG-funded survey of vernal pools catalogued some 2500 vernal pool amphibian breeding sites statewide.

Birds: Birds are the most studied and best monitored group of wildlife in Vermont—and nationally. A monumental advancement in bird monitoring during the past decade was the completion and publication of Vermont’s Second Breeding Bird Atlas (2013). A collaborative effort, the Atlas was developed with more than 350 citizen scientists contributing more than 50,000 hours over five years to document every bird species breeding across the state. The project was led by the Vermont Center for Ecostudies (VCE), the Fish & Wildlife Department and others, with significant SWG funding. Other important Vermont bird monitoring efforts include: the annual Breeding Bird Surveys, Mountain Birdwatch, Forest Bird Monitoring Project and LoonWatch (VCE); Common Tern and Peregrine Falcon monitoring programs (Audubon and VFWD); Bald Eagle, Wild Turkey, American Woodcock, waterfowl, and Double Crested Cormorants (VFWD). Regional and

national monitoring efforts include the Breeding Bird Survey and Atlantic Coast Joint Venture.

Fish: Historically referred to as "vermin fishes" (Greeley 1930) and "trash fish," the species not prized by anglers have not been the focus of fisheries biologists either. More recently VFWD, the Vermont Cooperative Fish and Wildlife Research Unit, and other partners have surveyed a broader range of fish including Round Whitefish, Stonecat, darters and Muskellunge. And, a robust, long-term monitoring program for Lake Sturgeon is now underway. [*Fishes of Vermont*](#) (Langdon et. al. 2006) is supported by VDEC's 9,000 record fish distribution database. Non-native invasive species, such as alewife and zebra mussel, are also the subject regular surveys.

Invertebrates: With significant advances were made in the past decade in assessing the status of Vermont's invertebrates—most projects were SWG funded. The Vermont Butterfly Atlas (2002-2007), a statewide survey and analysis of historic records and collections documented the distribution of 103 butterfly species, including 12 species new to Vermont. The first statewide dragonfly and damselfly survey focused on peatlands and large river habitat was completed in 2009 providing species distribution and occurrence information which has broadened our understanding of these rare habitat-specialists. The Cobblestone Tiger Beetle and the Hairy-necked Tiger Beetle were the focus of dedicated surveys from 2005-2010. The Vermont Bumble Bee Survey (2012-2013) led by VCE had biologists and trained citizen scientists searching more than 1,500 locations across the state and recording more than 10,000 individual bumble bee encounters. And, surveys were conducted to determine the status of Vermont's freshwater mussels, their habitat needs, fish hosts and to establish appropriate species population goals. Virtually all other invertebrate taxa remain largely uninvestigated and unknown in Vermont and basic background surveys to document the presence and distribution of major orders of insects in Vermont are needed.

Mammals: Deer, Moose, Black Bear and furbearing species are closely monitored by VFWD which also monitors the endangered American Marten, Canada Lynx and several bat species in both winter hibernacula and summer maternity colonies. The Small Mammals Atlas (2007-2009) compiled historic documents and museum collections and conducted field surveys documenting 2,844 small mammal captures from 47 sites and created distribution maps for all 23 small mammal species in Vermont. Additionally, Keeping Track, Inc. has citizen monitoring teams in many sections of the state and region collecting long-term data on Black Bear, Bobcat, Moose, Fisher, River Otter, and Mink. Numerous individual localized surveys also occur but most are not ongoing, repeatable monitoring efforts.

Habitat and Vegetation Surveys and Monitoring

The Vermont Fish & Wildlife Department VFWD conducts ongoing natural community inventories to identify and map natural community types statewide. A survey and report on the distribution, ecology, classification of hardwood swamps was completed in 2004. VFWD also updates and maintains data on known and mapped significant natural communities, maps natural communities on state land and works with non-governmental organization partners to map or identify significant natural communities on their lands.

The [Biomonitoring and Aquatic Studies program](#) established by DEC's Watershed Management Division 1985 monitors the biological integrity of Vermont's lakes, wetlands, rivers, and streams and tracks long-term trends in water quality through changes over time to ambient fish

The [Lake Champlain Long-term Water Quality and Biological Monitoring Project](#) began in 1992. A joint effort shared by DEC and the New York State Department of Environmental Conservation, the primary purpose of the project is to detect long-term environmental change in the lake.

The [Forest Inventory and Analysis](#) (FIA) is a recurring inventory conducted by the US Forest Service's FIA Unit of the [Northeastern Research Station](#) in conjunction with the [Vermont Department of Forests, Parks & Recreation](#). The inventory provides data for measuring changes and trends in the extent and condition of forest land, associated timber volumes, and rates of timber growth, mortality, and removal (Wharton et. al 2003). Though this information is developed primarily for timber management and does not track old-growth forests it does provide important information to wildlife managers.

The [National Resources Inventory program](#) of the [National Resource Conservation Service](#) (NRCS) collects and distributes data on a state, regional and national level about the status, condition, and trends of soil, water, and related resources. The focus is primarily on agricultural lands with data includes available land-use types and land-use changes, erosion, and wetlands.

The [Vermont Monitoring Cooperative](#) (VMC) coordinates numerous monitoring and survey operations in Vermont focusing primarily on forest health issues. VMC's databases house more than two decades of data gathered for five main components of the forested ecosystem: air, water, forest, wildlife and soil resources.

Meeting the Congressional Requirements for Monitoring and Adaptive Management

Monitoring is clearly a linchpin in the adaptive management process. Monitoring is also a complex, demanding and expensive task that never ends. Monitoring was also the weakest link of virtually every Wildlife Action Plan in 2005. The reason is that with the funding and staffing resources available to states, monitoring the status of all SGCN and their habitat and threats was simply not possible. For the 2015-2025 Vermont will attempt the following:

1. **Monitoring of Threatened & Endangered Species and other select SGCN:** Monitoring programs are in effect for many of the state and federally listed species in Vermont, including, for example, the Spiny Softshell Turtle, Timber Rattlesnake and Eastern Ratsnake, Bald Eagle and Common Tern, Lake Sturgeon, Canada Lynx and Little Brown Bat, and Jesup's Milk-vetch and Northeastern Bulrush. Monitoring programs remain in effect for Peregrine Falcon and Common Loon which were both removed from Vermont's endangered species list in 2005.
2. **Taxa-wide surveys:** The Vermont Breeding Bird Atlas is the state's most comprehensive bird survey. The first Atlas (published in 1981) documented the distribution of every bird species breeding in the state. The second Atlas (published

- 2013) repeated this effort and in doing so documented changes in species distributions. The best data we have for many species. To complete this exhaustive, statewide survey took monumental effort with hundreds of volunteers donating thousands of hours from 2003-2007. Similar efforts were initiated with the Vermont Butterfly Atlas (2007) and the Vermont Bumble Bee Survey (2013) providing our first statewide snapshots of these groups where the data was collected in a rigorous, repeatable manner. While it may be another 20 years before they are repeated they can provide vital information regarding species status and trends. Additional statewide surveys may be initiated for other taxonomic groups in the coming years.
3. **Other species surveys:** Many of the surveys described above in the section titled “Current Survey and Monitoring of Vermont’s Wildlife and Habitats” will continue, and additional surveys for other SGCN will be initiated as staffing and budgets allow. Regional species assessments will also be supported through a collaboration of the fish and wildlife agencies of the 13 northeastern states (Northeast Association of Fish & Wildlife Agencies) which together fund projects through the [Regional Conservation Needs program](#) such as the [Wood Turtle Status Assessment](#) and Dragonfly and [Damselfly Status Assessment](#).
 4. **Development of baseline distribution and abundance estimates:** Little data exists for most SGCN (lack of data was one criterion for selection as a Species of Greatest Conservation Need). Determining SGCN distribution and abundance is needed in order to establish meaningful baseline data which then can be used to determine measurable goals and objectives that are the foundation of monitoring priorities.
 5. **Survey/Monitoring Protocol Development:** Rigorous protocols for surveying, monitoring and data analysis for many species do not exist, or are not applied consistently throughout a species’ range to provide robust data. VFWD will continue to collaborate with partners on the development of survey protocols through programs such as the Regional Conservation Needs program (e.g., [A Framework for Coordinated Bird Monitoring in the Northeast](#) and [Development of Regional Analysis for Frog Call Survey Data from the North American Amphibian Monitoring Program](#)). To be successful, any Action Plan monitoring program will need to address these four challenges. It is hoped that the Action Plan and SWG funds will help direct future research and development efforts, facilitate the integration of existing monitoring projects across organizations and improve collaboration.
 6. **Regional Monitoring of SGCN and Habitats:** The Northeast states collaborated to develop the [Northeast Regional Monitoring and Performance Measures Framework](#) (NEAFWA 2008) which identified representative habitats and species groups and proposed indicators of status and trends. States then tasked The Nature Conservancy to test the Framework with a GIS-based evaluation of target habitats conditions, population trends, and land protection status. The final report, [Conservation Status of Fish, Wildlife, and Natural Habitats in the Northeast Landscape](#) (Anderson and Olivero 2011), provides baseline measures at the regional level from which to gauge changes and progress on conservation efforts. Repeating

this evaluation every 5-10 years could provide regional-scale measures of SGCN and habitat status.

7. **Threat Monitoring:** Vermont will continue to monitor for diseases such as White-nose Syndrome (bats), Snake Fungal Disease (rattlesnakes), Chronic Wasting Disease (deer), Viral Hemorrhagic Septicemia (fishes), *Batrachochytrium salamandrivorans* (salamanders) and Avian Influenza (birds) and many other diseases affecting SGCN that have been found or could be introduced to the state. Vermont has also programs in effect to monitor for the eventual arrival of forest pests, including the Asian long-horned beetle and emerald ash borer, in order to prevent the establishment and/or limit their spread within the state. Vermont lacks rigorous method for tracking the loss of habitat across the state.
8. **Effectiveness Monitoring:** [Wildlife TRACS](#) (Tracking and Reporting Actions for the Conservation of Species) is a database developed by the US Fish & Wildlife Service for tracking and reporting conservation activities funded through its Wildlife and Sport Fish Restoration Program. It will include an effectiveness tracking component intended to track and report project outputs, effectiveness measures, and species and habitat outcomes (based on [Measuring the Effectiveness of State Wildlife Grants Final Report](#) (AFWA 2008)). It should be noted that though TRACS will provide a consistent system for tracking projects effectiveness, it may take years to determine if a project is indeed effective (e.g., waiting for trees to grow to sufficient height to shade a stream) and validation of a strategy's success may be difficult to tease out from other problems impacting a species or a site (e.g., the strategy did produce more salmon fry but the results were masked two unseasonably hot summers and an accidental chemical spill).

Considerations for the development monitoring programs:

Before any new monitoring programs are initiated a review of existing efforts and careful planning are required. Such planning should take into account the following considerations:

Collaboration: Planning to develop and implement a Species of Greatest Conservation Need monitoring program should begin with collaboration. As with the design of actions in this report, successful monitoring of SGCN will require the help and cooperation of many partners. Many current survey and monitoring efforts are conducted by interagency and inter-organizational efforts locally, regionally and nationally. These collaborations share expertise, make the best of limited resources, prevent redundancies of effort, increase the level of expertise of volunteers and improve program quality and effectiveness.

How much collaboration is needed? As many entities as possible should be brought together to develop consistent monitoring protocols and systems for data collection and data sharing, identifying indicators for species and habitats and goals and objectives for SGCN conservation.

The need for collaborative fund raising efforts cannot be overstated. Sufficient funds are imperative for monitoring to be effective. The State Wildlife Grants program currently is not sufficiently funded to finance the monitoring needs outlined here. Even if it was, state-side match is insufficient. A collaborative effort of agencies, conservation partners, local, state

and federal elected officials, NGOs and private businesses and individuals is needed to develop adequate funding mechanisms at the state and federal levels.

Coordination: The coordination of monitoring programs, summarizing of results and sharing data with resources managers, researchers, local, state and national decision makers, educators, stakeholders and the general public will be essential to the success of a monitoring collaborative, to Action Plan efforts and to wildlife conservation in general. Solid coordination throughout the implementation phase will also make revisions of the Action Plan report straightforward and uncomplicated.

Indicators: Monitoring every SGCN, their habitats, problems and the effects of conservation actions is too costly and time-consuming to ever complete. Relevant indicators that are measurable, precise, consistent, and sensitive are needed as coarse filters to make monitoring useful and manageable. Indicators should also be of appropriate scale, easily obtained and obvious in meaning so that results can be supported by a broad array of users.

Citizen Science: Successful monitoring projects such as VINS' Bird Atlas, Butterfly Atlas and LoonWatch, the Vermont Reptile and Amphibian Atlas, Keeping Track Inc.'s big mammal monitoring, Audubon's Christmas Bird Counts, and Great Backyard Bird Count and VFWD's Big Game Report Stations provide multiple benefits that should be considered in the development of new monitoring efforts. In addition to the direct benefits—improved wildlife knowledge—citizen-based monitoring also provides wildlife education through active field work on local projects, boosts awareness of and involvement in natural resource protection at the community level, and can be highly cost-effective.

VMC as a Model for Coordination of Statewide SGCN Monitoring: The Vermont Monitoring Cooperative (VMC) (<http://vmc.snr.uvm.edu>) is a collaborative partnership that collects and pools information and data on Vermont's forested ecosystems. Participating cooperators from government, academic and private sectors, conduct research projects on a variety of topics including forest health, air quality and meteorology, wildlife and aquatic systems. The VMC makes the data and results from these projects available to other scientists, educators, resource managers and the general public through its online data library and card catalogue containing the data and metadata from more than 100 projects.

Data storage and data sharing: The volume of government (local, state, federal), NGO, and private sectors data available for plants, animals, ecosystems, climate, geology, hydrology, social and economic that could be used to conserve wildlife is simply huge. The management, storage and accessibility of monitoring data will be a significant issue for any coordinated monitoring efforts. VFWD's Natural Heritage Inventory manages much of the current data for rare wildlife in collected in Vermont but the program is already understaffed. The Natural Heritage Inventory is the Vermont affiliate to NatureServe (www.natureserve.org) an international network of biological inventories—known as natural heritage programs and conservation data centers—operating in all 50 U.S. states, Canada, Latin America and the Caribbean. NatureServe collects and manages data on rare, threatened and endangered plants, animals, and ecosystems, establishes scientific standards for biological inventory and biodiversity data management, and develops data management tools.

Adapting Conservation Actions and the Wildlife Action Plan in Response to New Information or Changing Conditions

The Little Brown and Northern Long-eared Bats underscore the need to adapt management to changing conditions and information. In 2003, VFWD established a Bat Conservation and Management Program determine the distribution and abundance of nine bat species and to identify conservation strategies for these bats. Then in 2008 White-Nose Syndrome (WNS) appeared in the Northeast and monitoring efforts revealed population declines in excess of 90% between 2008 and 2010 for Little Brown and Northern Long-eared Bats at many hibernacula. Several of the high priority conservation strategies identified in 2003 had to set aside as biologist scrambled to identify the cause and extent of WNS and to prevent the total extirpation of these species. Action Plan monitoring and review procedures will be the primary tool to identify new information, changing conditions and the need for adaptation. It will act at three scales—individual conservation projects, ongoing plan-wide adaptations (year-to-year), and 10-year plan review.

The iterative nature of adaptive management (plan→ implement→ monitor→ evaluate→ plan→...) builds opportunities to adapt directly into Action Plan project management activities. Project reporting, monitoring and the increased communication and coordination among conservation partners fostered by Action Plan implementation will feed into overall Action Plan management from year-to-year. The USFWS allows states to incorporate significant “Emerging Issues” into the Wildlife Action Plan without full plan-wide revisions to their Action Plans, as was done with bats and WNS in 2009. All this information will be used to formally review and revise the Action Plan on a 10-year cycle (see also Action Plan Review later in this chapter).

Action Plan: Implementation

Congressional intent for Wildlife Action Plans is for states to identify and address the needs of species that might require help in order to prevent their becoming threatened or endangered. The full import of the word “comprehensive” becomes overwhelmingly clear as numbers in this report are tallied (more than 2,000 threats and 1,000 conservation actions identified for 133 vertebrate species, 200 invertebrates, 645 plants and more than 100 habitat/community/landscape categories). The next steps, conducting the recommended research, setting species and habitat goals and objectives, implementing strategies and designing and implementing the monitoring programs outlined in this report requires the continued help and support of all conservation partners—those that participated in the Action Plan development and new partners as well.

Congress has designated state fish & wildlife agencies as Action Plan and State Wildlife Grants (SWG) custodians because these agencies are mandated by state law to manage and protect wildlife. Custodial responsibilities include not only delivering the completed Action Plan but also for regular review and updating of the Action Plan report and administering SWG funds. To carry out these responsibilities the VFWD will assign sufficient staff and resources to this program to manage projects, coordinate efforts and monitor overall program operations.

The VFWD will take the lead in coordinating the implementation of the research and monitoring recommendations and conservation strategies described in this report. While the Department may be responsible for implementing much of the research, monitoring and conservation strategies, Conservation partners may be the more logical and appropriate leaders for other research and strategy implementation, due to their skills and expertise, staffing, history, location, available resources and constituencies.

The Action Plan will remain a work in progress for many years, an experiment in long-term multi-species conservation on a broad scale. Much of the work in this document is ground breaking. Many of the species examined here have not received focused attention before. The next few cycles of implementation, review and updating of individual strategies and the Action Plan report overall will be the particularly important for working out kinks, testing methods, and improving aspects of the Action Plan.

Implementation and Participation

As a wildlife conservation plan for the entire state, the Wildlife Action Plan includes some strategies that almost any individual or organization can implement. And, any and all interested partners are encouraged to take part. Though many of these actions will not require the notification of VFWD, tracking the implementation and outcomes of actions is vital to the monitoring and adaptive management goals outlined elsewhere in this chapter. All participating partners are encouraged to consult with VFWD prior to taking action.

Impacts on other species, habitats and ecological processes and functions should always be considered when implementing conservation actions to benefit Species of Greatest Conservation Need (SGCN). Implementation may also be subject to changing conditions and regulatory review (where required) and should be conducted in cooperation with land

managers, land owners and key stakeholders. Large scale conservation efforts (e.g., broad scale monitoring) should be coordinated through VFWD, interagency workgroups and formal agreements where applicable.

Coordination and Collaboration

As noted throughout this report, coordination of efforts is vital to leveraging available resources to ensure maximum wildlife benefit. VFWD will take the lead in facilitating communications among conservation partners, including local, state and federal agencies.

Coordination between the 13 northeast states occurs through the Northeast Association of Fish & Wildlife Agencies (NEAFWA) and the regional office of the USFWS Division of Federal Assistance. NEAFWA established the [Regional Conservation Needs \(RCN\)](#) Program in 2008 to formalize a cooperative approach Action Plan implementation across multiple states. The purpose of the RCN program is to develop, coordinate, and implement conservation actions to address issues, threats, and opportunities that are most effectively tackled at a regional scale.

National coordination will be spearheaded by the Association of Fish & Wildlife Agencies and the US Fish & Wildlife Service.

Prioritizing Conservation Need

During the identification and assessment of SGCN our Action Plan technical teams began the process to prioritize conservation need through the following actions: SGCN were assigned either medium or high priority status (low priority species are deemed relatively secure for now, see Action Plan development), conservation actions, research and monitoring needs and habitat problems were similarly ranked.

We did not prioritize needs and actions beyond this. The Action Plan is a conservation guide for the state—not only VFWD or the Agency of Natural Resources. It is meant to provide guidance to organizations, agencies and individuals who wish to conserve wildlife. The goals and missions of the many and varied partners involved in the project span a broad spectrum of wildlife interests, skills and reach (some are very local, others are state, regional and federal entities). It was clear that there would be no prioritization that would satisfy all partners and that conservation need is so great that there is room for everyone to select the species and habitats they find most important and implement the strategies they are most capable of working on.

When it comes to allocating SWG funds to specific projects, further prioritization is required. Prioritization will take into account the goal of the SWG program—to keep wildlife populations from declining to the point that they require protection under the federal Endangered Species Act (ESA)—and Congressional intent—that SWG funds benefit wildlife that have not historically been the primary beneficiaries of the Federal Aid in Wildlife Restoration Act, Federal Aid in Sport Fish Restoration Program or the federal ESA. Prioritization will also be based on the impact of problems to SGCN and habitats, the project's ability to affect positive change, other conservation and social impacts and the availability of matching funds.

Action Plan Review

Element number six of the eight required elements for an Action Plan (see Chapter 1: Congressional Guidelines) requires that states provide “descriptions of procedures to review the strategy at intervals not to exceed ten years.”

Vermont will update its Action Plan on a 10-year cycle. Ten years will allow for planning, and implementation of actions and for detecting responses in at least some SGCN populations. Vermont’s adaptive management approach to Action Plan implementation, however, means that species and habitat monitoring, formal project reporting and financial tracking will be ongoing and will provide a constant flow of information during the intervening years. Managers, wildlife planners and biologists will use this data to hone strategies, fine tune operations and make mid-course corrections within each ten year cycle. Review activities will include:

- Twice yearly expenditure tracking for individual projects by SWG project managers.
- Annual financial reporting of all in-kind match for individual projects by SWG project managers.
- Full project reports due within 90 days of completion of individual SWG projects by SWG project managers.
- Providing regular Federal Assistance reports to the US Fish & Wildlife Service Division of Federal Assistance.

The process to review and update the Action Plan in 2025 should begin at least two years prior to the deadline. The current thinking is that the review process should mirror the current Action Plan revision process to update each of the eight elements from the original congressional guidelines as follows:

1. Revise the list of SGCN and update information on the distribution and abundance of SGCN. Which species can be removed from the list, which should be added?
2. Update information on the location and condition of key habitats. Describe key habitats of any new SGCN.
3. Describe threats and problems impacting SGCN and their habitats. Update research needs.
4. Review the success of conservation actions implemented to date. Identify conservation actions to conserve SGCN and their habitats.
5. Review Action Plan monitoring efforts to date. Describe plans to monitor species, habitats and conservation actions for the future.
6. Update and describe the process for the next plan review.
7. Review coordination efforts to date. Update plans to coordinate with other plans and planning entities.
8. Revise and describe plans to include the public in the design and implementation of the next Action Plan report.

Literature Cited

- Anderson, M.G. and A. Olivero Sheldon. 2011. [Conservation Status of Fish, Wildlife, and Natural Habitats in the Northeast Landscape: Implementation of the Northeast Monitoring Framework](#). The Nature Conservancy, Eastern Conservation Science
- Association of Fish and Wildlife Agencies (AFWA). 2011. [Measuring the Effectiveness of State Wildlife Grants Final Report](#). Washington, DC.
- Association of Fish and Wildlife Agencies (AFWA), Teaming With Wildlife Committee, State Wildlife Action Plan Best Practices Working Group. 2012. Best Practices for State Wildlife Action Plans—Voluntary Guidance to States for Revision and Implementation. Washington, DC: Association of Fish and Wildlife Agencies. <http://www.fishwildlife.org/files/SWAPBestPractices.pdf>
- Greeley, J. R. 1930. Fishes of the Champlain watershed, pages 44-87, in: A biological survey of the Champlain watershed. Supplemental to Nineteenth Annual Report, 1929. New York State Conservation Department, Albany.
- Langdon, R.W., M.T. Ferguson, and K.M. Cox. 2006. [Fishes of Vermont](#). Vermont Department of Fish and Wildlife, Waterbury, Vermont.
<http://www.vtfishandwildlife.com/cms/One.aspx?portalId=73163&pageId=216868>
- NEAFWA 2008. Monitoring the Conservation of Fish and Wildlife in the Northeast: A Report on the Monitoring and Performance Reporting Framework for the Northeast Association of Fish and Wildlife Agencies” <http://rcngrants.org/content/regional-monitoring-and-performance-framework>.
- Northeast Coordinated Bird Monitoring Partnership. 2007. A Framework for Coordinated Bird Monitoring in the Northeast. Northeast Coordinated Bird Monitoring Partnership Report.
<http://www.nebirdmonitor.org/framework>
- Noss, R. and A.Y. Cooperrider. 1994. Saving Nature’s Legacy: Protecting and Restoring Biodiversity. Island Press. Washington DC
- Salafsky, N, R. Margoluis, and K. Redford. 2001. Adaptive Management: A Tool for Conservation Practitioners. Biodiversity Support Program. www.fosonline.org/Adaptive_Management1.cfm

**Vermont's
Wildlife Action Plan**

Chapter 8

Glossary & Acronym Key

Draft • September 2015

Wildlife Action Plan Glossary.....	1
Acronym Key.....	4

Chapter 8: Glossary & Acronym Key

This glossary contains definitions to many of the terms used in this document.

Actively managed: For wildlife this means that a management plan for the species or a suite of species exists. (E.g. an osprey plan, waterfowl plan, spruce grouse plan.)

Anthropogenic: Conditions that result from human activities. “Anthropo-” meaning human and “-genic” meaning produced from.

Comprehensive Wildlife Conservation Strategy (CWCS): This Wildlife Action Plan was developed under the working title of Comprehensive Wildlife Conservation Strategy (CWCS). Upon federal approval the name was changed to Wildlife Action Plan (Action Plan). The Action Plan/CWCS sets a plan of action for conserving Vermont's wildlife by addressing conservation issues, management needs, and priorities. It is intended to be used by anyone with an interest in wildlife conservation.

Conservation: Plans and actions that will help restore and/or sustain Vermont's wildlife populations, with a focus on Species of Greatest Conservation Need (SGCN), and utilizing the full array of traditional conservation tools such as management (e.g. habitat manipulation, restoration (e.g. acquisition, fee-simple easements), landowner education and incentives.

Conservation Opportunity Areas: areas of land and water where the likelihood of successful conservation is strongest and the conservation needs of wildlife and their habitats would be best met.

Conservation Partners: The wildlife biologists, ecologists, sportsmen and other conservationists, non-governmental organizations, business leaders, colleges and universities and state and federal agencies representing more than 60 entities (table 1-1) that worked with the Vermont Fish & Wildlife Department to create Vermont's Comprehensive Wildlife Conservation Strategy. When implementation of the Action Plan begins, any and all individuals, organizations, agencies and other entities wishing to participate will be considered conservation partners.

Conservation Reserve Program (CRP): provides annual land rental payments up to 15 years and cost sharing assistance to install water quality enhancement practices on environmentally sensitive land.

Conservation Reserve Enhancement Program (CREP): State and Federal partnership allowing incentive payments to landowners who set aside environmentally sensitive land along streams or field boundaries.

Common Species: "Keeping Common Species Common" is a phrase Congress used to describe its goal for the SWG program and the Action Plan. Common in this situation refers to any species that is not on the federal Endangered Species List (Threaten or Endangered).

Contiguous Forest: An area of forested land with either no roads or low densities of class IV roads, and little or no human development (buildings, parking areas, lawns, gravel pits). Contiguous forest may have various age classes of forest cover and include other habitat types such as wetlands or grasslands that are part of the overall contiguous habitat complex.

Corridor: A route that permits the direct travel or spread of animals or plants from one area or region to another, either by the gradual spread of a species' population along the route or by the movement of individual animals, seeds, pollen, spores, or microbes.

Cultural Habitat: (sometimes referred to as anthropogenic habitat) communities and sites that are either created and/or maintained by human activities or are modified by human influence to such a degree that the physical condition is substantially different from what existed prior to human influence (e.g. old mines, hayfields used by grassland birds, buildings and structures used by bats).

Data Gap: A clear data need identified in the Action Plan as important to the conservation of a species or habitat.

Ecosystem: A complex array of organisms, their natural environment, the interactions between them, and the ecological processes that sustain the system. Ecosystems can be defined at any scale, from rotting logs, to Lake Champlain, to the Green Mountains.

Endangered Species: A species in danger of becoming extinct that is protected by either the federal Endangered Species Act or the Vermont Endangered Species Act.

Endemic species Found only in a certain place. For the purposes of this document endemic refers to species found *only* in Vermont. There are no known endemic species in Vermont. The most likely possibilities are invertebrates.

Exotic Invasive & Pest Species: An invasive species is defined by the as a species that is 1) non-native (or alien) to the ecosystem under consideration and 2) whose introduction causes or is likely to cause economic or environmental harm or harm to human health.

Environmental Quality Incentives Program (EQIP): provides cost sharing payments to participants who install enduring conservation practices to help control soil erosion and improve water quality.

Forest Cover Type: A descriptive classification of forestland based on present occupancy of an area by tree species (Society of American Foresters).

Game Species: Wildlife species that are subject to legal hunting, fishing or harvesting.

Habitat: A place where a plant or animal lives. A place where an organism lives. Habitat is generally thought of in terms of single species such as bear or calypso orchid habitat.

Herp: an abbreviation for herpetile, which includes both amphibian and reptile species.

Herptile: amphibian and reptile species

Hyporheic Zone: is the region beneath and adjacent to streams and rivers where surface and ground water mix. The hyporheic zone: links aquatic and terrestrial systems; serves as transition areas between surface water and groundwater systems; and, can contain species common to both surface and subsurface waters.

Indicator species: A species, or community whose presence in an area indicates the presence of certain environmental conditions.

Indicators: Indicators are measures that track inputs, outputs, and outcomes by stating them in specific and observable terms. They are also used to monitor natural resource conditions and the threats that can degrade natural ecosystems (e.g. the number lakes infested with Eurasian watermilfoil; the distribution of lakes infested with Eurasian watermilfoil # of boat checks conducted; % of boaters aware of exotic species laws)

Landscape: A heterogeneous area of land containing groups of natural communities and clusters of interacting ecosystems. These can be of widely varying scales, but normally include a range of elevations, bedrock, and soils.

Life-history traits: Examples include be species with low fecundity, that take a long time to reach sexual maturity, that take a long time between reproductive events (sturgeon, wood turtle)

Limiting factor: The factor limiting the growth, abundance, or distribution of a population of organisms or a habitat.

Metadata: Definitional information that provides information about or documentation of other data

Metapopulation: A small number of relatively isolated populations that may occasionally exchange individuals

Mosaic: A pattern of vegetation in which two or more different plant communities are interspersed in patches.

Natural Community: An interacting assemblage of plants and animals, their physical environment, and the natural processes that affect them.

Neotropical Migrants: Birds especially songbirds, that summer and breed in North America but migrate to the tropics for the winter. Neotropical refers to the region south of the Tropic of Cancer that includes southern Mexico, Central and South America, and the West Indies

Nongame Wildlife: Wildlife species that are not subject to legal hunting, fishing or harvesting.

Pathogen: Any disease producing microorganism or material

Problem: A force causing a negative impact at the species, population, habitat and landscape levels (e.g., habitat conversion, pollution, illegal pet trade). A problem can also be the lack of information or a data gap vital to the successful management of a species. Because this report addresses an extremely broad range of problems affecting species and their habitats, the term "problem" may not always be the most appropriate term: threat, stress, stressor, issue, concern and limiting factor may at times be more accurate.

Regulated Hunting/Fishing/Trapping: The harvest of wildlife under regulations stipulating setting of seasons, time frame of lawful harvest, open and closed zones, methods of take, bag limits, possession limits, and reporting or tagging of species.

Responsibility Species: Species for which Vermont has a long-term stewardship responsibility because they are not doing well regionally, even if populations are stable in Vermont. E.g. bobolink

SGCN: see Species of Greatest Conservation Need

Species of Greatest Conservation Need (SGCN): According to federal legislation and guidance from the USFWS on the development of Comprehensive Wildlife Conservation Strategies, "each State will determine these species in the context of developing its [Wildlife Conservation Strategy]. These species must be fauna, and not flora, and may include aquatic species and invertebrates. A State's list of "species of the greatest conservation need" may include currently listed Federal and State wildlife species and other species of concern. We anticipate that the composition of this list will change over time as the status and conservation need of species changes within the State." The term Species of Greatest Conservation Need is not a statutory designation similar to the terms "endangered" or "threatened" codified by federal and state Endangered Species Acts.

Take/Taking: "Take" and "Taking" mean pursuing, shooting, hunting, killing, capturing, trapping, disturbing, harrying, worrying, or wounding snaring and netting fish, birds and quadrupeds and all lesser acts including placing, setting, drawing or using any net or other device commonly used to take fish or wild animals, whether they result in taking or not. It includes every attempt to take and every act of assistance to another person in taking or attempting to take fish or wild animals.

Threatened Species: Any species which is likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range that is protected by either the federal Endangered Species Act or the Vermont Endangered Species Act

Wildlife: According to State Wildlife Grants legislation, wildlife is any species of wild, free-ranging fauna including fish, and invertebrates and also fauna in captive breeding programs the object of which is to reintroduce individuals of a depleted indigenous species in a previously occupied range.

Wildlife Action Plan (Action Plan) The Action Plan sets a plan of action for conserving Vermont's wildlife by addressing conservation issues, management needs, and priorities. It is intended to be used by anyone with an interest in wildlife conservation. It was developed under the working title of Comprehensive Wildlife Conservation Strategy (CWCS).

Acronym Key

This key includes many of the acronyms used in this document. Please let the authors know if additional entries are warranted.

AFS: American Fisheries Society
AFWA: Association of Fish and Wildlife Agencies
AMP: Acceptable Management Practice
ANR: Vermont Agency of Natural Resources (comprised of VFWD, DEC, FPR)
AOT: Vermont Agency of Transportation
AVCC: Association of Vermont Conservation Commissions
BBS: Breeding Bird Survey
BCR: Bird Conservation Region
BMP: Best management practice
CBC: Christmas Bird Count
CITES: Convention on International Trade in Endangered Species of Flora and Fauna
Cons Comms: Conservation Committees of towns
CRASC: Connecticut River Atlantic Salmon Commission
CRJC: Connecticut River Joint Commission
CRP: Conservation Reserve Program (a program of FSA)
CWCS: Comprehensive Wildlife Conservation Strategy, the original name for the Wildlife Action Plan.
DEC: Vermont Department of Environmental Conservation, also VDEC
DHCD: Vermont Department of Housing & Community Development
DJ: Dingell-Johnson Act of 1950, Federal Aid in Sport Fish Restoration Act
EO: Element Occurrence
EQUIP: Environmental Quality Incentives Program (a program of NRCS)
ESC: Endangered Species Committee of the Agency of Natural Resources
FIA: Forest Inventory Analysis
FIP: Forest Incentives Program (USFS)
FPR: Vermont Department of Forests, Parks & Recreation, also VFPR
FSA: Farm Service Agency (a USDA agency) (www.fsa.usda.gov/vt/)
FWD: Vermont Fish & Wildlife Department
GIS: Geographic Information System
GRP: Grassland Reserve Program
HAT: Hunters, Anglers & Trapper of Vermont
Herp Atlas: Vermont Reptile & Amphibian Atlas
IBA: Important Bird Areas
LCBP: Lake Champlain Basin Program
LCC: Lake Champlain Committee
LCI: Lake Champlain International
LCLT: Lake Champlain Land Trust
MBTA: Migratory Bird Treaty Act of 1940
NABCI: North American Bird Conservation Initiative
NALCC: North Atlantic Landscape Conservation Cooperative
NEPARC: Northeast Partners in Amphibian and Reptile Conservation
NEPCoP: New England Plant Conservation Program
Nongame Fund: Vermont Nongame Wildlife Fund

NHFGD: New Hampshire Fish & Game Department
NHI: Natural Heritage Inventory of the Vermont Fish & Wildlife Department
NRCS: U.S. Natural Resource Conservation Service (habitat programs include EQIP, CRP)
NWI: National Wetlands Inventory
NWR: National Wildlife Refuge
NWTF: National Wild Turkey Federation
OS: The Orianne Society
PARC: Partners in Amphibian and Reptile Conservation
PIF: Partners in Flight
PR: Pittman-Robertson Act of 1937, the Federal Aid to Wildlife Restoration Act
RGS: Ruffed Grouse Society
RPC: Regional Planning Commissions
SAF: Society of American Foresters
SAG: Scientific Advisory Group (advises the Agency of Natural Resources' Endangered Species Committee)
SCI: Staying Connected Initiative
SGCN: Species of Greatest Conservation Need
SWG: State Wildlife Grants program
TNC: The Nature Conservancy
TU: Trout Unlimited
USACE: United States Army Corps of Engineers
USDA: United States Department of Agriculture
USEPA: United States Environmental Protection Agency
USFS: United States Forest Service
USFWS: United States Fish & Wildlife Service
USGS: United States Geological Service
VASA: Vermont All-Terrain Vehicle Sportsman's Association
VCGI: Vermont Center for Geographic Information
VDHCD: Vermont Department of Housing & Community Development
VCE: Vermont Center for Ecostudies
VFWD: Vermont Fish & Wildlife Department
VFPR: Vermont Forests, Parks & Recreation Department
VHCB: Vermont Housing & Community Board
VINS: Vermont Institute of Natural Sciences
VLCT: Vermont League of Cities & Towns
VLT: Vermont Land Trust
VNRC: Vermont Natural Resources Council
VT Coop: Vermont Cooperative Fish and Wildlife Research Unit (of the US Geological Service)
VTa: Vermont Trappers Association
VTFSC: Vermont Federation of Sportsmen's Clubs
VWA: Vermont Woodlands Association
VTrans: Vermont Agency of Transportation (also AOT)
WCS: Wildlife Conservation Society
WHIP: Wildlife Habitat Improvement Program (a program of NRCS)
WMA: Wildlife Management Area (managed by VFWD)
WDP: Wildlife Diversity Program of the Vermont Fish & Wildlife Department (previously the Nongame & Natural Heritage Program)
WRP: Wetland Reserve Program (NRCS)

Vermont's Wildlife Action Plan

Chapter 9

Revising Vermont's Wildlife Action Plan

Draft • September 2015

Developing Vermont's Wildlife Action Plan.....	2
Threats, Problems and Species of Greatest Conservation Need	4
Defining Threats and Problems.....	4
Threat Categories	5
Conservation Action Development	6
Outreach and Public Involvement	7
Species & Habitat Conservation.....	8
Identifying Species of Greatest Conservation Need.....	8
Conservation of Species of Greatest Conservation Need	12
Literature Cited	13

Developing Vermont's Wildlife Action Plan

The revision Vermont's Wildlife Action Plan began in January 2013 when a Revision Team of Vermont Fish & Wildlife Department staff met to begin project scoping. Federal guidelines, planning literature and past planning efforts were reviewed and an organizational structure and revision process were subsequently developed. The identification of Species of Greatest Conservation Need (SGCN) occurred from July 2014 through January 2015. Habitat delineation for SGCN, problem assessment and strategy development occurred from October 2014 through June 2015. Integration and conservation planning ran from May through August 2015. Review and additional input by the Department, agencies and other stakeholders and the general public, will occur between September and November 2015. Final document preparation and editing will occur in December 2015. The anticipated submission date of the Action Plan is December 31, 2015.

The Planning Team reaffirmed five primary goals used to guide its first Wildlife Action Plan as the revision's guiding framework, and added two additional goals:

1. Conserve, enhance and restore Vermont's wildlife and wildlife habitat.
2. Represent good science and conservation planning.
3. Identify conservation priorities yet remain flexible and open to new opportunities.
4. Develop the Action Plan for the entire state; one that all agencies, organizations and individuals can find useful.
5. Build and support advocates for wildlife conservation.
6. Build on the good work of the first Wildlife Action Plan.
7. Develop the Action Plan in a manner that will support regional roll-up of Wildlife Action Plan information among member states of the Northeast Association of Fish & Wildlife Agencies per the Northeast Lexicon (Crisfield 2013) for improved regional conservation.

The Planning Team recognized that meeting these goals required the resources, participation and ingenuity of many conservation-minded individuals, organizations and agencies. This in turn required a development process that included conservation partners to the greatest extent possible. Six teams of taxonomic experts (Species Teams) and a Landscape Team and were created to develop the Wildlife Action Plan. Team members are listed in table 9.1.

Species Teams: (selected Fish and Wildlife staff and other taxonomic experts). Six Species Teams were created: Amphibian & Reptile (Herps), Bird, Fish, Invertebrate, Mammal, and Plant. These teams developed and refined lists of Species of Greatest Conservation Need; assessed species distribution and abundance, identified habitats, communities, threats and actions; developed monitoring and performance measures.

Landscape Team: (selected Fish and Wildlife staff and conservation partners with expertise in GIS, landscape assessment and conservation design). The Landscape Team was charged with developing a landscape-level conservation design for the state, one that would address the needs of most, if not all, Species of Greatest Conservation Need.

Table 9.1: Team and Committee Members, Wildlife Action Plan Revision

*Denotes team/committee chairpersons

Vermont Action Plan Revision Team		Bird Team	
Steve Parren*	VT Fish & Wildlife Dept	John Buck*	VT Fish & Wildlife Dept
Ken Cox	VT Fish & Wildlife Dept	Dr. William Barnard	Norwich University
Steve Gomez	VT Fish & Wildlife Dept	Chip Darmstadt	North Branch Nature Center
Jon Kart	VT Fish & Wildlife Dept	Margaret Fowle	Audubon VT
Eric Sorenson	VT Fish & Wildlife Dept	John Gobeille	VT Fish & Wildlife Dept.
Susan Warner	VT Fish & Wildlife Dept	Mark Labarr	Audubon VT
Lael Will	VT Fish & Wildlife Dept	Sally Laughlin	First VT Bird Atlas
		Dr. Rosalind Renfrew	VT Center for Ecostudies
Planning Team		David Sausville	VT Fish & Wildlife Dept
Steve Parren*	VT Fish & Wildlife Dept	Dr. Allan Strong	University of Vermont
Jon Kart	VT Fish & Wildlife Dept	Erin Talmadge	Birds of VT Museum
Christopher Hilke	National Wildlife Federation		
Landscape Steering Committee		Fish Team	
Eric Sorenson*	VT Fish & Wildlife Dept	Kenneth Cox*	VT Fish & Wildlife Dept
Jens Hilke*	VT Fish & Wildlife Dept	Dr. William Barnard	Norwich University
Bob Zaino*	VT Fish & Wildlife Dept	Dr. Douglas Facey	Saint Michael's College
Liz Thompson	Vermont Land Trust	Mark Ferguson	VT Fish & Wildlife Dept
John Austin	VT Fish & Wildlife Dept	Eric Howe	Lake Champlain Basin Program
Jayson Benoit	NorthWoods Stewardship Ctr	Richard Langdon	VT Dept of Environmental Conservation
Jeff Briggs	VT Forest Parks & Recreation Dept	Invertebrate Team	
Dan Farrell	The Nature Conservancy	Mark Ferguson*	VT Fish & Wildlife Dept
Jon Kart	VT Fish & Wildlife Dept	Steve Fiske	VT Dept of Environmental Conservation
Jane Lazorchak	VT Fish & Wildlife Dept	Trish Hanson	VT Forest Parks & Recreation Dept
Paul Marangelo	The Nature Conservancy	Kent McFarland	VT Center for Ecostudies
Doug Morin	VT Fish & Wildlife Dept	Bryan Pfeiffer	Consulting Entomologist
Steve Parren	VT Fish & Wildlife Dept		
Nancy Patch	VT Forest Parks & Recreation Dept	Mammal Team	
Rose Paul	The Nature Conservancy	Chris Bernier*	VT Fish & Wildlife Dept
Kim Royar	VT Fish & Wildlife Dept	Alyssa Bennett	VT Fish & Wildlife Dept
Mark Scott	VT Fish & Wildlife Dept	Dr. William Kilpatrick	University of Vermont
		Dr. James Murdoch	University of Vermont
Amphibian & Reptile Team		Dr. Peter Smith	Green Mountain College
Doug Blodgett*	VT Fish & Wildlife Dept	Christopher Spatz	Cougar Rewilding Foundation/NE Wolf Coalition
Jim Andrews	Middlebury College		
Steve Faccio	VT Center for Ecostudies	Plant Team	
Chris Slesar	VT Agency of Transportation	Bob Popp*	VT Fish & Wildlife Dept
		Everett Marshall*	VT Fish & Wildlife Dept
		Charlie Hohn	VT Fish & Wildlife Dept.
		Aaron Marcus	VT Fish & Wildlife Dept
		Eric Sorenson	VT Fish & Wildlife Dept
		Bob Zaino	VT Fish & Wildlife Dept

Threats, Problems and Species of Greatest Conservation Need

Defining Threats and Problems

Element number three of the eight congressionally required elements of a Wildlife Action Plan requires that states: describe the problems that may adversely affect Species of Greatest Conservation Need or their habitats and priority research and survey efforts needed to identify factors which may assist in restoration and improved conservation of these species and habitats. Problem and threats are defined as follows:

Problem: Something that is a concern and could cause a negative impact at the species, population, habitat and/or landscape levels (e.g., habitat conversion, pollution, illegal pet trade). A problem can also be the lack of information or a data gap vital to the successful management of a species.

Threat (direct): Processes or human activities “that have caused, are causing, or may cause the destruction, degradation, and/or impairment of biodiversity targets” (adapted from Salafsky et al. 2008).

Threat (indirect): The factors contributing to or enabling direct threats. Typically there is a chain of contributing factors behind any given direct threat. Synonyms include contributing factors, underlying factors, drivers, and root causes (adapted from Salafsky et al. 2008).

For the purposes of this report, problem and threat are used in a similar or related manner. For each Species of Greatest Conservation Need in the Action Plan we identified priority problems. Priority research needed to evaluate other potential problems was also identified. They are detailed in SGCN conservation summaries (Appendix A) and in habitat/community summaries (Appendix B).

Each of the threats and problems identified in the Action Plan was assigned to one of 24 categories roughly grouped into habitat-related factors and non-habitat-related factors. These categories make it possible to search our database for similar factors impacting other species. It also makes it easier to roll-up for broad scale conservation planning. The categories were cross-walked (Appendix G) with those developed by the International Union for Conservation of Nature and Natural Resources (IUCN) (Salafsky et al. 2008) to aid in the regional roll-up of Action Plan data as recommended by the Diversity Technical Committee of the Northeast Association of Fish & Wildlife Agencies (Crisfield 2013).

The categories are not mutually exclusive and threats can often logically be placed into more than one category depending on the particular stress it causes for a species or habitat. For example, a road can fragment the habitat of grassland nesting birds, cars traveling the road can injure or kill amphibians that were crossing the road to mate in an adjacent pool, and salt spread on the road to prevent icing can wash into a stream impacting its population of Brook Trout. In this example the threats stemming from the road would be recorded in the "Habitat Fragmentation," "Impacts of Roads & Transportation Systems," and "Pollution" categories.

Threats are often species and/or habitat specific. What may negatively impact one species may benefit another. For example, if a cold water stream with a healthy Brook Trout population was dammed it might no longer support Brook Trout. That impact to the dam would be described as the "conversion of habitat" category. However, the reservoir created by the dam might make it more suitable for a warm water fish species.

Clearly life is too complex to be placed into any one box. Therefore it is important to read the full description of a factor affecting a species or habitat in the appropriate species or habitat summary. Definitions can be found in Appendix G.

Threat Categories

See Appendix G for definitions of each category. For context, see Appendix A for SGCN conservation summaries and Appendix B for habitat/community summaries.

Habitat-Related Threat/Problem Categories

- Climate Change
- Habitat Alteration/ Degradation
- Habitat Conversion
- Habitat Fragmentation
- Hydrologic Alteration
- Impacts of Roads & Transportation Systems
- Impacts of Energy Infrastructure & Development
- Inadequate Distribution of Successional Stages
- Inadequate Disturbance Regime
- Invasion by Exotic Species
- Parcelization
- Sedimentation

Non-Habitat-Related Threat/Problem Categories

- Competition
- Disease
- Genetics
- Harvest or Collection
- Incompatible Recreation
- Loss of Food Base or Prey Base
- Loss of Relationship with Other Species
- Parasitism
- Pollution
- Predation or Herbivory
- Reproductive Traits
- Trampling & Direct Impacts

Conservation Action Development

Element number four of the eight congressionally required elements of a Wildlife Action Plan requires that states describe “conservation actions proposed to conserve the identified species and habitats and priorities for implementing such actions.”

We identified actions to address the threats and problems impacting each of Species of Greatest Conservation Need (SGCN) and their habitats. Selected actions are based on the best science available today as well as a strategic assessment of needs and priorities of all wildlife species. In the coming years, as monitoring data on SGCN and conservation actions becomes available, as priorities change, or new threats or opportunities arise, actions may need to be revisited. Not every action in this report will be eligible for State Wildlife Grant funding. Furthermore, it may not be suitable, or feasible, for the Vermont Fish & Wildlife Department to implement some of the actions in this report, however, some conservation partners may find them fitting and practical.

Actions are described in short narratives in each SGCN conservation summary and in each habitat, community and landscape summary. Actions are intentionally broad, directional, and nonspecific so as not to constrain our selection of procedures for implementing them. For example, an action such as “provide technical assistance to landowners to maintain or improve riparian habitat for Species of Greatest Conservation Need” allows for different approaches to providing that assistance and leaves the door open to a variety of providers to implement. Where action implementation is to be funded by the State Wildlife Grant program the approach should be consistent with the Department’s mission and strategic plan, and precise procedures will be detailed in operational plans once the Action Plan is finalized.

Vermont’s Action Plan was designed for the state, not just the Fish & Wildlife Department. While the VFWD may be responsible for implementing many of the actions in this report, it could be conservation partners that are the more logical and appropriate leaders for others, due to their skills and expertise, staffing, history, location, available resources and constituencies.

Each of the actions identified in this report were assigned to one of 27 categories in six major classes. The categories were developed by the Conservation Measures Partnership (Salafsky 2005) as a means of standardizing terminology (not practices) among conservation practitioners worldwide. Many states have used these same categories to organize the strategies and actions in their Action Plan. They have also been incorporated into Wildlife TRACS (Tracking and Reporting Actions for the Conservation of Species) the US Fish & Wildlife Services’ system for tracking and reporting conservation activities. States, including Vermont, will use TRACS for all work funded through the USFWS once it is fully operational.

The action categories are used solely for the purpose of organizing and grouping strategies developed by Action Plan teams and committees. It was not our goal to create strategies for every category. A few categories were not applicable to the species or habitats in Vermont whereas others were deemed not as effective. Definitions for each strategy can be found in Appendix G.

Outreach and Public Involvement

The Vermont Fish & Wildlife Department recognized that to meet our Action Plan revision goals that we needed the resources, participation and ingenuity of our conservation partners. More than 30 partners representing 20 different organizations and agencies participated on the landscape team or one of the taxonomic teams.

Additional outreach and public involvement efforts focused on the following groups:

This section is incomplete. It will be completed after the draft Action Plan is reviewed by conservation partners and the general public.

Species & Habitat Conservation

Identifying Species of Greatest Conservation Need

Congress created the State Wildlife Grants program (SWG) in 2001 with the goal of preventing wildlife populations from declining to the point of requiring Endangered Species Act protections. To receive SWG funds, state and tribal fish and wildlife agencies agreed to develop statewide Wildlife Action Plans. Congress directed that the Action Plan identify and be focused on the "Species of Greatest Conservation Need."

Congress left it up to each state to identify their Species of Greatest Conservation Need (SGCN). The State Wildlife Grants program defines wildlife as "any species of wild, free-ranging fauna including aquatic species and invertebrates as well as native fauna in captive breeding programs intended for reintroduction within its previously occupied range." Furthermore, it was Congress' intent that SWG assist wildlife that "have not previously benefited from other federal wildlife conservation and management programs" (e.g., Federal Aid to Wildlife Restoration Act, Federal Aid in Sport Fish Restoration Act, or the Endangered Species Act). In Vermont, SGCN include:

- Species with declining populations;
- Species threatened or potentially threatened; and,
- Species that are so little known in the state that experts cannot yet ascertain status.

Though plants are not eligible for State Wildlife Grants Program funding, Vermont's Action Plan does include plant SGCN. Plant-specific conservation strategies, if and when they are implemented, will be funded through mechanisms other than SWG. Several game and sportfish species are identified here as SGCN. Other established funding programs for the conservation of these species may be used before using SWG.

Vermont began its process of identifying Species of Greatest Conservation Need (SGCN) with a systematic review of all its known wildlife. The review considered both the well-known wildlife species supported by large datasets and poorly understood species.

The six Species Teams (Amphibian & Reptile, Bird, Fish, Invertebrate, Mammal and Plant) conducted the reviews and selected SGCN using the review criteria in table 9.2. They were provided lists of species found in Vermont within their respective taxa (the Invertebrate team received the most up-to-date invertebrate list available, but it is widely accepted that a complete list of the estimated 21,000 invertebrates in Vermont may never be possible. The lists and supporting information were developed by the VFWD's Wildlife Diversity Program using its Natural Heritage Database and augmented with other databases, records and information from NatureServe, universities and research facilities, regional and national monitoring efforts, published literature and the knowledge of technical experts. The following groups had major, taxon-wide State rarity rank reviews: Amphibians & Reptiles (2007), Bumble Bees (2014), Birds (2010), Fishes (2005), Bats (2011), Other small mammals (2008), moths and butterflies (2010), dragonflies and damselflies (2008) and Vascular Plants (2014). Ranks for individual species were updated as needed.

Table 9.2: Review Criteria for Identifying Species of Greatest Conservation Need

Category	Criterion	Allowed Response	Definition/example
Species that are rare or declining	State and/or Federally listed Threatened or Endangered species	Endangered, Threatened, Special Concern [See Appendix I for definitions of T& E status and ranks]	E: Endangered: in immediate danger of becoming extirpated in the state T: Threatened: with high possibility of becoming endangered in the near future. SC: Special Concern: rare; status should be watched
	Rare and very rare species	S-Ranks S1,S2 [See appendix I for definitions of T& E status and ranks]	S1: Critically imperiled (very rare): At very high risk of extinction or extirpation due to extreme rarity (often 5 or fewer populations), very steep declines, or other factors. S2: Imperiled (rare): At high risk of extinction or extirpation due to very restricted range, very few populations (often 20 or fewer), steep declines, or other factors
	State Trend	Stable, Fluctuating, Declining, Increasing, Unknown	Based on research data such as BBS routes, other monitoring and best judgment of experts
	Regionally Rare	Yes/No/ Unknown	Based on regional and national research, BBS routes, other monitoring and consensus within technical teams.
	Extirpated in Vermont	Yes/No/ Unknown	
Vulnerable species at risk due to any of the following	Habitat Loss/Conversion/fragmentation	Yes-development, Yes-succession, Yes-natural causes, No, Unknown	Species negatively affected by habitat conversion, degradation, fragmentation or succession
	Life-history traits making the species vulnerable	Yes/No/ Unknown	Species with low fecundity, that take a long time to reach sexual maturity, that take a long time between reproductive events (e.g., sturgeon, wood turtle)
	Species vulnerable to taking	Yes-Regulated, Yes-Unregulated, No, Unknown	Hunting, trapping or collection, legal or otherwise.
	Species vulnerable to other deadly contact with humans	Yes/No/ Unknown	Road kill (bobcat, turtles), wind turbines (birds, bats) contaminates (fish) etc.
	Species w/ limited, localized at-risk populations	Yes/No/ Unknown	Populations that cannot or do not intermix with the meta-population. E.g., non-vagile invertebrates in a sandplain community and perhaps spruce grouse.
	Species significantly impacted by exotics	Yes/No/ Unknown	Impact may lead to elimination of populations, limits to long-term stability, extirpation
Species or species groups w/ unknown status or taxonomy	Unknown status-more data is needed	Yes/No/ Unknown	
	Species w/ taxonomic uncertainties	Yes/No/ Unknown	

Category	Criterion	Allowed Response	Definition/example
Other factors to consider	Keystone species	Yes/No/ Unknown	Species with a disproportionately strong influence on ecosystem functioning and diversity (Power et al.1996).
	Responsibility species	Yes/No/ Unknown	Species for which Vermont has a long-term stewardship responsibility because they are not doing well regionally, even if populations are stable in Vermont (e.g., Bobolink)
	Endemic species	Yes/No/ Unknown	Species found only in Vermont
	Relationship to core population	central peripheral, disjunct, unknown	
	Requires rare or specialized habitats	Yes/No/ Unknown	A species with a very narrow niche, e.g., a species requiring a host plant found only in a handful of serpentine rock outcrops.
	Species with limited dispersal capability	Yes/No/ Unknown	Non-vagile species in dispersed habitats.
	Requires key Vermont migration stopover points	Yes/No/ Unknown	
	Species selected based on expert opinion	Yes/No	Combined opinion of the team.
	Actively managed? (if so list applicable plan(s))	Yes-Mgt plan exists, Yes-regulated, No	Does a management plan exist for the species or species group? (E.g., an osprey plan, waterfowl plan, species recovery plan.)
Secure?	Species Secure	Yes/No/ Unknown	Combined opinion of the team
	Final Assessment	High, Medium, Low Priority	

Once the reviews were complete the Species Team selected SGCN using selection criteria found in Table 9.3. Species were assigned conservation priorities of high, medium or low. Species ranked medium and high constitute Vermont's Species of Greatest Conservation Need. Low priority species were considered secure. There were a few cases where a specific Species Team approached their tasks differently:

Bird Team: An unusually rich collection of data and prior conservation planning efforts are available for bird conservation—far more than is available for other taxa, including the second Vermont Breeding Bird Atlas (2013), the USFWS Breeding Bird Surveys and information from Partners-In-Flight, North American Bird Conservation Initiative, National Audubon Society's Watch List, and the American Bird Conservancy's Green List.

Invertebrate Team: It is estimated that Vermont is home to approximately 21,000 invertebrate species (McFarland, pers comm). The vast majority are un-cataloged, unstudied and just plain unknown. Application of the review criteria to invertebrates on a species-by-species basis would be unproductive. Instead the Invertebrate Team interviewed additional experts within Vermont, regionally and nationally to help in the identification of species and Species Groups of Greatest Conservation Need. The team also took advantage of several significant advances made since (and because of) the adoption of Vermont's first Wildlife Action Plan in 2005, including: the Vermont

Butterfly Atlas, a Peatland and Large River Odonate Survey and the Vermont Bumble Bee Survey.

Plant Team: The Plant Team also had to contend with a huge list of species—more than 2000 vascular plants (Flora 1993) and 600 bryophytes (Allard 2004). The team took advantage of plant conservation assessments previously conducted by the Agency of Natural Resources' Endangered Species Committee to create its list of Species of Greatest Conservation Need. All species ranked S1 (critically imperiled) and S2 (imperiled) became SGCN. Those SGCN also on the New England Plant Conservation Program list of regionally rare plants were then ranked High Priority. All others were ranked medium priority.

Table 9.3: Criteria for Selecting Vermont's Species of Greatest Conservation Need

Because the circumstances, issues and problems impacting each species differ, teams were given some flexibility in assigning ranks to species.

Species (and Species Groups) of Greatest Conservation Need	High Priority	Species that are vulnerable (rarity is an aspect of vulnerability).
		Species with immediate limits to its survivability based on known problems and/or known impacts to the population
		Species exhibit negative population trends.
		Species may be extirpated locally (Vermont) but still exist regionally.
	Medium Priority	Species may be well distributed and even locally abundant, but populations are challenged by factors that increase mortality or habitat loss and therefore threaten the species in Vermont.
		Consider what is known about the species regionally.
		Since this may be the most difficult category to assign species to, there should be a consensus among group members.
Common Species	Low Priority	Species is secure for the immediate future.
		Species may be vulnerable to some mortality and/or problems (e.g., habitat degradation) but population is abundant enough to tolerate negative forces

The list of Species of Greatest Conservation Need includes 132 vertebrate species (out of a total of 468), 200 invertebrate species or groups (out of an estimated 21,000) and 645 plant species out of approximately 2000 vascular and non-vascular species. Table 9.4 provides summary statistics.

Table 9.4: Summary Statistics for Vermont's Species of Greatest Conservation Need

High and medium priority-ranked species constitute Vermont's SGCN.

*21,400 is the estimated number of Vermont invertebrates

** This low percentage reflects the large number of invertebrates whose conservation status is unknown

	Total species in VT	High Priority SGCN	Medium Priority SGCN	Total SGCN	% SGCN of total VT Species
Amphibians & Reptiles	42	12	7	19	45%
Birds	269	29	22	51	19%
Fish	94	12	13	27	29%
Invertebrates*	21,400*	141	59	200	0.93%**
Mammals	63	17	18	35	57%
Plants	2000	221	424	645	32%
Total	23,866	432	543	977	2.87%

Conservation of Species of Greatest Conservation Need

Fine Filter-Species

Once Species of Greatest Conservation Need were identified, Taxa Teams developed conservation summaries each SGCN. Reports identified species distribution, habitat needs, problems affecting species and their habitats, research and monitoring needs and conservation strategies for each SGCN (Congressionally required elements #1-#5). Invertebrate SGCN were addressed in groups rather than as individual species. Fifteen invertebrate groups were created based on taxonomy (e.g., Bumble Bees, Crustaceans, Tiger Beetles) and habitat use (e.g., freshwater, grasslands, hardwood forests). Individual conservation summaries were not developed for plant SGCN but a taxon-wide summary is provided in chapter 5. All data was entered into the Action Plan database.

Distribution for all SGCN was identified by biophysical region (Girton & Capen 1997) using terminology consistent with VFWD's element occurrence tracking procedures. Distribution of fish SGCN and some additional aquatic SGCN were also identified by 8-digit watershed unit (NRCS 2009). Historic occurrence was noted in a narrative for some of the rarer and extirpated SGCN.

Habitat descriptions for SGCN include a narrative, elevation preferences, migrant status, home range and patch size requirements and landscape requirements (e.g., corridor needs, habitat mosaics or wetland complexes, preference for managed or passively managed forest, large grasslands or developed landscapes).

Research and monitoring were also identified and prioritized for each animal SGCN.

Priority threats and potential risks to Species of Greatest Conservation Need were enumerated for each species. These were not exhaustive lists of all possible problems. Teams identified only those factors posing significant and potentially significant threats for a species. A narrative description was entered into the database. Species teams also assigned each problem to one of 24 habitat related and non-habitat related problem categories (Appendix G). These categories have been cross-walked with those developed by the International Union for Conservation of Nature and Natural Resources (IUCN) (Salafsky et al. 2008) to aid in the regional roll-up of Action Plan data as recommended by the Diversity Technical Committee of the Northeast Association of Fish & Wildlife Agencies (Crisfield 2013).

Species specific conservation actions were also developed by the Species Teams. Actions were designed to address identified threats. Actions were assigned either a "medium" or "high" priority status (low priority actions are not included in the Action Plan) and each strategy was also assigned to a category (Salafsky 2004) to aid in organizing and review of actions (Appendix G).

Actions were not prioritized beyond this step. As a conservation guide for the state, Vermont's Action Plan is meant to provide guidance to organizations, agencies and individuals wishing to conserve wildlife. The varied goals and missions of the partners involved in the Action Plan span a broad spectrum of wildlife interests, skills and reach (some are local; others are state, regional and federal entities). No prioritization was found to satisfy all partners, however, the conservation need is deemed so great that there is room for

everyone to select the species and habitats they find most important and implement the actions they are most capable of working on.

Coarse Filter-Conservation at Multiple Scales

To aid in the development of community and landscape level conservation actions, each SGCN was assigned to at least one of more than 100 habitat types (natural communities, aquatic habitats, cultural habitats and or landscapes). These habitats were grouped into 24 major categories (Chapter 4. table 4.1) and conservation summaries were developed for each. The summaries include descriptions and general locations; current conditions; desired conditions based on the needs of associated SGCN; prioritized threats and conservation actions, potential conservation partners and funding sources for action implementation; and, a listing of other relevant plans and planning processes.

Threats and problems described in the habitat summaries (and in species summaries) are not comprehensive. Only those problems ranked as medium and high are included in this report. This was a strategic decision to focus attention on those threats and problems determined or perceived to be most important. If additional problem(s) are later identified as significantly impacting a species or habitat it will be incorporated into the Action Plan database during project review and reporting. Actions and actions to address additional problem(s) will also be eligible for SWG funding.

Habitat Classification & Ecological Divisions

"Wetland, Woodland, Wildland - A guide to the natural communities of Vermont" (2000) by Thompson and Sorenson was used as the basis for terrestrial natural communities. Forest cover types (Eyre 1980) and U.S Forest Service Forest Inventory & Analysis types (USDA 2003) were used for early successional and managed forests. "A Classification of the Aquatic Communities of Vermont" by Langdon et. al. (1998) was used as the basis for aquatic habitat designations and Reschke (1990) was adapted for cultural habitats.

SGCN distribution was identified to biophysical region (Girton & Capen 1997) and 8-digit watersheds (NRCS 2003). These landscape units were selected in part because they will integrate well with other conservation efforts within the state and regionally. Biophysical regions can be considered a sub-unit of the Bailey's section (Bailey 1995, Bailey 1998) providing finer grain detail. Data can be integrated into Bailey's sections to aide in regional, national and international conservation efforts.

Literature Cited

- Allard, D. 2004. A Preliminary Recovery Plan for the Bryophytes of Vermont.
- Association of Fish and Wildlife Agencies (AFWA). 2011. Measuring the Effectiveness of State Wildlife Grants Final Report
- Association of Fish and Wildlife Agencies (AFWA), Teaming With Wildlife Committee, State Wildlife Action Plan Best Practices Working Group. 2012. Best Practices for State Wildlife Action Plans—Voluntary Guidance to States for Revision and Implementation. Washington, DC: Association of Fish and Wildlife Agencies. <http://www.fishwildlife.org/files/SWAPBestPractices.pdf>
- Association of Fish & Wildlife Agencies. 2009. Voluntary Guidance for States to Incorporate Climate Change into State Wildlife Action Plans & Other Management Plans. http://www.fishwildlife.org/files/AFWA-Voluntary-Guidance-Incorporating-Climate-Change_SWAP.pdf

- Bailey, R.G. 1995. Description of the Ecoregions of the United States, 2d ed., USDA-Forest Service Miscellaneous Publication 1391
- Bailey, R.G. 1998. Ecoregions Map of North America: Explanatory Note. Prepared in Cooperation with The Nature Conservancy and the US Geological Survey. USDA Forest Service, Miscellaneous Publication Number 1548. Washington, DC.
- Crisfield, E. and the Northeast Fish and Wildlife Diversity Technical Committee (NFWDTTC). 2013. The Northeast Lexicon: Terminology Conventions and Data Framework for State Wildlife Action Plans in the Northeast Region. A report submitted to the Northeast Fish and Wildlife Diversity Technical Committee. Terwilliger Consulting, Inc., Locustville, VA.
- Eyre, F. H. ed. 1980. Forest Cover Types of the United States. Society of American Foresters. Washington, DC.
- Flora of North America Editorial Committee, eds. 1993+. Flora of North America North of Mexico. 7+ vols. New York and Oxford.
- Girton, P. and D. Capen. 1997. A report on biophysical regions in Vermont. Unpublished report prepared for the Vermont EcoMapping roundtable.
- Langdon, R., J. Andrews, K. Cox, S. Fiske, N. Kamman, and S. Warren. 1998. A classification of the aquatic communities of Vermont. The Nature Conservancy and the Vermont Biodiversity Project, Montpelier, Vermont.
- Natural Resources Conservation Service, U.S. Department of Agriculture. 20009. WaterHydro_WBD8VT. VT Center for Geographic Information. Vermont.
- NEAFWA 2008. Monitoring the Conservation of Fish and Wildlife in the Northeast: A Report on the Monitoring and Performance Reporting Framework for the Northeast Association of Fish and Wildlife Agencies” <http://rcngrants.org/content/regional-monitoring-and-performance-framework>.
- Salafsky, N. D. Salzer, A. J. Stattersfield, C. Hilton-Taylor, et al. 2008. [A Standard Lexicon for Biodiversity Conservation: Unified Classifications of Threats and Actions](#). Conservation Biology 22 (4)
- Salafsky, N., D. Salzer, J. Ervin, T. Boucher, and W. Otlie. 2003. Conventions for defining, naming, measuring, combining, and mapping threats to conservation: an initial proposal for a standard system, December 2003 Draft. Bethesda, MD.
- Salafsky N., D. Salzer. 2005. Proposed Taxonomy of Conservation Actions Draft 5. January, 11 2005. Bethesda, MD.
- Thompson, E. H., and E. R. Sorenson. 2000. Wetland, woodland, wildland - A guide to the natural communities of Vermont. University Press of New England, Hanover
- U.S. Department of Agriculture, Forest Service. 2003. Forest inventory and analysis national core field guide, volume 1: field data collection procedures for phase 2 plots, version 1.7. U.S. Department of Agriculture, Forest Service, Forest Inventory and Analysis, 201 14th St., Washington, D.C., 20250

Vermont's Wildlife Action Plan

Amphibian & Reptile SGCN Conservation Reports

Appendix A1

Draft 10/1/2015

Species	Page
Jefferson Salamander	1
Blue-spotted Salamander	8
Spotted Salamander	15
Four-toed Salamander	22
Mudpuppy	29
Fowler's Toad	38
Boreal Chorus Frog	47
Spotted Turtle	54
Wood Turtle	60
Eastern Musk Turtle (Stinkpot)	67
Spiny Softshell Turtle	71
Five-lined Skink	79
North American Racer	85
Eastern Ratsnake	91
Common Watersnake	98
DeKay's Brownsnake	104
Eastern Ribbonsnake	109
Smooth Greensnake	114
Timber Rattlesnake	120

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Jefferson Salamander**
Scientific Name: **Ambystoma jeffersonianum**
Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: S2

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Jefferson Salamander is rare in Vermont (S2, SC), and is considered a species of conservation concern within the region due to evidence of population declines and the fact that a high proportion of the global population occurs within the Northeast (Terres). Jefferson Salamander breeding habitat is limited almost exclusively to temporary woodland pools surrounded by relatively large stands of mature hardwoods. There is evidence that the species may be sensitive to forest fragmentation, and in southern New England, some populations appear to have been outcompeted by its congener, the Blue-spotted Salamander (Klemens, personal communication). In addition, unisexual female hybrid populations exist that introduce uncertainties about the species' taxonomy, population biology, persistence, and long-term viability.

Distribution

The distribution of this rare woodland species is widespread but scattered in Vermont; rare or absent from higher elevations of the Green Mountains.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Probable
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Middle Connecticut
Lake Champlain
Middle Connecticut
West
Upper Connecticut-Mascoma
Black-Ottawquechee
Hudson-Hoosic
Otter Creek
White
Winooski River

Probable Watersheds

Lamoille River
Waits
Deerfield
Passumpsic



Common Name: **Jefferson Salamander**
Scientific Name: **Ambystoma jeffersonianum**
Species Group: **Herp**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Habitat mosaics that include breeding pools embedded in closed canopy forest stands is most critical. Maintaining forested corridors connected to other suitable habitat patches with breeding pools is important for juvenile dispersal and to maintain metapopulation dynamics. Requires well-shaded, relatively mature hardwood/hemlock forest stands with abundant coarse woody debris, leaf litter, and underground refugia (small mammal tunnels, rock crevices, etc.) surrounding temporary woodland (vernal) pools (Faccio 2003). May also use semi-permanent pools. Most commonly found in ridgetop, mid-elevation Northern Hardwood forests in the foothills of the Green Mountains.

Habitat Types:

Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Seeps and Pools
Northern Hardwood
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Sedimentation
Habitat Fragmentation
Impacts of Roads or Transportation Systems
Climate Change

Description of habitat threat(s): Any habitat conversion, alteration, or fragmentation that disrupts species' ability to move between breeding and terrestrial sites, disrupts connectivity between breeding sites/metapopulations (Compton et al 2007), changes water/soil chemistry, temperature, pool hydroperiod, humidity, etc, may have negative effects. Road mortality can have major impacts on migrating adults and dispersing juveniles, especially when located between terrestrial and breeding habitats. Climate change that affects hydroperiod and/or water temperature of breeding pools could have significant impacts on productivity (Rowe and Dunson 1995).

Non-Habitat Threats:

Genetics
Competition
Disease
Pollution

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Jefferson Salamander**
Scientific Name: **Ambystoma jeffersonianum**
Species Group: **Herp**

Reproductive Traits

Trampling or Direct Impacts

Description of non-habitat threat(s): Unisexual, female hybrid populations reproduce via gynogenesis (requires sperm from diploid males which is not incorporated into the genome of embryos; Petranks 1998), resulting in female-biased sex ratios. Since hybrid females require males in the population for successful breeding, but do not produce males to replace those that have been lost, they may have the potential to dilute genetic variability of diploid populations. Competition from the Blue-spotted Salamander, which is more tolerant of disturbed habitats (Klemens 1993), may be a problem in areas where both species are found together (Champlain Valley, southern NE). Two emerging diseases (Ranavirus and Batrachochytrium salamandrivorans (Bs)) could present a threat in the future. Ranavirus was recently shown to be widely distributed in vernal pools in six mid-Atlantic states (Scott A. Smith pers. Comm.), while the fungal disease Bs, which originated in Asia, was recently detected in Europe (Stokstad 2014). Loss of metapopulation structure due to fragmentation of suitable habitats by roads or other non-permeable development is problematic. Jefferson Salamander is more sensitive to acidification of breeding pools compared to other Ambystoma species (Petranks 1998). Complete egg mortality occurs in pools with low pH, and water with a pH <4.5 is often lethal to larvae. Also, widespread treatment of breeding pools to control West Nile Virus would likely have negative effects on amphibians. This is a long-lived amphibian that may not breed every year and produces relatively few eggs per breeding cycle.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Continue to field-verify mapped potential pools and other breeding sites statewide.
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	1) Identify distribution and relative abundance of VT populations 2) Identify significant breeding sites. Large numbers (e.g., >25 egg masses) of breeders or evidence of use by any SGCN. 3) Continue to field-verify mapped potential vernal pools statewide.
Research	Threats and Their Significance	High	1) Conduct sampling to determine if Ranavirus is present in the state, and if so, determine its distribution and which species are affected. 2) Identify sites where road mortality is high annually and evaluate mitigation methods. 3) Identify and evaluate limiting factors.
Research	Population Genetics	Medium	Determine population genetics statewide and monitor populations for changes in sex ratios.
Research	Taxonomy	Medium	
Monitoring	Population Change	High	Implement a statistically robust monitoring protocol to track breeding phenology, population trends, and productivity, as well as changes in breeding site variables (e.g. hydroperiod, water chemistry, etc.).
Monitoring	Habitat Change	High	
Monitoring	Range Shifts	Medium	
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Jefferson Salamander**
 Scientific Name: **Ambystoma jeffersonianum**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	High	Determine presence/absence and distribution of Ranavirus in VT, and which species are most affected.	Number and distribution of sites sampled	VCE, Reptile & Amphibian Atlas, ANR Wetlands Division	SWG, Private Grants
Standards		Timber harvest should be conducted on frozen ground and rutting of ground that could change hydrology of an area or intercept amphibians should be avoided.	Number of operations that did not disturb ground	landowners, FPR, USFS, consulting foresters	State Lands Mgmt, GMNF, EQIP, Current Use
Standards		Maintain 75% mature canopy cover beyond the 30-meter buffer in the terrestrial habitat.	Sites with canopy cover	landowners, consulting foresters	State Lands Mgmt funds, EQIP, Current Use
Standards		Maintain 30-meter water quality buffer around entire perimeter of the pool. Timber harvesting, roads and any ground disturbing activities to be excluded within this buffer.	Number of sites having water quality protected	landowners, consulting foresters	EQIP, Current Use
Compatible Resource Use		Maintain permeable forested habitat matrix between and among breeding populations so that individuals can be exchanged among populations.	Number of pools with forest surrounding	landowners, consulting foresters	EQIP, Current Use
Technical Assistance, Training, Learning Networks		Identify significant road crossings and develop safe road crossings to address roadkill.	Number of sites reported	VTrans, Reptile & Amphibian Atlas, VCE	VTrans, FHWA
Habitat Restoration		When existing road traffic is impacting migration of animals from terrestrial habitat to aquatic pool habitat, identify problem areas and redesign roads with crossing structures when roads are being upgraded.	Number of structures installed	VTrans	VTrans, FHWA
Standards		Maintain habitat mosaic and connectivity between breeding pools. If two large sites are separated beyond the dispersal distance of a species it might be helpful to create or enhance pools that would link the two large sites.	Number of pools within habitat matrix	landowners, consulting foresters	State Lands Mgmt fund, EQIP, Current Use
Compatible Resource Use		Site all permanent roads more than 200 meters from a breeding pool, downslope of the pool if possible.	Number of roads sited to minimize impacts to pools	VTrans, developers	VTrans, development conditions
Habitat Restoration		When feasible restore deciduous or mixed forest surrounding breeding pool.	Number of sites restored	landowners, consulting foresters	Current Use, EQIP

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Jefferson Salamander**
 Scientific Name: **Ambystoma jeffersonianum**
 Species Group: **Herp**

Awareness Raising and Communications		Help people understand the essential needs of all life stages, especially upland habitat in proximity to breeding pool.	Number of people exposed to conservation message	VFWD Outreach Division, media, Reptile & Amphibian Atlas, VCE	marketing funds
Compatible Resource Use		Maintain breeding pools and needed terrestrial habitat--usually directly adjacent to pool perimeter out to 200m but could be the equivalent area along a portion of the perimeter while minimizing edge.	Pools with upland habitat.	landowners, consulting foresters	Current Use, EQIP
Compatible Resource Use		Maintain corridor connections between upland and pool habitat.	Number of sites with upland connections	landowners, consulting foresters	Current Use, EQIP
Habitat Restoration		If loss of important sites is likely due to development, consider creating or enhancing other pools that might allow some adults to transfer to the new site if they encounter it or develop a new breeding population from dispersal of colonizers.	Number of utilized pools.	VTrans, Towns	VTrans, development conditions
Technical Assistance, Training, Learning Networks	High	Implement vernal pool management guidelines as described by VFWD.	Number of trainings offered. Number of entities adopting the guidelines.	VFWD, FPR, Coverts, VWA, VT Family Forests, SAF, Land Trusts, Consulting Foresters	SWG

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Jefferson Salamander
Scientific Name: *Ambystoma jeffersonianum*
Species Group: Herp

Bibliography

- Andrews, J. 2001. The Vermont Reptile and Amphibian Atlas. <http://cat.middlebury.edu/herpatlas/>
- Bishop, S.C. 1941 (June). The salamanders of New York. New York State Museum bulletin No. 324. The University of the State of New York, Albany, New York 365 pp
- Bishop, S.C. 1994. Handbook of salamanders: The salamanders of the United States, of Canada, and of lower California. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 555 pp. (contains as much information on each species as The Salamanders of New York.)
- Bogart, J.P. and M. W. Klemens. 1997. Hybrids and genetic interactions of mole salamanders (*Ambystoma jeffersonianum* and *A. laterale*) in New York and New England. *American Museum Novitates* 3218: 1-78.
- Bogart, J.P. and Licht, L.E. 1986. Reproduction and the origin of polyploids in hybrid salamanders of the genus *Ambystoma*. *Canadian Journal of Genetic Cytology* 28:605-617.
- Brodman, R. 2002. Twelve-year study monitoring two species of pond-breeding salamanders in Northeast Ohio. *Ohio Journal of Science* 102: 123-127.
- Calhoun, A.J.K. and M. W. Klemens. 2002. Best Development Practices: Conserving pool-breeding amphibians in residential and commercial developments in the Northeastern United States. MCA Technical Paper No. 5, Metropolitan Conservation Alliance, Wildlife Conservation Society, Bronx, New York. 57 pp. (Call 924-925-9175 to order.)
- Calhoun, A.J.K. and P.G. deMaynadier. 2002. Forestry Habitat Management Guidelines for Vernal Pool Wildlife in Maine. A cooperative publication of the University of Maine, Maine Department of Inland Fisheries and Wildlife, Maine Audubon, and Maine Department of Conservation, Orono.
- Calhoun, A.J.K. and P.G. deMaynadier. 2008. Science and Conservation of Vernal Pools in Northeastern North America. CRC Press, Boca Rotan, FL, USA.
- Compton, B.W., K. McGarigal, S.A. Cushman, L.R. Gamble. 2007. A resistant-kernel model of connectivity for amphibians that breed in vernal pools. *Conservation Biology* 21:788-799
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Duellman, W.E. and L. Trueb. 1994. Biology of amphibians. The Johns Hopkins University Press, Baltimore, Maryland 670 pp. (The standard text for amphibians.)
- Epple, A.O. 1983. The amphibians of New England. Down East Books, Camden, Maine 138 pp.
- Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.
- Faccio, S.D. 2003. Post-breeding emigration and habitat use by Jefferson and Spotted salamanders in Vermont. *Journal of Herpetology* 37:479-489.
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)
- Heyer, W.R., M.A. Donnelly, R.W. McDiarmid, L-A. C. Hayek, and M.S. Foster. 1994. Measuring and monitoring biological diversity: standard methods for amphibians. Smithsonian Institution Press, Washington 364 pp.
- Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)
- Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Jefferson Salamander
Scientific Name: *Ambystoma jeffersonianum*
Species Group: Herp

Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp. (Visit the PARC website for more information: www.parcplace.org.)

Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)

Martel, A., et al. 2014. Recent introduction of a chytrid fungus endangers Western Palearctic salamanders. *Science* 346, 630; DOI: 10.1126/science.1258268.

Mitchell, J.C. 2000. Amphibian monitoring methods and field guide. Smithsonian National Zoological Park's Conservation & Research Center, Front Royal, Virginia 56 pp. (Very accessible, designed for citizen scientists.)

NEPARC. 2010. Northeast Amphibian and Reptile Species of Regional Responsibility and Conservation Concern. Northeast Partners in Amphibian and Reptile Conservation (NEPARC). Publication 2010-1.

Peterman, W.E., J.A. Crawford, and A.R. Kuhns. 2013. Using species distribution and occupancy modeling to guide survey efforts and assess species status. *Journal for Nature Conservation* 21: 114-121.

Petranka, J.W. 1998. Salamanders of the United States and Canada. Smithsonian Institution Press, Washington, DC 587 pp

Pfingsten, R.A. and F.L. Downs. 1989. Salamanders of Ohio. Bulletin of the Ohio Biological Survey Vol. 7, No. 2. College of Biological Sciences, The Ohio State University, Columbus, Ohio 315 pp, 29 plates.

Rowe, C.L., and W.A. Dunson. 1993. Relationships among biotic parameters and breeding effort by three amphibians in temporary wetlands of central Pennsylvania. *Wetlands* 13:237-246

Stebbins, R.C. and N.W. Cohen. 1995. A natural history of amphibians. Princeton University Press, Princeton, New Jersey 316 pp

Stokstad, E. 2014. The coming salamander plague. *Science* 346: 530-531.

Terres, G.D. 1999. Wildlife species of regional conservation concern in the Northeastern United States. *Northeast Wildlife* 54:93-100.

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp. (A standard text.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Blue-spotted Salamander**
Scientific Name: **Ambystoma laterale**
Species Group: **Herp**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S3

State Trend: Unknown

Extirpated in VT?

Regional SGCN? Yes

Assessment Narrative:

Blue-spotted Salamander is rare in Vermont (S3, SC), and is considered a species of conservation concern within the region due to its unknown population status, and taxonomic uncertainty (Terres 1999). The Blue-spotted Salamander is dependent on habitat mosaics consisting of lowland forest adjacent to fishless wetlands suitable for breeding. In addition, unisexual female hybrid populations exist that introduce uncertainties about the species' taxonomy, population biology, persistence, and long-term viability.

Distribution

Well distributed in the Champlain Valley and southern Champlain Hills, with scattered populations elsewhere; rare or absent from higher elevations of the Green Mountains.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Unknown	Southern Green Mtns	Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Known Watersheds

Metawee River
Passumpsic
West
Hudson-Hoosic
Lake Champlain
Lamoille River
Missiquoi River
Otter Creek
St. Francois River
Upper Connecticut
Winooski River

Probable Watersheds

Middle Connecticut
Waits
White



Common Name: **Blue-spotted Salamander**
Scientific Name: **Ambystoma laterale**
Species Group: **Herp**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Habitat mosaics that include breeding sites embedded in, or adjacent to, closed canopy forest stands are most critical. Maintaining forested corridors connected to other suitable habitat patches with breeding pools is important for juvenile dispersal and to maintain metapopulation dynamics. Requires hardwood/mixed forest stands with abundant coarse woody debris, leaf litter, and underground refugia (small mammal tunnels, rock crevices, etc.) surrounding a variety of wetland types, including red maple swamps, fens, marshes, temporary woodland (vernal) pools, etc. Most commonly found at lower elevations (<350m), in woodlands adjacent to forested wetlands with sufficient cover for breeding. More tolerant of disturbed habitats and smaller patch size than Jefferson Salamander (Klemens 1993).

Habitat Types:

Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Hardwood Swamps
Softwood Swamps
Seeps and Pools
Northern Hardwood
Open Peatlands
Marshes and Sedge Meadows
Shrub Swamps
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Habitat Fragmentation
Impacts of Roads or Transportation Systems
Climate Change

Description of habitat threat(s): Any habitat conversion, alteration, or fragmentation that disrupts species' ability to move between breeding and terrestrial sites, disrupts connectivity between breeding sites/metapopulations (Compton et al 2007), changes water/soil chemistry, temperature, pool hydroperiod, humidity, etc, may have negative effects. Road mortality can have major impacts on migrating adults and dispersing juveniles, especially when located between terrestrial and breeding habitats. Climate change that affects hydroperiod and/or water temperature of breeding pools could have significant impacts on productivity (Rowe and Dunson 1995).

Non-Habitat Threats:



Common Name: **Blue-spotted Salamander**
 Scientific Name: **Ambystoma laterale**
 Species Group: **Herp**

Genetics

Disease

Pollution

Reproductive Traits

Trampling or Direct Impacts

Description of non-habitat threat(s): Unisexual, female hybrid populations reproduce via gynogenesis requiring sperm from diploid males which is not incorporated into the genome of embryos (Petranka 1998), resulting in female-biased sex ratios. Since hybrid females require males in the population for successful breeding, but do not produce males to replace those that have been lost, they may have the potential to dilute genetic variability of diploid populations. Two emerging diseases (Ranavirus and Batrachochytrium salamandrivorans (Bs)) could present a threat in the future. Ranavirus was recently shown to be widely distributed in vernal pools in six mid-Atlantic states (Scott A. Smith pers. Comm.), while the fungal disease Bs, which originated in Asia, was recently detected in Europe (Stokstad 2014). Loss of metapopulation structure leading to genetic isolation due to fragmentation of suitable habitats by roads or other non-permeable development is problematic. Acid precipitation that lowers pH of breeding pools may reduce productivity. Also, widespread treatment of breeding pools to control West Nile Virus would likely have negative effects on amphibians. This is a long-lived amphibian that may not breed every year and produces relatively few eggs per breeding cycle.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Continue to field-verify mapped potential pools and other breeding sites statewide.
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	1) Identify distribution and relative abundance of VT populations. 2) Identify significant breeding sites. Large numbers (e.g., >25 egg masses) of breeders or evidence of use by any SGCN. 3) Continue to field-verify mapped potential vernal pools statewide.
Research	Threats and Their Significance	High	1) Conduct sampling to determine if Ranavirus is present in the state, and if so, determine its distribution and which species are affected. 2) Identify sites where road mortality is high annually and evaluate mitigation methods. 3) Identify and evaluate limiting factors.
Research	Population Genetics	Medium	Determine population genetics statewide and monitor populations for changes in sex ratios.
Research	Taxonomy	Medium	
Monitoring	Population Change	High	Implement a statistically robust monitoring protocol to track breeding phenology, population trends, and productivity, as well as changes in breeding site variables (e.g. hydroperiod, water chemistry, etc.).
Monitoring	Habitat Change	High	
Monitoring	Range Shifts	Medium	
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Blue-spotted Salamander**
 Scientific Name: **Ambystoma laterale**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks	High	Identify significant road crossings and develop safe road crossings to reduce roadkill.	Number of reported crossings.	VTrans, VCE, Town highway crews & Cons Comms	Vtrans, FHWA
Standards	Medium	Maintain 30-meter water quality buffer around entire perimeter of the pool. Timber harvesting, roads and any ground disturbing activities to be excluded within this buffer.	Number of pools with protected buffers.	DEC Wetlands	EPA
Habitat Restoration	Medium	If loss of important sites is likely due to development, consider creating or enhancing other pools that might allow some adults to transfer to the new site if they encounter it or develop a new breeding population from dispersal of colonizers.	Connectivity among clusters of pools. No net loss of functional breeding pools.	DEC Wetlands	EPA
Standards	Medium	Maintain habitat mosaic and connectivity between breeding pools. If two large sites are separated beyond the dispersal distance of a species it might be helpful to create or enhance pools that would link the two large sites.	Connectivity among clusters of pools. No net loss of functional breeding pools.	DEC Wetlands	EPA
Technical Assistance, Training, Learning Networks	High	Identify significant road crossings and develop safe road crossings to reduce roadkill.	Numbers of sites identified and addressed.	VTrans, FHWA	VTrans, FHWA
Awareness Raising and Communications	Medium	Help people understand the essential needs of all life stages, especially upland habitat in proximity to breeding pool.	Number of programs and individuals reached with message.	Reptile & Amphibian Atlas, VFWD Outreach Division, VCE	Private Foundation grants
Technical Assistance, Training, Learning Networks	High	Implement vernal pool management guidelines as described by VFWD.	Number of trainings offered. Number of entities adopting the guidelines.	VFWD, FPR, Coverts, VWA, VT Family Forests, SAF, Land Trusts, Consulting Foresters	SWG
Compatible Resource Use	High	Maintain corridor connections between upland and pool habitat.	Number of pools with secure travel paths to and from upland habitat. No net loss.	Consulting Foresters, USFS	Land Mgmt Agency operating budgets

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Blue-spotted Salamander**
 Scientific Name: **Ambystoma laterale**
 Species Group: **Herp**

Research	High	Determine presence/absence and distribution of Ranavirus in VT, and which species are most affected.	Number and distribution of sites sampled.	VCE, Reptile & Amphibian Atlas, DEC Wetlands	SWG, Private grants
Habitat Restoration	High	When existing road traffic is impacting migration of animals from terrestrial habitat to aquatic pool habitat, identify problem areas and redesign roads with crossing structures when roads are being upgraded.	Number of structures installed.	VTrans, FHWA, Towns	FHWA
Standards	High	Timber harvest should be conducted on frozen or dry ground to avoid rutting of ground that could change hydrology of an area or intercept amphibians should be avoided.	Number of harvests conducted on frozen ground.	FPR, Consulting Foresters, USFS	timber sale
Standards	High	Maintain 75% mature canopy cover beyond the 30-meter buffer in the terrestrial habitat.	Number and percentage of habitats with adequate canopy.	FPR, Consulting Foresters, USFS	
Compatible Resource Use	High	Maintain breeding pools and needed terrestrial habitat--usually directly adjacent to pool perimeter out to 200m but could be the equivalent area along a portion of the perimeter while minimizing edge).	Number of pools with adjacent upland habitat that is permeable for salamanders moving to and from pool.	Landowners	
Compatible Resource Use	Medium	Site all permanent roads more than 200 meters from a breeding pool, downslope of the pool if possible.	Number/ percentage of roads sited appropriately.	Developers, Act 250 Commission s	Planning grants, development costs
Compatible Resource Use	High	Maintain permeable forested habitat matrix between and among breeding populations so that individuals can be exchanged among populations.	Number of linkages between populations.	Reptile & Amphibian Atlas, Conservation Organization s, USFS	Current Use Program, EQIP
Habitat Restoration	Low	When feasible restore deciduous or mixed forest surrounding breeding pool.	Number of pools restored.	Vermont Family Forests	Current Use



Common Name: Blue-spotted Salamander
Scientific Name: *Ambystoma laterale*
Species Group: Herp

Bibliography

- Andrews, J. 2001. The Vermont Reptile and Amphibian Atlas. <http://cat.middlebury.edu/herpatlas/>
- Bishop, S.C. 1941 (June). The salamanders of New York. New York State Museum bulletin No. 324. The University of the State of New York, Albany, New York 365 pp
- Bishop, S.C. 1994. Handbook of salamanders: The salamanders of the United States, of Canada, and of lower California. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 555 pp. (contains as much information on each species as The Salamanders of New York.)
- Bogart, J.P. and M. W. Klemens. 1997. Hybrids and genetic interactions of mole salamanders (*Ambystoma jeffersonianum* and *A. laterale*) in New York and New England. *American Museum Novitates* 3218: 1-78.
- Bogart, J.P. and Licht, L.E. 1986. Reproduction and the origin of polyploids in hybrid salamanders of the genus *Ambystoma*. *Canadian Journal of Genetic Cytology* 28:605-617.
- Brodman, R. 2002. Twelve-year study monitoring two species of pond-breeding salamanders in Northeast Ohio. *Ohio Journal of Science* 102: 123-127.
- Calhoun, A.J.K. and M. W. Klemens. 2002. Best Development Practices: Conserving pool-breeding amphibians in residential and commercial developments in the Northeastern United States. MCA Technical Paper No. 5, Metropolitan Conservation Alliance, Wildlife Conservation Society, Bronx, New York. 57 pp. (Call 924-925-9175 to order.)
- Calhoun, A.J.K. and P.G. deMaynadier. 2002. Forestry Habitat Management Guidelines for Vernal Pool Wildlife in Maine. A cooperative publication of the University of Maine, Maine Department of Inland Fisheries and Wildlife, Maine Audubon, and Maine Department of Conservation, Orono.
- Calhoun, A.J.K. and P.G. deMaynadier. 2008. Science and Conservation of Vernal Pools in Northeastern North America. CRC Press, Boca Rotan, FL, USA.
- Compton, B.W., K. McGarigal, S.A. Cushman, L.R. Gamble. 2007. A resistant-kernel model of connectivity for amphibians that breed in vernal pools. *Conservation Biology* 21:788-799
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Duellman, W.E. and L. Trueb. 1994. Biology of amphibians. The Johns Hopkins University Press, Baltimore, Maryland 670 pp. (The standard text for amphibians.)
- Epple, A.O. 1983. The amphibians of New England. Down East Books, Camden, Maine 138 pp.
- Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.
- Faccio, S.D. 2003. Post-breeding emigration and habitat use by Jefferson and Spotted salamanders in Vermont. *Journal of Herpetology* 37:479-489.
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)
- Heyer, W.R., M.A. Donnelly, R.W. McDiarmid, L-A. C. Hayek, and M.S. Foster. 1994. Measuring and monitoring biological diversity: standard methods for amphibians. Smithsonian Institution Press, Washington 364 pp.
- Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)
- Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Blue-spotted Salamander
Scientific Name: *Ambystoma laterale*
Species Group: Herp

Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp. (Visit the PARC website for more information: www.parcplace.org.)

Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)

Martel, A., et al. 2014. Recent introduction of a chytrid fungus endangers Western Palearctic salamanders. *Science* 346, 630; DOI: 10.1126/science.1258268.

Mitchell, J.C. 2000. Amphibian monitoring methods and field guide. Smithsonian National Zoological Park's Conservation & Research Center, Front Royal, Virginia 56 pp. (Very accessible, designed for citizen scientists.)

NEPARC. 2010. Northeast Amphibian and Reptile Species of Regional Responsibility and Conservation Concern. Northeast Partners in Amphibian and Reptile Conservation (NEPARC). Publication 2010-1.

Peterman, W.E., J.A. Crawford, and A.R. Kuhns. 2013. Using species distribution and occupancy modeling to guide survey efforts and assess species status. *Journal for Nature Conservation* 21: 114-121.

Petranka, J.W. 1998. Salamanders of the United States and Canada. Smithsonian Institution Press, Washington, DC 587 pp

Pfingsten, R.A. and F.L. Downs. 1989. Salamanders of Ohio. Bulletin of the Ohio Biological Survey Vol. 7, No. 2. College of Biological Sciences, The Ohio State University, Columbus, Ohio 315 pp, 29 plates.

Rowe, C.L., and W.A. Dunson. 1993. Relationships among biotic parameters and breeding effort by three amphibians in temporary wetlands of central Pennsylvania. *Wetlands* 13:237-246

Stebbins, R.C. and N.W. Cohen. 1995. A natural history of amphibians. Princeton University Press, Princeton, New Jersey 316 pp

Stokstad, E. 2014. The coming salamander plague. *Science* 346: 530-531.

Terres, G.D. 1999. Wildlife species of regional conservation concern in the Northeastern United States. *Northeast Wildlife* 54:93-100.

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp. (A standard text.



Common Name: **Spotted Salamander**
Scientific Name: **Ambystoma maculatum**
Species Group: **Herp**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5

State Trend:

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

Spotted Salamander is found in pools adjacent to woodlands. It is widespread and abundant, but is killed in large numbers when it migrates across roads to and from its breeding pools, and is therefore a SGCN. Conservation status could change over time with increased human pressure that impacts the species in Vermont. For now we consider it to be a medium priority species.

Distribution

The Spotted Salamander is distributed widely in the state of Vermont, including the Green Mountains.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Known Watersheds

Black - Ottauquechee
Deerfield
Hudson-Hoosic
Lake Champlain
Lamoille River
Metawee River
Middle Connecticut
Missiquoi River
Otter Creek
Passumpsic
St. Francois River
Upper Connecticut
Upper Connecticut - Mascoma



Common Name: **Spotted Salamander**
Scientific Name: **Ambystoma maculatum**
Species Group: **Herp**

Waits

West

White

Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Spotted Salamanders must have both ephemeral breeding pools and upland habitat in proximity. They may also breed in semi-permanent or permanent wetlands where there is suitable cover to avoid predation by fish. Upland, non-breeding habitat requirements include hardwood/mixed forest stands with abundant coarse woody debris, leaf litter and underground refugia (e.g., small mammal tunnels, rock crevices) surrounding suitable breeding pools (Faccio 2003).

Habitat Types:

Oak-Pine Northern Hardwood

Hardwood Swamps

Softwood Swamps

Seeps and Pools

Spruce Fir Northern Hardwood

Northern Hardwood

Floodplain Forests

Open Peatlands

Marshes and Sedge Meadows

Shrub Swamps

Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Alteration

Habitat Fragmentation

Impacts of Roads or Transportation Systems

Climate Change

Description of habitat threat(s): Habitat problems for the Spotted Salamander include loss of upland and pool habitat, loss of movement between habitats, road mortality and less water as a result of climate change.



Common Name: **Spotted Salamander**
Scientific Name: **Ambystoma maculatum**
Species Group: **Herp**

Non-Habitat Threats:

Pollution

Reproductive Traits

Trampling or Direct Impacts

Description of non-habitat threat(s): Two emerging diseases (Ranavirus and Batrachochytrium salamandrivorans (Bs)) could present a threat in the future. Ranavirus was recently shown to be widely distributed in vernal pools in six mid-Atlantic states (Scott A. Smith pers. comm.), while the fungal disease Bs, which originated in Asia, was recently detected in Europe (Stokstad 2014). Loss of metapopulation structure leading to genetic isolation due to fragmentation of suitable habitats by roads or other non-permeable development is problematic. Acid precipitation that lowers pH of breeding pools may reduce productivity. Also, widespread treatment of breeding pools to control West Nile Virus would likely have negative effects on amphibians. This is a long-lived amphibian that may not breed every year and produces relatively few eggs per breeding cycle.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Continue to field-verify mapped potential pools and other breeding sites statewide.
Research	Basic Life History	Medium	Need to determine travel distances to pools
Research	Distribution and Abundance	Medium	1) Identify significant breeding sites. Large numbers (e.g., >25 egg masses) of breeders or evidence of use by any SGCN. 2) Continue to field-verify mapped potential vernal pools statewide
Research	Threats and Their Significance	High	1) Conduct sampling to determine if Ranavirus is present in the state, and if so, determine its distribution and which species are affected. 2) Identify sites where road mortality is high annually and evaluate mitigation methods. 3) Identify and evaluate limiting factors.
Monitoring	Population Change	High	Implement a statistically robust monitoring protocol to track breeding phenology, population trends, and productivity, as well as changes in breeding site variables (e.g. hydroperiod, water chemistry, etc.).
Monitoring	Habitat Change	High	It is important to understand the existing habitat base and track trends.
Monitoring	Monitor Threats	High	We should both monitor and manage limiting factors.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Spotted Salamander**
 Scientific Name: **Ambystoma maculatum**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	High	Determine presence/absence and distribution of Ranavirus in VT, and which species are most affected.	Number and distribution of sites sampled	VCE, Reptile & Amphibian Atlas, DEC Wetlands	SWG, Private grants
Standards		Maintain 75% mature canopy cover beyond the 30-meter buffer in the terrestrial habitat	Number of sites where canopy cover is retained	landowners state and federal lands managers	state lands Mgmt funds, federal lands Mgmt, EQIP
Technical Assistance, Training, Learning Networks		Identify significant road crossings and develop safe road crossings to address roadkill.	Number of sites identified and crossings developed.	Reptile & Amphibian Atlas, VTrans	VTrans, FHWA
Habitat Restoration		When feasible restore deciduous or mixed forest surrounding breeding pool	Number of sites with restored forest cover	landowners, consulting foresters, EQIP biologists	EQIP, Current Use
Standards		Maintain 30-meter water quality buffer around entire perimeter of the pool. Timber harvesting, roads and any ground disturbing activities to be excluded within this buffer	Number of sites where water quality is protected	DEC Water Quality/Wetlands Office, consulting foresters, EQIP biologists	EQIP
Standards		maintain habitat mosaic and maintain connectivity between breeding pools. If two large sites are separated beyond the dispersal distance of a species it might be helpful to create or enhance pools that would link the two large sites.	Number of areas linked.	FPR and VFWD lands managers, USFS, Federal Refuges, private landowners	State lands Mgmt funds, EQIP
Compatible Resource Use		Site all permanent roads more than 200 meters from a breeding pool, downslope of the pool if possible	Number of roads sited that minimize impacts.	VTrans, developers, Towns	Vtrans
Awareness Raising and Communications		Help people understand the essential needs of all life stages, especially upland habitat in proximity to breeding pool.	Number of people who are exposed to message	Reptile & Amphibian Atlas, VCE, SAG-Herps	private grants
Compatible Resource Use		Maintain corridor connections between upland and pool habitat	Number of sites with connections between pool and upland.	Landowners, consulting foresters, EQIP biologists	EQIP, Current Use

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Spotted Salamander**
 Scientific Name: **Ambystoma maculatum**
 Species Group: **Herp**

Habitat Restoration		When existing road traffic is impacting migration of animals from terrestrial habitat to aquatic pool habitat, identify problem areas and redesign roads with crossing structures when roads are being upgraded.	Number of structures installed	VTrans	VTrans, FHWA
Habitat Restoration		If loss of important sites is likely due to development, consider creating or enhancing other pools that might allow some adults to transfer to the new site if they encounter it or develop a new breeding population from dispersal of colonizers.	Number of sites enhanced.	VTrans	VTrans
Compatible Resource Use		Maintain breeding pools and needed terrestrial habitat--usually directly adjacent to pool perimeter out to 200m but could be the equivalent area along a portion of the perimeter while minimizing edge.	Number of sites with both pools and upland habitat intact.	Landowners, consulting foresters, Reptile & Amphibian Atlas	EQIP, Partners in Wildlife
Standards		Timber harvest should be conducted on frozen ground and rutting of ground that could change hydrology of an area or intercept amphibians should be avoided.	Number of harvest operations that do not disturb ground.	Consulting foresters, FPR	Current Use
Technical Assistance, Training, Learning Networks	High	Implement vernal pool management guidelines as described by VFWD.	Number of trainings offered. Number of entities adopting the guidelines.	VFWD, FPR, Coverts, VWA, VT Family Forests, SAF, Land Trusts, Consulting Foresters	SWG

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Spotted Salamander
Scientific Name: *Ambystoma maculatum*
Species Group: Herp

Bibliography

- Andrews, J. 2001. The Vermont Reptile and Amphibian Atlas. <http://cat.middlebury.edu/herpatlas/>
- Bishop, S.C. 1941 (June). The salamanders of New York. New York State Museum bulletin No. 324. The University of the State of New York, Albany, New York 365 pp
- Bishop, S.C. 1994. Handbook of salamanders: The salamanders of the United States, of Canada, and of lower California. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 555 pp. (contains as much information on each species as The Salamanders of New York.)
- Calhoun, A.J.K. and M. W. Klemens. 2002. Best Development Practices: Conserving pool-breeding amphibians in residential and commercial developments in the Northeastern United States. MCA Technical Paper No. 5, Metropolitan Conservation Alliance, Wildlife Conservation Society, Bronx, New York. 57 pp. (Call 924-925-9175 to order.)
- Calhoun, A.J.K. and P.G. deMaynadier. 2002. Forestry Habitat Management Guidelines for Vernal Pool Wildlife in Maine. A cooperative publication of the University of Maine, Maine Department of Inland Fisheries and Wildlife, Maine Audubon, and Maine Department of Conservation, Orono.
- Calhoun, A.J.K. and P.G. deMaynadier. 2008. Science and Conservation of Vernal Pools in Northeastern North America. CRC Press, Boca Rotan, FL, USA.
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- Compton, B.W., K. McGarigal, S.A. Cushman, L.R. Gamble. 2007. A resistant-kernel model of connectivity for amphibians that breed in vernal pools. *Conservation Biology* 21:788-799.
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Duellman, W.E. and L. Trueb. 1994. Biology of amphibians. The Johns Hopkins University Press, Baltimore, Maryland 670 pp. (The standard text for amphibians.)
- Epple, A.O. 1983. The amphibians of New England. Down East Books, Camden, Maine 138 pp.
- Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.
- Faccio, S.D. 2003. Post-breeding emigration and habitat use by Jefferson and Spotted salamanders in Vermont. *Journal of Herpetology* 37:479-489.
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)
- Heyer, W.R., M.A. Donnelly, R.W. McDiarmid, L-A. C. Hayek, and M.S. Foster. 1994. Measuring and monitoring biological diversity: standard methods for amphibians. Smithsonian Institution Press, Washington 364 pp.
- Homan, R. B. Windmiller, and J. Reed. 2004. Critical thresholds associated with habitat loss for two vernal pool-breeding amphibians. *Ecological Applications* 14: 1547-1553.
- Homan, R., J. Regosin, D. Rodrigues, J. Reed, B. Windmiller, and L. Romero. 2003. Impacts of varying habitat quality on the physiological stress of spotted salamanders (*Ambystoma maculatum*). *Animal Conservation* 6: 11-18.
- Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)
- Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.
- Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp. (Visit the PARC website for more information: www.parcplace.org.)

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Spotted Salamander
Scientific Name: *Ambystoma maculatum*
Species Group: Herp

Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)

Martel, A., et al. 2014. Recent introduction of a chytrid fungus endangers Western Palearctic salamanders. *Science* 346, 630; DOI: 10.1126/science.1258268.

Mitchell, J.C. 2000. Amphibian monitoring methods and field guide. Smithsonian National Zoological Park's Conservation & Research Center, Front Royal, Virginia 56 pp. (Very accessible, designed for citizen scientists.)

NEPARC. 2010. Northeast Amphibian and Reptile Species of Regional Responsibility and Conservation Concern. Northeast Partners in Amphibian and Reptile Conservation (NEPARC). Publication 2010-1.

Petranka, J.W. 1998. Salamanders of the United States and Canada. Smithsonian Institution Press, Washington, DC 587 pp

Pfingsten, R.A. and F.L. Downs. 1989. Salamanders of Ohio. Bulletin of the Ohio Biological Survey Vol. 7, No. 2. College of Biological Sciences, The Ohio State University, Columbus, Ohio 315 pp, 29 plates.

Regosin, J. B. Windmiller, J. Reed. 2003. Influence of abundance of small mammal burrows and conspecifics on the density and distribution of spotted salamanders in terrestrial habitats. *Canadian Journal of Zoology* 81:591-605.

Semlitsch, R.D. and J.R. Bodie. Are small wetlands expendable? *Conservation Biology* 12(5):1129-1133.

Semlitsch, R.D. and J.R. Bodie. 2003. Biological criteria for buffer zones around wetlands and riparian habitats for amphibians and reptiles. *Conservation Biology* 17(5):1219-1228.

Stebbins, R.C. and N.W. Cohen. 1995. A natural history of amphibians. Princeton University Press, Princeton, New Jersey 316 pp

Stokstad, E. 2014. The coming salamander plague. *Science* 346: 530-531.

Tyning, T.F. 1990. A guide to amphibians and reptiles. Little, Brown and Company. Boston Massachusetts 400 pp.

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp. (A standard text.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Four-toed Salamander**
Scientific Name: **Hemidactylium scutatum**
Species Group: **Herp**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S2

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

The Four-toed Salamander is rare in Vermont (S2, SC) and the region. Although the species is small and secretive, its distribution appears to be limited to low elevations of the Champlain Valley, Taconics, and probably the lower Connecticut River Valley. The species has specialized breeding requirements and is dependent on habitat mosaics consisting of lowland forest adjacent to suitable wetlands for breeding.

Distribution

The distribution of the Four-toed Salamander appears to be limited to low elevations of the Champlain Valley, Taconics, and probably the lower Connecticut River Valley.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

West
Upper Connecticut-Mascoma
Mettawee River
Lake Champlain
Middle Connecticut
Mettawee River
Otter Creek
Winooski River

Probable Watersheds

Lamoille River
Missisquoi River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Moss or sedge hummocks or moss mats overhanging (often sphagnum) standing water is a critical feature of Four-toed Salamander breeding sites. Eggs are deposited in moss mats and, upon hatching, larvae wiggle



Common Name: Four-toed Salamander
Scientific Name: *Hemidactylium scutatum*
Species Group: Herp

through moss and enter pools. Habitat mosaics that include these breeding sites embedded in, or adjacent to, closed canopy forest stands are critical. Maintaining forested corridors connected to other suitable habitat patches with breeding sites is important for juvenile dispersal and to maintain metapopulation dynamics. Eggs have also been reported in the literature in rotting logs or leaf litter. Requires relatively mature, moist hardwood/mixed forest stands with abundant coarse woody debris, leaf litter, and underground refugia (small mammal tunnels, rock crevices, etc.) in close proximity to suitable breeding sites. Breeding sites include a variety of mossy wetlands, including red maple swamps, bogs, fens, temporary woodland (vernal) pools, etc. Most commonly found in bottomland forests adjacent to shallow, mossy wetlands with pools and sufficient cover for breeding. In surveys in Massachusetts and Maine, species appears to drop-out of suitable sites above 300 m elevation (A. Richmond, pers. Comm.)

Habitat Types:

Hardwood Swamps
Softwood Swamps
Seeps and Pools
Northern Hardwood
Oak-Pine Northern Hardwood
Open Peatlands
Marshes and Sedge Meadows
Wet Shores
Shrub Swamps
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Impacts of Roads or Transportation Systems
Habitat Fragmentation
Climate Change

Description of habitat threat(s): The Four-toed Salamander has specialized breeding requirements that make it vulnerable to habitat disturbance. Any habitat conversion, alteration, or fragmentation that disrupts species' ability to move between breeding and terrestrial sites, changes water/soil chemistry, temperature, pool hydroperiod, humidity, etc, may have negative effects. Road mortality can negatively impact migrating adults and dispersing juveniles, especially when located between terrestrial and breeding habitats. Climate change that affects hydroperiod and/or water temperature of breeding pools could have significant impacts on productivity.

Non-Habitat Threats:

Genetics
Disease



Common Name: **Four-toed Salamander**
Scientific Name: **Hemidactylium scutatum**
Species Group: **Herp**

Pollution

Trampling or Direct Impacts

Description of non-habitat threat(s): Loss of metapopulation structure leading to genetic isolation due to fragmentation of suitable habitats by roads or other non-permeable development is problematic. Widespread treatment of breeding pools to control West Nile Virus would likely have negative effects on many amphibians, including Four-toed Salamanders. Two emerging diseases (Ranavirus and Batrachochytrium salamandrivorans (Bs)) could present a threat in the future. Ranavirus was recently shown to be widely distributed in vernal pools in six mid-Atlantic states (Scott A. Smith pers. comm.), while the fungal disease Bs, which originated in Asia, was recently detected in Europe (Stokstad 2014).

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Gain better understanding of breeding habitat requirements in Vermont and upper elevational limits.
Research	Basic Life History	Medium	
Research	Distribution and Abundance	High	1) Identify significant breeding sites. Large numbers (e.g., >25 egg masses) of breeders or evidence of use by any SGCN. 2) Determine distribution and relative abundance of species in southern Connecticut River Valley, Vermont Valley, Champlain Hills, and Taconics. 3) Confirm whether the disjunct population represented by the near-historic record from 1989 in Fairlee is extant.
Research	Threats and Their Significance	High	1) Conduct sampling to determine if Ranavirus is present in the state, and if so, determine its distribution and which species are affected. 2) Identify sites where road mortality is high annually and evaluate mitigation methods. 3) Identify and evaluate limiting factors.
Research	Population Genetics	Low	
Monitoring	Population Change	High	
Monitoring	Habitat Change	High	It is important to understand the existing habitat base and track trends.
Monitoring	Range Shifts	Medium	
Monitoring	Monitor Threats	High	We should both monitor and manage limiting factors.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Four-toed Salamander**
 Scientific Name: **Hemidactylium scutatum**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	When existing road traffic is impacting migration of animals from terrestrial habitat to aquatic pool habitat, identify problem areas and redesign roads with crossing structures when roads are being upgraded.	Number of redesigns of roads.	VTrans	VTrans, FHWA
Technical Assistance, Training, Learning Networks	High	Identify significant road crossings and develop safe road crossings to address roadkill.	Number of crossing identified and structures installed	VTrans	VTrans
Compatible Resource Use	High	Maintain forested habitat matrix around breeding sites.	Number of pools within forested matrix	Landowners	Current Use, EQIP
Standards	Medium	Maintain 30-meter water quality buffer around entire perimeter of the pool. Timber harvesting, roads and any ground disturbing activities to be excluded within this buffer.	Number of pool sites managed for good water quality	Consulting Foresters, private and public landowners	Current Use, EQIP
Standards	High	Maintain 75% mature canopy cover beyond the 30-meter buffer in the terrestrial habitat.	Number of pools with canopy cover retained	Consulting Foresters, private and public landowners	Current Use, EQIP
Compatible Resource Use	High	Maintain corridor connections between upland and pool habitat.	Number of breeding sites with secure upland connection	Landowners, consulting foresters	volunteer compliance, Current Use, EQIP
Technical Assistance, Training, Learning Networks	High	Implement vernal pool management guidelines as described by VFWD.	Number of trainings offered. Number of entities adopting the guidelines.	VFWD, FPR, Coverts, VWA, VT Family Forests, SAF, Land Trusts, Consulting Foresters	SWG
Habitat Restoration	Medium	If loss of important sites is likely due to development, consider creating or enhancing other pools that might allow some adults to transfer to the new site if they encounter it or develop a new breeding population from dispersal of colonizers.	Number of utilized pools created	Landowners, VTrans	VTrans, volunteer effort
Research	High	Determine presence/absence and distribution of Ranavirus in VT, and which species are most affected.	Number and distribution of sites sampled.	VCE, Reptile & Amphibian Atlas, DEC Wetlands	SWG, private grants

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Four-toed Salamander**
 Scientific Name: **Hemidactylium scutatum**
 Species Group: **Herp**

Standards	High	Maintain habitat mosaic and maintain connectivity between breeding pools. If two large sites are separated beyond the dispersal distance of a species it might be helpful to create or enhance pools that would link the two large sites.	Number of sites enhanced.	Landowners, consulting foresters	Current Use, EQIP
Standards	High	Timber harvest should be conducted on frozen ground and rutting of ground that could change hydrology of an area or intercept amphibians should be avoided.	Number of operations that avoid ground alteration.	FPR, Consulting Foresters	State Lands Mgmt, Current Use
Awareness Raising and Communications	Medium	Help people understand the essential needs of all life stages, especially upland habitat in proximity to breeding pool.	Number of people who receive message.	Reptile & Amphibian Atlas, VCE	private grant
Compatible Resource Use	High	Maintain breeding pools and needed terrestrial habitat--usually directly adjacent to pool perimeter out to 200m but could be the equivalent area along a portion of the perimeter while minimizing edge.	Number of pools with needed upland	landowners, consulting foresters	State Lands Mgmt, Current Use, EQIP
Habitat Restoration	Low	When feasible restore deciduous or mixed forest surrounding breeding pool.	Sites with restored forested habitat.	Landowners	volunteer effort, EQIP
Compatible Resource Use	Medium	Site all permanent roads more than 200 meters from a breeding pool, downslope of the pool if possible.	Number of roads sited so that pool impacts are minimized.	VTrans, Towns	VTrans, FHWA
Technical Assistance, Training, Learning Networks	High	Identify significant road crossings and develop safe road crossings to reduce roadkill.	Number of structures installed.	VTrans, VCE	VTrans

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Four-toed Salamander
Scientific Name: *Hemidactylium scutatum*
Species Group: Herp

Bibliography

Andrews, J. 2001. The Vermont Reptile and Amphibian Atlas. <http://cat.middlebury.edu/herpatlas/>

Bishop, S.C. 1941 (June). The salamanders of New York. New York State Museum bulletin No. 324. The University of the State of New York, Albany, New York 365 pp

Bishop, S.C. 1994. Handbook of salamanders: The salamanders of the United States, of Canada, and of lower California. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 555 pp. (contains as much information on each species as The Salamanders of New York.)

Calhoun, A.J.K. and M. W. Klemens. 2002. Best Development Practices: Conserving pool-breeding amphibians in residential and commercial developments in the Northeastern United States. MCA Technical Paper No. 5, Metropolitan Conservation Alliance, Wildlife Conservation Society, Bronx, New York. 57 pp. (Call 924-925-9175 to order.)

Calhoun, A.J.K. and P.G. deMaynadier. 2002. Forestry Habitat Management Guidelines for Vernal Pool Wildlife in Maine. A cooperative publication of the University of Maine, Maine Department of Inland Fisheries and Wildlife, Maine Audubon, and Maine Department of Conservation, Orono.

Chalmers, Rebecca J.; Loftin, Cynthia S. 2006. Wetland and microhabitat use by nesting four-toed salamanders in Maine. *Journal of Herpetology*. 40(4): 478-485.

Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.

Corser, Jeffrey D.; Dodd, C. Kenneth, Jr. 2004. Fluctuations in a metapopulation of nesting four-toed salamanders, *Hemidactylium scutatum*, in the Great Smoky Mountains National Park, USA, 1999-2003. *Natural Areas Journal*. 24(2): 135-140.

DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.

Duellman, W.E. and L. Trueb. 1994. Biology of amphibians. The Johns Hopkins University Press, Baltimore, Maryland 670 pp. (The standard text for amphibians.)

Epple, A.O. 1983. The amphibians of New England. Down East Books, Camden, Maine 138 pp.

Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.

Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)

Harris, R. N., and D. E. Gill. 1980. Communal nesting, brooding behavior, and embryonic survival of the four-toed salamander *Hemidactylium scutatum*. *Herpetologica* 36 :141-144.

Harris, R. N., and P. M. Ludwig. 2004. Resource level and reproductive frequency in female four-toed salamander, *Hemidactylium scutatum*. *Ecology* 85:1585-1590.

Heyer, W.R., M.A. Donnelly, R.W. McDiarmid, L-A. C. Hayek, and M.S. Foster. 1994. Measuring and monitoring biological diversity: standard methods for amphibians. Smithsonian Institution Press, Washington 364 pp.

Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)

Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.

Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp. (Visit the PARC website for more information: www.parcplace.org.)

Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Four-toed Salamander
Scientific Name: *Hemidactylium scutatum*
Species Group: Herp

Martel, A., et al. 2014. Recent introduction of a chytrid fungus endangers Western Palearctic salamanders. *Science* 346, 630; DOI: 10.1126/science.1258268.

Mitchell, J.C. 2000. Amphibian monitoring methods and field guide. Smithsonian National Zoological Park's Conservation & Research Center, Front Royal, Virginia 56 pp. (Very accessible, designed for citizen scientists.)

Natural Heritage and Endangered Species Program. 2007. Massachusetts Forestry Conservation Management Practices for Four-toed Salamanders. Draft (August 2007). Natural Heritage and Endangered Species Program, Massachusetts Division of Fisheries and Wildlife, Westborough, Massachusetts, USA. <http://www.mass.gov/eea/docs/dfg/nhesp/regulatory-review/fourtoed-salamander-cmp.pdf>

NEPARC. 2010. Northeast Amphibian and Reptile Species of Regional Responsibility and Conservation Concern. Northeast Partners in Amphibian and Reptile Conservation (NEPARC). Publication 2010-1.

Petranka, J.W. 1998. Salamanders of the United States and Canada. Smithsonian Institution Press, Washington, DC 587 pp

Pfingsten, R.A. and F.L. Downs. 1989. Salamanders of Ohio. Bulletin of the Ohio Biological Survey Vol. 7, No. 2. College of Biological Sciences, The Ohio State University, Columbus, Ohio 315 pp, 29 plates.

Semlitsch, R.D. and J.R. Bodie. Are small wetlands expendable? *Conservation Biology* 12(5):1129-1133.

Stebbins, R.C. and N.W. Cohen. 1995. A natural history of amphibians. Princeton University Press, Princeton, New Jersey 316 pp

Stokstad, E. 2014. The coming salamander plague. *Science* 346: 530-531.

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp. (A standard text.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Mudpuppy**
Scientific Name: **Necturus maculosus**
Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The Mudpuppy is native to Lake Champlain where it appears to be restricted to shoals and shallows and the lower reaches of lake's tributaries. The Burlington area is the source of the type specimen from the late 1700s. There is also a Connecticut River population that is generally agreed to be from an introduced source. Recent analysis of genetic samples from Lake Champlain and the Connecticut River (Chellman 2011) confirmed the affinities of the Lake Champlain Mudpuppy with populations of the Ohio River drainage, whereas, the affinities of the Mudpuppy in the Connecticut River drainage are not with populations sampled from the Northeast, further supporting the likelihood that the Connecticut River population was introduced. The Mudpuppy is a long-lived species that does not reach reproductive maturity until about six years old. Survival of breeding adults is very important to the maintenance of populations.

Surveying for the Mudpuppy has proven to be difficult and therefore our understanding of abundance, and even distribution, remains uncertain. U.S. Fish & Wildlife Service attempts to survey for Mudpuppy in and near Lake Champlain met with little success. Two recommendations to list this species as threatened were not accepted by the Agency of Natural Resources.

Widespread lampricide applications in many Lake Champlain tributaries is a risk to the Mudpuppy with large losses documented in some treated rivers. Evidence indicates possible losses of some smaller populations. The largest known kill from lampricide applications took place in the Lamoille River during its first treatment (2009) where 528 dead Mudpuppies were found along the shoreline and shallows of the treated area that was searched for non-target mortality. Rescue operations for non-target species affected by lampricide during the second (2013) application of the same river found fewer than 10 Mudpuppies, strongly suggesting a very significant decline in that river. Specimens in some cage studies during lampricide treatments have survived and we do not fully understand why the effects have been so different among treatments. A smaller population in Lewis Creek appears to have declined to the point where the Mudpuppy is no longer detectable. No specimens have been located in Lewis Creek during non-target post-treatment surveys for the last two decades.

Distribution

Primarily the shoals and shallows of Lake Champlain and lower reaches of its tributaries (to the first impassable falls or dam), and the Connecticut River and its tributaries north to Waterford. In addition, it is reported from some waterbodies easily accessed (no barriers) from the Lake Champlain and Connecticut River populations (for e.g., Lake Morey). This species has also been reported from Otter Creek in Brandon but not since 1998. Populations in and near the Connecticut River appear to be introduced from a source other than our native population in Lake Champlain and its tributaries.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Confident	Taconic Mtns	Not Probable

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Mudpuppy**
Scientific Name: **Necturus maculosus**
Species Group: **Herp**

Northeastern Highlands Probable

Distribution by Watershed:

Known Watersheds

Probable Watersheds

Lamoille River

Metawee River

Upper Connecticut - Mascoma

Upper Connecticut

Waits

Winooski River

Middle Connecticut

West

Black-Ottawquechee

Lake Champlain

Missiquoi River

Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

The Mudpuppy is a generalist, occupying myriad habitats throughout its range including lakes, rivers, small creeks, canals, irrigation ditches and reservoirs (Escleshymer 1906; Hamilton 1932; Bishop 1941; Harris 1959a, b). Waters inhabited may be clear or turbid; still, slow or rapid flowing; cold or warm and may have gravel, cobbled, or muddy bottoms (Harris 1959a). Adults need suitable habitat with cover (for nesting, protection from predators, and shelter from light) such as flat rocks, slabs, logs or planks (Escleshymer 1906; Pearse 1910; Bishop 1941; Harris 1959a). Juveniles are usually found in greater numbers in the substrate of pools where silt and organic debris have accumulated to a minimal thickness of several mm (Matson 1990).

In one study the maximum distance that a Mudpuppy moved was 256m in its aquatic environment (Shoop and Gunning 1967). There is some evidence of short distance seasonal migrations from shallows to pools or river channels.

Habitat Types:

Aquatic: Lacustrine

Aquatic: Fluvial

Aquatic: Lower CT River

Aquatic: Large Lake Champlain Tribs Below Falls

Aquatic: Man-Made Water Bodies

Aquatic: Lake Champlain



Common Name: **Mudpuppy**
Scientific Name: **Necturus maculosus**
Species Group: **Herp**

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Sedimentation
Inadequate Disturbance Regime
Habitat Fragmentation

Description of habitat threat(s): The amount of quality habitat for this species at present is likely less than that of 200 years ago but how much less is unknown. Upstream movement for feeding and spawning has probably been limited by dam building. Pollution and sedimentation of nesting sites have caused declines in populations in Ohio (Pfingsten and White 1989) and is likely to have limited appropriate habitat availability in the Lake Champlain drainage system. Removal of snags and other navigational hazards on rivers and lakeshores may have decreased nesting sites.

Non-Habitat Threats:

Disease
Pollution
Reproductive Traits
Trampling or Direct Impacts

Description of non-habitat threat(s): A large percentage of the area of known distribution for the native Mudpuppy in Vermont coincides with the lampricide-treated spawning habitat for Sea Lamprey. Significant mortality has been documented due to TFM lampricide applications in NY and VT. The Mudpuppy can be negatively affected by chemical pesticides (Bonin et al. 1995; Gendron et al. 1997), especially those used in the control of parasitic Sea Lamprey (*Petromyzon marinus*) such as 3-trifluoromethyl-4-nitrophenol (TFM) and TFM/2', 5-dichloro-4-nitrosalicylanide (Niclosamide) mixtures (Matson 1990; Boogaard et al. 2003). Acute toxicity studies of TFM and of TFM/Niclosamide mixtures have been conducted on adult Mudpuppies (Boogaard et al. 2003; Boogaard et al. 2008) and juvenile Mudpuppies (Neuderfer 2002; Neuderfer et al. 2004; Boogaard et al. 2008) including one-year-old Mudpuppies (Durfey and Neuderfer 2009). All of these studies found that TFM and TFM/Niclosamide mixtures caused mortality in Mudpuppies but that the sensitivity to these lampricides was age specific (Neuderfer 2002; Boogaard et al. 2003). Boogaard et al. (2003, 2008) found that adult Mudpuppies were sufficiently more resistant to the lampricides than Sea Lamprey and suggested that there was a sufficient safety margin for selective control of Sea Lamprey in the presence of adult Mudpuppies (i.e., <10% expected mortality). Whereas the information on the acute toxicity of juvenile Mudpuppies between 2 and 5 years of age is limited (Neuderfer 2002), the studies (Neuderfer et al. 2004; Boogaard et al. 2008; Durfey and Neuderfer 2009) on juvenile mudpuppies (young of the year or 1 year of age) has found that this life stage is at risk of substantial treatment related mortality at lampricide concentrations at those needed to control Sea Lampreys. These toxicity studies (Neuderfer 2002; Boogaard et al. 2003; Neuderfer et al. 2004; Boogaard et al. 2008; Durfey and Neuderfer 2009) predict the greatest mortality to young of the year and 1 year old Mudpuppies and little mortality to adult Mudpuppies from exposure to lampricides. The field data from the 2009 Lamoille River lampricide treatment reveal that 19.3% of the dead Mudpuppies detected were adults.

Some individuals are caught on hooks. A large Mudpuppy die off due to botulism was reported in the Great Lakes in 2003.



Common Name: **Mudpuppy**
Scientific Name: **Necturus maculosus**
Species Group: **Herp**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Determine how the mudpuppy is using Vermont's lake and river habitat and when. Gather data on egg-laying sites, instream shelter, and seasonal movement patterns.
Research	Basic Life History	Medium	Determine if and when mudpuppies are entering or leaving Vermont rivers would to help us understand the potential impacts of TFM treatments.
Research	Distribution and Abundance	High	1) Develop survey techniques to effectively sample the Mudpuppy. 2) Mudpuppy distribution is not well known in VT and we do not have a good idea of population size or trend. Monitor the size and determine the sustainability of existing populations. eDNA methods are currently being tested to evaluate Mudpuppy distribution in the Champlain basin.
Research	Threats and Their Significance	High	We need to better understand the impacts of TFM applications, fragmentation and changed river flows due to dams, and sedimentation.
Research	Population Genetics	Medium	Determine the genetic source or at least the most closely related population to our Connecticut River Mudpuppy population.
Monitoring	Population Change	High	
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	High	Range shift may be a useful index of population change.
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Mudpuppy**
 Scientific Name: **Necturus maculosus**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications	Medium	Help Vermonters assign value to this species through educational programs, printed material, web site information, field trips, TV and video information.	Number of programs and individuals reached with message.	ECHO, VFWD Outreach Division	Corporate Sponsors, Lake Champlain Basin Program
Habitat Restoration	Medium	Consider removal of stream barriers that restrict upstream movement.	Number of miles of streams reopened.	FERC, Trout Unlimited, VTrans, towns	Power Companies, VTrans, municipalities
Habitat Restoration	High	Maintain adequate stream flows to support mudpuppy populations.	Number of streams	DEC Water Quality, Power Generation companies, FERC	Power Generation companies
Awareness Raising and Communications	High	Encourage reports of sightings to the Vermont Natural Heritage Inventory and the VT Reptile & Amphibian Atlas.	Numbers of reports received	Reptile & Amphibian Atlas, volunteers	Nongame Wildlife Fund, SWG
Invasive Species Control & Prevention	High	Investigate and implement alternative lamprey control methods that have less impacts to mudpuppy populations.	Number of times alternative methods used.	TNC, USFWS, VFWD Fisheries	DJ, USFWS, SWG
Habitat Restoration	Medium	Prevent sedimentation that degrades mudpuppy habitat.	Improvements in water quality. Number of miles of buffer strips established. Turbidity measures	DEC Water Quality, Dam operators, Dept. of Agriculture, VTrans, NRCS	Dam operators, Dept. of Agriculture, VTrans, EQIP
Planning & Zoning	Medium	Implement stream management to promote better water quality in mudpuppy waters.	Streams with improved water quality	USFWS, VFWD Fisheries, DEC Water Quality	DJ, USFWS



Common Name: Mudpuppy
Scientific Name: *Necturus maculosus*
Species Group: Herp

Bibliography

- Allran, J. W., and W. H. Karasov. 2001. Effects of Atrazine on embryos, larvae and adults of Anuran amphibians. *Environmental Toxicology and Chemistry*, 20:769-775.
- Andrews, J. S. 2013. The Vermont Reptile and Amphibian Atlas. VtHerpAtlas.org
- Beebee, T. J. C., and R. A. Griffiths. 2005. The amphibian decline crisis: a watershed for conservation biology? *Biological Conservation*, 125:271-285.
- Blaustein, A. R., and J. M. Kiesecker. 2002. Complexity in conservation: lessons from the global decline of amphibian populations. *Ecological Letters*, 5:597-608.
- Bishop, S. C. 1926. Notes on the habitats and development of the mudpuppy, *Necturus maculosus* (Rafinesque). *New York State Museum Bulletin*, 268:5-60.
- Bishop, S. C. 1941. The salamanders of New York. *New York State Museum Bulletin*, 324:1-365.
- Bonin, J., J. L. DesGranges, C. A. Bishop, J. Rodrigue, A. Gendron, and J. E. Elliot. 1995. Comparative study of contaminants in the mudpuppy (Amphibia) and the common snapping turtle (Reptilia), St. Lawrence River, Canada. *Archives of Environmental Contaminant Toxicology*, 28:184-194.
- Boogaard, M. A., T. D. Bills, and D. A. Johnson. 2003. Acute toxicity of TFM and a TFM/Niclosamide mixture to selected species of fish, including lake sturgeon (*Acipenser fulvescens*) and mudpuppies (*Necturus maculosus*), in laboratory and field exposures. *Journal of Great Lakes Research* 29 (Supplement 1):529-541.
- Boogaard, M. A., M. P. Gaikowski, J. E. Rivera, and T. D. Hubert. 2008. Relative toxicity of the lampricides TFM and a TFM:1% niclosamide mixture to adult and juvenile mudpuppies (*Necturus maculosus*) in comparison to sea lampreys (*Petromyzon marinus*). Report for the Great Lake Fishery Commission, Ann Arbor, Michigan, 22 pp.
- Boone, M. D., and R. D. Semlitsch. 2002. Interactions of an insecticide with competition and pond drying in amphibian communities. *Ecological Applications*, 12:307-316.
- Breisch, A. 1996. Effects of lampricides of amphibians: Little Ausable and Ausable Rivers and Deltas, Lake Champlain, NY. Unpublished report New York Department of Environmental Conservation, Delmar.
- Breisch, A. 1999. Non-target amphibian mortality associated with the 1999 lampricide treatment. Unpublished report New York Department of Environmental Conservation, Delmar.
- Breisch, A. 2000. Non-target amphibian mortality associated with the 2000 lampricide treatment. Unpublished report New York Department of Environmental Conservation, Delmar.
- Cagle, F. R. 1954. Observations on the life history of the salamander *Necturus louisianensis*. *Copeia*, 1954:257-260.
- Chellman, I. C. 2011. Population demographics and genetic diversity of mudpuppies (*Necturus maculosus*) in Vermont. Unpublished M.S. thesis, University of Vermont, Burlington, 92 pp.
- Chellman, I. C., and D. L. Parrish 2010. Developing methods for sampling mudpuppies in Vermont tributaries of Lake Champlain. Final Report, Vermont Fish and Wildlife Department, 71 pp.
- Collins, J. P., and A. Storfer. 2003. Global amphibian declines: sorting the hypotheses. *Diversity and Distribution*, 9:89-98.
- Crocker, D. W. 1960. Mudpuppies in Maine. *Maine Field Naturalist*, 16:14-17.
- DeKay, J. E. 1842. The New York Fauna: Part III Reptiles and Amphibia. White, W., A. White and J. Visscher, Albany.
- Du Preez, L. H., N. Kunene, R. Hanner, J. P. Gisey, K. R. Solomon, A. Hosmer, and G. J. Van Der Kraak. 2009. Population-specific incidence of testicular ovarian follicles in *Xenopus laevis* from South Africa: a potential issue in endocrine testing. *Aquatic Toxicology*, 95:10-16.
- Durfey, L., and G. Neuderfer. 2009. Acute toxicity of the lampricide mixture TFM/1% Niclosamide to one-year-old mudpuppies (*Necturus maculosus*). New York State Department of Environmental Conservation, Ray Brook, NY. 12 pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Mudpuppy
Scientific Name: *Necturus maculosus*
Species Group: Herp

- Eycleshymer, A. C. 1906. The habits of *Necturus maculosus*. *American Naturalist*, 40:123-136.
- Fitch, K. L. 1959. Observations on the nesting habits of the mudpuppy, *Necturus maculosus* (Rafinesque). *Copeia*, 1959:339-340.
- Gendron, A. D. 1999. Status report on the mudpuppy, *Necturus maculosus* (Rafinesque), in Canada. Report to Reptile and Amphibian Subcommittee, Committee on the Status of Endangered Wildlife in Canada, Wildlife Canada, Quebec.
- Gendron, A. D., C. A. Bishop, R. Fortin, and A. Hontela. 1997. In vivo testing of the functional integrity of the corticosterone-producing in mudpuppy (Amphibia) exposed to chlorinated hydrocarbons in the wild. *Environmental Toxicology and Chemistry*, 16:1694-1706.
- Gibbons, J. W., and S. Nelson, Jr. 1968. Observations of the mudpuppy, *Necturus maculosus*, in a Michigan lake. *American Midland Naturalists*, 80:562-564.
- Gilderhaus, P.A. and B.G. H. Johnson. 1980. Effects of sea lamprey (*Petromyzon marinus*) control in the Great Lakes on aquatic plants, invertebrates, and amphibians. *Canadian Journal of Fisheries and Aquatic Sciences* 37:1895-1905.
- Green, N. B., and T. K. Pauley. 1987. *Amphibians and Reptiles of West Virginia*. University of Pittsburg Press, Pittsburg, PA.
- Hayes, T. B. 2004. There is no denying this: defusing the confusion about Atrazine. *BioScience*, 54:1138-1149.
- Hayes, T. B., K. Haston, M. Tsui, A. Hoang, C. Haeffele, and A. Vonk. 2003. Atrazine-induced hermaphroditism at 0.1 ppb in American leopard frogs (*Rana pipiens*): laboratory and field evidence. *Environmental Health Perspective*, 2003:568-575.
- Hayes, T. B., V. Khoury, A. Narayan, M. Nazir, A. Park, T. Brown, L. Adame, E. Chan, D. Buchholz, T. Stueve, and S. Gallipeau. 2010. Atrazine induces complete feminization and chemical castration in male African clawed frogs (*Xenopus laevis*). *Proceedings of the National Academy of Sciences, USA*, 107:4612-4617.
- Hamilton, W. J., Jr. 1932. The food and feeding habits of some eastern salamanders. *Copeia*, 1932:83-86.
- Harris, J. P., Jr. 1959a. The natural history of *Necturus*: I. Habitats and habits. *Field and Laboratory*, 27:11-20.
- Harris, J. P., Jr. 1959b. The natural history of *Necturus*: III. Food and feeding. *Field and Laboratory*, 27:105-111.
- Harris, J. P., Jr. 1959c. The natural history of *Necturus*: II. Respiration. *Field and Laboratory*, 27:71-77.
- Henshaw, S. 1904. *Fauna of New England*. 2. List of the Batrachia. *Occasional Paper of the Boston Society of Natural History*, 7:1-10.
- Hosmer, A. J. 2009. Response from Syngenta: the need for objective, science based advocacy. *Environmental Science and Technology*, October 2009:7595.
- Hutchison, V. H., and J. P. Ritchart. 1989. Annual cycle of thermal tolerance in the salamander, *Necturus maculosus*. *Journal of Herpetology*, 23:73-76.
- Klemens, M. W. 1993. *Amphibians and reptiles of Connecticut and adjacent regions*. *State Geological and Natural History Survey of Connecticut Bulletin*, 112:1-318.
- Lagler, K. F., and K. E. Goellner. 1941. Notes on *Necturus maculosus* (Rafinesque), from Evans Lake, Michigan. *Copeia*, 1941:96-98.
- Lytle, M., and N. Staats. 2003. Mudpuppy (*Necturus maculosus*) sampling in the Lake Champlain drainage. Progress Report. U.S. Fish and Wildlife Service, Essex Junction, VT. 15 pp.
- Lytle, M., and N. Staats. 2004. Mudpuppy (*Necturus maculosus*) sampling in the Lake Champlain drainage. Progress Report. U.S. Fish and Wildlife Service, Essex Junction, VT. 5 pp.
- Mairs, D. F. 1999. Common mudpuppy. Pp. 33-36, in *Maine Amphibians and Reptiles* (M. L. Hunter, A. J. K. Calhoun, and M. McCollough, eds.). University of Maine Press, Orono.
- Matson, T. O. 1990. Estimation of numbers for a riverine *Necturus* population before and after TFM lampricide exposure. *Kirtlandia*, 45:33-38.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Mudpuppy
Scientific Name: *Necturus maculosus*
Species Group: Herp

- Matson, T.O. 1998. Evidence for home ranges in mudpuppies and implications for impacts due to episodic applications of the lampricide TFM. In Status and Conservation of Midwestern amphibians. Ed. M.J. Lannoo Univ. of Iowa Press. 507 pp.
- McDaniel, T. V., P. A. Martin, G. C. Barrett, K. Hughes, A. D. Gendron, L. Shirose, and C. A. Bishop. 2009. Relative abundance, age structure, and body size in mudpuppy populations in southwestern Ontario. *Journal of Great Lakes Research*, 35:182-189.
- Mitchell, J.C. 2000. Amphibian monitoring methods and field guide. Smithsonian National Zoological Park's Conservation & Research Center, Front Royal, Virginia 56 pp. (Very accessible, designed for citizen scientists.)
- Neuderfer, G. N. 2002. Draft summary of acute toxicity of TFM to mixed-aged juvenile mudpuppies (*Necturus maculosus*). New York State Department of Environmental Conservation, Avon, NY. 32 pp.
- Neuderfer, G. N., B. D. Chipman, and L. Durfey. 2004. Draft acute toxicity of TFM and a TFM-1% Niclosamide mixture to juvenile mudpuppies. New York State Department of Environmental Conservation, Avon, NY. 17 pp.
- Petranka, J. W. 1998. Salamanders of the United States and Canada. Smithsonian Institutional Press, Washington D.C.
- Pearse, A. S. 1921. Habits of the mud-puppy, *Necturus*, an enemy of food fishes. Bureau of Fisheries Economic Circular, 29:161-208.
- Pfingsten, R. A., and A. M. White. 1989. *Necturus maculosus* (Rafinesque). Pp. 71-78, in Salamanders of Ohio (R. A. Pfingsten and F. L. Downs, eds.). Ohio Biological Survey, Columbus.
- Pope, C. H. 1947. Amphibians and Reptiles of the Chicago Area. Chicago Natural History Museum Press, Chicago, IL.
- Reptile and Amphibian Scientific Advisory Group to the Vermont Endangered Species Committee. 2011. Narrative, supporting information, and bibliography for the proposed listing of the mudpuppy (*Necturus maculosus*) as a state threatened species in Vermont.
- Relyea, R. A. 2005. The impacts of insecticides and herbicides on the biodiversity and productivity of aquatic communities. *Ecological Applications*, 15:618-627.
- Richmond, A. M. 2000. The mudpuppy: New England's largest salamander. *Massachusetts Wildlife*, 45:9-15.
- Rohr, J. R., and K. A. McCoy. 2010a. A qualitative meta-analysis reveals consistent effects of Atrazine on freshwater fish and amphibian. *Environmental Health Perspective*, 118:20-32.
- Rohr, J. R., and K. A. McCoy. 2010b. Preserving environmental health and scientific credibility: a practical guide to reducing conflicts of interest. *Conservation Letters*, 3:143-150.
- Sajdak, R. A. 1982. Seasonal activity, habitat selection, and population structure of the mudpuppy, *Necturus maculosus*, in a Wisconsin stream. Unpubl. M.S. thesis, University of Wisconsin, Milwaukee.
- Schmidt, R. E. 2004. Mudpuppy (*Necturus maculosus*) in the tidal Hudson River, with comments on its status as native. *Northeastern Naturalists*, 11:179-188.
- Shoop, C. R. 1965. Aspects of reproduction in Louisiana *Necturus* populations. *American Midland Naturalist*, 74:357-367.
- Shoop, C. R., and G. E. Gunning. 1967. Seasonal activity and movements of *Necturus* in Louisiana. *Copeia*, 1967:732-737.
- Solomon, K. R., J. A. Carr, L. H. Du Preez, J. P. Giesy, R. J. Kendall, E. E. Smith, and G. J. Van Der Kraak. 2008. Effects of Atrazine on fish, amphibian, and aquatic reptiles: a critical review. *Critical Reviews in Toxicology*, 38:721-772.
- Stejneger, L., and T. Barbour. 1933. A Checklist of North American Amphibians and Reptiles. Third edition, Cambridge.
- Stuart, S. N., J. S. Chanson, N. A. Cox, B. E. Young, A. L. Rodrigues, D. L. Fischman, and R. W. Waller. 2004. Status and trends of amphibian declines and extinctions worldwide. *Science*, 306:1783-1786.
- Thompson, D. G., K. R. Solomon, B. F. Wojtaszek, A. N. Edginton, G. R. Stephenson, and R. A. Relyea. 2006. The impacts of insecticides and herbicides on the biodiversity and productivity of aquatic communities: [with response]. *Ecological Applications*, 16:2022-2034.
- Thompson, Z. 1853. Natural History of Vermont. Charles E. Tuttle Co., Rutland, VT.



Common Name: **Mudpuppy**
Scientific Name: **Necturus maculosus**
Species Group: **Herp**

Wake, D. B., and V. T. Vredenburg. 2008. Are we in the midst of the sixth mass extinction? A view from the world of amphibian. *Proceedings of the National Academy of Sciences, USA*, 105:11466-11473.

Warfel, H. E. 1936. Notes on the occurrence of *Necturus maculosus* (Rafinesque) in Massachusetts. *Copeia*, 1936:237.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Fowler's Toad**
Scientific Name: **Anaxyrus fowleri**
Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The Fowler's Toad is extremely rare in Vermont (S1) and was listed as state-endangered in 2015. The Fowler's Toad has specialized habitat requirements and depends on habitat mosaics consisting of disturbed shorelines and uplands adjacent to shallow breeding sites. Habitat loss due to succession is also a problem for this species. The last documented sighting in Vermont was in 2007 in Vernon. However, recent reports of calling from islands within the Connecticut River of NH suggest that populations may still exist in Vermont.

Distribution

The Fowler's Toad is primarily a species of the Atlantic Coastal Plain, its distribution in Vermont is limited to the lower Connecticut River valley, with populations documented at one site each in Vernon and White River Junction (Andrews 2001, Barker and Caduto 1984).

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Middle Connecticut

Black-Ottawquechee

Probable Watersheds

NA

NA

NA

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

The most critical habitat features include dry, sandy woodlands near shallow water for breeding. Breeding sites may include river edges, pond or lake margins, shallow wetlands (forested or emergent), vernal pools, roadside ditches, etc. Fowler's Toads are tolerant of and dependent upon warmer temperatures than American Toads (Frost and Martin, 1971). Along the north shore of Lake Erie all Fowler's Toad reports are within ½ kilometer of the shore and the toads require habitat in the early stages of ecological succession. At those sites they require five habitat types in close proximity to sustain a population (COSEWIC, 2010):
--Hibernation habitat (sandy dunes)



Common Name: **Fowler's Toad**
Scientific Name: **Anaxyrus fowleri**
Species Group: **Herp**

- Breeding, egg-laying habitat (sparsely vegetated still-water ponds, sandy bottom pools, shallow rocky shoals, or rocky pools).
- Feeding and hydration habitat (sandy riverside and lakeshore habitats with bare to sparse vegetation cover)
- Daytime retreat and aestivation habitat (sandy beaches and shoreline debris), and
- Dispersal corridor habitat.

Overwintering habitat is mentioned as a potential limiting factor in Canada (COSEWIC, 2010). Burrows must be deep enough for the toads to avoid freezing, close enough to the water table to be damp, but not so deep as to be flooded. Toads are not tolerant of freezing or of long-term submergence while over wintering.

Stille (1952) reported small home ranges with most toads emerging from the ground within 60-210 meters of the water's edge. In Canada (COSEWIC, 2010) Fowler's Toads (nocturnal) spend days buried in soil up to 400 m from the waters edge but they must move to the water as soon as they emerge to replace moisture lost while in the soil.

Along Lake Erie, Fowler's Toads depend upon breeding sites that are continually created or maintained by disturbance.

Breeding habitat in Vermont appears to be the disturbed margins of the Connecticut River and its tributaries in Windham and Windsor Counties, and perhaps shorelines of other water bodies near sandy soils in those floodplains. Terrestrial habitat appears to be largely open areas of adjacent floodplains and lower-elevation uplands within a few hundred meters of those breeding sites, particularly those with sandy or gravelly soils. This includes yard edges and moderately developed residential or agricultural areas. According to Klemens (1993) the species prefers well-drained sand and gravel habitat in Connecticut. Wright and Wright (1949) state "wherever Fowler's Toads are sympatric with American Toads (as they are anywhere in Vermont), Fowler's Toads occur in rivers, streams, or lake beaches" and American Toads in the uplands. This appears to be the case in Vermont. Soil maps show large deposits of sand in the Vernon area.

Habitat Types:

Upland Shores

Oak-Pine Northern Hardwood

Hardwood Swamps

Seeps and Pools

Northern Hardwood

Floodplain Forests

Marshes and Sedge Meadows

Early Succession Pine and Hemlock

Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Lawns, Gardens, and Row Crops

Wet Shores

Shrub Swamps

Early Succession Northern Hardwoods



Common Name: **Fowler's Toad**
Scientific Name: **Anaxyrus fowleri**
Species Group: **Herp**

Early Succession Upland Oak
Aquatic: Lacustrine
Aquatic: Lower CT River
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development
Habitat Alteration
Habitat Fragmentation
Inadequate Disturbance Regime
Impacts of Roads or Transportation Systems
Climate Change

Description of habitat threat(s): The Fowler's Toad has specialized habitat requirements that make it vulnerable to disturbance. Any habitat conversion, alteration, or fragmentation that disrupts species' ability to move between breeding and terrestrial sites may have negative effects. Road mortality can negatively impact migrating adults and dispersing juveniles, especially when located between terrestrial and breeding habitats. Climate change that affects hydroperiod and/or water temperature of breeding pools could have significant impacts on productivity.

Early successional habitat in sandy soils within 400 meters of the Connecticut River has probably been reduced significantly with the development of an extensive series of flood control dams in the Connecticut River drainage. In addition, sandy and gravelly soils in the floodplain have been desirable sites for shoreline development and agriculture. Some types of low-density development and agriculture (pasture, some crops, new farm ponds) may have created open early-successional foraging habitat or breeding habitat for this species; however, high-density development with heavy road traffic (toads suffer high road mortality), row crops and intensive pesticide or herbicide use (atrazine) are probably not consistent with continued Fowler's Toad use. Bank stabilization activities would also limit the amount of potential habitat for this species.

This floodplain area has also seen significant road building. Routes 91 and 5 both parallel the river within the floodplain on the Vermont side as well as numerous smaller roads such as 142 in Vernont.

Breeding habitat of this species seems to be concentrated on islands and along the shorelines of the Connecticut River, within the disturbance zone. Regular removal of vegetation, new deposits of sand and gravel, and creation of small pools may be necessary to maintain breeding habitat. Dams along the Connecticut River are built to control flooding. It is possible that they control flooding to an extent that limits the amount of breeding habitat for this species. In addition, subdivision and development within the narrow Connecticut River valley may be limiting useful habitat and increasing mortality on roads and by machinery.

This species regularly undergoes large population changes. The existence of nearby healthy populations to recolonize previously occupied areas is essential. In addition, the colonizers within those populations need to be able to safely traverse the landscape along the river for some distance as populations rebuild. Given



Common Name: **Fowler's Toad**
Scientific Name: **Anaxyrus fowleri**
Species Group: **Herp**

distances between populations that may be larger than the dispersal range of juvenile toads, all five required habitat types will need to be located fairly regularly (~every 8 miles) along the shore of the Connecticut River in order for recolonization to take place from a distant source. Impediments to travel exist in increased road traffic, more intensive or chemical dependent agricultural methods, and intensive development such as in the towns along the river.

Non-Habitat Threats:

Genetics

Disease

Pollution

Trampling or Direct Impacts

Description of non-habitat threat(s): Known populations of this species occur in close proximity to humans, and in a region of the state that continues to see significant development pressure. Fragmentation of suitable habitats by roads or other non-permeable development may result in loss of metapopulation structure and leading to genetic isolation, especially considering the limited and localized populations of this species. Widespread treatment of breeding pools to control West Nile Virus would likely have negative effects on many amphibians, including Fowler's Toad.

According to Freda and Dunson (1986) this species shows decreased larval growth rates with increased acidity (lowered pH) due to acid rain. It is also less tolerant than most amphibians to atrazine (Birge et al., 2000), and is particularly sensitive to the insecticide azinphos-methol (Guthion; Mayer and Ellersieck, 1986). The organochlorides endrin, toxaphene, dieldrin, toxaphene, DDT, and lindane are also highly toxic to larval Fowler's Toads (Sanders, 1970). Adults were also highly sensitive to organochlorides (Ferguson and Gilbert, 1968) as well as pyrethroid insecticides (Bennett et al., 1983) and the metals chromium, gallium, titanium, and aluminum (Birge et al., 2000). In southwestern Ontario, agricultural chemicals were listed as a possible contributing factor to Fowler's Toads declines. The herbicide Trifluralin and the insecticide Endrin were reported to be particularly toxic to toads (COSEWIC, 2010). The disappearance of Fowler's Toads from many of the Massachusetts islands was thought to be the result of DDT use according to Lazell (1976). DDT is also suspected of eliminating populations on Point Pelee in Canada (COSEWIC, 2010). We have not looked at the available data on the level of any of these substances in the Connecticut River or on surrounding lands, although we expect atrazine is widely used on corn crops along the Connecticut River.

Fowler's Toads are susceptible to mycobacterial (Shively et al., 1981) and parasitic infections (Jilek and Wolff, 1978; Ashton and Rabalais, 1978; McAllister et al., 1989; and Vences et al., 2003). Botulism is also considered a potential threat to Fowler's Toads (COSEWIC, 2010). Along the north shore of Lake Erie it was noticed that shoreline mats of algae created the anaerobic conditions that allow *Clostridium botulinum* to survive.

Toads overwinter and avoid predation and desiccation during the day and during dry periods by digging into sandy or loose soil (Harding and Holman, 1992). By the end of the winter they have burrowed to depths of up to 15-30 cm (R. Latham quoted in Oliver, 1955). Tilling of the soil in late fall or early spring may disturb or kill overwintering Fowler's Toads. Tilling during other times of the year could have the same impact on toads underground for the daytime hours or when aestivating to escape dehydration.



Common Name: **Fowler's Toad**
Scientific Name: **Anaxyrus fowleri**
Species Group: **Herp**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Due to the likely spotty distribution of the rare Fowler's Toad in Vermont, it is important that we document and map habitat including connectivity of patches.
Research	Distribution and Abundance	High	1) There is a need to better document the distribution of the Fowler's Toad in Vermont, which will require dedicated searches during the calling period. 2) Continue to document species distribution in Connecticut River Valley with targeted searches of potential sites, and sites where previously reported.
Research	Threats and Their Significance	High	
Monitoring	Population Change	High	Monitor known population annually and attempt to determine population size and demographics.
Monitoring	Habitat Change	High	The habitat of the Fowler's Toad is likely vulnerable to human development and fragmentation of its habitat, including breeding pools. Monitoring the amount and regularity of habitat creation though flooding would provide better insight into the potential impacts of flood control dams.
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Fowler's Toad**
 Scientific Name: **Anaxyrus fowleri**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Protected Area Management	Medium	Eliminate the use of pesticides and herbicides within protected areas and maintain early successional habitat patches in a way that minimizes direct mortality.	Development and implementation of management plans.	Easement holders, VFWD; VFPR	VHCB, CNWR, Carbon tax revenues
Policy & Regulations	High	Through the FERC dam relicensing process insure adequate water releases to create and maintain breeding habitat.	Increased scouring, vegetation removal, and sand and gravel deposits along Connecticut River.	NH F&W, TNC	
Easements	Medium	1) Purchase land or easements along margin of Connecticut River to the 100 year flood high water line to allow safe flooding and scouring. 2) Purchase land or easements for known terrestrial habitat along Stebbins Road in Vernon.	Change in the acreage of conserved land	VLT, Conte NWR, local and national land trusts, RPCs	VHCB, CNWR, Carbon tax revenues
Awareness Raising and Communications	High	Encourage reports of sightings to the VT Heritage Inventory and the VT Reptile & Amphibian Atlas.	Number of reports received	Reptile & Amphibian Atlas	Nongame Wildlife Fund, private grants
Policy & Regulations	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	ANR, Health Dept, USFWS	
Technical Assistance, Training, Learning Networks	High	Implement vernal pool management guidelines as described by VFWD.	Number of trainings offered. Number of entities adopting the guidelines.	VFWD, FPR, Coverts, VWA, VT Family Forests, SAF, Land Trusts, Consulting Foresters	SWG

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Fowler's Toad**
Scientific Name: **Anaxyrus fowleri**
Species Group: **Herp**

Bibliography

Andrews, J. 2013. The Vermont Reptile and Amphibian Atlas. VtHerpAtlas.org

Ashton, A.D. and F.C. Rabalais. 1978. Helminth parasites of some anurans of northwestern Ohio. Proceedings of the Helminthological Society of Washington 45:141-142.

Barker, M.W. and M.J. Caduto. 1984. Geographic distribution: Bufo woodhousii fowleri. SSAR Herpetological Review 15:51-52.

Bennett, R.S., E.E. Klaas, J.R. Coats, M.A. Mayse and E.J. Kolbe. 1983. Fenvalerate residues in nontarget organisms from treated cotton fields. Bulletin of Environmental Toxicology 31:61-65.

Birge, W.J., A.G. Westerman and J.A. Spromberg. 2000. Comparative toxicology and risk assessment of amphibians. Pp. 727-791. In Sparling, D.W., G. Linder and C.A. Bishop (Eds.), Ecotoxicology of amphibians and reptiles. Society for Environmental Toxicology and Contaminants (SETAC) Press, Pensacola, Florida.

Breden, F. 1987. Population structure of Fowler's Toads Bufo woodhousei fowleri. Copeia 1987:386-395.

Breden, F. 1988. The natural history and ecology of Fowler's Toad, Bufo woodhousei fowleri (Amphibia: Bufonidae) in the Indiana Dunes National Lakeshore. Fieldiana Zoology 49:1-16.

Clarke, R.D. 1977. Postmetamorphic survivorship of Fowler's Toads Bufo woodhousei fowleri. Copeia 1977 594-597.

Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.

Committee on the Status of Endangered Wildlife in Canada (COSEWIC). 2010. COSEWIC assessment and status report on the Fowler's Toad Anaxyrus fowleri in Canada. Ottawa. Vii + 58 pp.

DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.

Dickerson, M.C. 1969. The frog book: North American toads and frogs, with a study of the habits and life histories of those of the northeastern states. Dover Publications, Inc., New York 253 pp.

Dodd, C.K. 2013. Frogs of the United States and Canada. Johns Hopkins University Press, Baltimore, Maryland, 982 pp. in two volumes.

Duellman, W.E. and L. Trueb. 1994. Biology of amphibians. The Johns Hopkins University Press, Baltimore, Maryland 670 pp. (The standard text for amphibians.)

Epple, A.O. 1983. The amphibians of New England. Down East Books, Camden, Maine 138 pp.

Ferguson, D.E. and C.C. Gilbert. 1968. Tolerances of three species of anuran amphibians to five chlorinated hydrocarbon insecticides. Journal of the Mississippi Academy of Sciences 13:135-138.

Freda, J. and W.A. Dunson. 1985. The influence of external cation concentration on the hatching of amphibian embryos in water of low pH. Canadian Journal of Zoology 63:2649-2656.

Frost, J.S. and E.W. Martin. 1971. A comparison of distribution and high temperature tolerance in Bufo americanus and Bufo woodhousii fowleri. Copeia 1971:750-751.

Green, D.M. 1992. Fowler's Toads Bufo woodhousii fowleri, at Long Point, Ontario: Changing abundance and implications for conservation. Pp. 37-45. In Bishop, C.A. and K.E. Pettit (Eds.), Declines in Canadian amphibian populations: designing a national monitoring strategy. Canadian Wildlife Service, Occasional Publications, Number 76, Environment Canada, Ottawa, Ontario, Canada.

Green, D.M. 1997. Temporal variation in abundance and age structure in Fowler's Toads, Bufo fowleri at Long Point, Ontario. Pp. 45-56. In Green D.M. (Ed.), Amphibians in decline: Canadian studies of a global problem. Herpetological Conservation, Number 1, Society for the Study of Amphibians and Reptiles, St. Louis, Missouri.

Green, D.M., A.R. Yagi, and Stewart E. Hamill. 2011. Recovery Strategy for the Fowler's Toad, (Anaxyrus fowleri) in Ontario. Ontario Recovery Strategy Series. Prepared for the Ontario Ministry of Natural Resources, Peterborough, Ontario. Vi + 21 pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Fowler's Toad**
Scientific Name: **Anaxyrus fowleri**
Species Group: **Herp**

Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)

Hranitz, J.M., T.S. Klinger, F.C. Hill, R.G. Sagar, T. Mencken and J. Carr. 1993. Morphometric variation between Bufo woodhousii fowleri Hinckley (Anura: Bufonidae) on Assateague Island, Virginia and the adjacent mainland. Brimleyana 19:65-75.

Heyer, W.R., M.A. Donnelly, R.W. McDiarmid, L-A. C. Hayek, and M.S. Foster. 1994. Measuring and monitoring biological diversity: standard methods for amphibians. Smithsonian Institution Press, Washington 364 pp.

Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp.

IUCN. 2012. International Union for the Conservation of Nature online map of Fowler's Toad worldwide distribution <http://maps.iucnredlist.org/map.html?id=54640>

Jackson, S.D., R.M. Richmond, T.F. Tynning and C.W. Leahy (eds). 2010. Massachusetts Herpetological Atlas 1992-1998, Massachusetts Audubon Society & University of Massachusetts (www.massherpatlas.org).

Jilek, R. and R. Wolff. 1978. Occurrence of Spinitectus gracilis Ward Magath 1916 (Nematoda: Spiruroidea) in the toad (Bufo woodhousii fowleri) in Illinois. Journal of Parasitology 64:619.

Kellner, A. and D.M. Green. 1995. Age structure and age of maturity in Fowler's Toads Bufo woodhousii fowleri at their northern range limit. Journal of Herpetology 29:417-421.

Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp. (Visit the PARC website for more information: www.parcplace.org.)

Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)

Lannoo, M. (Ed.) 2005. Amphibian declines: the conservation status of United States species. University of California Press, Berkeley and Los Angeles, California 926 pp.

Lazell, J.D. Jr. 1976. This broken archipelago. Cape Cod and the islands, amphibians and reptiles. Demeter Press, New York. 260 pp.

Mayer, F.L. and M.R. Ellersieck. 1986. Manual of acute toxicity: interpretation and database for 410 chemicals and 66 species of freshwater animals. U.S. Fish and Wildlife Service, Resource Publication, Number 160, Washington, D.C.

McAllister, C.T., S.J. Upton and D.B. Conn. 1989. A comparative study of endoparasites in three species of sympatric Bufo (Anura: Bufonidae), from Texas. Proceedings of the Helminthological Society of Washington 56:162-167.

Mitchell, J.C. 2000. Amphibian monitoring methods and field guide. Smithsonian National Zoological Park's Conservation & Research Center, Front Royal, Virginia 56 pp. (Very accessible, designed for citizen scientists.)

Oliver, J.A. 1955. The natural history of North American amphibians and reptiles. D. Van Nostrand Company, Princeton, New Jersey.

Quinn, H.R. and C. Scott. 2005. Critical areas, In Lannoo, M. (Ed.). Amphibian declines: the conservation status of United States species. University of California Press, Berkeley and Los Angeles, California 926 pp.

Reptile and Amphibian Scientific Advisory Group to the Vermont Endangered Species Committee. 2013. Narrative, supporting information, and bibliography for the proposed listing of the Fowler's Toad (Anaxyrus fowleri) as a state endangered species in Vermont.

Sanders, H.O. 1970. Pesticide toxicities to tadpoles of the Western Chorus Frog Pseudacris maculate and Fowler's Toads Bufo woodhousii fowleri. Copeia 1970: 246-251.

Shively, J.N., J.G. Songer, S. Prchal, M.S. Keasey III, and C.O. Thoen. 1981. Mycobacterium marinum infection in Bufonidae. Journal of Wildlife Diseases 17:3-7.



Common Name: **Fowler's Toad**
Scientific Name: **Anaxyrus fowleri**
Species Group: **Herp**

Smith, M. A. and D.M. Green. 2006. Sex isolation and fidelity: unbiased long distance dispersal in a terrestrial amphibian. *Ecography* 29:649-658.

Stebbins, R.C. and N.W. Cohen. 1995. A natural history of amphibians. Princeton University Press, Princeton, New Jersey 316 pp.

Stille, W.T. 1952. The nocturnal amphibian fauna of the southern Lake Michigan beach. *Ecology* 33:149-162.

Vences, M., M.W. Penuel-Matthews, D.R. Vieites and R. Altig. 2003. Natural history notes: *Rana temporaria* (Common Frog) and *Bufo fowleri* (Fowler's Toad). Protozoan infestation. *Herpetological Review* 34:237-238.

Wright, A.H. and A.A. Wright. 1995. Handbook of frogs and toads of the United States and Canada. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 640 pp.

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp



Common Name: **Boreal Chorus Frog**
Scientific Name: **Pseudacris maculata**
Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Recent genetic research by Moriarty Lemmon et al. (2007) indicated that the chorus frog of northeastern North America previously considered to be *P. triseriata*, which includes Vermont's population, is actually *P. maculata*. This small eastern portion of the species' range is separated from the larger western distribution by a minimum of 200 miles.

The Boreal Chorus Frog previously was considered a western species, present in Ontario and west to the Northwest Territories and eastern British Columbia, and south into the United States from Minnesota to Montana into northern New Mexico. *P. triseriata* was considered to occur from the Saint Lawrence Valley in Quebec westward to Minnesota and South Dakota, and south to Oklahoma, Indiana, and western New York. Genetic evidence presented by Moriarty Lemmon et al. (2007) redefined the geographic boundaries for several members of this genus, including these two species. According to this work, *P. triseriata* is more limited, occurring from western Kentucky and southern Illinois, northeastward through Michigan, southern Ontario, and western New York. *P. maculata* is more widespread, occurring from Illinois north to Ontario and westward, with a disjunct population occurring in southeastern Ontario, and within the Saint Lawrence Valley of Quebec and New York and the northern Lake Champlain Valley.

P. triseriata used to be listed as an SGCN species in Vermont. Based on the study above which is supported by additional anecdotal information (responses to tapes) and gaps in distribution, we feel the best current science suggests that the species we thought was *P. triseriata* is actually *P. maculata*. This taxonomy has not been accepted by all authorities, but seems to be the most supported at present. As a result, in 2011 *P. triseriata* was removed from our list of Vermont endangered species and *P. maculata* was added. We are now removing *P. triseriata* from our list of SGCN and replacing it with *P. maculata*. The supporting information here is an updated version of the same information previously presented but under a corrected name.

This species was located in townships along northern Lake Champlain from Swanton/Alburt to Georgia in the 1970s. Searches in 1988, 1996, and 1997 at the original site and along roads in the northwestern corner of the state were unsuccessful. A novel occurrence of singing male chorus frogs was observed in 1998 and 1999 in Alburt, with only two heard during each visit. This breeding site is located less than one mile from the Quebec border. Annual searches in 2000-2009 have not located chorus frogs at either of the two last known sites or other locations. It has also disappeared from the NY portion of the Lake Champlain Basin (see Corser et al. 2012) and many of its historic locations in Quebec and eastern Ontario. The COSEWIC website states "populations in Quebec are documented to have declined at a rate of 37% over 10 years and are expected to continue to decline. Despite there being some areas where chorus frogs remain evident, surveys of populations in Ontario indicate a significant decline in abundance of 30% over the past decade". In Vermont this species is state-listed as endangered.

Habitat loss due to development, succession, and drainage of pools in agricultural fields are all reported problems for this species. Species may also be vulnerable to taking by mowing of agricultural fields and newer chemicals (herbicides) used in agriculture. Recent studies also show this genus to be sensitive to chemicals produced by the invasive plant species European Buckthorn (*Rhamnus cathartica*). This plant is currently widespread in the Lake Champlain basin. Habitat in Vermont seems to be flooded or ponded pasture with shrubs. There is evidence to support calling competition between this species and the Spring Peeper (*Pseudacris crucifer*), so calling times and perhaps localized distribution may need to be isolated from each

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Boreal Chorus Frog**
Scientific Name: **Pseudacris maculata**
Species Group: **Herp**

other. There appears to be appropriate habitat remaining within this species historic distribution within Vermont, leading this writer to suspect that changes other than or in addition to habitat loss and fragmentation are causing declines.

Distribution

This species was located in townships along northern Lake Champlain from Swanton/Alburg to Georgia in the 1970s. Searches in 1988, 1996, and 1997 at the original site and along roads in the northwestern corner of the state were unsuccessful. A novel occurrence of singing male chorus frogs was observed in 1998 and 1999 in Alburg, with only two heard during each visit. This breeding site is located less than one mile from the Quebec border. Annual searches in 2000-2009 have not located Chorus Frogs at either of the two last known sites or other locations.

Distribution by Biophysical Region:

Champlain Valley	Probable	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

The Boreal Chorus Frog (*P. maculata*) is primarily a terrestrial species, occurring in moist, marshy, vegetated environments near breeding sites (NatureServe 2011, IMNH 2000). It may be found in grassy and shrubby areas, meadows, open woodlands, fens, bogs, or similar low habitats (NatureServe 2011, IMNH 2000, Bider and Matte 1996) not far from open ponds, ditches, marshes, temporary pools, or other wetlands (NatureServe 2011, IMNH 2000, HerpNet). The species is generally not found in forested areas (Bider and Matte 1996). *P. maculata* breeds in temporary to semi-permanent waters of no current, often quite small with no or few fish predators (HerpNet, Lannoo 2005). These are typically open habitats (no canopy) with abundant emergent vegetation which provides both cover and egg attachment sites (IMNH 2000, Lannoo 2005). Boreal chorus frogs are considered poor dispersers, though individual studies have found adults traveling 275 (Quebec) and 685 m (Colorado) from breeding sites (HerpNet, Whiting 2004). Outside the breeding season, this is generally a nocturnal species, being inactive during the day, and is active into September or October (Lannoo 2005, NatureServe 2011). Overwintering occurs under protective cover or underground in the upland areas surrounding a breeding site (NatureServe 2011). Some indirect evidence suggests limited possible hibernation within the breeding pool (Whiting 2004).

In Vermont, the breeding sites observed in 1985 and 1998 were both small, open cattail ponds adjacent to agricultural lands and larger wetland systems. No habitat analyses were undertaken, but the available terrestrial



Common Name: Boreal Chorus Frog
Scientific Name: *Pseudacris maculata*
Species Group: Herp

and aquatic habitat types available agreed with those described in literature for *P. maculata*. The field notes taken by Fred Schueler during his 1975 survey indicate the Vermont chorus frog to be somewhat of a generalist, using a variety of open habitats. He observed breeding in ditches, pools, potholes, and ponds with grasses and cattails, located within or adjacent to meadows, hayfields, woods, pastures, alder swamps, and "cattail swamps."

Habitat Types:

Hardwood Swamps
Seeps and Pools
Marshes and Sedge Meadows
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Shrub Swamps
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Impacts of Roads or Transportation Systems
Climate Change
Habitat Fragmentation
Invasion by Exotic Species

Description of habitat threat(s): Shallow wetland habitat appears to be limited and no strong population center known that would provide colonizers. Appears to be limited to NW Vermont. Known distribution retracted from northeastern shore of Lake Champlain (Canadian border to Georgia) to isolated site in Alburg. Also declines in Quebec and eastern Ontario. Needs early successional, open or edge habitat, does not compete well with Wood Frogs, genus has shown sensitivity to a chemical (emodin) released by invasive buckthorn (*Rhamnus cathartica*). Shallow open seasonal wetlands in the Lake Champlain Basin are particularly likely to be drained for agriculture. These lands are also exposed to regularly chemical use and mechanized cutting.

Non-Habitat Threats:

Genetics
Competition
Disease
Pollution
Harvest or Collection
Reproductive Traits
Predation or Herbivory



Common Name: **Boreal Chorus Frog**
Scientific Name: **Pseudacris maculata**
Species Group: **Herp**

Trampling or Direct Impacts

Description of non-habitat threat(s): Pesticides, herbicides, and runoff. may be inadvertently collected when Leopard Frogs are collected by sweep net or drift fence. Chytrid fungus or ranavirus may play a role in declines either alone or synergistically. Any remaining populations are likely genetically isolated. Call does not compete well with Spring Peeper (*Pseudacris crucifer*). More successful if isolated in time or space from their calls. Does not coexist well with Wood Frogs (*Lithobates sylvaticus*) perhaps as a result of predation. Modern agricultural harvesting equipment is probably too fast and efficient to coexist with populations.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Determine habitat needs.
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Thoroughly survey for this species in Vermont.
Research	Threats and Their Significance	High	Determine the primary limiting factors for this species.
Research	Population Genetics	Medium	Confirm species, Determine if genetically isolated.
Monitoring	Population Change	High	
Monitoring	Habitat Change	High	
Monitoring	Range Shifts	High	Determine how Vermont fits in with regional population change.
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Boreal Chorus Frog**
 Scientific Name: **Pseudacris maculata**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	Medium	Determine which species of Chorus Frog we have/had, how genetically isolated/inbred it is/was and what can be done to address issues if needed.	Genetic assessments	Academic Community, Province of Quebec, NY DEC	SWG, Nongame Fund, private grants
Research	Medium	Determine the presence/absences of amphibian diseases in the historic area of the Chorus Frog and to control their spread and impact.	Disease surveys	Regional Wildlife Health Lab, Academic community	SWG, Nongame Fund, Private grants
Invasive Species Control & Prevention	Medium	Examine presence/absence of invasives in historic habitat of this species. Conduct lab experiments of impacts of Emodin on this species.		Academic community, TNC	SWG, Nongame Fund, private grants
Protected Area Management	Medium	Manage selected areas for early succession, work with agriculturalists to manage lands in a way that maintains early successional habitat in a way that is amphibian friendly. Encourage light pasturing rather than harvesting and cultivating near wetlands.	Change in the acres of wetland/grassland mosaics	Sportsmen, Audubon, groups interested in early successional bird species	SWG, VHCB, DU,
Policy & Regulations	Medium	Improve agricultural practices that degrade or remove wetlands and surrounding habitat and replace and restore wetlands altered or lost by agriculture.	Wetland Acreage Conserved, Created, or Recreated	DEC, Environmental Organizations, LCI, Lake Champlain protection groups	Private Foundations, water quality grants
Policy & Regulations	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	ANR, Health Dept, USFWS	
Awareness Raising and Communications	Medium	Encourage reports of sightings to the VT Wildlife Diversity Program and the VT Reptile & Amphibian Atlas	Number of training sessions, person-nights surveyed, Number of reports received	Reptile & Amphibian Atlas	SWG, Nongame Wildlife Fund
Easements		Attempt to protect (through easement or purchase) any sites found with breeding populations and adjacent terrestrial habitat.	Acreage and number of sites conserved	Reptile & Amphibian Atlas, Lake Champlain Land Trust, VLT	VHCB

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Boreal Chorus Frog**
Scientific Name: **Pseudacris maculata**
Species Group: **Herp**

Bibliography

- Andrews, J. 2013. The Vermont Reptile and Amphibian Atlas. VtHerpAtlas.org
- Bider, J. R., and S. Matte. 1996. An atlas of reptiles and amphibians of Québec. St. Lawrence Valley Natural History Society and Ministère de l'Environnement et de la Faune, Québec. Montreal. 110 pages.
- Calhoun, A.J.K. and M. W. Klemens. 2002. Best Development Practices: Conserving pool-breeding amphibians in residential and commercial developments in the Northeastern United States. MCA Technical Paper No. 5, Metropolitan Conservation Alliance, Wildlife Conservation Society, Bronx, New York. 57 pp.
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- Committee on the Status of Endangered Wildlife in Canada (COSEWIC). 2014. Boreal Chorus Frog. [Ttp://www.cosewic.gc.ca](http://www.cosewic.gc.ca)
- Corser, J. D., K. J. Roblee, and G. Johnson. 2012. Shifting status and distribution of range margin chorus frog (*Pseudacris*) populations in eastern Great Lakes watersheds. *Journal of Great Lakes Research*, 38(4) 806-811.
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Dickerson, M.C. 1969. The frog book: North American toads and frogs, with a study of the habits and life histories of those of the northeastern states. Dover Publications, Inc., New York 253 pp.
- Dodd, C.K. 2013. Frogs of the United States and Canada. Johns Hopkins University Press, Baltimore, Maryland, 982 pp. in two volumes.
- Duellman, W.E. and L. Trueb. 1994. Biology of amphibians. The Johns Hopkins University Press, Baltimore, Maryland 670 pp. (The standard text for amphibians.)
- Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp.
- HerpNet: Amphibians and Reptiles of Minnesota. No date. Boreal chorus frog - *Pseudacris maculata*. Available <http://herpnet.net/Minnesota-Herpnetology/>.
- Heyer, W.R., M.A. Donnelly, R.W. McDiarmid, L-A. C. Hayek, and M.S. Foster. 1994. Measuring and monitoring biological diversity: standard methods for amphibians. Smithsonian Institution Press, Washington 364 pp.
- IMNH. 2000. *Pseudacris maculata* (Boreal Chorus Frog), Idaho Museum of Natural History. Available <http://imnh.isu.edu/digitalatlas/bio/amph/anurans/psma/Psmafra.htm>.
- Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp.
- Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp.
- Lannoo, M. 2005. Amphibian declines: the conservation status of United States species. Berkeley, CA Univ. of California Press. 1094 pages.
- Mitchell, J.C. 2000. Amphibian monitoring methods and field guide. Smithsonian National Zoological Park's Conservation & Research Center, Front Royal, Virginia 56 pp.
- Moriarty Lemmon, E., A. R. Lemmon, J. A. Lee-Yaw, J. T. Collins, and D. C. Cannatella. 2007. Phylogeny-based delimitation of species boundaries and contact zones in the trilling chorus frogs (*Pseudacris*). *Molecular Phylogenetics and Evolution* 44:1068-1082.
- NatureServe. 2011. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.0. NatureServe, Arlington, VA. U.S.A. Available <http://www.natureserve.org/explorer>. (Accessed: 10 June 2011)



Common Name: **Boreal Chorus Frog**
Scientific Name: **Pseudacris maculata**
Species Group: **Herp**

Sacerdote, A. B., and R. B. King. 2014. Direct effects of an invasive European buckthorn metabolite on embryo survival and development in *Xenopus laevis* and *Pseudacris triseriata*. *Journal of Herpetology* 48(1):51-58.

Semlitsch, R.D. and J.R. Bodie. Are small wetlands expendable? *Conservation Biology* 12(5):1129-1133.

Semlitsch, R.D. and J.R. Bodie. 2003. Biological criteria for buffer zones around wetlands and riparian habitats for amphibians and reptiles. *Conservation Biology* 17(5):1219-1228.

Stebbins, R.C. and N.W. Cohen. 1995. A natural history of amphibians. Princeton University Press, Princeton, New Jersey 316 pp

Wright, A.H. and A.A. Wright. 1995. Handbook of frogs and toads of the United States and Canada. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 640 pp.

Whiting, A. 2004. Population ecology of the Western chorus frog (*Pseudacris triseriata*). Master's Thesis, McGill University,

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp



Common Name: **Spotted Turtle**
Scientific Name: **Clemmys guttata**
Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The state-endangered Spotted Turtle is known from three locations in Vermont and is a species of regional conservation concern. It is on Appendix II of CITES, the IUCN lists it as endangered on its Red List, and is a SGCN in Vermont. Two Spotted Turtle sites in Vermont are compromised by surrounding landuse. At one site a RR bisects the wetland. Twenty under rail crossings have been installed to allow safe passage of the turtles. A road parallels much of the other compromised site. Both are believed to have very small populations. The third known site is more robust and found within a large wetland complex, which provides some buffering from surrounding landuse.

Distribution

There are three widely separated known populations in Vermont. The disjunct nature of the three known occurrences means that each needs to maintain itself with little likelihood of rescue from nearby populations. It is possible that some migrants from Massachusetts could reach one of our southern populations. The species is not a migrant but some individuals do wander. Based on limited telemetry work in Vermont it may be young adult males are more prone to dispersal. Because of the vulnerability of this species to illegal collection and the protections of our state law, specific locations are not being shared in this summary.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Hudson-Hoosic
Lake Champlain
Middle Connecticut

Probable Watersheds

NA

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Based on field inventory and monitoring, the Spotted Turtle uses red maple swamps and other wetland habitats in Vermont. In Massachusetts it has been documented to patch together woodland vernal pools in some



Common Name: Spotted Turtle
Scientific Name: *Clemmys guttata*
Species Group: Herp

localities and can also use ponds. The two smaller Vermont sites are linear riparian wetlands. The third and larger population is part of a large wetland complex. Nesting by this species is reported in the literature to include sphagnum moss and at the larger site a hatchling was documented on a large sphagnum mat. Some sphagnum occurs at one of the smaller sites, but nesting has been documented along a railroad bed. At the other smaller site two female Spotted Turtles have been located near a dirt road during the nesting season.

Potential wetland habitats include Hardwood Swamps, Marshes and Sedge Meadows, Shrub Swamps, Open Peatlands. We have located this species in red maple swamps, shrub swamps, and a bog/fen wetland as well as an adjacent emergent swamp. In Vermont we have found this species at low elevation sites less than 100 m in elevation. Home range is often reported as less than 5ha, but linear movements can be quite long. At one riparian wetland a female Spotted Turtle moved over 300 m. We believe large wetland complexes provide the best habitat by buffering from outside influences and providing options during different moisture regimes. This may become more important with weather patterns linked to climate change. Movement corridors are important in some situations where Spotted Turtles piece together needed resources within a habitat mosaic and this has been documented in Massachusetts with turtles traveling between a large wetland and dispersed vernal pools in surrounding uplands. Long-distance movements are unlikely and assisted migration may need to be considered in the future.

Habitat Types:

Oak-Pine Northern Hardwood
Hardwood Swamps
Softwood Swamps
Seeps and Pools
Northern Hardwood
Open Peatlands
Marshes and Sedge Meadows
Wet Shores
Shrub Swamps
Aquatic: Lacustrine
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Habitat Fragmentation
Impacts of Roads or Transportation Systems
Climate Change

Description of habitat threat(s): Habitat threats for the Spotted Turtle include loss of wetland overwintering and foraging habitat, loss of nesting areas, fragmentation isolating populations and separating needed seasonal habitats, as well as road and railroad mortality. Climate change could affect the Spotted



Common Name: **Spotted Turtle**
 Scientific Name: **Clemmys guttata**
 Species Group: **Herp**

Turtle if a change in water regime results in wetlands and pools drying.

Non-Habitat Threats:

Genetics

Harvest or Collection

Reproductive Traits

Predation or Herbivory

Trampling or Direct Impacts

Description of non-habitat threat(s): The Spotted Turtle is vulnerable to collection, nest predation, road mortality, and population rescue is unlikely. If close to humans, risks increase.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	1) Given the rarity of this species, examine critical habitat (e.g., wintering sites). 2) Gather data on specific habitat requirements of Vermont populations: denning sites, egg-laying sites, foraging areas, overwintering areas and movement corridors.
Research	Distribution and Abundance	High	Pursue further surveys for Spotted Turtles in Vermont. Need to document all populations and their abundance, as well as the spatial relationship of populations.
Research	Threats and Their Significance	High	Need to identify and resolve potential limiting factors.
Research	Population Genetics	Medium	We don't know if population genetics is a problem or not, but species is isolated in Vermont.
Monitoring	Population Change	High	1) Track population size, age and sex distribution. 2) Monitor the size and determine the sustainability of existing populations through age-class or genetic analysis.
Monitoring	Habitat Change	High	It would be important to track changes in quality and quantity of habitat, as well as connectivity between habitats.
Monitoring	Monitor Threats	High	We need to determine if we are making progress alleviating limiting factors.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Spotted Turtle**
 Scientific Name: **Clemmys guttata**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks		With staff turnover try assigning responsibility for species recovery to new VFWD employees to address limited staffing realities.	Recovery plans written and implemented, actions taken, results documented	ESC, Orriane Society	SWG, Nongame Fund, Orriane Society
Compatible Resource Use		Control ATV and other off road usage in known habitat to avoid impacts.	Frequency of ATV use	Railroad, Game Wardens, VASA	operating funds
Compatible Resource Use		Continue to work with landowners and users of area to protect known habitat.	Number of cooperating landowners	local game warden, landowners, country forester	SWG, Nongame Wildlife Fund
Habitat Restoration		Create nesting sites and passages connecting wetland habitats. (e.g. Railroad used American Recovery and Reinvestment Act funds to construct 20 passages in 2012).	Number of enhancements (e.g., 20 passages under rails and 13 nesting pits created at one site).	railroad, local warden	railroad, Nongame Wildlife Fund
Standards		Develop land management guidelines for owners and managers of appropriate habitat and make them readily available through multiple media, including print and the web.	Number of landowners and managers who receive and use guidelines	landowners, consulting foresters, EQIP biologists	SWG, EQIP, Current Use
Awareness Raising and Communications		Encourage reports of sightings to the VT Natural Heritage Inventory and the VT Reptile & Amphibian Atlas.	Number of reports received annually	Reptile & Amphibian Atlas	Nongame Wildlife Fund
Species Restoration		Consider reintroduction or augmentation from closest healthy source. Maintaining and enhancing extant populations is always a priority and should be continued.	number of populations.	Bonnyvale Environmental Center	SWG
Species Restoration		Protect nests and adults by predator trapping and removal.	number of raccoons harvested per year.	Trappers Assoc., landowners	Trappers Assoc members
Easements		Conserve known habitat through fee simple purchase, development rights or easements, management agreements, and education of private landowners and managers regarding appropriate management.	acres of land conserved.	VHCB	VHCB
Awareness Raising and Communications		Help Vermonters assign value to this species through educational programs, printed material, web site information, field trips, TV and video information.	Number of people exposed to message	VFWD Outreach division, Bonnyvale Environmental Center	marketing funds, private grants

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Spotted Turtle**
 Scientific Name: **Clemmys guttata**
 Species Group: **Herp**

Habitat Restoration	Work to maintain connectivity with populations to the south in Massachusetts.	Number of potential connections to populations south of Vermont	Mass Fish and Wildlife, private landowners, VLT	VHCB
Compatible Resource Use	Review all roadway projects in appropriate habitat, check against known crossing areas VTRANS, VFWD, VT Reptile & Amphibian Atlas, survey appropriate habitat when unknown.	Number of projects reviewed with spotted turtle planning information	Reptile & Amphibian Atlas, VTrans	VTrans
Awareness Raising and Communications	Encourage reports of road-killed specimens, road crossing, and road basking areas to VFWD, VTRANS, and the VT Reptile & Amphibian Atlas.	Number of reports.	Reptile & Amphibian Atlas, VTrans	VTrans, SWG



Common Name: **Spotted Turtle**
Scientific Name: **Clemmys guttata**
Species Group: **Herp**

Bibliography

- Andrews, J. 2014 or most recent. The Vermont Reptile and Amphibian Atlas. www.VTHerpAtlas.org
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- Carr, A. 1995. Handbook of turtles: The turtles of the United States, Canada, and Baja California. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 542 pp. (A reprint of an old classic.)
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Ernst, C.H., J.E. Lovich, and R.W. Barbour. 1994. Turtles of the United States and Canada. Smithsonian Institution Press, Washington D. C. 578 pp. (The best current source for detailed information.)
- Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)
- Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)
- Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.
- Joyal, L.A., M. McCollough, and M.L. Hunter 2001 Landscape ecology approaches to wetland species conservation: a case study of two turtle species in southern Maine. *Conservation Biology* 15(6):1755-1762.
- Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)
- Klemens, M. (ed.) 2000. Turtle conservation. Smithsonian Institution Press. Washington 334 pp. (A current discussion of conservation challenges.)
- Litzgus, J.D. and R.J. Brooks. 2000. Habitat and temperature selection of *Clemmys guttata* in a northern population.
- Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp. (This book provides excellent information on the species of reptile that we share with Virginia; most of our species are found in this book.)
- Semlitsch, R.D. and J.R. Bodie. Are small wetlands expendable? *Conservation Biology* 12(5):1129-1133.
- Semlitsch, R.D. and J.R. Bodie. 2003. Biological criteria for buffer zones around wetlands and riparian habitats for amphibians and reptiles. *Conservation Biology* 17(5):1219-1228.
- Terres, G.D. 1999. Wildlife species of regional conservation concern in the Northeastern United States. *Northeast Wildlife* 54:93-100.
- Tyning, T.F. 1990. A guide to amphibians and reptiles. Little, Brown and Company. Boston Massachusetts 400 pp

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Wood Turtle**
Scientific Name: **Glyptemys insculpta**
Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: S3

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

With more than 50% of its distribution in the Northeast the Wood Turtle was selected as a Regional Species of Greatest Conservation Need (RSGCN) among the 13 Northeastern states. Vermont is at the core of its range. The Wood Turtle is listed as an Appendix II species by Convention on International Trade in Endangered Species (CITES), meaning that it not necessarily threatened with extinction, but in which trade must be controlled in order to avoid utilization incompatible with their survival. It has been petitioned in the past for listing as a federally listed species. The Wood Turtle is believed to still be widely distributed, but potentially declining throughout its range due to its long age to maturity and increasing losses of long-lived adults. It is still encountered over a wide region, which may be masking the conservation peril this species faces. Two populations have been monitored in Vermont (Parren 2013) and Vermont was included in a Northeast range-wide assessment (Jones et al. 2014). This species is getting regional recognition as an RCN species.

The Wood Turtle is found in riparian habitats and upland habitats adjacent to streams, providing it is not heavily developed or intensively used for agriculture. It is vulnerable to mowing of agricultural fields and to road mortality. Habitat loss due to succession is also a challenge for this species, but it seeks out and uses even small suitable patches within woodlands. It is also at risk to collection.

Distribution

The Wood Turtle is well distributed throughout Vermont, but not at very high elevations and rarely encountered at the lowest levels of the Champlain Valley. A suitable medium gradient river or large creek is required.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Known Watersheds

Middle Connecticut
West
Upper Connecticut-Mascoma
Black-Ottawquechee
Hudson-Hoosic

Probable Watersheds

Waits
Upper Connecticut-Mascoma
Passumpsic
St. Francois River
Upper Connecticut



Common Name: **Wood Turtle**
Scientific Name: **Glyptemys insculpta**
Species Group: **Herp**

Mettawee River
Lake Champlain
Lamoille River
Missiquoi River
Otter Creek
White
Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

The Wood Turtle is found in a landscape of rolling hills. Upland habitat adjacent to streams needs to be permeable to Wood Turtle terrestrial wanderings. It uses shrub swamps, alder swamps, and can use human-altered landscapes if not too severely changed, but it must have a suitable home stream. Home range can be calculated as less than a hectare if long-distant movements for females to nest or for males to patrol a river or stream to breed females (1.5 km) are not included. Micro Habitat: moderate gradient streams with refuge sites, sand gravel & rock streambeds.

Habitat Types:

Oak-Pine Northern Hardwood
Hardwood Swamps
Softwood Swamps
Seeps and Pools
Spruce Fir Northern Hardwood
Northern Hardwood
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Early Succession Pine and Hemlock
Early Succession Upland Oak
Early Succession Other Types
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Wet Shores
Shrub Swamps
Early Succession Northern Hardwoods
Aquatic: Fluvial



Common Name: **Wood Turtle**
Scientific Name: **Glyptemys insculpta**
Species Group: **Herp**

Aquatic: Large Lake Champlain Tribs Below Falls

Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Succession

Habitat Alteration

Habitat Fragmentation

Incompatible Recreation

Description of habitat threat(s): Loss of upland habitat adjacent to Wood Turtle home stream is real problem. It can tolerate a fair amount of succession providing some suitable patches remain. Although the Wood Turtle can use an agricultural landscape, too intensive a use such as row crops, is unsuitable. Mowing and driving over fields can directly impact Wood Turtles found in agricultural fields (Erb and Jones 2011). Habitat is being broken up by development and roads. Trails can bring more people into contact with Wood Turtles and their populations are often impacted (Graber and Burger 1995). Roads paralleling Wood Turtle streams directly impact Wood Turtle populations and can over time reduce the use of habitat by removal of turtles (Parren 2013). Wood Turtles seem well adapted to finding food resources within a broad habitat matrix, but this requires wandering a large area with associated risks.

Non-Habitat Threats:

Harvest or Collection

Predation or Herbivory

Trampling or Direct Impacts

Description of non-habitat threat(s): Wood Turtles are susceptible to collection as pets and we have detected for profit illegal collection in Vermont. Nest depredation is a threat all turtles face and Wood Turtles also suffer limb and tail loss to predators. In some cases direct mortality occurs. Road traffic causes Wood Turtle mortality and the roads themselves can be thought of as a landscape feature limiting Wood Turtle habitat (see above). Proximity to human habitation and our land use is a risk to Wood Turtle survival due to collection, direct mortality from vehicles and other equipment, as well as increased predators such as skunks and raccoons.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Wood Turtle**
 Scientific Name: **Glyptemys insculpta**
 Species Group: **Herp**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	1) Identification and protection of communal wintering and nesting sites is critical. 2) Gather data on specific habitat requirements of Vermont populations: denning sites, egg-laying sites, foraging areas, and movement corridors.
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Identify distribution and relative abundance of populations in Vermont as good knowledge of current distribution and abundance is essential for good monitoring.
Research	Threats and Their Significance	High	It is important to have a solid understanding of limiting factors and how they impact populations of Wood Turtles.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	High	It is essential to monitor population change for this vulnerable species, consider doing so by monitoring the size and determine the sustainability of existing populations through age-class or genetic analysis.
Monitoring	Habitat Change	High	Since habitat loss/change affects the resilience of Wood Turtle populations, it is important to monitor habitat change.
Monitoring	Range Shifts	High	The VT Reptile & Amphibian Atlas has provided a pretty good baseline of Wood Turtle distribution by township. Failure to detect Wood Turtles in these same townships in the future will be cause for alarm.
Monitoring	Monitor Threats	High	Road mortality, collection, and nest success should be monitored.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Wood Turtle**
 Scientific Name: **Glyptemys insculpta**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Compatible Resource Use		Encourage holding off field mowing until October in Wood Turtle habitat or setting mowing bar at 6 inches or higher.	Number of areas adhering to mowing guidance	Corps of Engineers, EQIP biologists	Corps of Engineers, EQIP
Compatible Resource Use		Encourage land-use practices on private lands that continue to allow the Wood Turtle to maintain itself in Vermont.	Number of enrolled landowners	FPR, EQIP, USFWS, private landowners	Current Use Program, EQIP, Partners in Wildlife
Easements		Conserve known habitat through fee simple purchase, development rights or easements, management agreements, and education of private landowners and managers regarding appropriate management.	Number of sites protected	VLT, local land trust, towns	VHCB
Compatible Resource Use		Direct trail development away from streams to avoid impacts to Wood Turtle populations.	Number of trails sited in a way to avoid impacts	recreation planners, developers, regulators	private grants, SWG, Nongame Wildlife Fund, Technical Assistance
Habitat Restoration		Develop, implement, and monitor, road crossing structures and barriers for this species.	Effectiveness of crossing structures	VTrans	VTrans, FHWA
Easements		Use conservation easements to protect suitable habitat on privately owned land.	Number of sites protected	VLT, local land trusts	VHCB, private funds
Species Restoration		If populations limited, consider reintroduction or augmentation from closest healthy source. Maintaining and enhancing extant populations is always a priority and should be continued.	Number of reintroduced or augmented populations	Landowners, Corps of Engineers, SAG-Herps	SWG
Compatible Resource Use		Site new roads 1000' away from Wood Turtle streams and avoid parallel roads.	Number of new roads sited away from streams	VTrans	VTrans
Publically-Owned Protected Areas		Protect suitable habitat on publicly owned land.	Number of sites on public land	ANR, TNC, USFS	state lands Mgmt funds, TNC, GMNF
Awareness Raising and Communications		Encourage reports of sightings to the VT Natural Heritage Inventory and the VT Reptile & Amphibian Atlas.	Number of reports	Reptile & Amphibian Atlas Project, Orriane Society	Nongame Wildlife Fund, SWG, Orriane Society

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Wood Turtle**
 Scientific Name: **Glyptemys insculpta**
 Species Group: **Herp**

Planning & Zoning	Include Wood Turtle habitat in town zoning to limit impacts from development.	Number of towns considering wood turtle habitat in zoning	Cons Comms, VLCT	SWG
Standards	Develop land management guidelines for owners and managers of appropriate habitat and make them readily available through multiple media, including print and the web.	Number of landowners and managers who receive and use guidelines	Reptile & Amphibian Atlas, EQUIP biologists	SWG, EQUIP
Compatible Resource Use	Manage ATV and other off road usage in known habitat to avoid impacts to Wood Turtles.	Number of sites where ATV use is managed	landowners, state and federal lands managers, VASA	state lands Mgmt funds, landowner decisions
Awareness Raising and Communications	Encourage reports of road-killed specimens, road crossing, and road basking areas to VFWD, VTRANS, and the VT Reptile & Amphibian Atlas.	Number of reports	VTrans, Reptile & Amphibian Atlas Project	VTrans

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Wood Turtle**
Scientific Name: **Glyptemys insculpta**
Species Group: **Herp**

Bibliography

- Andrews, J. 2014 or latest. The Vermont Reptile and Amphibian Atlas. www.VTHerpAtlas.org
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- Carr, A. 1995. Handbook of turtles: The turtles of the United States, Canada, and Baja California. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 542 pp. (A reprint of an old classic.)
- Compton, B.W., J.M. Rhymer, and M. McCollough. 2002. Habitat selection by wood turtles (*Clemmys insculpta*): an application of paired logistic regression. *Ecology*. 83(3):833-843.
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Erb, L., and M.T. Jones. 2011. Can turtle mortality be reduced in managed fields? *Northeastern Naturalist* 18:489–496. Ernst, C.H., J.E. Lovich, and R.W. Barbour. 1994. Turtles of the United States and Canada. Smithsonian Institution Press, Washington D. C. 578 pp. (The best current source for detailed information.)
- Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.
- Fowle, S. 2001. Guidelines for protecting wood turtles and their habitats in Massachusetts. Mass Div, Fish and Wildlife, Natural Heritage and Endangered Species Program.
- Garber, S.D. and J. Burger. 1995. A 20-yr study documenting the relationship between turtle decline and human recreation. *Ecological Applications*. 5(4): 1151-1162.
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)
- Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)
- Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.
- Jones, M. T., L. L. Willey, P. R. Sievert, and T. S. B. Akre. 2014. Status and Conservation of the Wood Turtle in the Northeastern United States. Final Report to the Regional Conservation Needs (RCN) Program. 289 pp.
- Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)
- Klemens, M. (ed.) 2000. Turtle conservation. Smithsonian Institution Press. Washington 334 pp. (A current discussion of conservation challenges.)
- Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp. (This book provides excellent information on the species of reptile that we share with Virginia; most of our species are found in this book.)
- Parren, S.G. 2013. A Twenty-Five Year Study Of The Wood Turtle *Glyptemys insculpta* In Vermont: Movements, Behavior, Injuries, And Death. *Herpetological Conservation and Biology* 8(1):176–190.
- Saumure, R.A. and J.R. Bider. 1998. Impact of agricultural development on a population of wood turtles (*Clemmys insculpta*) in southern Quebec, Canada. *Chelonian Conservation and Biology*. 3(1):37-45.
- Terres, G.D. 1999. Wildlife species of regional conservation concern in the Northeastern United States. *Northeast Wildlife* 54:93-100.
- Tyning, T.F. 1990. A guide to amphibians and reptiles. Little, Brown and Company. Boston Massachusetts 400 pp.
- Walde, A.D., J.R. Bider, C. Daugkem D. Masse, J. Bourgeois, J. Jutras, and R.D. Titman. 2003. Ecological aspects of a wood turtle, *Glyptemys insculpta*, population at the northern limit of its range in Quebec. *Canadian Field-Naturalist* 117(3):377-388.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Musk Turtle (Stinkpot)**
Scientific Name: **Sternotherus odoratus**
Species Group: **Herp**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S2

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

The Musk Turtle is only recorded from the Champlain Valley of Vermont in about ten townships. It is small with limited dispersal capabilities. Home range is less than 2ha. Water chestnut control harvesting is a potential limiting factor to this species.

This species might be impacted by shoreline development that leads to loss of aquatic vegetation and suitable nesting sites (Harding 1997). In Vermont, ANR staff involved in water chestnut harvesting are aware of the potential to impact Musk Turtles that might be collected with the vegetation, and operate the harvester slowly and are on the lookout for turtles.

It is very easy to overlook Musk Turtles and they can be mistaken for Painted Turtles when observed on a road.

Distribution

Musk turtle reports are clustered in Colchester-Milton-Grand Isle, Ferrisburgh, and West Haven, Benson, Orwell, Castleton, Hubbardton, Sudbury.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Unknown
Champlain Hills	Unknown	Vermont Valley	Probable
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Unknown
Northeastern Highlands	Unknown		

Distribution by Watershed:

Known Watersheds

Lake Champlain

Probable Watersheds

Otter Creek

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Shallow permanent water of lakes and large ponds with aquatic vegetation, and large slow rivers. Shallow permanent water of lakes and large ponds with aquatic vegetation, and large slow rivers. This species is sometimes referred to as a bottom walker. It nests under logs and at the debris line of shorelines. It is largely restricted to the lowest elevations in Vermont within the Champlain Valley (<60m elevation).



Common Name: **Eastern Musk Turtle (Stinkpot)**
Scientific Name: **Sternotherus odoratus**
Species Group: **Herp**

Habitat Types:

Marshes and Sedge Meadows
Shrub Swamps
Aquatic: Lacustrine
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat

Description of habitat threat(s): Development of shoreline that impacts nesting sites and aquatic vegetation.

Non-Habitat Threats:

Trampling or Direct Impacts

Description of non-habitat threat(s): Mechanical harvesting of water chestnut. Literature suggests anglers sometimes persecute. Might be subject to some collection as pets but it is not well documented.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Gather data on specific habitat requirements of Vermont populations: nesting sites, foraging areas, over wintering sites.
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Conduct statewide survey of Musk Turtle in Vermont. Identify distribution and relative abundance of populations in Vermont.
Research	Threats and Their Significance	Medium	Consider impacts of chestnut harvesting and shoreline development.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	High	Once baseline is established it would be helpful to monitor population over time.
Monitoring	Habitat Change	High	It would be good to detect habitat change while there is still time to act.
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Musk Turtle (Stinkpot)**
 Scientific Name: **Sternotherus odoratus**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks		Encourage observation and rescue of Musk Turtles removed from lakes by mechanical weed harvesting. Develop a training program to train weed harvesters how to do this.	Number of harvest operations that properly screen for turtles.	DEC Water Chestnut Program, Towns that manage aquatic weeds, lake association	
Standards		Develop land and water management guidelines for owners and managers of appropriate habitat and make them readily available through multiple media, including print and the web.	Number of owners and managers who receive information.	ECHO Leahy Center for Lake Champlain	Lake Champlain Basin Program
Awareness Raising and Communications		Encourage reports of Musk Turtle sightings to the VT Natural Heritage Inventory and the VT Reptile & Amphibian Atlas.	Number of reports received.	Reptile & Amphibian Atlas	Nongame Wildlife Fund

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Eastern Musk Turtle (Stinkpot)
Scientific Name: *Sternotherus odoratus*
Species Group: Herp

Bibliography

- Andrews, J. 2014 or most recent. The Vermont Reptile and Amphibian Atlas. www.VTHerpAtlas.org
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- Booms, T.L. 1999. Vertebrate s removed by mechanical weed harvesting in Lake Keesus, Wisconsin. *Journal of Aquatic Plant Management* 37:34-36.
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Ernst, C.H., J.E. Lovich, and R.W. Barbour. 1994. Turtles of the United States and Canada. Smithsonian Institution Press, Washington D. C. 578 pp. (The best current source for detailed information.)
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)
- Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)
- Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.
- Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)
- Klemens, M. (ed.) 2000. Turtle conservation. Smithsonian Institution Press. Washington 334 pp. (A current discussion of conservation challenges.)
- Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp. (This book provides excellent information on the species of reptile that we share with Virginia; most of our species are found in this book.)
- Tennant, A. 2003. Snakes of North America: eastern and central regions. Lone Star Books, Lanham, Maryland. 605 pp. (One of a two excellent new snake resources.)
- Tyning, T.F. 1990. A guide to amphibians and reptiles. Little, Brown and Company. Boston Massachusetts 400 pp.
- Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp. (A standard text.).



Common Name: **Spiny Softshell Turtle**
 Scientific Name: **Apalone spinifera**
 Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The Spiny Softshell Turtle is restricted to Lake Champlain in VT and Province of Quebec, and is not found elsewhere in New England as wild populations. It is not known from the NY side of Lake Champlain. All other historical locations in Quebec have been lost and the Winooski River population in Vermont has been lost. Shoreline development has limited nesting and basking areas for this species.

The Softshell is a very strong swimmer, but it is restricted to habitats in or near Lake Champlain and is unlikely to gain immigrants from outside Lake Champlain. Habitat loss due to succession is also a problem for this species, but can be corrected with vegetative management and natural processes such as ice scour of nesting beaches. Increasing lake use is limiting this species directly by boat strikes and limiting habitat through lakeshore development, especially sea walls and riprap. Human disturbances limit basking and can chase nesting females away from nesting beaches. Egg predator populations are abnormally high due to lack of larger predators and trapping pressure, and are subsidized by humans in the form of increased food supplies (garbage, pet food, corn).

A sustained nest management effort appears to be reducing nest depredation and many hatchlings have been documented to have emerged from nests and others have been taken into captive care and then released. It is hoped that these efforts will be successful in enhancing the population of this state-threatened species in Lake Champlain. A recovery plan has been written and accepted by the Agency of Natural Resources (VFWD 2009). The ongoing nest management actions include monitoring of nest clutch size which should detect recruitment of surviving females as they enter the breeding population at about age 14.

Galois et al. (2002) reported home range size to be 3,200 ha (12.8 sq. miles) for females and 275 ha for the smaller males. Individual adult females are known to travel from the Bay Bridge to the Pike River, a distance of ~19km and some females move upstream to nest. The large home range estimates of adult females is a result of lengthy movements between seasonal habitats.

Distribution

The Softshell is currently restricted to Missisquoi Bay and surrounding areas of shore and lower reaches of rivers and creeks from Pike River to St. Alban's Bay and a smaller subpopulation is associated with the lower Lamoille River and surrounding lake.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		



Common Name: **Spiny Softshell Turtle**
Scientific Name: **Apalone spinifera**
Species Group: **Herp**

Distribution by Watershed:

Known Watersheds

Lake Champlain

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

The following is based on general and regional literature as well as extensive local knowledge. The Spiny Softshell Turtle is a highly aquatic species inhabiting lakes, larger rivers, and associated wetlands. In general, important habitat features include a soft bottom with some aquatic vegetation for foraging and escape cover and fallen trees with underwater limbs, sandbars, and mudflats for basking. Individual or group basking sites such as rocks, logs, mud, sandbanks, or floating debris are also considered important (Ernst et al. 1994). Graham and Graham (1997) found partially submerged dead tree trunks to be the preferred basking substrate in the Lamoille River. Nesting sites are generally free of vegetation, have adequate solar exposure, are well drained throughout the nesting and incubation periods and occur on open sand and gravel/shale pebble deposits. Hatchlings prefer small shallow puddles or shallow waters on the lee end of sandbars that provide warmer and quieter water than surrounding areas (Plummer 1977a). This preference may be attributed to one or a combination of the following: food resources, swimming ability, differences in thermal preferences, social interactions, and predator avoidance (Congdon et al. 1992). Hibernacula must provide well oxygenated water and be free of ice scour, human disturbance, and predators. Turtles during hibernation are particularly vulnerable as they live under the ice for several months with low metabolism and reduced activity. The importance of adequate overwintering sites cannot be overstated, particularly at our latitude.

Habitat Types:

Upland Shores

Marshes and Sedge Meadows

Wet Shores

Shrub Swamps

Aquatic: Lacustrine

Aquatic: Large Lake Champlain Tribs Below Falls

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Succession

Habitat Alteration

Habitat Fragmentation

Inadequate Disturbance Regime

Description of habitat threat(s): Shoreline development is the biggest habitat threat. Both nesting and



Common Name: **Spiny Softshell Turtle**
Scientific Name: **Apalone spinifera**
Species Group: **Herp**

basking substrates are lost. Natural processes that create and clear nesting areas along shoreline are now impaired. Dams can cut off upstream populations or even divide populations. Marinas or other deep water development can impact hibernacula.

Non-Habitat Threats:

Pollution

Predation or Herbivory

Trampling or Direct Impacts

Description of non-habitat threat(s): Softshell Turtles are subject to intense nest depredation due to high levels of nest predators and concentrated nesting in a few locations. The nesting sites are impacted by human disturbance during nesting by shoreline and water recreation and nearby camps. Activity on nesting beaches can directly impact eggs and we have documented one case of equipment being driven over nests and causing damage. Human disturbance limits basking opportunities and duration. Pollution may have played a role in the decline of the Winooski River population and there remains concern about contaminants in Lake Champlain and possible impacts from toxic blue-green algal blooms.



Common Name: **Spiny Softshell Turtle**
 Scientific Name: **Apalone spinifera**
 Species Group: **Herp**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	1) Determine feasibility of a Winooski River softshell population restoration. 2) Conduct habitat surveys and assessments that provide useful information about distribution, quality, and level of disturbance by humans.
Research	Basic Life History	Low	1) Continue studies monitoring individuals via radio-tagging in an effort to document habitat utilization and movements between those habitats among seasons and years. Movements of radio-tagged individuals will aid in our understanding of the extent of interchange between populations. 2) Develop emergence estimates based on the number of hatchlings produced from each nest, through either direct observation or the counting of eggshell fragments. 3) Nest success can be documented by monitoring nests and calculating the proportion of nests that successfully hatch young by the end of the nesting season.
Research	Distribution and Abundance	Medium	Have a pretty good handle on adults but not on juveniles. Document that recruitment of young into the breeding population is occurring.
Research	Threats and Their Significance	High	1) Assess recovered dead specimens for size, weight, length, age estimate, sex, and the cause of death determined. 2) Employ tracking boards and camera sets to determine what species are predated at a nesting site. 3) Investigate sensitivity to environmental contaminants.
Research	Population Genetics	Medium	Investigate if Lake Champlain softshells are genetically distinct as there is interest in the uniqueness of Lake Champlain turtle population. Some work has been conducted by UVM researchers.
Research	Other Research	High	All individuals captured for research should be measured, age estimated, sexed, and possibly marked via pit tags which would provide long-term information.
Monitoring	Population Change	High	Consider the use of genetic methods for investigating populations.
Monitoring	Habitat Change	High	
Monitoring	Range Shifts	High	
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Spiny Softshell Turtle**
 Scientific Name: **Apalone spinifera**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Alliance Development		Collaborate with other American and Canadian investigators, organizations, and agencies.	Number of interactions with partners.	Société de la faune et des parcs du Québec, Société d'Histoire Naturelle de la Vallée du St-Laurent	
Compliance & Enforcement		Protect high-use basking areas from human disturbance via on and offshore signage and law enforcement.	Number of sites with signage and patrol.	VFWD Outreach Div., game wardens	Nongame Wildlife Funds
Compliance & Enforcement		Protect nesting beaches from human disturbance during nesting season via on and offshore signage, law enforcement, and, if appropriate, visual screens.	Number of sites managed.	USDA Wildlife Service, game wardens	SWG
Technical Assistance, Training, Learning Networks		Encourage softshell habitat landowners to become monitors and land stewards of that habitat for the purpose of softshell conservation.	Numbers of cooperating landowners.	Lake Champlain Land Trust	Lake Champlain Land Trust, Nongame Wildlife Fund
Awareness Raising and Communications		Develop and place signage along important habitat areas.	Number of sites with signage.	USDA Wildlife Service, State Parks staff	SWG, Nongame Wildlife Fund
Species Restoration		Consider recruiting volunteers to monitor potential nesting sites during the nesting season in an effort to identify previously undocumented nesting sites.	Number of trained volunteers and hours expended.	Lake Champlain Land Trust, Audubon VT	volunteer effort
Species Restoration		Create nesting habitat in suitable areas close to water.	Number of created nesting area	Corps of Engineers, Missisquoi National Wildlife Refuge	SWG
Species Restoration		Explore other deterrents such as fencing (chain link/floppy), electric wire, discouraging winter denning near nesting sites, night shooting, and night patrols with a trained dog to lessen predation.	Number of sites where alternative methods employed.	USDA Wildlife Services	SWG

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Spiny Softshell Turtle
Scientific Name: *Apalone spinifera*
Species Group: Herp

Awareness Raising and Communications	Develop and place informational brochures at fishing license agents, marinas, fishing derbies, and State Parks and camping areas.	Brochures distributed	State Parks, VFWD Outreach Div.	Marketing funds, Nongame Wildlife Fund
Species Restoration	Removing debris and large rocks, provide sand-shale substrate, trim or remove shading brush, and dig out encroaching vegetation in old shale deposits at nesting beaches.	Area improved for nesting.	volunteers, Audubon VT, UVM students	volunteer time
Species Restoration	Consider headstarting young if their survival in the nest is compromised.	Number of young salvaged.	ECHO Center for Lake Champlain, Ecomuseum (Montreal)	volunteer effort
Species Restoration	Employ mammalian predator trapping programs at nesting beaches that exhibit a relatively high concentration of nests to reduce the number of nests predated.	Number of predators removed.	USDA Wildlife Services	SWG
Market Forces	Develop an incentives program for dairy farmers to halt the access and trampling of sandy shorelines by cows (i.e., provide farmers with large water tanks and electric fencing).	Number of sites where livestock trampling of shoreline controlled.	NRCS, Farm Bureau	NRCS
Species Restoration	Basking habitat could be created via floating platforms or permanent structures.	Number of sites benefiting from basking structures.	Normandea u Inc	VTrans bridge project
Awareness Raising and Communications	A Spiny Softshell Turtle reporting program that encourages the public to document softshell sightings should be part of the overall public outreach effort for this species.	Number of reports received.	Reptile & Amphibian Atlas project	Nongame Wildlife Fund
Awareness Raising and Communications	Incorporate softshells into existing Vermont Agency of Natural Resources fish and wildlife publications (i.e., law digest and fishing guide).	Number of times message is carried in ANR publications.	VFWD outreach division and commissioner	marketing funds
Species Restoration	Protect known habitats from disturbance: nesting, wintering, basking, foraging.	Number of sites protected.	Missisquoi National Wildlife Refuge, Société de la Faune et Des Parcs du Québec, TNC of Canada, FPR, VFWD	Lake Champlain Mgmt funds, Lake Champlain Basin Program funds, SWG
Awareness Raising and Communications	Recruit local volunteers for the purpose of monitoring nesting beaches and increasing the general public's awareness of Lake Champlain Spiny Softshells.	Number of volunteers and effort expended	Audubon VT	Nongame Fund, Audubon VT

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Spiny Softshell Turtle
Scientific Name: *Apalone spinifera*
Species Group: Herp

Species Restoration		When feasible, protect nests with ½" x ½" hardware cloth or vinyl-coated wire mesh cages by state and federal biologists in an effort to reduce the number of depredated nests.	Number of successful nests protected,	USDA Wildlife Services	SWG
Awareness Raising and Communications		Organize workshops at boat-ramps to educate anglers on turtle identification and fishing hook removal.	Number of workshops held and numbers of people who attend.	angler organizations, Lake Champlain Committee, LCI	LCI
Compliance & Enforcement		Identify areas that provide critical foraging habitat, particularly for juveniles, and protect vulnerable areas from human disturbance via on and offshore signage and law enforcement if the areas are reasonably small and identifiable.	Number of areas documented.	TNC of Canada, Normandeau Associates	VTrans bridge monitoring
Compliance & Enforcement		Develop and maintain internal communications with law enforcement and biologists to build awareness and support for turtle protection.	Number/frequency of exchanges of information.	Game wardens, fisheries biologists	SWG, Nongame Wildlife Fund
Habitat Restoration		When feasible, basking areas will be enhanced via natural (e.g., tree limbs and trunks) structures in an effort to increase basking surface area.	Number of basking areas	Missisquoi National Wildlife Refuge	refuge operating budget
Habitat Restoration		Improve water quality in Lake Champlain by reducing sources of existing pollution and prevent future pollution impacts.	Improvements in water quality	DEC, Farmers, Towns	Clean and Clear Program
Technical Assistance, Training, Learning Networks	Low	Identify significant road crossings and develop safe road crossings to reduce the potential for roadkill. Softshells rarely venture far from water so are less vulnerable to road mortality than other turtles.	Number of sites identified and crossings developed.	Reptile & Amphibian Atlas,	VTrans, FHWA
Species Restoration		Trap nesting areas that exhibit a relatively high concentration of nests in an effort to reduce the number of predated nests.	Number of areas trapped	USDA Wildlife Services	SWG
Compliance & Enforcement		Further develop program by which softshell sightings and/or harassment can be reported to Vermont's Wildlife Diversity Program.	Number of reports received	Reptile & Amphibian Atlas	Nongame Wildlife Fund
Policy & Regulations		Explore and implement legal protection to benefit the Spiny Softshell Turtle, including the establishment of a legal means of designating and protecting habitats critical for softshells, both on land and water.	Number of legal tools provided.	Game wardens	state general

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Spiny Softshell Turtle
Scientific Name: *Apalone spinifera*
Species Group: Herp

Publically-Owned Protected Areas	Pursue acquisition of those areas identified as important for maintaining and enhancing Spiny Softshell Turtles.	Number of sites and acreage conserved.	Lake Champlain Land Trust, Nature Conservancy of Canada	VHCB funds
Awareness Raising and Communications	Develop and distribute information to landowners of current and potential riverine and lakeside softshell habitat.	Numbers of landowners/camp owners contacted.	Lake Champlain Land Trust	Lake Champlain Land Trust, Nongame Wildlife Fund
Compliance & Enforcement	Monitor hibernacula when Spiny Softshells have congregated (September – May) to ensure disturbance is minimal.	Frequency of monitoring	Société de la faune et des parcs du Québec	Québec grant
Technical Assistance, Training, Learning Networks	Inform state biologists of potential problem for hibernacula (e.g., potential marina development) and take appropriate actions when a hibernaculum's physical characteristics and/or hibernating individuals are limited.	Number of actions taken to protect turtles and their habitat.	Act 250 coordinator, game warden, Missisquoi National Wildlife Refuge	

Bibliography

- Andrews, J. 2014 or most recent. The Vermont Reptile and Amphibian Atlas. www.VTHerpAtlas.org
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- Carr, A. 1995. Handbook of turtles: The turtles of the United States, Canada, and Baja California. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 542 pp. (A reprint of an old classic.)
- Congdon, J.D., S.W. Gotte, and R.W. McDiarmid. 1992. Ontogenetic changes in habitat use by juvenile turtles, *Chelydra serpentina* and *Chrysemys picta*. Canadian Field-Naturalist 106:241-248. DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Ernst, C.H., J.E. Lovich, and R.W. Barbour. 1994. Turtles of the United States and Canada. Smithsonian Institution Press, Washington D. C. 578 pp. (The best current source for detailed information.)
- Galois, P., M. Levéillé, L. Bouthillier, C. Daigle, and S. Parren. 2002. Movement patterns, activity, and home range of the eastern spiny softshell turtle (*Apalone spinifera*) in northern Lake Champlain, Québec, Vermont. Journal of Herpetology 36:402-411.
- Graham, T.E. and A.A. Graham. 1997. Ecology of the eastern spiny softshell, *Apalone spinifera spinifera*, in the Lamoille River, Vermont. Chelonian Conservation and Biology 2:363-369. Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)
- Klemens, M. (ed.) 2000. Turtle conservation. Smithsonian Institution Press. Washington 334 pp. (A current discussion of conservation challenges.)
- Plummer, M.V. 1977. Activity, habitat, and population structure in the turtle, *Trionyx muticus*. Copeia 1977:431-440.
- VFWD (Wells, D. and S. Parren) 2009 Vermont Spiny Softshell Turtle Recovery Plan (final)



Common Name: **Five-lined Skink**
Scientific Name: **Plestiodon fasciatus**
Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

The Five-lined Skink is an S1, state-endangered species, that is known from only three locations in West Haven, VT. All Five-lined Skink sites in Vermont feature talus and exposed rock within a mile of Lake Champlain. Total records of individual sightings number approximately 40. We have almost no data on their abundance and natural history in VT. We would benefit from data on distribution, behavior, seasonal movements, egg-laying sites, predators, food, population size, genetic heterozygosity and microhabitat requirements.

Distribution

The Five-lined Skink is known from three locations in West Haven, VT.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Mettawee River

Lake Champlain

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

The Five-lined Skink is known currently only on talus slopes and nearby cliff faces, exposed rocky ridges, and rocky shorelines. The ridges are composed off a mixture of ledge, broken rock, and scattered juniper or hardwoods. All known sites have a south or southwestern exposure, low elevation, nearby water, and relatively warm climates for Vermont. Anecdotal historic reports mention the use of exposed faces of old buildings near the above habitat and old mining areas. A skink sighting of this nature was recently documented in an abandoned shanty.

Known in Vermont only from West Haven/Benson along Lake Champlain where talus slopes and exposed cliff faces and ridges of low elevation are close to water. Critical habitat includes leaf litter and coarse woody debris mixed with exposed broken rock/ledge seems to be preferred.



Common Name: **Five-lined Skink**
Scientific Name: **Plestiodon fasciatus**
Species Group: **Herp**

Habitat Types:

Outcrops and Alpine
Cliffs and Talus
Northern Hardwood
Oak-Pine Northern Hardwood
Subterranean
Building or Structure
Mine
Other Cultural

Current Threats

Habitat Threats:

Habitat Succession
Habitat Alteration
Impacts of Roads or Transportation Systems
Habitat Fragmentation

Description of habitat threat(s): Five-lined Skinks need solar exposure, rock slides and fire may play a roll in keeping talus and ledges exposed. Development of lake-shore areas where the skink is found, removal of coarse woody debris, and introduction of cats could prove to eliminate local populations. Skinks may move from denning sites on talus to feeding areas nearby. Short-distance seasonal movements seem likely. They do not seem to move across open field but rather short distances from talus to cliff, ridge, field edge, or lake-shore. Moderate traffic, wide roads and large agricultural fields could limit movements. Future sources of coarse woody debris need to be maintained (old snags, large dead trees, etc.).

Non-Habitat Threats:

Genetics
Predation or Herbivory
Loss of Prey Base
Trampling or Direct Impacts

Description of non-habitat threat(s): Genetic isolation of very localized populations are potentially a problem. Predation by cats or other introduced or subsidized predators could be a problem. Insecticide use could impact their prey base. When cold, lizards move slowly. Excessive trampling (intensive agricultural, residential, or recreational use could be a problem). Sites may have become isolated by large agricultural fields. Continuous sunny and rocky edge habitat may connect one large meta-population (Bald Mountain, Austin Hill, and adjacent rocky shorelines and talus slopes). Two other known locations (Dresden Narrows, Benson) may be isolated.



Common Name: **Five-lined Skink**
Scientific Name: **Plestiodon fasciatus**
Species Group: **Herp**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	1) Gather data on specific habitat requirements of Vermont populations: denning sites, egg-laying sites, foraging areas, and movement corridors. 2) Identify critical habitat that includes basking sites.
Research	Basic Life History	Medium	
Research	Distribution and Abundance	High	1) Determine distribution and abundance in Vermont. 2) Survey anthropogenic sites such as old mines and talus piles in western Rutland County for this species. 3) Identify appropriate habitat in Western Rutland and Addison Counties from maps and photos.
Research	Threats and Their Significance	High	Recover any dead specimens or use other means to obtain genetic tissue samples as the basis for genetic assessment/demographic info.
Research	Population Genetics	High	Recover any dead specimens or use other means to obtain genetic tissue samples as the basis for genetic assessment/demographic info.
Monitoring	Population Change	High	Monitor the size and determine the sustainability of existing populations through age-class and/or genetic analysis.
Monitoring	Habitat Change	High	
Monitoring	Range Shifts	Medium	
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Five-lined Skink**
 Scientific Name: **Plestiodon fasciatus**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks		Train Rattlesnake researchers and game wardens to keep data on the sightings and habits of this species.	Number of cooperators who gather information on skinks.	game wardens, volunteers	SWG, TNC
Standards		Develop land management guidelines for owners and managers of appropriate habitat and make them readily available through multiple media, including print and the web.	Number of landowners and managers who receive and use guidelines	VFWD district biologists, consulting foresters	SWG
Technical Assistance, Training, Learning Networks	Medium	Train Rattlesnake "Responders" to collect and submit data on the sightings and habits of the five-lined skink.	Number of cooperators who gather information on skinks. Number of skink tissue samples collected.	Game Wardens, volunteers, VFWD district biologists	SWG, TNC
Awareness Raising and Communications		Help Vermonters assign value to this species through educational programs, printed material, web site information, field trips, TV and video information.	Number of people who receive message.	VFWD Outreach Division, TNC	Outreach marketing funds
Easements		Conserve known habitat through fee simple purchase, development rights or easements, management agreements and education of private landowners and managers regarding appropriate management.	Number of sites and acreage conserved.	TNC, VLT	VHCB funds
Awareness Raising and Communications		Encourage reports of sightings to the VFWD Wildlife Diversity Program and the Vermont Reptile and Amphibian Atlas.	Number of reports received	Reptile & Amphibian Atlas, TNC	Private grants, Nongame Wildlife Fund, SWG
Species Restoration		Work to maintain connectivity with populations to the west in New York State and between the two known populations. Collect tissue sample when/where possible for genetic assessment.	Quality and quantity of connecting habitat.	New York DEC, TNC, VLT	VHCB, TNC, SWG
Habitat Restoration		Experiment with habitat enhancement such as creating small openings in heavily shaded areas along the top of cliffs and talus slopes, dropping logs onto the talus, maintaining coarse woody debris and scattered cover.	Number of sites with active management that have been monitored.	TNC	TNC, SWG
Alliance Development		Continue to work cooperatively with important landowners such as the Nature Conservancy. Develop and maintain allies in local government and private citizens.	Number of joint meetings with partners.	TNC, landowners	various

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Five-lined Skink**
 Scientific Name: **Plestiodon fasciatus**
 Species Group: **Herp**

Compatible Resource Use	Manage ATV and other off road usage in known habitat to avoid impacts.	Number of sites where ATV use is controlled.	Landowners, TNC, Game Wardens, VASA	Land Mgmt funds
Awareness Raising and Communications	Keep cats away from known habitat. Discourage or re-direct residences away from known habitats.	Number of areas fenced or otherwise protected	Landowners, TNC	TNC
Awareness Raising and Communications	Encourage reports of road-killed specimens, road crossing, and road basking areas to VFWD, VTRANS, and the VT Reptile & Amphibian Atlas.	Number of sites reported	VTrans, Reptile & Amphibian Atlas, volunteers TNC	Vtrans, TNC
Species Restoration	If local populations are determined to be unsustainable, consider reintroduction or augmentation from closest healthy source. Maintaining and enhancing extant populations is always a priority and should be continued.	Number of extant sites	TNC, NYDEC	Private grant, SWG



Common Name: **Five-lined Skink**
Scientific Name: **Plestiodon fasciatus**
Species Group: **Herp**

Bibliography

Andrews, J. 2013. The Vermont Reptile and Amphibian Atlas. <http://cat.middlebury.edu/herpatlas/>

Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.

DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.

Gibbs, J.P., A. Breisch, P. Ducey, G. Johnson, J. Behler, and R.C. Bothner. 2007. The reptiles and Amphibians of New York State, Identification, Natural History, and Conservation. Oxford University Press, Oxford, NY 422pp.

Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)

Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)

Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp. (Visit the PARC website for more information: www.parcplace.org.)

Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)

Lougheed. 2004. The importance of cover rock in northern populations of the five-lined skink (*Eumeces fasciatus*). *Herpetologica* 60(3): 287-294.

Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp. (This book provides excellent information on the species of reptile that we share with Virginia; most of our species are found in this book.)

Smith, H.M. 1995. Handbook of lizards: lizards of the United States and Canada. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 557 pp. (A reprint of an old classic.)

Tyning, T.F. 1990. A guide to amphibians and reptiles. Little, Brown and Company. Boston Massachusetts 400 pp.

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp. (A standard text)



Common Name: **North American Racer**
 Scientific Name: **Coluber constrictor**
 Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1

State Trend:

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The North American Racer is currently known from only one site in Vermont. It had not been documented in Vermont since 1985 and had been feared extirpated until relocated in 2003. It is a very rare species in Vermont (S1) and was listed as threatened in 2005. Anecdotal historic reports in the southern Connecticut River Valley and on nearby ridges repeatedly speak of a North American Racer that was commonly seen in this area twenty-five or more years ago. Recent Reptile & Amphibian Atlas records include racer reports in a six-town area in this region although the species is currently known from only one site in VT. Focused Racer research from 2004-2007 identified eight individuals in Vernon/Gilford which were PIT-tagged and monitored. Subsequent survey efforts occurred annually but no animals were seen between 2008 - 2014 sparking added fears of extirpation. Two sightings were most recently documented in 2014 in Gilford. Habitat loss due to succession is likely negatively affecting this species. Since 2007, habitat improvement efforts directed specifically towards Racers have occurred at the Gilford I-91 weigh-station site.

Historically, this species probably expanded in numbers as Vermont's forests were cleared. Open pasture, fields mowed by hand or horse, or fields not mechanically mowed probably provided expanded habitat. Currently, with farm loss/abandonment over the past several decades, habitat favorable to Racers has declined significantly.

Distribution

The only known population of the North American Racer in Vermont uses early successional open ledge, grass, fern, and other herbaceous cover exclusively during the summer. It does move through short (30m) sections of woodlands between patches. It may move larger distances through woodlands to denning sites. Telemetry research on this species in 2004-2007 identified long, narrow home range movements covering some three miles, tightly associated with a major powerline and grass margins of the Interstate right-of-way. The animal is known to den along ledges with talus slopes and exposed rock in other northern locations. One former denning site in Vermont has been located. Recent racer habitat improvement efforts have focused on creating early successional (grass/shrub) habitat, creation of experimental hibernacula, egg-laying substrate and basking habitat at the animal's last known occurrence site in Gilford.

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Confident
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Historic Records Only
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Middle Connecticut

Probable Watersheds

West



Common Name: **North American Racer**
Scientific Name: **Coluber constrictor**
Species Group: **Herp**

Deerfield
Black-Ottawquechee
Hudson-Hoosic
Mettawee River
Lake Champlain
Otter Creek

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

The North American Racer is currently known only from a very limited portion of a two-town area in the Connecticut River Valley. The species primarily uses early successional and sunny habitat along low rocky ridges in warm portions of the state. It probably moves from ridges to adjacent open areas at lower elevations to feed during the summer months. It needs to get below frost line for denning. Overwintering mortality has been documented for this species. Connected mosaics of early successional habitat and rocky exposed ledges is probably required.

Habitat Types:

Outcrops and Alpine
Cliffs and Talus
Oak-Pine Northern Hardwood
Early Succession Pine and Hemlock
Subterranean
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Early Succession Upland Oak

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Habitat Fragmentation
Impacts of Roads or Transportation Systems

Description of habitat threat(s): This species probably expanded in numbers as early Vermont was cleared. Open pasture, fields mowed by hand or horse, or fields not mechanically baled probably provided expanded habitat. Loss of early successional habitat including small farms, increased row cropping, increasing speed and mechanization of mowing and bailing and increased parcelization are believed to have limited appropriate habitat.

Non-Habitat Threats:



Common Name: **North American Racer**
Scientific Name: **Coluber constrictor**
Species Group: **Herp**

Genetics

Disease

Reproductive Traits

Trampling or Direct Impacts

Description of non-habitat threat(s): Isolation of appropriate habitat patches and small population size may have led to genetic isolation. Snake Fungal Disease is a potential threat but as yet unknown in VT racers. Increasing mechanization and speed of mowing and bailing causes direct mortality. Increasing ATV use in rural areas is a direct problem. Direct persecution from encounters with humans and possibly dogs needs to be addressed.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	While the Racers' general requirements are known from literature, we lack full understanding of VT habitat utilization. Gather specific habitat requirement data for VT populations: denning sites, egg-laying sites, foraging areas, and movement corridors.
Research	Distribution and Abundance	High	1) Continue racer surveys at known sites and for additional populations. Identify distribution and relative abundance of populations. Look for/examine evidence of Rutland/Bennington county populations. 2) Identify appropriate southeastern VT habitat from maps, photos, aerial surveys, and ground survey and interviews in likely areas.
Research	Threats and Their Significance	High	Opportunistically capture racers and recover any dead specimens to assess health status, obtain demographic information and genetic tissue samples for analysis.
Monitoring	Population Change	High	1) Monitor the size and determine the sustainability of existing populations through age-class or genetic analysis. 2) Review pertinent literature to investigate/inform the possibility of augmentation from closest, healthy source population.
Monitoring	Habitat Change	High	Continue on-going habitat improvement/maintenance schedules and monitor weigh station sites for evidence of racer use/occupation.
Monitoring	Monitor Threats	High	Watch for Snake Fungal Disease in Racer populations.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **North American Racer**
 Scientific Name: **Coluber constrictor**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration		Continue to maintain and create early successional habitat. Create/enhance denning, basking, egg-laying habitat if limited. Monitor existing improvements for evidence of use. Educate private landowners about maintaining habitat in a snake friendly manner.	Number of acres and specific sites maintained or enhanced.	VELCO, local landowners, VFWD district biologists, VTrans	VELCO, VFWD, VTrans, SWG
Alliance Development	Medium	Continue to work cooperatively with organizations and individuals in southeastern Vermont. Develop and maintain allies.	Number of partners.	Bonnyvale Environmental Center, local conservation commission, landowners, consulting foresters, VTrans, Reptile & Amphibian Atlas	SWG
Awareness Raising and Communications		Educate landowners in area about snakes in general and encourage coexistence with snakes. Inform them about the identification, natural history, and conservation problems and needs of this species.	Number of people who receive message.	VFWD Outreach Division	marketing funds, SWG
Species Restoration		If local populations are determined to be unsustainable, consider augmentation from closest healthy source. Maintaining and enhancing extant populations is always a priority and should be continued.	Number of extant populations.	Mass F&W, NH F&W	SWG
Planning & Zoning		Review all roadway projects in appropriate habitat, check against known crossing areas VTRANS, VFWD, VT Reptile & Amphibian Atlas, survey appropriate habitat when unknown.	Number of projects that utilize racer information.	VTrans, Reptile & Amphibian Atlas	FHWA, SWG
Awareness Raising and Communications	Medium	Encourage reports of sightings to the VT Heritage Inventory and the VT Reptile & Amphibian Atlas.	Number of reports received.	Reptile & Amphibian Atlas	Nongame Wildlife Fund
Awareness Raising and Communications		Help Vermonters assign value to this species through educational programs, printed material, web site information, field trips, TV and video information.	Number of people who received message.	VFWD Outreach Division	Nongame Wildlife Fund, SWG

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: North American Racer
Scientific Name: Coluber constrictor
Species Group: Herp

Standards	Develop land management guidelines for owners and managers of appropriate habitat and make them readily available through multiple media, including print and the web.	Number of managers and landowners who receive message.	Consulting foresters, VELCO, VTrans	SWG
Technical Assistance, Training, Learning Networks	Establish a web site with conservation information on this species and trained local contacts who can relocate snakes.	Establishment of web site containing information on racer.	VFWD Outreach Division, VTrans, Bonnyvale Environmental Center, Reptile & Amphibian Atlas	VTrans, VFWD marketing funds
Habitat Restoration	Develop, implement, and monitor, road crossing structures and barriers for this species.	Number of structures installed.	VTrans	FHWA
Easements	Conserve know habitat through fee simple purchase, development rights or easements, management agreements, and education of private landowners and managers regarding appropriate management.	Number of areas conserved.	consulting foresters, landowners, VLT, Vtrans, local land trusts	VHCB, FHWA
Species Restoration	Work to maintain connectivity with populations to the south in Massachusetts.	Maintenance of connectivity.	VTrans, VFWD, Mass Highway Dept. Town Planning Commission	FHWA, Vtrans
Awareness Raising and Communications	Encourage reports of road-killed specimens, road crossing, and road basking areas to VFWD, VTRANS, and the VT Reptile & Amphibian Atlas.	Number of reports received.	VTrans, Reptile & Amphibian Atlas	Nongame Wildlife Fund
Compatible Resource Use	Manage ATV and other off road usage in know habitat to avoid impacts.	Number of areas where ATV use is controlled.	Landowners, VELCO, VTrans, VASA	VELCO, VTrans
Awareness Raising and Communications	Quickly and thoroughly, counter myths and misinformation appearing in the press that may limit this species.	Number of press articles. Numbers of individuals who received message	Bonnyvale Environmental Center, Reptile & Amphibian Atlas	Private funds and grants

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: North American Racer
Scientific Name: *Coluber constrictor*
Species Group: Herp

Bibliography

Andrews, J. 2013. The Vermont Reptile and Amphibian Atlas. <http://cat.middlebury.edu/herpatlas/>

----2012. "Rare Snakes of Vermont- Census/Monitoring/Tagging of Rare Snakes in Rutland and Windham Counties, VT". Final Report- April 27, 2012. State Wildlife Grant Award #06120FY10183.

----2008. "North American Racer (*Coluber constrictor*) use of state lands in Guilford, VT, 2008". Final Report - Project Guilford IR 091-1(25). Vtrans.

Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.

DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.

Ernst, C.H. and R.W. Barbour. 1989. Snakes of eastern North America. George Mason University Press, Fairfax, Virginia. 282 pp. (An excellent source for detailed information.)

Ernst, C. H., and E. Ernst. 2003. Snakes of the United States and Canada. Smithsonian Institution Press, Washington D.C. 668 pp. (The latest and most complete source for snakes.)

Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.

Fitch, H..S. 1963. Natural History of the Racer. University of Kansas Publications, Museum of Natural History, Lawrence Kansas Vol 15, No. 8; pp. 351-468

Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)

Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp. (Visit the PARC website for more information: www.parcplace.org.)

Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)

Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp. (This book provides excellent information on the species of reptile that we share with Virginia; most of our species are found in this book.)

Tennant, A. 2003. Snakes of North America: eastern and central regions. Lone Star Books, Lanham, Maryland. 605 pp. (One of a two excellent new snake resources.)

Tyning, T.F. 1990. A guide to amphibians and reptiles. Little, Brown and Company. Boston Massachusetts 400 pp.

Wright, A.H. and A.A. Wright. 1994. Handbook of snakes of the United States and Canada, volumes 1 and 2. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 1105 pp. (A reprint of an old classic.)

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp. (A standard text)



Common Name: **Eastern Ratsnake**
 Scientific Name: **Pantherophis alleghaniensis**
 Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2

State Trend: Declining

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

The Eastern Ratsnake is a S2 species that is listed as state-threatened. Development, habitat fragmentation, road mortality, and direct persecution limit both Ratsnake populations in Vermont. The northern population appears to be entirely isolated. Anecdotal reports strongly suggest that both populations are declining.

Distribution

The Eastern Ratsnake is known from only two regions of VT. One meta-population can be found in western Rutland County and extending into southwestern Addison County. The second population is very localized on the border of Monkton, Bristol, and New Haven. The southern population is essentially bounded on the south by Route 4, on the west by Route 30, and on the north by Route 73 with an extension on Bald Hill in Sudbury reaching across Otter Creek into Leicester. The northern population is essentially bordered on the south by Plank Road, on the west by North Street (and the adjoining wooded swamp) on the north by Piney Hill Road and on the East by the Monkton-Bristol Road. Recent work in Vermont (Andrews 2012) has shown that individual snakes migrate at least 1.5 miles to and from hibernacula. Many, but not all, of these dens are on or immediately below south or southwest facing cliffs along the shore of Lake Champlain or similar but more interior cliffs and talus slopes. At least one population must use less-exposed ledges due to an apparent lack of cliffs within known migration distances.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Probable Watersheds

Lake Champlain
 Metawee River
 Otter Creek



Common Name: **Eastern Ratsnake**
Scientific Name: **Pantherophis alleghaniensis**
Species Group: **Herp**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

This species dens in cracks and caves on cliff faces, in rocky talus slopes often at the base of cliffs, and possibly in rocky woodlands and along ledges at low elevation (<400m) with a southern or southwestern exposure. Many of these dens are along the shore of Lake Champlain or similar but more interior cliffs and talus slopes are also used. At least one population must use less-exposed but deeply eroded ledges due to an apparent lack of cliffs within known migration distances.

From these dens individuals travel distances of 1.5 miles or more to summer foraging areas that consist primarily of cliff tops, field edges, old fields, old orchards, abandoned or seldom used buildings and barns, or wetland margins. Large exposed dead or partly dead hollow trees along field edges, cliff tops, on talus slopes, or along margins or wetlands and water bodies are often used as refuges, as well as basking and feeding areas. Habitat mosaics including rocky slopes, cliffs, large dead and hollow trees, old fields, wetlands, and old buildings are ideal. Areas that also include abundant amphibians such as Northern Leopard Frogs seem to be most frequented. Along Lake Champlain while traveling between dens and foraging areas, ratsnakes spend most of their travel time within the thickly-vegetated shoreline margin of the lake. In other areas they appear to use densely vegetated corridors and hedgerows as their travel corridors. They also return to favored barns and protected refugia year after year.

Appropriate denning areas (talus, rock crevices) with spring basking opportunities, low elevation warm rocky woodlands, and safe connectivity to woodland or edge foraging areas with low human and road density appear to be important. Standing hollow snags with good sun exposure provide feeding, protection, and basking.

Habitat Types:

Upland Shores
Outcrops and Alpine
Cliffs and Talus
Hardwood Swamps
Oak-Pine Northern Hardwood
Marshes and Sedge Meadows
Early Succession Upland Oak
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Building or Structure
Mine
Other Cultural

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession



Common Name: **Eastern Ratsnake**
Scientific Name: **Pantherophis alleghaniensis**
Species Group: **Herp**

Habitat Alteration

Impacts of Roads or Transportation Systems

Habitat Fragmentation

Climate Change

Description of habitat threat(s): This species often travels relatively long distances from den sites for foraging areas. As development continues, this travel may require movement through increasingly mechanized agricultural areas, and across increasingly busy roads. This movement in combination with their affinity for edge habitat, old buildings, barns, and outbuildings often brings them into contact with humans. Increasing development brings snakes into contact with more humans, and their machines.

Non-Habitat Threats:

Genetics

Disease

Reproductive Traits

Trampling or Direct Impacts

Description of non-habitat threat(s): Some populations appear to be completely isolated, others may become so. This is a long-lived species, consequently direct persecution from humans and increased road mortality of adult breeders can outpace production. Roads attract and hold cold snakes as basking areas. Increasing road density and traffic are a problems. Increased ATV use in and near woodland fragments is known to cause mortality to snakes basking in trails. Increasing efficiency and speed of farm equipment for the planting and harvest of crop lands increases mortality from this source. Snake fungal disease has recently been identified in this area and is known to cause mortality in ratsnakes. It may have impacts on our populations but the extent of and impacts from the disease in Vermont are currently unknown.



Common Name: **Eastern Ratsnake**
 Scientific Name: **Pantherophis alleghaniensis**
 Species Group: **Herp**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Determine food requirements, denning sites, nesting locations, foraging areas, movement corridors, annual range, and other important natural history information that can be used to better protect and/or enhance habitat.
Research	Basic Life History	Medium	
Research	Distribution and Abundance	High	1) Identify distribution and relative abundance (population sizes) of populations in Vermont. 2) Survey all areas from which reports have originated in the last ten years.
Research	Threats and Their Significance	High	Is this species susceptible to or carrying Snake Fungal Disease or other diseases in Vermont? Are there specific locations where snake fencing and underpasses would benefit this species?
Research	Population Genetics	Medium	Is the northern population in danger due too limited genetic diversity?
Monitoring	Population Change	High	Are populations associated with any of the dens declining? If so, at what rate?
Monitoring	Habitat Change	High	How and at what rate is habitat changing? Is traffic increasing?
Monitoring	Range Shifts	High	Has this species disappeared from any historic range. Is it moving into any previously uninhabited range?
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Ratsnake**
 Scientific Name: **Pantherophis alleghaniensis**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications	High	Educate landowners and residents within the snake's range to encourage coexistence with snakes.	Number of programs and number of people who receive message.	landowners, Cons Comms	SWG, Private grants
Technical Assistance, Training, Learning Networks	High	Work with VTrans crew and other land managers to raise awareness of conservation need and implement conservation actions that benefit snakes.	Number of crew members who receive training	VTrans	VTrans training funding
Alliance Development	High	Continue working cooperatively with important landowners such as The Nature Conservancy. Develop and maintain allies in local government and private citizens.	Number of joint meetings held with partners.	TNC, landowners, towns	SWG, Nongame Wildlife Fund
Natural Processes Restoration	Medium	Develop, implement, and monitor, road crossing structures and barriers for this species.	Number of structures installed	VTrans, Academic community	SWG, VTrans, FHWA
Species Restoration	Medium	Work to maintain connectivity with populations to the west in New York State and between known populations.	Quantity (acorage) and quality of connective habitat.	NY DEC, VLT, Champlain Land Trust	VHCB
Standards	Medium	Review all roadway projects in appropriate habitat, check against known crossing areas VTRANS, VFWD, VT Reptile & Amphibian Atlas, survey appropriate habitat when unknown.	Number of projects where ratsnake information was used for planning	VTrans, Reptile & Amphibian Atlas	VTrans, FHWA
Compatible Resource Use		Manage selected areas for early succession, work with agriculturalists to manage lands to maintain early successional habitat does not cause direct mortality (e.g., light pasturing rather than harvesting/cultivation near wetlands and denning habitat.	Acres of wetland/grassland mosaics within safe travel distance of known denning habitat.	Sportsmen, Audubon VT	SWG, VHCB
Planning & Zoning	High	Identify important denning areas and movement corridors and minimize development, clearing, road building and increased traffic in these areas. Maintain low density human use in mosaics in known areas.	Number of specific sites identified. Number of sites with compatible land use.	TNC, landowners, land managers, VTrans, town government s, Town/ Regional Planners	SWG, TNC, VTrans
Compatible Resource Use	Medium	Manage ATV and other off road usage in known habitat to avoid impacts.	Number of sites where ATV use is controlled	Landowners, Game Wardens, TNC, VASA	ATV License fees, SWG

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Ratsnake**
 Scientific Name: **Pantherophis alleghaniensis**
 Species Group: **Herp**

Research	Medium	Determine the presence/absences of reptile diseases in the historic area of the Chorus frog and to control their spread and impact.	Disease surveys	Regional Wildlife Health Lab, Academic community	SWG, Nongame Fund, Private grants
Awareness Raising and Communications	High	Quickly and thoroughly, counter myths and misinformation appearing in the press that may limit this species.	Number of press responses carried by media.	SAG-Herps, Reptile & Amphibian Atlas, VFWD Outreach Division	SWG, Marketing funds
Technical Assistance, Training, Learning Networks		Maintain and support the network of trained snake relocators for this species as well as Rattlesnakes. Put information about Ratsnakes and this service on the same materials and website as for rattlesnake.	Number of requests for assistance.	Volunteers, Reptile & Amphibian Atlas, TNC, Orianne Society	TNC, Orianne Society
Awareness Raising and Communications	Medium	Encourage reports of sightings to the VT Wildlife Diversity Program and the VT Reptile & Amphibian Atlas.	Number of reports received	Reptile & Amphibian Atlas	Nongame Wildlife Fund, SWG
Standards	Medium	Develop land management guidelines for owners and managers of appropriate habitat and make them readily available through multiple media, including print and the web.	Numbers of landowners and managers who become aware and use guidelines	VFWD district biologists, consulting foresters	SWG
Awareness Raising and Communications	Medium	Help Vermonters assign value to this species through educational programs, printed material, web site information, field trips, TV and video information.	Number of people who receive information	ECHO Center for Lake Champlain, VFWD Outreach Division	Corporate Sponsors, Lake Champlain Basin Program, Marketing funds
Awareness Raising and Communications	Medium	Encourage reports of road-killed specimens, road crossing, and road basking areas to VFWD, VTrans, and the VT Reptile & Amphibian Atlas.	Number of sites reported	VTrans, Reptile & Amphibian Atlas.	VTrans, FHWA, SWG
Publically-Owned Protected Areas	High	Protect through easement or purchase or collaboration, denning sites, travel corridors and foraging areas.	Acreage and number of sites conserved	Reptile & Amphibian Atlas, Lake Champlain Land Trust, VLT	VHCB



Common Name: **Eastern Ratsnake**
Scientific Name: ***Pantherophis alleghaniensis***
Species Group: **Herp**

Bibliography

- Andrews, J. 2012. Rare snakes of Vermont-census/monitoring/tagging of rare snakes in Rutland and Windham Counties, Vermont, final report for State Wildlife Grant Award #06120FY10183
- Andrews, J. 2013. The Vermont Reptile and Amphibian Atlas. VtHerpAtlas.org
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Ernst, C. H., and E. Ernst. 2003. Snakes of the United States and Canada. Smithsonian Institution Press, Washington D.C. 668 pp.
- Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp.
- Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp.
- Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp.
- Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp.
- Krulikowski, L. 2007 Snakes of New England: A Photographic and Natural History Study. Self-published 320 pp.
- Lillywhite, H.B. 2014. How snakes work: structure, function, and behavior of the world's snakes. Oxford University Press, NY, NY 241 pp.
- Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp.
- Mullin, S.J. and R.A. Seigel eds. 2009. Snakes: ecology and evolution, Cornell University Press, Ithaca, NY 365 pp.
- Tennant, A. 2003. Snakes of North America: eastern and central regions. Lone Star Books, Lanham, Maryland. 605 pp.
- Wright, A.H. and A.A. Wright. 1994. Handbook of snakes of the United States and Canada, volumes 1 and 2. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 1105 pp.
- Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Common Watersnake**
Scientific Name: **Nerodia sipedon**
Species Group: **Herp**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S3

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

The Common Watersnake (previously known as the Northern Watersnake) is mainly a Champlain Valley species in Vermont, but also found in SE Vermont. This species is relatively large and aggressive, so is sometimes killed by humans. This species does suffer some road mortality and is purposely killed by some. It is usually associated with large wetlands of the Champlain Valley. Shoreline development may increase negative impacts from humans. Literature suggests pollution may be a problem (Harding 1997; Hunter, Calhoun, McCollough 1999). In some locations, the Northern Water Snake can be found in large numbers (e.g., Bristol Pond).

Distribution

The Common Watersnake is known from the Champlain Valley, Shaftsbury, and Vernon.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Certain
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Middle Connecticut
Hudson-Hoosic
Mettawee River
Lake Champlain
Missiquoi River
Otter Creek

Probable Watersheds

West
Lamoille River
Missisquoi River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

Wetlands associated with permanent water bodies. Also used flooded meadows. Avoids deeply shaded areas (Hunter et al. 1999). Uses overwintering sites in upland rock outcrops with cracks. Basking sites near water.



Common Name: **Common Watersnake**
Scientific Name: **Nerodia sipedon**
Species Group: **Herp**

Habitat Types:

Marshes and Sedge Meadows
Shrub Swamps
Aquatic: Lacustrine
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Impacts of Roads or Transportation Systems
Climate Change
Habitat Fragmentation

Description of habitat threat(s): Loss and conversion of wetlands, and shoreline development

Non-Habitat Threats:

Disease
Pollution
Trampling or Direct Impacts

Description of non-habitat threat(s): Northern Water Snakes may be impacted by pollution of their aquatic habitat. They are sometimes persecuted by people and are run over when crossing roads. Northern Water Snakes have become entangled in plastic erosion control/landscape netting. Snake fungal disease has recently been identified in this area and may potentially cause mortality in this species. Ranavirus may also impact population.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Common Watersnake**
 Scientific Name: **Nerodia sipedon**
 Species Group: **Herp**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	1) Identification of wintering sites would be important. 2) Gather data on specific habitat requirements of Vermont populations: denning sites, birthing sites, foraging areas, and movement corridors.
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	1) Need to develop a good baseline for this species. Identify distribution and relative abundance of populations in Vermont. 2) Target some surveys along the Connecticut River Valley.
Research	Threats and Their Significance	High	1) It would be helpful to know the level of mortality due to human activity. 2) Investigate water quality and human impacts to snakes.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	High	Need to monitor population and distribution change in order to take action while there is still time.
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	High	Monitor for changes in Vermont distribution and manage accordingly.
Monitoring	Monitor Threats	High	1) Monitor for snake fungal disease. 2) It is important to monitor limiting factors to gauge impacts to the species.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Common Watersnake**
 Scientific Name: **Nerodia sipedon**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Compatible Resource Use		Maintain or regain water quality in known use areas.	Maintenance or improvement in water quality.	Wetlands managers, farmers, Towns	Clean and Clear funding
Compatible Resource Use		Manage ATV and other off road usage in known habitat to lessen impacts.	Number of areas where ATV use is controlled.	Missisquoi National Wildlife Refuge, landowners, state lands managers, VASA	Refuge Mgmt funds, State Lands Mgmt
Awareness Raising and Communications		Quickly and thoroughly, counter myths and misinformation appearing in the press that may limit this species.	Number of media outlets that carry rebuttal of myths.	Reptile & Amphibian Atlas, SAG-Herps, media, VFWD Outreach Division	Marketing funds, volunteer efforts
Species Restoration		Reexamine species status at regular intervals (no longer than every 10 years) to determine if listing is appropriate.	Frequency of reviews.	SAG-Herps	volunteer effort
Species Restoration		Protect denning areas.	Number of sites protected.	landowners, managers	EQIP
Policy & Regulations		Establish and maintain 100-foot buffers of natural vegetation along water bodies in known habitat.	Number of sites with protected buffer habitat.	landowners, wetland managers	state lands Mgmt funds, EQIP, Partners in Wildlife
Planning & Zoning		Review all roadway projects in appropriate habitat, check against known crossing areas VTRANS, VFWD, VT Reptile & Amphibian Atlas, survey appropriate habitat when unknown.	Number of projects that use watersnake information for planning.	VTrans	VTrans FHWA
Standards		Develop management guidelines for owners and managers of appropriate habitat and make them readily available through multiple media, including print and the web.	Numbers of landowners and managers who receive and use guidelines.	Wetlands Managers, landowners	SWG
Habitat Restoration		Develop, implement, and monitor, road crossing structures and barriers for this species.	Number of structures installed.	VTrans	VTrans, FHWA

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Common Watersnake**
 Scientific Name: **Nerodia sipedon**
 Species Group: **Herp**

Standards	Medium	Encourage compliance with ANR Stormwater Guidelines regarding erosion control matting, in both regulated and non-regulated contexts.	Number of contractors and homeowners who use non-plastic EC and landscape matting. Number of retailers stocking and selling non-plastic matting.	VT Stormwater Office, VT Association of General Contractors. Product manufacturers and distributors.	SWG
Awareness Raising and Communications		Put information about watersnakes on the web.	Number of sites with posting.	Lake Champlain Committee, ECHO Leahy Center for Lake Champlain, Lake Champlain Basin Program	Lake Champlain Basin Program funds
Awareness Raising and Communications		Place informational posters at access areas where this species is known (Button Bay, Shelburne Pond, Bristol Pond, Vernon Pond).	Number of sites with signage.	game wardens	Nongame Wildlife Fund
Awareness Raising and Communications		Encourage reports of sightings to the VT Heritage Inventory and the VT Reptile & Amphibian Atlas.	Number of reports received.	Reptile & Amphibian Atlas	private grants, Nongame Wildlife Fund
Awareness Raising and Communications		Encourage reports of road-killed specimens, road crossing, and road basking areas to VFWD, VTRANS, and the VT Reptile & Amphibian Atlas.	Number of sites reported.	VTrans, Reptile & Amphibian Atlas	VTrans
Easements		Conserve known habitat through fee simple purchase, development rights or easements, management agreements and education of private landowners and managers regarding appropriate management.	Number of sites and acreage conserved	VLT, Missisquoi National Wildlife Refuge	VHCB funds, refuge acquisition
Awareness Raising and Communications		Help Vermonters assign value to this species through educational programs, printed material, web site information, field trips, TV and video information.	Number of people exposed to message.	Reptile & Amphibian Atlas, VFWD Outreach Division	marketing funds
Awareness Raising and Communications		Educate anglers regarding the conservation needs, habits of this species, and inform them of the protected status of this species.	Number of anglers exposed to message.	VFWD Outreach Division, angler groups, refuge staff	Marketing funds

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Common Watersnake
Scientific Name: *Nerodia sipedon*
Species Group: Herp

Easements	Consider creation of basking, denning, and refuge areas (rock piles) near appropriate foraging habitat.	Number of sites created and used.	Wetland managers, refuge staff	state land Mgmt funds, refuge operating budget

Bibliography

Andrews, J. 2013. The Vermont Reptile and Amphibian Atlas. VTHerpAtlas.org

Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.

DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.

Ernst, C.H. and R.W. Barbour. 1989. Snakes of eastern North America. George Mason University Press, Fairfax, Virginia. 282 pp.

Ernst, C. H., and E. Ernst. 2003. Snakes of the United States and Canada. Smithsonian Institution Press, Washington D.C. 668 pp.

Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.

Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp.

Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp.

Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.

Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp.

Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)

Krulikowski, L. 2007. Snakes of New England: A Photographic and Natural History Study. Self-published 320 pp

Lillywhite, H.B. 2014. How snakes work: structure, function, and behavior of the world's snakes. Oxford University Press, NY, NY 241 pp.

Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp. (This book provides excellent information on the species of reptile that we share with Virginia; most of our species are found in this book.)

Mullin, S.J. and R.A. Seigel eds. 2009. Snakes: ecology and evolution, Cornell University Press, Ithaca, NY 365 pp.

Tennant, A. 2003. Snakes of North America: eastern and central regions. Lone Star Books, Lanham, Maryland. 605 pp.

Tyning, T.F. 1990. A guide to amphibians and reptiles. Little, Brown and Company. Boston Massachusetts 400 pp.

Wright, A.H. and A.A. Wright. 1994. Handbook of snakes of the United States and Canada, volumes 1 and 2. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 1105 pp.

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp.



Common Name: **DeKay's Brownsnake**
 Scientific Name: **Storeria dekayi**
 Species Group: **Herp**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S4

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The DeKay's Brownsnake is an S4 species in Vermont, but populations are small and highly localized. The Brown Snake reaches its ecological limit across northern New England (Hunter et al. 1999), where it is less tolerant of disturbed sites and dependent upon habitat mosaics consisting of wetlands or riparian margins adjacent to upland forest overwintering sites.

Distribution

The Brown Snake is primarily found in the Champlain Valley, Taconics, and a few scattered records from the southern CT River Valley. It is widespread and more common in southern New England (Klemens 1993).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Probable
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Probable	Southern Green Mtns	Probable
Northern VT Piedmont	Probable	Taconic Mtns	Confident
Northeastern Highlands	Unknown		

Distribution by Watershed:

Known Watersheds

Middle Connecticut
 West
 Upper Connecticut-Mascoma
 Black-Ottawquechee
 Deerfield
 Lake Champlain
 Lamoille River
 Otter Creek
 Winooski River

Probable Watersheds

Hudson-Hoosic
 Missiquoi River



Common Name: **DeKay's Brownsnake**
Scientific Name: **Storeria dekayi**
Species Group: **Herp**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

DeKay's Brown Snake primarily occupies wet woods and fields, sedge meadows, seeps, and wetland or stream margins adjacent to upland forest. They are typically found under a variety of cover objects, including logs, stones, brush piles, leaf litter, etc. Critical habitat for this snake includes lowland wetlands or riparian margins adjacent to upland forest where it overwinters.

Habitat Types:

Hardwood Swamps
Seeps and Pools
Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops
Shrub Swamps
Early Succession Northern Hardwoods
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Impacts of Roads or Transportation Systems
Habitat Fragmentation

Description of habitat threat(s): Any habitat conversion, alteration, or fragmentation that disrupts species' ability to move between foraging and overwintering sites may have negative effects. Road mortality can negatively impact migrating adults and dispersing juveniles, especially when located between hibernaculum and foraging habitats. In Vermont this species appears less tolerant of disturbed habitats than in southern New England near the core of its range.

Non-Habitat Threats:

Trampling or Direct Impacts

Description of non-habitat threat(s): This species often occurs in close proximity to humans, and its distribution is primarily in a region of the state that continues to see significant development pressure. Fragmentation of suitable habitats by roads or other non-permeable development may result in loss of



Common Name: **DeKay's Brownsnake**
Scientific Name: **Storeria dekayi**
Species Group: **Herp**

metapopulation structure leading to genetic isolation and prevention of recolonization, especially considering the limited and localized populations of this species.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	1) Gather data on specific habitat requirements of Vermont populations: denning sites, birthing sites, foraging areas, and movement corridors. 2) Determine if and how habitat differs in Vermont compared to the core of the Brown Snake range.
Research	Distribution and Abundance	High	Determine species statewide distribution and relative abundance with emphasis in Taconics and southern CT River Valley.
Research	Threats and Their Significance	High	
Research	Other Research	Medium	Develop enhancement techniques for birthing and overwintering habitat.
Monitoring	Population Change	High	Monitor population sizes and distribution changes.
Monitoring	Habitat Change	High	
Monitoring	Range Shifts	High	Range distribution monitoring may be how we are able to track population change in Vermont (maintenance or loss of populations).
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **DeKay's Brownsnake**
 Scientific Name: **Storeria dekayi**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications		Encourage reports of sightings to the VT Heritage Inventory and the VT Reptile & Amphibian Atlas.	Number of reports received. Geographic coverage of reports.	Reptile & Amphibian Atlas	Nongame Wildlife Fund
Compatible Resource Use		Review all roadway projects in appropriate habitat, check against known crossing areas VTRANS, VFWD, VT Reptile & Amphibian Atlas, survey appropriate habitat when unknown.	Number of sites where information on crossing areas	VTrans, Reptile & Amphibian Atlas	VTrans planning funds
Compatible Resource Use		Manage ATV and other off road usage in known habitat to lessen impacts.	Number of brownsnake areas with restricted or managed ATV use.	Land managers, private landowners, VASA	
Awareness Raising and Communications		Encourage reports of road-killed specimens, road crossing, and road basking areas to VFWD, VTRANS, and the VT Reptile & Amphibian Atlas.	Number of areas reported.	VTrans, Reptile & Amphibian Atlas	VTrans funds
Habitat Restoration		Maintain habitat mosaic and connectivity necessary for this species, particularly in Champlain Valley.	Number of intact habitats and connections	Consulting Foresters, landowners, Cons Comms	Current Use, EQIP
Standards		Develop land management guidelines for owners and managers of appropriate habitat and make them readily available through multiple media, including print and the web.			
Easements		Conserve known habitat through fee simple purchase, development rights or easements, management agreements and education of private landowners and managers regarding appropriate management.	Number of known sites conserved.	Consulting foresters, local Cons Comms	VHCB funds
Species Restoration	High	Reexamine species status at regular intervals (no longer than every 10 years) to determine if Endangered Species Act listing is appropriate.	Frequency of review	SAG-Herps	donated time
Awareness Raising and Communications		Help Vermonters assign value to this species through educational programs, printed material, web site information, field trips, TV and video information.	Number of people exposed to message.	VFWD Outreach Division	Marketing funds
Species Restoration		Develop, install, and monitor, road crossing structures and barriers for this species.	Number of structures installed.	VTrans	FHWA



Common Name: **DeKay's Brownsnake**
Scientific Name: **Storeria dekayi**
Species Group: **Herp**

Bibliography

- Andrews, J. 2013. The Vermont Reptile and Amphibian Atlas. <http://cat.middlebury.edu/herpatlas/>
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Ernst, C.H. and R.W. Barbour. 1989. Snakes of eastern North America. George Mason University Press, Fairfax, Virginia. 282 pp. (An excellent source for detailed information.)
- Ernst, C. H., and E. Ernst. 2003. Snakes of the United States and Canada. Smithsonian Institution Press, Washington D.C. 668 pp. (The latest and most complete source for snakes.)
- Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)
- Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)
- Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.
- Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp. (Visit the PARC website for more information: www.parcplace.org.)
- Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)
- Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp. (This book provides excellent information on the species of reptile that we share with Virginia; most of our species are found in this book.)
- Tennant, A. 2003. Snakes of North America: eastern and central regions. Lone Star Books, Lanham, Maryland. 605 pp. (One of a two excellent new snake resources.)
- Wright, A.H. and A.A. Wright. 1994. Handbook of snakes of the United States and Canada, volumes 1 and 2. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 1105 pp. (A reprint of an old classic.)
- Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp. (A standard text.)



Common Name: **Eastern Ribbonsnake**
 Scientific Name: **Thamnophis sauritus**
 Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The Eastern Ribbonsnake is a rare species in Vermont (S2) and is considered a species of special concern in Vermont. The Eastern Ribbonsnake is one of the rarest of snakes in Vermont based on the number of known current sites. It seems to depend on a combination of a relatively warm, undeveloped lowland site and wetlands.

Distribution

It is currently documented from only six locations in Vermont: five in western Rutland County and one along the southern Connecticut River valley. A handful of historic records and sightings come from further north in the Lake Champlain basin and the Connecticut River Valley.

Distribution by Biophysical Region:

Champlain Valley	Probable	Southern VT Piedmont	Confident
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

West
 Mettawee River
 Lake Champlain

Probable Watersheds

Middle Connecticut
 Hudson-Hoosic
 Otter Creek
 Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

This species requires wetland edges with sunny exposed basking sites in warm, low-elevation, largely undeveloped, areas. The presence of nearby rocky woodlands and talus seems to increase the chances of finding this species.



Common Name: **Eastern Ribbonsnake**
Scientific Name: **Thamnophis sauritus**
Species Group: **Herp**

Habitat Types:

Cliffs and Talus
Seeps and Pools
Oak-Pine Northern Hardwood
Marshes and Sedge Meadows
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Shrub Swamps
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Habitat Alteration
Impacts of Roads or Transportation Systems
Habitat Fragmentation

Description of habitat threat(s): Conversion or drainage of wetlands, shoreline development, and fragmentation due to road density could all be problems.

Non-Habitat Threats:

Genetics
Loss of Prey Base
Trampling or Direct Impacts

Description of non-habitat threat(s): Some populations may be genetically isolated and others are becoming more so as a result of development. This species may be dependent on local amphibian populations that are known to vary annually. ATV use, increased traffic, cutting and bailing, and lawn mowing could all increase mortality significantly. It has not been located in moderately or heavily developed areas.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Ribbonsnake**
Scientific Name: **Thamnophis sauritus**
Species Group: **Herp**

Research and Monitoring Needs

<i>Type</i>	<i>Need</i>	<i>Priority</i>	<i>Description</i>
Research	Habitat Requirements	High	This species may use a wider variety of habitats in Vermont than is currently known. 1) Gather data on specific habitat requirements of Vermont populations: denning sites, birthing sites, foraging areas, overwintering sites and movement corridors.
Research	Distribution and Abundance	High	Identify distribution and relative abundance of populations in Vermont. Search for ribbonsnakes in areas of open talus in the Champlain, Connecticut River valley, and other relatively warm valleys, especially if adjacent to wetland foraging areas.
Research	Threats and Their Significance	High	Monitor for signs of Snake Fungal Disease in this species.
Research	Other Research	Medium	Develop enhancement techniques to improve for birthing and overwintering habitat.
Monitoring	Population Change	High	Monitor population sizes and distribution changes.
Monitoring	Habitat Change	High	
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Ribbonsnake**
 Scientific Name: **Thamnophis sauritus**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications		Quickly and thoroughly, counter myths and misinformation appearing in the press that may limit this species.	Number of response carried by media.	Media, SAG-Herps, Reptile & Amphibian Atlas, VFWD Outreach Division	volunteer, marketing funds
Publically-Owned Protected Areas		Locate populations on public lands and manage some specifically for this species.	Number of sites managed for ribbonsnake	FPR, USFS, VFWD	State Lands Mgmt funds, GMNF funds
Awareness Raising and Communications	Medium	Encourage reports of sightings to the VT Heritage Inventory and the VT Reptile & Amphibian Atlas.	Number of reports received.	Reptile & Amphibian Atlas	Nongame Wildlife Fund, private grants
Awareness Raising and Communications		Encourage reports of road-killed specimens, road crossing, and road basking areas to VFWD, VTrans, and the VT Reptile & Amphibian Atlas.	Number of sites reported.	VTrans, volunteers	VTrans
Compatible Resource Use		Review all roadway projects in appropriate habitat, check against known crossing areas VTrans, VFWD, VT Reptile & Amphibian Atlas, survey appropriate habitat when unknown.	Number of projects reviewed with planning information on snakes.	VTrans, Reptile & Amphibian Atlas	VTrans
Compatible Resource Use		Manage ATV and other off road usage in known habitat to lessen impacts.	Number of sites where ATV use is controlled.	Land managers, landowners, VASA	
Easements		Conserve known habitat through fee simple purchase, development rights or easements, management agreements and education of private landowners and managers regarding appropriate management.	Number of sites conserved.	Vermont Land Trust	VHCB funds
Awareness Raising and Communications		Help Vermonters assign value to this species through educational programs, printed material, web site information, field trips, TV and video information.	Number of people who receive message.	VFWD Outreach Division	Marketing funds
Standards		Develop land management guidelines for owners and managers of appropriate habitat and make them readily available through multiple media, including print and the web.	Number of landowners and managers who receive and use guidelines.	VFWD district biologists, consulting foresters	SWG
Habitat Restoration	Medium	Develop, implement, and monitor, road crossing structures and barriers for this species.	Number of structures installed.	VTrans	FHWA

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Eastern Ribbonsnake
Scientific Name: *Thamnophis sauritus*
Species Group: Herp

Species Restoration	High	Reexamine species status at regular intervals (no longer than every 10 years) to determine if ESA listing is appropriate.	Number of years since last review.	SAG-Herps	volunteer

Bibliography

Andrews, J. 2013. The Vermont Reptile and Amphibian Atlas. <http://cat.middlebury.edu/herpatlas/>

Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.

DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.

Ernst, C.H. and R.W. Barbour. 1989. Snakes of eastern North America. George Mason University Press, Fairfax, Virginia. 282 pp. (An excellent source for detailed information.)

Ernst, C. H., and E. Ernst. 2003. Snakes of the United States and Canada. Smithsonian Institution Press, Washington D.C. 668 pp. (The latest and most complete source for snakes.)

Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.

Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)

Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.

Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp. (Visit the PARC website for more information: www.parcplace.org.)

Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)

Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp. (This book provides excellent information on the species of reptile that we share with Virginia; most of our species are found in this book.)

Tennant, A. 2003. Snakes of North America: eastern and central regions. Lone Star Books, Lanham, Maryland. 605 pp. (One of a two excellent new snake resources.)

Wright, A.H. and A.A. Wright. 1994. Handbook of snakes of the United States and Canada, volumes 1 and 2. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 1105 pp. (A reprint of an old classic.)

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp. (A standard text.)



Common Name: **Smooth Greensnake**
 Scientific Name: **Opheodrys vernalis**
 Species Group: **Herp**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S4

State Trend:

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Few people encounter the Smooth Greensnake and it is thought to be found less frequently than in the past. Little is known about its distribution in Vermont and its considered uncommon in Vermont (S3). It's conservation status is uncertain and its considered a medium-priority SGCN.

Habitat loss due to development is also a problem for this species, especially in the lowlands. In past large beaver meadows may have been connected. Mechanization of agriculture, lawn mowing, and roads all are likely impacts. In the southern Great Lakes Basin it is reported to be decreasing due to intensive conversion of its habitat to agricultural uses and pesticides (Harding 1997. The Amphibians and Reptiles of the Great Lakes Region).

Distribution

Primarily at mid-elevational levels. Missing from Northeastern Vermont

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills		Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Unknown		

Distribution by Watershed:

Known Watersheds

Middle Connecticut

West

Waits

Upper Connecticut-Mascoma

Deerfield

Hudson-Hoosic

Mettawee River

Lake Champlain

Lamoille River

Missiquoi River



Common Name: **Smooth Greensnake**
Scientific Name: **Opheodrys vernalis**
Species Group: **Herp**

Otter Creek

White

Winooski River

Black-Ottawquechee

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Greensnakes use sedge meadows, marsh borders, pastures, powerlines, shrub areas, and early successional habitat not mowed regularly. Micro Habitat: dense annual vegetation. Critical habitat includes overwintering habitat (ant mounds), early successional foraging habitat, and areas for egg laying.

Habitat Types:

Open Peatlands

Marshes and Sedge Meadows

Early Succession Upland Oak

Early Succession Other Types

Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Wet Shores

Shrub Swamps

Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Succession

Habitat Alteration

Impacts of Roads or Transportation Systems

Habitat Fragmentation

Description of habitat threat(s): Mowing and pesticides are limiting factors for this species. Baling also does impact snakes.

Non-Habitat Threats:

Disease

Pollution

Trampling or Direct Impacts

Description of non-habitat threat(s): Mowing of habitat, road traffic, and pesticide use. Snakes have become entangled in plastic erosion control/landscape netting. Snake fungal disease has recently been



Common Name: **Smooth Greensnake**
Scientific Name: ***Opheodrys vernalis***
Species Group: **Herp**

identified in this area and may potentially cause mortality in this species. Ranavirus may also impact populations.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Gather data on specific habitat requirements of Vermont populations: denning sites, egg-laying sites, foraging areas, overwintering sites and movement corridors.
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Identify distribution and relative abundance of populations in Vermont.
Research	Threats and Their Significance	High	
Research	Population Genetics	Low	
Research	Other Research	Medium	Gather data from known habitat on how it is kept in early succession and apply this knowledge.
Monitoring	Population Change	High	Monitor population sizes and distribution changes.
Monitoring	Habitat Change	High	
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	High	Monitor for Snake Fungal Disease.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Smooth Greensnake**
 Scientific Name: **Opheodrys vernalis**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Standards	Medium	Encourage compliance with ANR Stormwater Guidelines regarding erosion control matting, in both regulated and non-regulated contexts.	Number of contractors and homeowners who use non-plastic EC and landscape matting. Number of retailers stocking and selling non-plastic matting.	VT Stormwater Office, VT Association of General Contractors.	SWG
Compatible Resource Use		Maintain connectivity between areas of appropriate early successional habitat.	Number of acres linked through connectivity	landowners	EQUIP, Current Use
Habitat Restoration		Develop, implement, and monitor, road crossing structures and barriers for this species.	Number of structures installed	VTrans	VTrans, FHWA
Awareness Raising and Communications		Encourage reports of sightings to the VT Heritage Inventory and the VT Reptile & Amphibian Atlas.	Number of reports received	Reptile & Amphibian Atlas	private grant, Nongame Wildlife Fund
Easements		Conserve known habitat through fee simple purchase, development rights or easements, management agreements and education of private landowners and managers regarding appropriate management.	Number of sites and acreage conserved	VLT, local land trusts	VHCB funds
Compatible Resource Use		Manage ATV and other off road usage in known habitat to lessen impacts.	Number of sites where ATV use is controlled	landowners, land managers, VASA	EQUIP, state lands Mgmt funds
Standards		Develop land management guidelines for owners and managers of appropriate habitat and make them readily available through multiple media, including print and the web.			
Awareness Raising and Communications		Encourage reports of road-killed specimens, road crossing, and road basking areas to VFWD, VTRANS, and the VT Reptile & Amphibian Atlas.	Number of reports received	VTrans	VTrans
Planning & Zoning		Review all roadway projects in appropriate habitat, check against known crossing areas VTRANS, VFWD, VT Reptile & Amphibian Atlas, survey appropriate habitat when unknown.	Number of projects reviewed using green snake information	VTrans, Reptile & Amphibian Atlas	VTrans, FHWA
Species Restoration		Reexamine species status at regular intervals (no longer than every 10 years) to determine if listing is appropriate.	Frequency of reviews	SAG-Herps	volunteer effort

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Smooth Greensnake**
 Scientific Name: **Opheodrys vernalis**
 Species Group: **Herp**

Awareness Raising and Communications	Help Vermonters assign value to this species through educational programs, printed material, web site information, field trips, TV and video information.	Number of people exposed to message.	Reptile & Amphibian Atlas, VFWD Outreach Div	Marketing funds
Publically-Owned Protected Areas	Locate populations on public lands and manage some specifically for this species.	Number of sites managed for green snake.	district foresters and wildlife managers	state lands Mgmt funds
Standards	Work with power companies, airports, horse farmers, and other landowners that provide large areas of early successional habitat to maintain it in a manner safe for this species.	Number of sites maintained in a safe manner.	Managers of powerlines, airport staff, landowners	VELCO, VTrans, EQIP

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Smooth Greensnake
Scientific Name: *Opheodrys vernalis*
Species Group: Herp

Bibliography

- Andrews, J. 2013. The Vermont Reptile and Amphibian Atlas. VTHerpAtlas.org
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Ernst, C.H. and R.W. Barbour. 1989. Snakes of eastern North America. George Mason University Press, Fairfax, Virginia. 282 pp.
- Ernst, C. H., and E. Ernst. 2003. Snakes of the United States and Canada. Smithsonian Institution Press, Washington D.C. 668 pp.
- Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp.
- Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp.
- Hunter, M.L., A. Calhoun, and M. McCullough (eds.). 1999. Maine amphibians and reptiles. The University of Maine Press, Orono, Maine 272 pp.
- Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp.
- Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)
- Krulikowski, L. 2007. Snakes of New England: A Photographic and Natural History Study. Self-published 320 pp
- Lillywhite, H.B. 2014. How snakes work: structure, function, and behavior of the world's snakes. Oxford University Press, NY, NY 241 pp.
- Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp. (This book provides excellent information on the species of reptile that we share with Virginia; most of our species are found in this book.)
- Mullin, S.J. and R.A. Seigel eds. 2009. Snakes: ecology and evolution, Cornell University Press, Ithaca, NY 365 pp.
- Tennant, A. 2003. Snakes of North America: eastern and central regions. Lone Star Books, Lanham, Maryland. 605 pp.
- Tyning, T.F. 1990. A guide to amphibians and reptiles. Little, Brown and Company. Boston Massachusetts 400 pp.
- Wright, A.H. and A.A. Wright. 1994. Handbook of snakes of the United States and Canada, volumes 1 and 2. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 1105 pp.
- Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp.



Common Name: **Timber Rattlesnake**
 Scientific Name: **Crotalus horridus**
 Species Group: **Herp**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: S1

State Trend:

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The Timber Rattlesnake is a state-endangered S1 species that was historically known from a much wider range in VT and the region. Current estimates of population and species distribution hover near some 75-85% loss and range contraction from historic levels. Recent telemetry research in Vermont further refined local home ranges by sex and range distribution of one population and concomitantly highlighted strategic rattlesnake habitats for future conservation efforts. Genetic analysis indicated relatively low heterozygosity in this cohort although random mating still appears to be occurring. Roads with high traffic levels appeared to present significant barriers to movement and dispersal. (Spear et.al. 2013).

Extant rattlesnake populations are known only from two denning areas in Vermont. Since it is a venomous species, it is more widely feared and persecuted—a significant threat to the species in Vermont. The species had a bounty on it until 1971. The bounty was lifted in 1987, but direct persecution and occasional takings still occur. Its habit of denning communally at sites which are now quite widely known make it unusually vulnerable to takings. This behavior also provides opportunities for geographically targeted protection. In 2012, a novel and lethal snake fungal disease, *Ophidiomyces ophidicola*, was isolated and identified in Vermont's Timber Rattlesnake populations and has since been found in various snake species in a dozen eastern states. This finding has prompted an in-depth, regional investigation of this disease in which Vermont is participating. Known as Snake Fungal Disease (SFD) it imposes an unknown but potentially devastating threat to the species. As the animals' name implies, the Timber Rattlesnake depends on warm low-elevation woodlands that are sparsely populated. Habitat fragmentation and concomitant increases in roads/traffic, human interactions present an increasing threat of mortalities.

Distribution

Two isolated populations of the Timber Rattlesnake are restricted to areas near the southern portion of Lake Champlain in western Rutland County below 1000 ft. in elevation. Research conducted in 2011-2012 indicates that genetic exchange between the two populations is likely non-existent. Populations in other parts of the state have been lost.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Historic Records Only
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Historic Records Only
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Mettawee River



Common Name: **Timber Rattlesnake**
Scientific Name: **Crotalus horridus**
Species Group: **Herp**

Lake Champlain

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

The Timber Rattlesnake in Vermont is documented only from western Rutland County at elevations below 1000 feet. In this region snakes den communally on south or southwest facing talus slopes which are near rocky ridges with exposed ledge and large undeveloped or sparsely developed areas of oak-hickory and maple-ash-hickory-oak vegetative communities. Males range annually about 2.5 miles from the natal den with females ranging up to 1.5 miles and gravid females utilizing still smaller ranges. Small, scattered canopy openings and forest wetlands along with their buffers are readily utilized and enhance habitat for foraging rattlesnakes. In Vermont, forested lands of approximately 5500 acres adjacent to the dens, sparsely developed and largely unfragmented by roads, support Vermont's discrete rattlesnake populations. (Spear et. al. 2013) Habitual movement corridors between dens and summer range are utilized annually. Roads present significant barriers and are deleterious to seasonal movement and dispersing snakes. Successful long distance movements between extant dens are now highly unlikely and may require human intervention/assistance in the future. Rattlesnakes maintain strict annual fidelity to their traditional, communal den sites. In Vermont's harsh climate, overwintering, frost-free den site requirements are exacting and thus, extremely limiting.

Critical habitat features are the rocky talus slopes with traditional dens, nearby ridges with exposed ledges, and extensive associated woodlands of oak-hickory and Pennsylvania sedge.

Habitat Types:

Outcrops and Alpine
Cliffs and Talus
Oak-Pine Northern Hardwood
Hardwood Swamps
Subterranean
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops
Shrub Swamps
Early Succession Northern Hardwoods
Early Succession Upland Oak
Mine

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Habitat Fragmentation



Common Name: **Timber Rattlesnake**
Scientific Name: **Crotalus horridus**
Species Group: **Herp**

Impacts of Roads or Transportation Systems

Description of habitat threat(s): The most immediate threat to Timber Rattlesnakes with potential population consequences is the documented presence of Snake Fungal Disease (SFD) in extant populations. The lethality and severity of SFD has the potential to overwhelm other efforts on behalf of the species. The rattlesnake uses large contiguous woodland areas adjacent to their dens. Heavy agricultural or residential use, or conversion to open land are all problems. Direct loss of habitat, increased habitat fragmentation and road-density with higher traffic levels result higher road mortality, population isolation and increased snake/ human interactions. Roads present a highly fragmenting landscape feature, heightened chances of direct mortality and formidable barriers to successful snake movement, dispersal and genetic exchange. Lowland wooded patches are popular building sites and thus are becoming increasingly fragmented. The resultant direct persecution stemming from human fear and intolerance especially towards adult rattlesnakes can profoundly impact a population.

Non-Habitat Threats:

Genetics

Disease

Harvest or Collection

Reproductive Traits

Trampling or Direct Impacts

Description of non-habitat threat(s): Snake Fungal Disease is an immediate and potentially overwhelming threat to Vermont's population. Genetic exchange between Vermont's two populations is likely non-existent. Exchange with New York's nearest rattlesnake population is unknown. Although protected, this venomous species is still illegally collected for various purposes and snakes that have been killed are occasionally discovered. Known traditional den sites and predictable patterns of behavior make the species very vulnerable to collection and persecution. Birthing sites also appear to be limited and traditional. This is a long-lived, K-selected species that can successfully reproduce only every 4-5 years. Consequently, loss of breeding adults, particularly adult females is a problem to the sustainability of the species. Heavy ATV use, increased traffic, and heavy recreational use along ridges during key time-periods is also a problem. Since this species is venomous, it is often feared and killed when found near residences.



Common Name: **Timber Rattlesnake**
 Scientific Name: **Crotalus horridus**
 Species Group: **Herp**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Better determine range and habitat usage and protect critical areas.
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Determine Population Status: continue monitoring during ingress/egress periods. Focus on adult females/reproductive status/litters as an index.
Research	Threats and Their Significance	High	Monitor Snake Fungal Disease Status (SFD), continue surveillance, disease testing in cooperation with Regional SFD Investigation.
Research	Population Genetics	Medium	Conduct periodic genetic assessment to inform genetic exchange/variability. Consider techniques to facilitate gene flow (translocation/captive rearing/headstarting).
Monitoring	Population Change	High	Monitor the size and determine the sustainability of existing populations through age-class and/or genetic analysis.
Monitoring	Habitat Change	Medium	Continue vigilance with regulatory habitat protections (Act 250, CPG's).
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	High	Continue the Rattlesnake Responder Program which 1) protects both snakes and residents and provides opportunity for outreach/education. 2) continue law enforcement efforts. Also See Threats research regarding SFD.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Timber Rattlesnake**
 Scientific Name: **Crotalus horridus**
 Species Group: **Herp**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Alliance Development	Medium	Continue to work cooperatively with important landowners such as the Nature Conservancy. Develop and maintain allies in local government and private citizens.	Number of partner contacts made annually.	TNC, towns, landowners	SWG
Technical Assistance, Training, Learning Networks	High	Maintain and support the network of trained Snake Responders. Put information about Rattlesnakes and this service on the web. Educate local landowners. Distribute Responder Team refrigerator magnets.	Number of times public receives technical assistance.	Volunteers, local warden, town officials, TNC, Reptile & Amphibian Atlas	TNC
Awareness Raising and Communications	High	Continue the Rattlesnake Responder Program to protect people and animals and take advantage of "teachable moments." Patrol denning and birthing areas during necessary times to protect all life stages and send an important message to the public.	Number of animals safely moved. Number of sites that are patrolled.	Rattlesnake responder team, Game Wardens	SWG
Habitat Restoration	Medium	Develop, implement, and monitor, road crossing structures and barriers for this species.	Number of structures installed.	VTrans, Consulting herpetologists	VTrans
Species Restoration	Medium	Work to maintain connectivity with populations to the west in New York State and between the two known populations.	Quantity and quality of landscape connections.	New York DEC, TNC, VLT	VHCB
Awareness Raising and Communications	Medium	Quickly and thoroughly, counter myths and misinformation appearing in the press that may limit this species.	Number of press responses carried by media.	SAG-Herps, Reptile & Amphibian Atlas, VFWD Outreach Div., TNC	SWG, volunteer efforts, marketing funds
Easements	Medium	Conserve known habitat through fee simple purchase, development rights or easements, management agreements, and education of private landowners and managers regarding appropriate management.	Number of sites conserved.	landowners, TNC, VLT, TPL, Orianne Society	VHCB, Orianne Society
Standards	Medium	Develop land management guidelines for owners and managers regarding appropriate management and make them readily available through multiple media, including print and the web.	Number of landowners and managers who receive and use the guidelines.	landowners and land managers, TNC, Towns	SWG, TNC

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Timber Rattlesnake
Scientific Name: *Crotalus horridus*
Species Group: Herp

Planning & Zoning	Medium	Review all roadway projects in appropriate habitat, check against known crossing areas VTRANS, VFWD, VT Reptile & Amphibian Atlas, survey appropriate habitat when unknown.	Number of projects reviewed using rattlesnake planning information.	VTrans, Reptile & Amphibian Atlas, Towns	VTrans,
Research	High	Monitor Snake Fungal Disease status, continue surveillance, disease testing in cooperation with Regional SFD Investigation.	Relative prevalence of SFD in populations.	TNC, Orianne Society, Veterinarians, Castleton State College	SWG, other research grants
Species Restoration	High	If local populations are determined to be unsustainable, consider augmentation from closest healthy source. Maintaining and enhancing extant populations is always a priority and should be continued.	Number of successful reintroductions or augmentations.	New York DEC	SWG
Technical Assistance, Training, Learning Networks	Medium	Work with District 3 VTrans crew and other land managers to raise awareness of conservation need and implement conservation actions that benefit snakes.	Number of VTrans and others managers cooperating.	VTrans	VTrans
Compatible Resource Use	High	Protect known denning areas and adjacent ledges and woodlands from incompatible development and heavy use during critical time periods. Protect foraging land from development.	Number of sites conserved.	TNC, landowners, VLT, Act 250 Staff	VHCB
Awareness Raising and Communications	Medium	Encourage reports of road-killed specimens, road crossing, and road basking areas to VFWD, VTRANS, and the VT Reptile & Amphibian Atlas.	Number of sites reported.	VTrans, Reptile & Amphibian Atlas, TNC	VTrans, Nongame Wildlife Fund
Compatible Resource Use	Medium	Manage ATV and other off road usage in known habitat to avoid impacts, including foraging habitat.	Number of sites where ATV use is controlled.	Landowners, land managers, VASA, other ATV user groups	land Mgmt funds/decisions
Awareness Raising and Communications	Medium	Encourage reports of sightings to the Vermont Wildlife Diversity Program and the VT Reptile & Amphibian Atlas.	Number of reports received.	Reptile & Amphibian Atlas	SWG, Nongame Wildlife Fund
Awareness Raising and Communications	Medium	Help Vermonters assign value to this species through educational programs, printed material, web site information, field trips, TV and video information.	Number of people exposed to message	VFWD outreach division, Reptile & Amphibian Atlas	marketing funds

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Timber Rattlesnake**
Scientific Name: **Crotalus horridus**
Species Group: **Herp**

Bibliography

- Andrews, J. 2001. The Vermont Reptile and Amphibian Atlas. <http://cat.middlebury.edu/herpatlas/>
- Brown, W.S. 1993. Biology, status, and management of the Timber Rattlesnake (*Crotalus horridus*): A guide for conservation. Herpetological Circular No. 22. Society for the Study of Amphibians and Reptiles 78 pp
- 1991. Female Reproduction ecology in a northern population of the timber rattlesnake *Crotalus horridus*. *Herpetologica* 47:101-115.
- 1995. The female timber rattlesnake: A key to conservation. *Reptile and Amphibian Magazine* Sept/Oct:12-19.
- Cavanaugh, C.J. 1994 *Crotalus horridus* (Timber Rattlesnake) Longevity. *Herpetological Review* 25:70.
- Clark, R. W., Brown, W. S., Stechert, R., & Zamudio, K. R. (2010). Roads, interrupted dispersal, and genetic diversity in timber rattlesnakes. *Conservation Biology*, 24(4), 1059-1069.
- Clark, R. W., M. N. Marchand, B. J. Clifford, R. Stechert, and S. Stephens. 2011. Decline of an isolated timber rattlesnake (*Crotalus horridus*) population: Interactions between climate change, disease, and loss of genetic diversity. *Biological Conservation* 144:886-891.
- Conant, R., and J.T. Collins. 1998. A field guide to reptiles and amphibians of eastern and central North America. Third Edition, expanded, Houghton Mifflin Company, Boston Massachusetts 616 pp.
- DeGraaf, R.M., and D.D. Rudis. 1983. Amphibians and reptiles of New England. The University of Massachusetts Press, Amherst, Massachusetts 85 pp.
- Ernst, C.H. and R.W. Barbour. 1989. Snakes of eastern North America. George Mason University Press, Fairfax, Virginia. 282 pp. (An excellent source for detailed information.)
- Ernst, C. H., and E. Ernst. 2003. Snakes of the United States and Canada. Smithsonian Institution Press, Washington D.C. 668 pp. (The latest and most complete source for snakes.)
- Evink, G. 2002. National Cooperative Highway Research Program Synthesis 305, Interaction between roadways and wildlife ecology, A synthesis of highway practice. Transportation Research Board, Washington D.C. 78 pp.
- Harding, J.H. 1997. Amphibians and reptiles of the Great Lakes Region. The University of Michigan Press, Ann Arbor, Michigan 378 pp. (All our species are included.)
- Hulse, A., C. J. McCoy, and E. Censky. 2001. Amphibian and reptiles of Pennsylvania and the Northeast. Cornell University Press, Ithaca, New York 419 pp. (Most of our species are included.)
- Kingsbury, B. and J. Gibson. 2002. Habitat management guidelines for amphibians and reptiles of the Midwest. Midwest Partners in Amphibian and Reptile Conservation (Midwest PARC). 57 pp. (Visit the PARC website for more information: www.parcplace.org.)
- Klauber, L.M. 1982. Rattlesnakes: their habits, life histories, & influence on mankind, abridged edition. University of California Press, Berkeley and Los Angeles, California 350 pp. (An edited reprint of an old classic.)
- Klemens, M.K. 1993. Amphibians and reptiles of Connecticut and adjacent regions. State Geological and Natural History Survey of Connecticut, Bulletin No. 112 318 pp. (Unfortunately this is currently out of print.)
- Mace, Georgina M., Nigel J. Collar, Kevin J. Gaston, C. R. A. I. G. HILTON-TAYLOR, H. Resit Akçakaya, N. I. G. E. L. LEADER-WILLIAMS, E. J. MILNER-GULLAND, and Simon N. Stuart. "Quantification of extinction risk: IUCN's system for classifying threatened species." *Conservation Biology* 22, no. 6 (2008): 1424-1442.
- Martin W.H. 1992a. The timber rattlesnake: its distribution and natural history. In T.F. Tynning (ed.), *Conservation of the Timber Rattlesnake in the northeast*, pp. 13-22. Massachusetts Audubon Society, Lincoln, Massachusetts.
- Mitchell, J.C. 1994. The reptiles of Virginia. Smithsonian Institution Press, Washington 352 pp. (This book provides excellent information on the species of reptile that we share with Virginia; most of our species are found in this book.)
- Reinert, H.K. and R.R. Rupert, Jr. 1999. Impacts of Translocation on Behavior and Survival of Timber Rattlesnakes, *Crotalus*

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Timber Rattlesnake
Scientific Name: *Crotalus horridus*
Species Group: Herp

horridus. Journal of Herpetology 33:45-61.

Rubio, M. 1998. Rattlesnake: Portrait of a predator. Smithsonian Institution Press, Washington D.C. 266 pp.

Spear, S., J. Bauder, D. Blodgett, C. Jenkins, K. Briggs. 2013. The Ecology of Timber Rattlesnakes (*Crotalus horridus*) in Vermont. Vermont Fish and Wildlife Department, Rutland, Vermont, USA.

Tennant, A. 2003. Snakes of North America: eastern and central regions. Lone Star Books, Lanham, Maryland. 605 pp. (One of two excellent new snake resources.)

Therres, G.D. 1999. Wildlife species of regional conservation concern in the Northeastern United States. Northeast Wildlife 54:93-100.

Thompson, E.H. and E.R. Sorenson. 2000. Wetland, Woodland, Wildland: A Guide to the Natural Communities of Vermont. Vermont Department of Fish and Wildlife and The Nature Conservancy. University Press of New England, Hanover, NH.

Tyning, T.F. 1990. A guide to amphibians and reptiles. Little, Brown and Company. Boston Massachusetts 400 pp.

Victoria, J. 2000. Status Assessment of the Timber Rattlesnake, Report for the Northeast Endangered Species Committee (DRAFT).

Wright, A.H. and A.A. Wright. 1994. Handbook of snakes of the United States and Canada, volumes 1 and 2. Comstock Publishing Associates, A Division of Cornell University Press, Ithaca, New York 1105 pp. (A reprint of an old classic.)

Zappalorti, R.T. and H.K. Reinert. 1992. Distribution and habitat utilization of the timber rattlesnake, *Crotalus horridus* (Linnaeus), in southern New Jersey with notes on hibernation. In T.F. Tyning (ed.) Conservation on the timber rattlesnake in the northeast, pp 1-2. Massachusetts Audubon Society, Lincoln, Massachusetts.

Zug, G.R. 1993. Herpetology: an introductory biology of amphibians and reptiles. Academic Press, A Division of Harcourt Brace & Company, San Diego, California 527 pp. (A standard text.)

Vermont's Wildlife Action Plan

Bird SGCN Conservation Reports

Appendix A2

Draft 10/1/2015

Species	Page
Common Loon	1
Pied-billed Grebe	5
American Bittern	8
Least Bittern.....	11
Great Blue Heron	14
Black-crowned Night-heron	17
American Black Duck.....	20
Bald Eagle	24
Northern Harrier	28
Northern Goshawk	32
Red-shouldered Hawk.....	35
American Kestrel.....	38
Peregrine Falcon.....	41
Spruce Grouse.....	45
Ruffed Grouse.....	49
Sora	53
Lesser Yellowlegs	55
Upland Sandpiper	58
American Woodcock	61
Common Tern.....	66
Black Tern.....	69
Black-billed Cuckoo.....	72
Short-eared Owl.....	75
Common Nighthawk.....	78
Whip-poor-will	82

Species	Page
Chimney Swift	86
Black-backed Woodpecker.....	89
Olive-sided Flycatcher.....	92
Purple Martin.....	94
Gray Jay.....	97
Sedge Wren	100
Bicknell's Thrush	103
Wood Thrush.....	108
Brown Thrasher.....	111
Blue-winged Warbler	114
Golden-winged Warbler.....	118
Chestnut-sided Warbler.....	121
Black-throated Blue Warbler.....	124
Prairie Warbler	127
Bay-breasted Warbler	130
Blackpoll Warbler	133
Cerulean Warbler	137
Canada Warbler	140
Eastern Towhee	146
Field Sparrow.....	149
Vesper Sparrow	152
Grasshopper Sparrow	155
Bobolink	158
Eastern Meadowlark.....	161
Rusty Blackbird	164

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Common Loon**
Scientific Name: **Gavia immer**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S2B,S4N

State Trend: Stable

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Officially designated as Endangered in VT in 1978, statewide population has steadily rebounded from a low of 8 nesting pairs in 1983 and 1984 to 43 pairs in 2004.

Common Loon was designated as Endangered in VT in 1978, and removed from the list in 2005. The statewide population has steadily rebounded from a low of 8 nesting pairs in 1983 and 1984, to 301 adults and 66 chicks in 2014. Sustained management and monitoring has continued due to the Vermont Loon Conservation Project: a partnership with Vermont Center for Ecostudies and the Vermont Fish and Wildlife Department.

Distribution

Breeding is concentrated in northeastern and north-central VT, with confirmed nests found in the southern Green Mountains. Breeding probable in Champlain Valley. Loons are a medium-distance migrant. After leaving Vermont Loons head to coastal marine wintering locations along the entire eastern seaboard of the United States

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Unknown
Champlain Hills	Confident	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Territories range from 9 to 161 ha, prefers freshwater lakes larger than 24 ha, particularly those containing small islands and coves. Lakes with undisturbed islands or marshy shorelines, adequate fish and crayfish prey base, and clear water to a depth of at least 3 m.

Habitat Types:

Aquatic: Lacustrine

Aquatic: Lake Champlain

Aquatic: Man-Made Water Bodies



Common Name: **Common Loon**
Scientific Name: **Gavia immer**
Species Group: **Bird**

Current Threats

Habitat Threats:

Habitat Alteration

Climate Change

Incompatible Recreation

Description of habitat threat(s): Lakeshore development and subsequent loss of shoreline habitat is a threat to the breeding population at this time. Effects of climate change may impact this species.

Non-Habitat Threats:

Competition

Pollution

Trampling or Direct Impacts

Description of non-habitat threat(s): Interference competition from extraterritorial loons has caused some territory and nest abandonments, as well as direct killing of chicks and adults, in recent years. Vermont Loon Conservation Program participated in a study looking at mercury in Loons. They found that mercury bioaccumulates, resulting in a negative effect on adult behavior and chick productivity. Recreational activities and direct human disturbance of nesting or nursery sites also a serious problem on more heavily used lakes. Lead fishing gear likely killed three adult Loons in 2014 in Vermont. Maine and New Hampshire have also reported deaths due to lead poisoning

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	Low	This is well known overall, due to sustained annual monitoring since 1978, and the Vermont Breeding Bird Atlas, but changes due to loss of nesting areas should be documented quickly.
Research	Threats and Their Significance	High	1) Collaborative research on extent and possible effects of mercury contamination in VT loons should be continued. 2) Patterns of shoreline development and ownership of current and recent nest sites needs to be documented. 3) Effects of climate change may impact population and should be documented.
Monitoring	Population Change	High	Sustained monitoring is crucial to documenting population trends
Monitoring	Habitat Change	Low	Changes of suitable nesting habitats, and use of rafts.
Monitoring	Monitor Threats	Medium	Monitoring of all limiting factors goes hand-in-hand with population monitoring and is critical to evaluate long-term viability and management needs of statewide population. Sick, weak, and dead loons should be collected and sent to wildlife health facilities for determination of cause of death, including interference competition from other loons (trauma), lead, and other contaminants. Annual summaries of known causes of deaths should be completed and disseminated. Results should be evaluated for management applications.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Common Loon**
 Scientific Name: **Gavia immer**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications	High	Continue concerted public education effort targeting landowners and lake users. Volunteers should continue to receive training, toward long-term goal of having loon monitoring and management be largely volunteer-based.	Public presentations, informational signs at VFWD lake access areas, media articles, and informal meetings with lakeshore residents and recreationists are all crucial to continued public awareness. Platforms and sign buoys must be used as necessary.	VFWD, VCE	Nongame Fund, SWG
Easements	High	Long-term protection of all current and recent (within past 5 years) loon nest sites should be secured through conservation easements and land acquisition.	Ownership of all current and recent nest sites should be documented. For those nest sites not currently protected, landowners should be contacted and protocols for securing protection should be developed.	VFWD, VCE, TNC, Lake Associations, power companies	SWG, PR
Species Restoration	High	Monitoring and management should be continued and supported.	Annual management should be continued as necessary. Annual LoonWatch should be continued indefinitely.	VFWD, VCE	VFWD, USFWS

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Common Loon**
Scientific Name: **Gavia immer**
Species Group: **Bird**

Bibliography

Borden, S.E. and C.C. Rimmer. 1998. Vermont Loon Recovery Plan. Unpubl. Report. Vermont Institute of Natural Science, Woodstock, VT and Vermont Fish and Wildlife Department, Waterbury, VT. 44 pp.

Gallo, S. 2013. Maine Loon Mortality: 1987 to 2012. Maine Audubon

Hanson, E.W., C.C. Rimmer, and S. Parren. 2003. The 2003 breeding status of Common Loons in Vermont. Unpubl. Report. Vermont Institute of Natural Science, Woodstock, VT and Vermont Fish and Wildlife Department, Waterbury, VT.

Hanson, E. 2013. Common Loon (*Gavia immer*). In: The Second Atlas of Breeding Birds of Vermont, (R. Renfrew, Ed.). University Press of New England, Hanover and London

Hanson, E. 2014. What have we learned from color marked Loons. Loon Caller. Vermont Loon Conservation Program, Vermont center for Ecostudies, Norwich, VT and Vermont Fish and Wildlife Department, Waterbury, VT

National Audubon Society. 2014. Audubon's Birds and Climate Change Report: A Primer for Practitioners. National Audubon Society, New York. Contributors: Gary Langham, Justin Schuetz, Candan Soykan, Chad Wilsey, Tom Auer, Geoff LeBaron, Connie Sanchez, Trish Distler. Version 1.2.

eBird. 2012. eBird: An online database of bird distribution and abundance [web application]. eBird, Cornell Lab of Ornithology, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: Date January 21, 2015).

Evers, David C., James D. Paruk, Judith W. McIntyre and Jack F. Barr. 2010. Common Loon (*Gavia immer*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/313>



Common Name: **Pied-billed Grebe**
Scientific Name: **Podilymbus podiceps**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2B,S3N

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

A sporadic breeder in Vermont, but believed to have been much more common historically. Loss of suitable wetlands since European settlement may have greatly reduced population. Development of wetland buffers and increased disturbance of nesting sites by recreationists and boat wakes may have helped continue the population decline. The second Atlas of Breeding Birds of Vermont showed little change of statewide distribution, but a slight decrease in occurrences was seen. The Breeding Bird Survey of Vermont showed a long term trend (1966 - 2012) of -8.6%. The 10-year trend from 2003-2012 (-7.99%) mirrors the long term loss.

Distribution

Most breeding was documented within wetlands located in the Champlain Valley and Lake Memphremagog regions. During the second Atlas breeding documentation was lost from the Green Mountains, Vermont Valley and Southern Piedmont biophysical regions.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Unknown
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Lakes, ponds, marshes, swamps, and slow-moving streams and rivers. Prefers shallow, permanent marshlands with stable water levels.



Common Name: **Pied-billed Grebe**
Scientific Name: **Podilymbus podiceps**
Species Group: **Bird**

Habitat Types:

Hardwood Swamps
Softwood Swamps
Seeps and Pools
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Wet Shores
Shrub Swamps
Aquatic: Lower CT River
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Invasion by Exotic Species
Incompatible Recreation

Description of habitat threat(s): Conversion of Habitat - loss of wetlands due to draining, filling for development and agriculture. Habitat Degradation - outdoor recreational activities disturb nesting. Invasion by exotic species - common reed and purple loosestrife compete with native vegetation for nesting and feeding sites.

Non-Habitat Threats:

Harvest or Collection
Trampling or Direct Impacts

Description of non-habitat threat(s): Harvest or Collection - sometimes mistakenly shot as waterfowl. Trampling or Direct Impacts; nests susceptible to damage by boating and associated wakes. Mercury accumulation. Human disturbance of nest sites. Disturbance by recreational water users

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Pied-billed Grebe**
 Scientific Name: **Podilymbus podiceps**
 Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	More surveys needed to determine distribution and abundance.
Research	Threats and Their Significance	Medium	Impact of recreational activities at known nest sites.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications	High	Protect nesting areas from destructive recreational activities through enforcement, signing, press releases, educational materials, television/radio commercials.	Presence/absence of nesting grebes, number of chicks surviving to fledgling stage.	Audubon-VT, VINS, lake associations, angler groups, boating organizations.	SWG, Nongame fund, USFWS
Publically-Owned Protected Areas	High	Protect large wetlands (>20 ac.) suitable as grebe nesting habitat, acquired in fee through purchase.	Number of acres conserved in fee.	USFWS, DU, TNC	PR, DU, VHCB, VLT, Lake Champlain Land Trust, VT Waterfowl Stamp Funds
Policy & Regulations	High	Protect potential nesting habitat (large wetlands) through regulatory process.	Number of wetland acres protected from development.	VT-DEC	PR, EPA

Bibliography

- The Atlas of Breeding Birds of Vermont. 1985. Laughlin, S.B. and D.P. Kibbe. University of New England Press. 456 p.
- Muller, M.J. and R.W. Storer. 1999. Pied-billed Grebe (*Podilymbus podiceps*). In: The Birds of North America, No. 410 (A. Poole and F. Gill, eds.). The Birds of North America, Inc., Philadelphia, PA.
- Kibbe, D. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- Sauer, J. R., J. E. Hines, J. E. Fallon, K. L. Pardieck, D. J. Ziolkowski, Jr., and W. A. Link. 2014. The North American Breeding Bird Survey, Results and Analysis 1966-2012. Version 02.19.2014 USGS Patuxent Wildlife Research Center, Laurel, MD



Common Name: **American Bittern**
Scientific Name: ***Botaurus lentiginosus***
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G4

Global Trend:

State Rank: S3B,S3N

State Trend: Stable

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The distribution and population status of this species of regional conservation concern are not well documented in VT. The Breeding Bird Survey indicates a 0.22% increase over the long term trend (1966-2012) and a 0.38% increase over the last 10-years, but this is based on a small sample of routes and low abundance, so cannot be considered reliable. The Second VT Breeding Bird Atlas showed an increase of presence in all biophysical regions, except the Southern Green Mountains and Southern Vermont Piedmont regions. Vermont recently delisted the species from the status of species of special concern.

Distribution

From first VT Breeding Bird Atlas, confirmed breeding in large wetland complexes in Champlain Valley, also in West Rutland Marsh, sites in lower Connecticut River Valley, and two sites in north-central VT. Probably breeds in other larger wetland complexes (e.g., Memphremagog) and scattered smaller wetlands throughout the state. The second VT Breeding Bird Atlas confirmed breeding in marsh systems located in the northern portions of the state. The southern portions showed a stable or slightly decreasing presence.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Probable
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Confident	Taconic Mtns	Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Primarily freshwater wetlands with tall, emergent vegetation. Inhabits wetlands of all sizes (0.1-1,000 ha), but more abundant on larger than smaller wetlands. Prefers impoundments and beaver-created wetlands to those of glacial origin. Also found in wet swale of poorly drained fields.

Habitat Types:

Marshes and Sedge Meadows

Grasslands, Hedgerows, Old Field, Shrub, or Orchard



Common Name: **American Bittern**
Scientific Name: ***Botaurus lentiginosus***
Species Group: **Bird**

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Invasion by Exotic Species

Description of habitat threat(s):); Loss or degradation of wetland habitats the primary problem to this species throughout its range. Changes in wetland isolation and water stabilization may erode habitat quality. Invasion of Phragmites and purple loosestrife a further problem to native wetland vegetation.

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Chemical contamination and human disturbance are identified problems.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	Medium	1) More robust data are needed on the statewide distribution and abundance of this species in wetlands of different sizes and vegetative composition. 2) A standardized, coordinated statewide survey of this and other wetland birds is needed to establish baseline information on distribution and abundance. An extensive, single-season survey could be followed by annual monitoring at a core number of wetlands. A volunteer-based survey that uses standardized, repeatable protocols could collect presence/absence and relative abundance data at a large number of wetland sites statewide. A core number (12-15) of sites could be annually monitored for long-term trends.
Research	Threats and Their Significance	Medium	
Monitoring	Population Change	High	A standardized monitoring program is needed for this and other wetland birds. Extensive sampling needs to be coordinated periodically, while a core sample of wetlands should be monitored annually.
Monitoring	Habitat Change	Medium	Important to monitor habitat quality and changes that may be occurring, e.g. from invasive plants like Phragmites and purple loosestrife.
Monitoring	Monitor Threats	Medium	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Bittern**
 Scientific Name: ***Botaurus lentiginosus***
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Invasive Species Control & Prevention	Medium	Assess whether wetland habitat quality is compromised by invasive species like Phragmites and purple loosestrife; implement control measures at selected sites to eliminate or minimize these species; evaluate success of measures for AMBI and other birds.	Correlate presence/absence and changes in relative abundance of AMBI and other wetland species, in relation to natural and manipulated changes in vegetation composition caused by increase or elimination of invasives.	VFWD, USFWS, TNC, VTDEC	Wetland Reserve Program, NFWF, SWG
Policy & Regulations	Medium	Ensure that wetlands inhabited by this species are well-protected, including a representative sample of smaller wetlands. Ensure that further wetland loss or degradation in VT is minimized.	Conduct a spatially explicit inventory and evaluation of wetlands in VT, and assess local regulations for protecting them. Involve local conservation commissions in wetlands inventories and protection, also monitoring.	VFWD, USFWS, TNC, local Cons Comms	Wetland Reserve Program
Compatible Resource Use	Medium	Monitor wetland habitat quality (sedimentation rates, nutrient fluxes, water quality, chemical contamination) and correlate with changes in AMBI relative abundance or presence/absence.	Correlate habitat parameters with standardized AMBI survey data, and changes in both over time.	VFWD, USFWS, TNC, local Cons Comms	Wetland Reserve program

Bibliography

- Gibbs, J.P. and S.M. Melvin. 1992. American Bittern in Migratory nongame birds of management concern in the northeastern United States (K. Schneider and D. Pence, Eds.). U.S. Fish and Wildlife Service, Newton Corner, MA.
- Gibbs, J.P., S.M. Melvin, and F.A. Reid. 1992. American Bittern in The Birds of North America, No. 18 (A. Poole, P. Stettenheim, and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union.
- Kibbe, D. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- Sauer, J. R., J. E. Hines, J. E. Fallon, K. L. Pardieck, D. J. Ziolkowski, Jr., and W. A. Link. 2014. The North American Breeding Bird Survey, Results and Analysis 1966-2012. Version 02.19.2014 USGS Patuxent Wildlife Research Center, Laurel, MD



Common Name: **Least Bittern**
Scientific Name: ***Ixobrychus exilis***
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2B,S2N

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

No state BBS data. BBS routes are not well-sited for monitoring marsh species. Status of species unknown in state; the species is regularly found in relatively few marshes in VT (Kibbe 1985).

Distribution

Primarily found in the deep water marshes of Lake Champlain Valley and more sparsely in other favorable habitats in other parts of the state.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Not Probable	Vermont Valley	Confident
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Found in freshwater and brackish marshes with dense, tall growths of emergent vegetation interspersed with woody vegetation and open water. Most abundant in hemi-marsh conditions with stable water levels, rarely found in areas without standing water. In freshwater marshes, generally prefers cattails (*Typha* spp.; Poole et al. 2009).

Habitat Types:

Marshes and Sedge Meadows

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Alteration

Invasion by Exotic Species

Description of habitat threat(s): Loss of wetlands will continue to limit the species. Invasion of



Common Name: **Least Bittern**
 Scientific Name: ***Ixobrychus exilis***
 Species Group: **Bird**

wetlands by loosestrife and phragmites will degrade habitat quality. Agricultural and urban runoff could reduce water quality and prey populations. Greater frequency of storm events could lead to variation in water levels in wetlands, particularly along Lake Champlain, leading to decreased nesting success.

Description of non-habitat threat(s):

Research and Monitoring Needs

<i>Type</i>	<i>Need</i>	<i>Priority</i>	<i>Description</i>
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	Information about the distribution of LEBI in VT is lacking. A state-wide (perhaps one-time) survey of all potential wetlands would also yield valuable information for other wetland-dependent species (SORA, VIRA, COGA, PBGR, AMBI, BLTE). Marshbird monitoring programs are limited in their spatial extent in VT. An extensive initial survey would provide baseline data for a long-term monitoring program that would lay the foundation for a more representative marshbird monitoring program.
Monitoring	Population Change	High	Improving the standardization and spatial extent of marsh monitoring programs for wetland birds would greatly help our understanding of the species' distribution and population status.
Monitoring	Habitat Change	Medium	Most wetlands on which LEBI are found are protected, but more information about wetland loss and degradation would be useful as loss of wetlands will continue to limit LEBI. Although regulations currently in place will likely protect most nesting sites, some research indicates that LEBI is not area-sensitive (Gibbs and Melvin 1990) and may be found on wetlands ≤ 0.4 ha.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Least Bittern**
 Scientific Name: ***Ixobrychus exilis***
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Invasive Species Control & Prevention	Medium	Prevent wetland invasions by Phragmites and Purple Loosestrife and remove these species where they have already invaded in order to maintain/improve habitat quality for LEBI.	Presence/absence of LEBI in relation to changes in vegetation composition.	VFWD, TNC, USFWS.	NFWF, Marsh bird monitoring groups, TNC, Wetland Reserve Program (NRCS).
Compatible Resource Use	Medium	Decrease sedimentation rates and nutrient influxes into marshes currently containing LEBI to maintain habitat quality.	Ideally, annual variation in abundance of LEBI could be correlated with changes in habitat quality. More realistically, survey results will need to be based on presence/absence in relation to changes in water quantity, quality and vegetation.	VFWD, TNC, USFWS.	NFWF, Marsh bird monitoring groups, TNC, Wetland Reserve Program (NRCS).

Bibliography

Gibbs, J. P., and S. M. Melvin. 1990. An assessment of wading birds and other wetlands avifauna and their habitat in Maine. Final Report, Maine Dep. Inland Fish. Wildl., Bangor, Maine.

Poole, A. F., P. Lowther, J. P. Gibbs, F. A. Reid, and S. M. Melvin. 2009. Least Bittern (*Ixobrychus exilis*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/017doi:10.2173/bna.17>

Kibbe, D. P. 1985. Least Bittern. Pages 36-37 in S. B. Laughlin and D. P. Kibbe, eds. The atlas of breeding birds of Vermont. University Presses of New England, Hanover, New Hampshire, USA.

Kibbe, D. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.



Common Name: **Great Blue Heron**
Scientific Name: **Ardea herodias**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S2S3B,S5N

State Trend: Stable

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

Population currently stable with numbers increasing at largest colony site, Missisquoi National Wildlife Refuge.

Largest breeding colony (350 pairs) currently located at Missisquoi National Wildlife Refuge, second largest at Porters Bay on Lake Champlain (~100 pairs). Smaller colonies located throughout the state. Missisquoi colony stable until 2000 when 600 pair colony failed due to disturbance early in the season. Has recovered to approx. 350 pairs.

Distribution

Widely distributed with the largest colonies located in Champlain Valley. Smaller colonies located throughout state.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills		Vermont Valley	Unknown
Northern Green Mtns	Probable	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Unknown
Northeastern Highlands	Probable		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☐

Colony nester, nesting in tall trees, usually in wooded swamps. Colony size ranges from a couple of pairs to more than 500 pairs. Inhabits marshes, swamps, streams and lakeshores.

Habitat Types:

Hardwood Swamps
Seeps and Pools
Floodplain Forests
Softwood Swamps



Common Name: **Great Blue Heron**
Scientific Name: **Ardea herodias**
Species Group: **Bird**

Current Threats

Habitat Threats:

Habitat Alteration

Incompatible Recreation

Description of habitat threat(s): Loss or degradation of nesting habitat at colony sites, especially larger colony sites directly impacts population

Non-Habitat Threats:

Competition

Predation or Herbivory

Description of non-habitat threat(s): Disturbance of nesting colony early in the season has lead to abandonment of nesting colonies. Increasing numbers of nesting Double-crested Cormorants at large colony sites results in competition for nesting space and habitat degradation

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Determine specific habitat requirements for nesting locations.
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	Document known nesting locations in the state, primarily smaller nesting colonies.
Research	Threats and Their Significance	High	Continue research efforts into competition with cormorants in breeding colonies unknown. More research is needed to better understand dynamics between these 2 species and effects on heron breeding colonies. Determine impacts of Double-crested Cormorants
Research	Population Genetics	Low	
Monitoring	Population Change	High	Annually monitor known nesting colonies.
Monitoring	Habitat Change	High	Monitor habitat changes at colony sites especially degradation of nesting trees due to the presence of Double-crested Cormorants.
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	High	Monitor disturbance and nest site competition at colony sites.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Great Blue Heron**
 Scientific Name: **Ardea herodias**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Protected Area Management	High	Reduce competition at nesting locations by Double-crested Cormorants	Minimize impacts of cormorants on nesting herons by limiting the number nesting cormorants at the colony site	USFWS, VTFWS, UVM, TNC	USFWS, TNC
Compatible Resource Use	Medium	Attempted predation by Bald Eagles is suspected of causing colony abandonment. Potential eagle nesting near colony sites could result in the loss of the colony	Maintain largest two colonies (Missisquoi and Porters Bay) in Vermont	USFWS, VT FWS, UVM, TNC	USFWS, SWG
Protected Area Management	High	Protect colony sites from human disturbance early in the nesting season to decrease chances of abandonment.	Increased education and awareness of individuals using the area (primarily boaters) through outreach efforts and signage.	USFWS, VTFWS, Audubon VT, TNC	USFWS, SWG
Habitat Restoration	High	Stop or reverse loss of vegetation used for nesting (trees) at Missisquoi NWR due to impacts of expanding Double-crested Cormorant colony. Suitable nesting structure needs to be maintained.	Maintenance of current nesting structure and identification of other suitable habitat at Missisquoi NWR	UFWS, UVM, Audubon	UFWS

Bibliography

Laughlin, S.B. and D. P. Kibbe, editors. 1985. The Atlas of Breeding Birds of Vermont. University Press of New England, Hanover, New Hampshire, USA.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Black-crowned Night-heron**
Scientific Name: **Nycticorax nycticorax**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S1B,S2N

State Trend:

Extirpated in VT?

Regional SGCN? Yes

Assessment Narrative:

Currently no documented nesting in Vermont.

Black-crowned Night-herons have been documented nesting on Lake Champlain, Vermont with the largest colony (30-50 pairs) having been on Young Island in Lake Champlain. Competition for nest sites and degradation of habitat by Double-crested Cormorants on Young Island resulted in the abandonment of that colony in the mid 1990's.

Distribution

Has nested at 2 sites along Lake Champlain with the largest being Young Island on the northern part of the lake. Nesting has not been documented in the state since the mid 1990's.

Distribution by Biophysical Region:

Champlain Valley	Unknown	Southern VT Piedmont	Unknown
Champlain Hills		Vermont Valley	Unknown
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Unknown
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☐

Prefers islands and wooded swamps for nesting locations. Feeds along shoreline and within marshes and swamps

Habitat Types:

Hardwood Swamps
Floodplain Forests
Marshes and Sedge Meadows
Shrub Swamps



Common Name: **Black-crowned Night-heron**
Scientific Name: **Nycticorax nycticorax**
Species Group: **Bird**

Current Threats

Habitat Threats:

Habitat Alteration

Description of habitat threat(s): Degradation of habitat as a result of nesting Double-crested Cormorants

Non-Habitat Threats:

Competition

Description of non-habitat threat(s): Competition for nest sites with Double-crested Cormorants

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Determine specific nesting habitat requirements
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Currently there is no documented breeding of this species in Vermont although breeding suspected. Surveys for breeding pairs and colonies should be undertaken to better assess status in Vermont
Research	Threats and Their Significance	Medium	Determine limiting factors to potential breeding locations.
Monitoring	Population Change	High	Determine presence/absence of species in the state.
Monitoring	Habitat Change	Medium	
Monitoring	Monitor Threats	Medium	If nesting sites located determine and monitor potential limiting factors.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Protected Area Management	Medium	Rapid increase in cormorant population of Young Island has displaced BCNH. Efforts to reduce cormorant numbers and restore areas of the island for BCNH nesting may result in BCNH nesting here in the future.	BCNH nesting on Young Island	VTWD, Wildlife Services	US government, USDA
Habitat Restoration	Medium	Restore nesting structure (trees and shrubs) on Young Island to enhance nesting opportunities.	BCNH nesting on Young Island	VTFW, UVM	USDA Wildlife Services, USFWS



Common Name: **Black-crowned Night-heron**
Scientific Name: **Nycticorax nycticorax**
Species Group: **Bird**

Bibliography

Laughlin, S. B. and D.P. Kibbe, editors. 1985. The Atlas of Breeding Birds of Vermont. University Press of New England, Hanover, New Hampshire, USA.



Common Name: **American Black Duck**
Scientific Name: **Anas rubripes**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S5B,S5N

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Population decline is readily apparent, but likely due to a number of factors including habitat loss, hybridization with mallard, and marine pollution that affects molluscs, an important winter food source. The second Atlas of Breeding Birds of Vermont showed a 32% decrease in distribution. The Breeding Bird Survey of Vermont showed a 5.9% decrease in occurrence in the long term trend, (1966-2012) and a 5.44% decrease during the 10- year period form 2003-2012. The Atlantic Flyway Breeding Waterfowl Plot Survey showed the 2014 breeding population 11.19% below the long term average, (1993-2014). The downward slope of the long term trends for the traditional and eastern survey areas mirror the decreasing population estimates of the Atlantic Flyway Breeding Waterfowl Plot Survey.

Distribution

Widespread across Vermont. Highest occurrences are found in the Champlain Valley, the Northern and Southern Vermont piedmonts and Southern Green Mountain biophysical regions.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Uses a variety of wetland habitats along the coast, in woodlands, boreal forest, mixed conifer-hardwoods, wherever there is water nearby. Nests in dense shrub vegetation usually near water but sometimes up to a mile or more away. Preferred wintering habitat includes brackish marshes bordering bays, estuaries, and agricultural areas, but also found on inland lakes, reservoirs, and marshes wherever ice-free conditions exist.



Common Name: **American Black Duck**
Scientific Name: **Anas rubripes**
Species Group: **Bird**

Habitat Types:

Hardwood Swamps
Open Peatlands
Marshes and Sedge Meadows
Floodplain Forests
Shrub Swamps
Aquatic: Lower CT River
Aquatic: Lacustrine
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Alteration

Description of habitat threat(s): (): Habitat conversion and degradation- conversion of wetlands to agriculture, and loss due to development including shoreline construction, ditching and other drainage methods; road building, alteration of wetland hydrology; invasive species such as purple loosestrife, common reed. On the wintering areas coastal erosion and filling of wetlands for development are a concern.

Description of non-habitat threat(s):

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Black Duck**
 Scientific Name: **Anas rubripes**
 Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	A breeding black duck survey is needed to determine where birds are breeding, by wetland or woodland type.
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Distribution and abundance of breeding black ducks are not well known in Vermont.
Research	Threats and Their Significance	High	The main limiting factor includes the loss of shrub wetlands from agriculture and residential and commercial development, including alteration and degradation of habitats over time. This duck is more susceptible to human disturbances than other duck species due to its shy nature and tendency to abandon nests when disturbed.
Research	Population Genetics	Low	This species hybridizes with the mallard but it is not believed to be a long term threat.
Research	Taxonomy	Low	
Monitoring	Population Change	High	Regional trends show a declining population in the St. Lawrence River Valley and northern New England.
Monitoring	Habitat Change	High	Wetland inventories should be updated periodically and analyzed for changes in wetland abundance by wetland type (i.e. scrub-shrub wetlands as potential black duck nesting). habitat.
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	High	Same as Habitat change.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Conservation Payments/Financial Incentives	High	Work with farmers and provide incentives for protection of wetlands from agricultural conversion.	Number of acres protected from conversion.	USFWS, USDA-NRCS, Ducks Unlimited	EQIP, NAWCA, PR, DU
Compliance & Enforcement	High	Better enforce state and federal wetland laws, including buffer zones.	Number of wetland acres and wetland buffer acres protected under state Conditional Use Determination regulatory process and federal Clean Water Act.	VT-DEC, EPA, US Army COE	EPA, PR
Protected Area Management	Medium	Attempted predation by Bald Eagles is suspected of causing colony abandonment. Potential eagle nesting near colony sites could result in the loss of the colony.	Maintain largest two colonies (Missisquoi and Porters Bay) in Vermont)	USFWS, VFWD, UVM, TNC	USFWS, SWG



Common Name: **American Black Duck**
Scientific Name: **Anas rubripes**
Species Group: **Bird**

Bibliography

Bellrose, F.C. 1976. Ducks, Geese, and Swans of North America. Harrisburg, PA: Stackpole Books. 544 p.

Coulter, M.C. and W.R. Miller. 1968. Nesting biology of black ducks in northern New England. VT Fish and Game Dept. Bull., No. 68-2. Montpelier, VT.

Longcore, J.R., D. G. McAuley, G. R. Hepp, and J. M. Rhymer. 2000. American Black Duck (*Anas rubripes*). In *The Birds of North America*, No. 481 (A. Poole and F. Gill, eds.) The Birds of North America, Inc., Philadelphia, PA. Renfrew, R.B. 2013. *The Second Atlas of Breeding Birds of Vermont*. Lebanon, NH: University Press of New England. 548 p.

Sauer, J. R., J. E. Hines, J. E. Fallon, K. L. Pardieck, D. J. Ziolkowski, Jr., and W. A. Link. 2014. *The North American Breeding Bird Survey, Results and Analysis 1966-2012*. Version 02.19.2014 USGS Patuxent Wildlife Research Center, Laurel, MD

Strong, A. 2013. *The Second Atlas of Breeding Birds of Vermont*. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

U.S. Fish and Wildlife Service. 2014. Atlantic Flyway Breeding Waterfowl Plot Survey report 2014. U.S. Department of Interior, Washington, D.C. USA.

U.S. Fish and Wildlife Service. 2014. Trends in Duck Breeding Populations 1955-2014 report 2014. U.S. Department of Interior, Washington, D.C. USA.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Bald Eagle**
Scientific Name: **Haliaeetus leucocephalus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1B,S2N

State Trend: Increasing

Extirpated in VT? Yes

Regional SGCN? No

Assessment Narrative:

State listed as endangered. Removed from the federal list of endangered and threatened species in 2007. 2014 population of 18 territorial pairs within Vermont's borders. Vermont population has seen a steady increase since the first pair nested successfully in 2008, with an 800% increase from the first to the second atlas (Renfrew 2013). Breeds in all adjacent states and Quebec. See: VT Bald Eagle Recovery Plan (VFWD 2010). Currently close to meeting downlisting goals of 19 pairs, 50% of which nest successfully in VT (VFWD, unpublished data).

Distribution

Breeding and wintering concentrated in the CT River and Lake Champlain watersheds. Numerous incidental sightings throughout the year on additional waterbodies throughout the state.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Unknown
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Confident	Taconic Mtns	Unknown
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Breeding: Lakes & rivers with large trees for nesting, perching and roosting. Prefers minimal human. Disturbance (USFWS 1999, DeGraaf & Yamasaki 2001). Wintering: Large waterbodies with open water or good supply of carion (USFWS 1999).



Common Name: **Bald Eagle**
Scientific Name: **Haliaeetus leucocephalus**
Species Group: **Bird**

Habitat Types:

Hardwood Swamps
Floodplain Forests
Marshes and Sedge Meadows
Aquatic: Lower CT River
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lake Champlain
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Impacts of Roads or Transportation Systems
Climate Change

Description of habitat threat(s): Habitat loss and human disturbance considered significant problems to breeding eagles. Roads and/or trails near nest site can be detrimental if human activity is not restricted. (Buehler 2000). Climate change has the potential to reduce food supply.

Non-Habitat Threats:

Disease
Pollution
Loss of Prey Base
Trampling or Direct Impacts

Description of non-habitat threat(s): Eagles are most vulnerable to toxic substances - lead, mercury, pesticides, and other toxic chemicals (DeSorbo and Evers 2007). Also vulnerable to collisions with vehicles and power lines and possibly to disease (USFWS 1999, Buehler 2000). Potentially threatened by wind turbines (Pagel et al. 2013)

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Bald Eagle**
 Scientific Name: **Haliaeetus leucocephalus**
 Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Follow up on reports of nesting pairs
Research	Threats and Their Significance	Medium	Monitor potential effects of climate change and wind turbines
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	High	Effects of chemical threats such as mercury not well-known in Vermont.
Monitoring	Population Change	High	Monitor population and productivity.
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	High	Monitor potential threats of habitat loss, human disturbance,

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Species Restoration	High	Protect nesting habitat of nesting pairs using predator guards, signage, and long-term protection as necessary	Monitoring productivity at active breeding sites	VFWD, Audubon VT, VINS	PR, SWG, private grants
Awareness Raising and Communications	Medium	Educate the public about eagle ecology and the importance of minimizing disturbance	Public presentations, informational signs, media articles are all necessary for increased public awareness.	VFWD, Audubon VT, VINS	PR, SWG, private grants
Species Restoration	High	Implement the VT Bald Eagle Recovery Plan, including breeding season monitoring, winter surveys, etc.	Monitoring population's distribution and productivity	VFWD, Audubon VT, VINS, NH Audubon, NY DEC	PR, SWG, private grants
Research	Medium	Send any deceased eagles for necropsies and toxic chemical testing	Determine any mortality caused by toxic chemicals – mercury, lead, etc.	VFWD, USFWS, NY DEC, Tufts Univ	USFWS



Common Name: **Bald Eagle**
Scientific Name: **Haliaeetus leucocephalus**
Species Group: **Bird**

Bibliography

Buehler, D. A. 2000. Bald eagle (*Haliaeetus leucocephalus*). In the Birds of North America, No. 506 (A. Poole & F. Gill, eds.). The Birds of North America, Inc., Philadelphia, PA. 40pp.

DeGraaf, R. M. & M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover, NH. 482 pp.

DeSorbo, C.R., and D.C. Evers. 2007. Evaluating exposure of Maine's bald eagle population to Mercury: assessing impacts on productivity and spatial exposure patterns. Report BRI 2007-02. Biodiversity Research Institute, Gorham, ME.

Fowle, M. R. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Laughin, S. B., and D. P. Kibbe, eds. 1985. The atlas of breeding birds of Vermont. University Press of New England, Hanover, NH. 456 pp.

US Fish & Wildlife Service. 1983. Northern states bald eagle recovery plan. US Fish and Wildlife Service, Twin Cities, MN. 76pp.

USFWS. 1999. Endangered and threatened wildlife and plant; proposed rule to remove the bald eagle in the lower 48 states from the list of endangered and threatened wildlife. Federal Register 64(128): 36454-36464.

VT Department of Fish & Wildlife (VFWD) 2010. Vermont Bald Eagle Recovery Plan. VT Fish and Wildlife Department, Waterbury, VT. 38 pp.



Common Name: **Northern Harrier**
Scientific Name: **Circus cyaneus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2B,S3S4N

State Trend: Increasing

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Recent positive population trends in Atlas (Renfrew 2013) and stable BBS data for VT, NH, and NY (Sauer et al. 2011). There is a documented decline of this species in some eastern states (PA), and there may be the suggestion of a decreasing range but increasing population density in appropriate habitat (Renfrew 2013). Listed as VT Species of Special Concern, and listed as endangered in CT and NH. Recent population increases may be due to the fact that harriers can nest in wetter grasslands that cannot be mowed early in the season. Abandoned farmland in the past 20 years may have increased the number of unused wet agricultural fields. Primary threats are incompatible agricultural management, wetland destruction, development grasslands, and succession.

Distribution

Distributed in all biophysical regions of the state, but most concentrated in the Champlain Valley (Renfrew 2013).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Open wetlands, marshy meadows, wet, lightly grazed pastures, old fields, marshes, upland prairies, mesic grasslands, drained marshlands, croplands, cold desert shrub-steppe, and riparian woodland. Densest populations typically associated with large tracts of undisturbed habitats dominated by thick vegetation (MacWhirter and Bildstein 1996).

Habitat Types:

Marshes and Sedge Meadows

Grasslands, Hedgerows, Old Field, Shrub, or Orchard



Common Name: **Northern Harrier**
 Scientific Name: **Circus cyaneus**
 Species Group: **Bird**

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Succession

Habitat Fragmentation

Description of habitat threat(s): Major problem is likely early mowing of hayfields. Other problems include heavy grazing rotations in pastures, especially wet pastures and wetland drainage or destruction. Additionally, hayfield abandonment (succession) and urban/suburban development are also problems (Renfrew 2013).

Non-Habitat Threats:

Loss of Prey Base

Trampling or Direct Impacts

Description of non-habitat threat(s): Early mowing can destroy nests and decreases rodent populations.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Wetlands are likely a safer nesting habitat for this species in VT. Proportion of birds nesting in wetlands versus hayfields would be helpful from a management standpoint.
Research	Basic Life History	Medium	Many studies on home range size, little on territory size. The disparity between the two (240 ha mean hr size versus 0.8 to 10 ha territory size) creates major variation in potential recommendations for habitat requirements for 500 pairs. Additionally, determining the causes of breeding failure and mortality are important.
Research	Distribution and Abundance	Low	
Research	Threats and Their Significance	Medium	Better information of timing of nesting in relation to hay harvest. Data from first breeding bird atlas suggests nestling dates are much later than necessary to fledge young prior to a Memorial Day cutting.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	High	Population would be relatively easy to monitor through roadside counts. Demographics would be more difficult to assess.
Monitoring	Habitat Change	Medium	It would be useful to know the proportion of grasslands lost to forest succession versus urban/suburban development.
Monitoring	Monitor Threats	High	Better information about the timing of hayfield cutting.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Northern Harrier**
 Scientific Name: **Circus cyaneus**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications	High	Education and outreach program that provides information about grassland dependent species and management options to enhance their populations in Vermont.	Number of landowners reached. Number of cooperating landowners who are maintaining grasslands by periodic late-summer mowing. Periodically assess (5yrs) grassland acreage in Vermont, through GIS analysis.	UVM, Audubon VT, VCE, NRCS.	NRCS, USDA, PR, SWG
Technical Assistance, Training, Learning Networks	High	Provide technical assistance to town and regional planning commissions to help conserve grassland habitats from development.	Number of town and regional planning commissions reached. # of cooperating landowners maintaining grasslands by periodic late-summer mowing. Periodically assess (5yrs) grassland acreage in Vermont, through GIS analyses.	UVM, NRCS.	USDA, NRCS.
Conservation Payments/Financial Incentives	High	Enroll farmers in NRCS funding programs (EQIP) for early/late or delayed mowing regimes	Number of cooperating farmers who are enrolled in EQIP. Additionally, estimates of foraging success rates of birds before and after hay harvest would help assess questions of abundance vs. availability of prey.	UVM, Audubon VT, VCE, NRCS.	NRCS, USDA.
Protected Area Management	High	Maintain nesting habitat throughout breeding season by following site specific conservation plans which include restricting mowing after July 15 on publicly owned lands (WMAs and state airports).	Maintain and increase current acreage under management on state lands	VFWD, NRCS, VTrans	VFWD
Habitat Restoration	High	Maintain grassland habitat in suitable locations through active management of woody vegetation within Grassland Bird Focus Areas.	Increase and maintain available habitat in suitable locations	VFWD, Audubon VT, NRCS, USFWS	PR



Common Name: **Northern Harrier**
Scientific Name: **Circus cyaneus**
Species Group: **Bird**

Bibliography

Ellison, W. G. 1985. Northern Harrier. Pages 72-73 in S. B. Laughlin and D. P. Kibbe, eds. The atlas of breeding birds of Vermont. University Presses of New England, Hanover, New Hampshire, USA.

MacWhirter, R. B., and K. L. Bildstein. 1996. Northern Harrier (*Circus cyaneus*). In The birds of North America, No. 210 (A. Poole and F. Gill, eds.). The birds of North America, Inc., Philadelphia, Pennsylvania, USA.

Perlut, N. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Sauer, J. R., J. E. Hines, J.E. Fallon, K.L. Pardieck, D.J. Ziolkowski, Jr., and W.A. Link. 2011. The North American breeding bird survey, results and analysis 1966-2009. Version 3.23.2011. USGS Patuxent Wildlife Research Center, Laurel, MD.



Common Name: **Northern Goshawk**
Scientific Name: **Accipiter gentilis**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S3S4B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Relatively abundant and widespread, Holarctic; population trends are difficult to determine; no hard evidence of a significant decline in recent decades, but probably declining in some areas primarily as a result of habitat alteration (natureserve.org). Formerly nested principally in Canada, but expanded breeding range south into northeastern North America beginning around 1950 (Laughlin & Kibbe 1985). In Vermont regions show a declining trend (21% statewide) in populations since the first Breeding Bird Atlas in 1985 (Renfrew 2013).

Distribution

Distribution info from VT Breeding Bird Atlas (Laughlin & Kibbe 1985, Renfrew 2013). NOGOs found in all regions of the state, with 22 confirmed breeding pairs mainly in the Green Mountain region of the state, and 1 in the lower Champlain Valley. Most sightings were in areas of medium - high elevation, with all but 1 in the Champlain Valley in the hilly areas on the periphery of the region. Most regions in VT show a declining trend (21% statewide) in populations since the first atlas (Renfrew 2013).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

Breeding: Forest interior habitats prefers mature forests with large trees and open understories. Found in all elevations up to treeline (DeGraaf & Yamasaki 2001). Nests usually in bottom of the canopy of a large hardwood tree in the East (Laughlin & Kibbe 1985). In Minnesota, 81% of 46 goshawks nests were in aspen trees, generally located in mature (>50 years) early successional upland hardwood stands (aspen and paper birch forest types) (Boal et al. 2001). Prey is primarily small to medium birds, but will also feed on small mammals. Preferred feeding habitats are openings in forests (DeGraaf & Yamasaki 2001).

Breeding: nest site must be away from human disturbance. Occupancy of nest sites positively correlated with stand size. Nests usually in bottom of the canopy of a large tree - goshawks tend to prefer hardwoods for nest trees in the East (Laughlin & Kibbe 1985). In Minnesota, 81% of 46 goshawks nests were in aspen trees,



Common Name: **Northern Goshawk**
Scientific Name: **Accipiter gentilis**
Species Group: **Bird**

generally located in mature (>50 years) early successional upland hardwood stands (aspen and paper birch forest types). Remaining nests were in paper birch, white pine or red oak (Boal et al. 2001).

Habitat Types:

Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Northern Hardwood
Early Succession Boreal Conifers
Early Succession Boreal Hardwoods
Early Succession Northern Hardwoods
Early Succession Spruce-Fir
Early Succession Pine and Hemlock
Early Succession Upland Oak

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Fragmentation
Impacts of Roads or Transportation Systems

Description of habitat threat(s): Loss of interior mature forest caused decline when Europeans settled New England. Does not nest in small forest tracks bounded by roads (DeGraaf & Yamasaki 2001).

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Highly sensitive to human presence (DeGraaf & Yamasaki 2001). Some sensitivity to pesticides/toxic chemicals.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Northern Goshawk**
 Scientific Name: **Accipiter gentilis**
 Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Basic Life History	Low	Estimate productivity of nesting pairs.
Research	Distribution and Abundance	Medium	Determine population status and trends in VT (locate nesting pairs).
Research	Threats and Their Significance	Medium	Need to determine significance of limiting factors to habitat in Vermont and whether active management/protection of this species is needed.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	Medium	Monitor trends in Vermont population
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	Medium	Monitor limiting factors to VT population

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Publically-Owned Protected Areas	High	Manage a portion of Vermont public lands with long rotations or as no-cut reserves.	Number of productive nests on conserved public lands.	VFWD, USFS, USFWS, USFS	SWG, PR
Habitat Restoration	Medium	Identify contiguous forests blocks w/mature components & encourage their conservation via easements or other financial incentives on private lands. Conserve contiguous forest blocks on public lands via appropriate long-range management plan designations.	Number and distribution of core forest blocks conserved on private and public lands	ANR, USFS, USFWS, VHCB, VLT, TNC	SWG, PR, VHCB

Bibliography

Boal, C. W., D. E. Anderson, and P. L. Kennedy. 2001. Home range and habitat use of northern goshawks (*Accipiter gentilis*) in Minnesota. Minnesota Cooperative Fish and Wildlife Research Unit, St. Paul, MN. 48 pp.

DeGraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover, NH. 482 pp.

Kibbe, D. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Laughlin, S. B., and D. P. Kibbe. 1985. The Atlas of Breeding Birds of Vermont. University Press of New England, Hanover, NH. 456 pp.

www.natureserve.org comprehensive report on the Northern Goshawk. Accessed: March 31, 2005.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Red-shouldered Hawk**
Scientific Name: **Buteo lineatus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S3S4B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Declined by 37% from first atlas to second atlas (Renfrew 2013). Appears to be stable or increasing in surrounding states and Ontario, as well as North America as a whole. BBS data shows increasing trend in the East, but current populations are thought to be far below historic levels (early 1900's).

Distribution

Reported in all biophysical regions (Renfrew 2013).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

Mature forested wetlands near natural openings for foraging and upland forests adjacent to wetlands (DeGraaf & Yamasaki 2001). Requires riparian woodlands with tall trees for nesting. Nest sites are often in the largest deciduous trees and always near water, such as river, pond or swamp. Prey upon herptiles, crustaceans, insects, and small mammals (Laughlin & Kibbe 1985, DeGraaf & Yamasaki 2001).



Common Name: **Red-shouldered Hawk**
Scientific Name: **Buteo lineatus**
Species Group: **Bird**

Habitat Types:

Oak-Pine Northern Hardwood
Hardwood Swamps
Seeps and Pools
Northern Hardwood
Floodplain Forests
Softwood Swamps
Open Peatlands
Marshes and Sedge Meadows
Wet Shores
Shrub Swamps

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration

Description of habitat threat(s): Declines in early 1900s due to loss of wetland habitat (DeGraaf & Yamasaki 2001). Shown to be vulnerable to habitat conversion, including deforestation, development, and draining of wetlands (Dykstra et al. 2008).

Non-Habitat Threats:

Pollution
Trampling or Direct Impacts

Description of non-habitat threat(s): RSHAs have been shown to accumulate pesticides such as PCBs (Laughlin & Kibbe 1985). May also be out-competed by more aggressive species such as Red-tailed Hawks and Great-horned Owls (Renfrew 2013).

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Red-shouldered Hawk**
 Scientific Name: **Buteo lineatus**
 Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	Determine population size and productivity in VT.
Research	Threats and Their Significance	High	Evaluate limiting factors to population in VT
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	Medium	Monitor population changes.
Monitoring	Habitat Change	High	Monitor loss of habitat
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	Medium	

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Species Restoration	Medium	Develop and implement protocols to monitor and manage population so as to minimize impacts.		VFWD, Audubon VT, VCE	
Habitat Restoration	High	Identify remaining blocks of contiguous forests w/mature components & encourage their conservation via easements or other financial incentives on private lands. Conserve these blocks on public lands via appropriate long-range management plan designations	Number and distribution of core forest blocks conserved on private and public lands	ANR, USFS, USFWS, VHCB, VLT, TNC	SWG, PR, VHCB

Bibliography

- Corwin, K. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- DeGraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover, NH. 482 pp.
- Dykstra, C.R., J. L. Hayes, and S. T. Crocoll. 2008. Red-shouldered Hawk (*Buteo lineatus*). In the birds of North America online (A. Poole, Ed.) Cornell Lab of Ornithology, Ithaca, NY. Retrieved from <http://bna.birds.cornell.edu/bna/species/107>.
- Laughlin, S. B., and D. P. Kibbe. 1985. The atlas of breeding birds of Vermont. University Press of New England, Hanover, NH. 456 pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Kestrel**
Scientific Name: **Falco sparverius**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Recent negative population trends in Atlas (Renfrew 2013) and BBS data for VT (NS) and survey-wide (Sauer et al. 2011). Listed as Species of Special Concern in 2009. Concern about population in the Northeast as a whole. Development of farmland, habitat succession, West Nile virus, and changes in farming practices are issues in the northeastern US.

Distribution

Distributed in all biophysical regions of the state, but most concentrated in the Champlain Valley and Northern Vermont Piedmont (Renfrew 2013).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Minimum area requirements appear to be ~25 ha grassland. Reports of home range size are variable. "Typical" densities are 0.11 to 1.74/100 ha (assuming peripatric home ranges, 57 to 909 ha). However, greater densities have been reported of 5.4 and 27.4/100 ha (3 -18 ha home range sizes). Kestrels are cavity nesters and use woodpecker holes, farm building crevices, or human-made nest boxes (Smallwood and Bird 2002).

Habitat Types:

Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Succession



Common Name: **American Kestrel**
 Scientific Name: **Falco sparverius**
 Species Group: **Bird**

Habitat Fragmentation

Description of habitat threat(s): Loss of grassland habitat to forest succession, conversion of agricultural areas to urban/suburban development, or changes in farming practices to eliminate hedgerows and trees for nesting (Renfrew 2013).

Non-Habitat Threats:

Disease

Predation or Herbivory

Trampling or Direct Impacts

Description of non-habitat threat(s): Automobile collisions are a concern where nest boxes have been placed on interstate highway signs (Smallwood and Bird 2002). There are also concerns with mortality from West Nile virus (Medica et al. 2007) and predation from increasing Cooper's Hawk populations (Farmer et al. 2006).

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Research should focus on relative importance of grassland habitat and cavity availability. If cavity availability is limiting, species could benefit from an active nest box placement program.
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	This species is relatively conspicuous and roadside counts could provide an excellent index of statewide population trends.
Research	Threats and Their Significance	High	Species is a useful model for environmental contaminants and climate change. May be a useful indicator species as they feed on herbivorous insects in agricultural habitats.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	Low	
Monitoring	Population Change	High	Nest monitoring in MA cranberry bogs shows 4 consecutive years of population declines. Peterson (2003) suggests the species is "quietly slipping away in New England."
Monitoring	Habitat Change	High	It would be helpful to know whether development or forest succession is more important to habitat loss.
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	High	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Kestrel**
 Scientific Name: **Falco sparverius**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Conservation Payments/Financial Incentives	High	Conserve grassland/shrubland habitats on private lands.	Number and total area of sites conserved.	USDA, NRCS, USFWS, VHCB	EQIP, SWG, PR, VHCB
Technical Assistance, Training, Learning Networks	High	Educate agricultural community and general public about grassland birds and management options to protect habitat	Enroll landowners into USDA habitat incentive programs	VFWD, Audubon VT, NRCS, UVM	PR, SWG, private funding sources
Habitat Restoration	High	Maintain grassland habitat in suitable locations through active management of woody vegetation within Grassland Bird Focus Areas.	Increase and maintain available habitat in suitable locations	VFWD, Audubon VT, NRCS, USFWS	USFWS, NRCS
Technical Assistance, Training, Learning Networks	Medium	Develop a nest box program for interested landowners.	Nest box occupancy rates as reported by landowners.	Audubon VT, VFWD	EQIP

Bibliography

- Clark, G.A. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- DeGraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University of New England Press, Hanover, New Hampshire, USA, and London, UK.
- Farmer, G.C, K. McCarty, S. Roberston, B. Roberston, and K.L. Bildstein. 2006. Suspected predation by accipiters on radio-tracked American Kestrels (*Falco sparverius*) in eastern Pennsylvania, USA. *Journal of Raptor Research* 40:294-297.
- Medica, D.L., R. Clauser, and K. Bildstein. 2007. Prevalence of West Nile Virus antibodies in a breeding population of American Kestrels (*Falco sparverius*) in Pennsylvania. *Journal of Wildlife Diseases* 43:538-541.
- Petersen, W. R. 2003. New England Region. *North American Birds* 57(4):465-469.
- Sauer, J. R., J. E. Hines, J.E. Fallon, K.L. Pardieck, D.J. Ziolkowski, Jr., and W.A. Link. 2011. The North American breeding bird survey, results and analysis 1966-2009. Version 3.23.2011. USGS Patuxent Wildlife Research Center, Laurel, MD.
- Smallwood, J. A., and D. M. Bird. 2002. American Kestrel (*Falco sparverius*). In *The birds of North America*, No. 602 (A. Poole and F. Gill, eds.). The birds of North America, Inc., Philadelphia, Pennsylvania, USA.
- William, W. G., and W. G. Ellison. 1985. American Kestrel. Pages 86-87 in S. B. Laughlin and D. P. Kibbe, eds. *The atlas of breeding birds of Vermont*. University Presses of New England, Hanover, New Hampshire, USA.



Common Name: **Peregrine Falcon**
Scientific Name: **Falco peregrinus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G4

Global Trend:

State Rank: S2B,S2N

State Trend: Stable

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Officially designated as Endangered in VT in 1972, statewide population has steadily recovered and surpassed recovery goals.

See: VT Peregrine Falcon Recovery Plan (Fowle et al. 2001) and Post-delisting Monitoring and Management Plan. Removed from VT Endangered Species List in 2005. Increases due in large part to concerted monitoring and management efforts; population continues to require monitoring and site protection to ensure successful nesting (Renfrew 2013).

Distribution

Breeding well distributed throughout the state. Nest sites known in all biophysical regions except the Vermont Valley.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Unknown
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Open areas for hunting, adequate food supply and steep rocky cliffs for nesting (Ratcliffe, 1993). Cliffs must be high enough (at least 30 m) to protect from terrestrial predators, have adequate horizontal nesting ledges (loose substrate for nest, protection from weather and predators), and have adequate perches and good views of the surrounding area for territorial defense and hunting (Hickey 1942, Ratcliffe 1993). Human disturbance on or above the nesting cliff must be minimal and limited to more than 1/4 mile from the nest site, but activity such as a road below the nesting cliff will not negatively affect the nesting birds (USFWS 1991, Ratcliffe 1993).



Common Name: **Peregrine Falcon**
Scientific Name: **Falco peregrinus**
Species Group: **Bird**

Habitat Types:

Upland Shores
Cliffs and Talus
Building or Structure
Lawns, Gardens, and Row Crops
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Aquatic: Lower CT River
Aquatic: Lacustrine
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lake Champlain
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession

Description of habitat threat(s):): Inappropriate development, poorly planned forest management and habitat changes on or near a cliff during the breeding season may disturb nesting peregrines and cause them to abandon their nest site. Any development on or near a cliff may be enough to cause a pair to abandon that nest site (USFWS 1991, Fowle et al. 2001).

Non-Habitat Threats:

Pollution
Predation or Herbivory
Loss of Prey Base
Trampling or Direct Impacts

Description of non-habitat threat(s): Pesticides and other toxic chemicals have shown negative effects in the past, and some of these chemicals persist today (Fowle et al. 2001, USFWS, unpublished data). Fire retardant chemicals have been found in elevated levels in Vermont and northeastern peregrine populations (Eriksson et al. 2004, Da Chen et al. 2008). Human disturbance on or near nesting cliffs is the greatest known problem to peregrines nesting in VT. Predation of young on the nest site has been an occasional problem in the past, as has adverse weather (Fowle et al. 2001, Ratcliffe 1993).

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Peregrine Falcon**
 Scientific Name: **Falco peregrinus**
 Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	
Research	Threats and Their Significance	Medium	Potential effects of toxic chemicals and ridgeline wind development
Research	Population Genetics	Low	
Monitoring	Population Change	High	Monitor breeding population and productivity annually
Monitoring	Habitat Change	Medium	Protect breeding habitat from human disturbance and development
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	Medium	Monitor human disturbance effects and protect nesting cliffs from disturbance. Where possible, monitor other potential threats on productivity such as predation and adverse weather

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	Medium	Monitor site occupancy and productivity, as well as monitor threats	Continue to evaluate the health of the breeding population annually	VFWD, Audubon VT, VINS, landowners, volunteers	PR, SWG, Private grants
Compatible Resource Use	High	Reduce/minimize human disturbance at nesting cliffs through access closures during the breeding season.		VFWD, Audubon VT, CRAG-VT	PR, SWG, Private grants
Awareness Raising and Communications	High	Continue concerted public education effort targeting climbers and recreational hikers to inform public of cliff closures. Continue volunteer-based monitoring efforts.	Public presentations, informational signs at climbing and hiking areas, and media articles and posts are all critical to maintaining public awareness.	VFWD, Audubon VT, CRAG-VT	PR, SWG, private grants



Common Name: **Peregrine Falcon**
Scientific Name: **Falco peregrinus**
Species Group: **Bird**

Bibliography

Fowle, M. R., S. G. Parren, S. D. Faccio, D.W. Blodgett, and J. C. Heintz. 2001. Vermont peregrine falcon recovery plan. National Wildlife Federation, unpublished document, Montpelier, VT. 43 pp.

Da Chen, M.J. La Guardia, E. Harvey, M. Amaral, K. Wohlfort, and R.C. Hale. 2008. Polybrominated diphenyl ethers in Peregrine Falcon (*Falco peregrinus*) eggs from the Northeastern U.S. *Environmental Science and Technology* 42:7594-7600.

Eriksson, P., E. Jakobsson, and A. Fredriksson. 2001. Brominated flame retardants: A novel class of developmental neurotoxicants in our environment? *Environmental Health Perspectives* 109:903-908.

Fowle, M.R. 2013. *The Second Atlas of Breeding Birds of Vermont*. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Hickey, J.J. 1942. Eastern populations of the duck hawk. *Auk* 59:176-204.

Ratcliffe, D. A. 1993. *The peregrine falcon*. Second edition. T. & A. D. Poyser, London. 454 pp.

US Fish and Wildlife Service. 1991. First update of peregrine falcon (*Falco peregrinus*) eastern population, revised recovery plan. Newton Corner, MA. 35 pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Spruce Grouse**
Scientific Name: **Falcipennis canadensis**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1B

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The Spruce Grouse is a state listed endangered species. The species is not listed in the Breeding Bird Survey data likely due to its interior forest nature and limited vocalizations render it difficult to detect. Draft Recovery Plan for Vermont. Spruce grouse inhabit the boreal forests of North America. Although considered common in Canada and in a few northern states, in Vermont the species is near the southern edge of its range. Historical accounts indicate the species was present in the northeastern counties of Orleans and Essex (Thompson 1853, Cutting 1884). Currently, breeding spruce grouse are restricted to a 62 km² (25 mi²) area of spruce-fir forest in northern Essex County (Royar and Alexander 1987). This breeding habitat is principally owned by the U.S. Fish and Wildlife Service (Nulhegan Division of the Silvio Conte Refuge) and the State of Vermont, Department of Fish and Wildlife (Wenlock Wildlife Management Area).

It is estimated that between 150 and 300 adult birds occur in this population, and periodic surveys since 1990 indicate a stable population. Although the future of this population would seem secure given the interest of the two public owners in conserving wildlife (assuming that vegetation management will continue to maintain and improve habitat and that disturbance from the potential increase in recreationists to this area can be controlled) a stochastic event such as a widespread fire or disease outbreak could prove disastrous.

Full recovery of Vermont spruce grouse, outlined in its recovery plan, will require the establishment of 2 additional sub-populations, most likely on the State Lands located in the southern Essex County towns of Victory and Granby, and in the northern Essex County town of Norton. During 2008 and 2009 a total of 136 spruce grouse (males, females, and chicks) from Maine and Quebec were translocated to the Victory Basin Wildlife Management Area. This was an effort to establish a second viable population as per goals of the recovery plan. Subsequent surveys from 2009 through 2014 indicate the translocated birds are not successfully reproducing and declining in number due to inherent mortality factors.

Distribution

Spruce grouse inhabit the boreal forests of North America and are considered common in much of Canada. PIF reports a continental estimate of 11,000,000 birds. In Vermont, Maine, and New Hampshire the species is at the southern edge of its range. Historical accounts indicate the species was present in the northeastern Vermont counties of Orleans and Essex (Thompson 1853, Cutting 1884). Currently, breeding spruce grouse are restricted to a 62 km² (25 mi²) area of spruce-fir forest in northern Essex County (Royar and Alexander 1987). This breeding habitat is principally owned by the U.S. Fish and Wildlife Service (Nulhegan Division of the Silvio Conte Refuge) and the State of Vermont, Department of Fish and Wildlife (Wenlock Wildlife Management Area).

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Probable	Taconic Mtns	Not Probable



Common Name: **Spruce Grouse**
Scientific Name: **Falcipennis canadensis**
Species Group: **Bird**

Northeastern Highlands Confident

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

The spruce grouse is a bird of the boreal forest. Highest grouse densities (40-80 adults in summer/100 ha) are found in young dense jack pine (*Pinus banksiana*) stands, where live branches occur from 4-8 meters (13-26 ft.) above ground (Szuba and Bendell 1983; Keppie 1995). Jack pine forests do not occur in Vermont, however, spruce-fir forests of similar structure provide suitable habitat throughout much of the species range. Keppie (1987) documented breeding densities of 9.8 - 21.9 adults/100 ha (0.25-0.55/ac) in a New Brunswick spruce-fir pine forest. Spruce (*Picea* spp.) is preferred over fir (*Abies balsamea*) because it develops and maintains better vertical stratification. A shrub layer of *Vaccinium* spp. or regenerating spruce-fir in low densities enhances habitat for spruce grouse (Robinson 1969). Larch (*Larix laricina*) in the overstory may provide a preferred fall food resource. Forest openings are important to female spruce grouse and their broods, as they provide greater abundance of accessible food resources for chicks than the dense forest (Allan 1985).

Habitat Types:

Spruce Fir Northern Hardwood
Softwood Swamps
Open Peatlands
Early Succession Boreal Conifers
Early Succession Spruce-Fir

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Habitat Fragmentation

Description of habitat threat(s): Two main causes of historic spruce grouse decline are habitat loss and human encroachment. As colonial settlements expanded from southern Vermont into northeastern Vermont forests, spruce fir forests were cleared and the relatively tame spruce grouse was undoubtedly taken for human consumption whenever the opportunity arose. As industrial timber companies were formed, vast areas of virgin spruce fir forest were cut, and 19th century loggers may have taken spruce grouse to supplement their daily fare. By 1980 spruce grouse apparently remained only in the Nulhegan Basin, which at the time was experiencing heavy cutting of its mostly mature forests.

With the dawn of the new millennium, much of Essex County forests are publicly owned or subject to conservation easements held by the State and/or private conservation organizations. As a result, Conservation and sustained forestry goals across much of Essex County forestland will likely maintain if



Common Name: Spruce Grouse
Scientific Name: *Falcipennis canadensis*
Species Group: Bird

not increase available spruce grouse habitat. Increasing human development, however, will no doubt continue to encroach on some peripheral habitats, and the forecast increases in outdoor recreationists to Essex County could disrupt breeding activities and/or increase susceptibility to predation or adverse weather conditions (especially if pets accompany their owners on excursions through grouse habitats)

Non-Habitat Threats:

Harvest or Collection

Predation or Herbivory

Description of non-habitat threat(s): Predation is likely the most common cause of spruce grouse mortality (Boag and Schroeder 1992) although no predator seems to depend on spruce grouse as a large part of its diet (Robinson 1980). A major predator of spruce grouse eggs is the red squirrel (*Tamiasciurus hudsonicus*) (Boag et al 1984, Naylor and Bendell 1987). Other potential predators in Vermont are the northern goshawk (*Accipiter gentilis*), barred owl (*Strix varia*), northern raven (*Corvus corax*), red fox (*Vulpes vulpes*), coyote (*Canis latrans*), bobcat (*Lynx rufus*), black bear (*Ursus americanus*), striped skunk (*Mephitis mephitis*), fisher (*Martes americana*) and ermine (*Mustela erminea*). On Mount Desert Island, 7 of 19 radio transmitter-carrying adult females (37%) were predated between April and late August (Whitcomb et.al.1996). Predators identified were a Red-tailed Hawk (*Buteo jamaicensis*), red fox and an unidentified raptor. A study in Ontario of 67 spruce grouse nests found 55% were depredated by red squirrels, red fox, black bear and striped skunk (D'Eon 1997).

Another potential problem, especially in small patches that might hold dispersing grouse, is accidental shootings by ruffed grouse hunters. A limited number of accidental shootings were documented during the 1980's in Ferdinand and Norton. The continuation of educational efforts aimed at grouse hunters in Essex County should help prevent this source of mortality from actually limiting the population.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	Vermont's triennial census within the Nulhegan Basin should be continued, and reported sighting from elsewhere during the breeding season should be investigated
Research	Threats and Their Significance	Low	
Research	Population Genetics	Low	Genetic comparisons between Vermont birds and potential sources for reintroduction should occur.
Monitoring	Population Change	Medium	Vermont's triennial census within the Nulhegan Basin should be continued, and reported sighting from elsewhere during the breeding season should be investigated
Monitoring	Habitat Change	Medium	Extent of spruce-fir forests in Northeastern Vermont should be periodically assessed (eg USFS Forest Survey).
Monitoring	Monitor Threats	Low	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Spruce Grouse**
 Scientific Name: **Falcipennis canadensis**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Implement the habitat management guidelines as detailed in VFWD's Spruce Grouse Recovery Plan.	Number of public land management plans which incorporate Spruce Grouse habitat management.	VFWD, USFWS	SWG, PR
Species Restoration	High	Establish a 2nd sub-population in Victory Basin to reduce likelihood of serious impact to the overall population. Enhance genetic exchange with NH s grouse with establishment of another population within dispersal distance of the Connecticut River.	Number of sub-populations established and maintained.	ANR	SWG
Awareness Raising and Communications	Medium	Continue with educational campaign to reduce accidental harvest by ruffed grouse hunters.	Number of accidentally-shot spruce grouse	VFWD, USFWS	SWG, PR

Bibliography

Alexander, C. E. 2013. Second Atlas of Breeding Birds of Vermont. Rosalind Renfrew, ed. 547 pp. University Press of New England

Alexander, C.E. and J.J. Horton. 1986. Wenlock wildlife management area plan. VT Fish and Wildlife Department, St. Johnsbury. 54 pp.

Alexander, C.E. and R.B. Chipman. 1993. Model habitat guidelines for Spruce Grouse (*Dendragapus canadensis*) in Vermont. Vt. Fish and Wildl. Dept., St. Johnsbury. 7 pp.

Alexander, C. and S. Parren. 2012. Vermont Spruce Grouse Recovery Plan. Vermont Fish and Wildlife Department. Montpelier, VT 05620.

Allan, T.A. 1985. Seasonal changes in habitat use by Main spruce grouse *Dendragapus canadensis*. Can. J. Zool. 63(12):2738-2742.

Boag, D.A., S.G. Reebbs, and M.A. Schroeder. 1984. Egg loss among Spruce Grouse inhabiting lodge pole pine forests. Can. J. Zool. 62:1034-1037.

Cutting, H. 1884. Catalogue of the birds of Vermont. Eighth Vt. Agric. Rep. Vt. Dept. of Agriculture.

D'Eon, R.G. 1997. Vegetative concealment, proximity to trails, and predator activity as relative factors affecting nest success and egg loss in Spruce Grouse, *Dendragapus canadensis*. Canadian Field-Naturalist 111(3): 399-402.

Keppie, D.M. 1987. Impact of demographic parameters upon a population of Spruce Grouse in New Brunswick. J. Wildl. Manage. 51:771-777.

Keppie, D.M. 1995. Spruce Grouse Habitat. Paper presented at New York State Tree Farm Committee Forest Stewardship workshop, Paul Smith. 8 p.

Laughlin, S. B., and D. P. Kibbe. 1985. The Atlas of Breeding Birds of Vermont. University Press of New England, Hanover, NH. 456 pp.

LeMay, Y. and J. Ferron. 1987. Évaluation d'une technique de recensement pour le Tétrás des Savanes (*Dendragopus canadensis*) et proposition d'une méthode de mesure de l'expansion de la population introduite sur L'île d'Anticosti. Rapport technique présenté au Ministère du Loisir, del a Chasse et de la Pêche. Québec. 25 pp.

Naylor, B.J. and J.F. Bendell. 1987. Egg loss and nest site selection in Spruce Grouse. World Pheasant Association, IV International Grouse Symposium. Lam, West Germany. Pp 272-283.

Robinson, W.L. 1980. Fool Hen: the Spruce Grouse on the Yellow Dog Plains. Univ. Wisconsin Press, Madison.

Royar, K.J. and C.E. Alexander. 1987. Spruce grouse project performance report, 1987. Unpublished report, Vt. Fish and Wildl. Dept., Springfield. 15 pp.

Szuba, K.J. and J.F. Bendell. 1983. Population densities and habitats of spruce grouse in Ontario, pp. 199-213 in Resources and dynamics of the boreal zone, Proceedings of a Conference, Thunder Bay, Ont., August 1982. (R.W. Wein, R.R. Riewe, and I.R. Methuen, Eds). Association of Canadian Universities for Northern Studies, Ottawa.

Thompson, Z. 1853. Natural history of Vermont. Charles E. Tuttle Co., Rutland. 286 pp.

Whitcomb, S.D., A.F. O'Connell, Jr., and F.A. Servello. 1996. Productivity of the Spruce Grouse at the southeastern limit of its range. J. Field Ornithol., 67(3):422-427.



Common Name: **Ruffed Grouse**
Scientific Name: **Bonasa umbellus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

While no trend data is collected in Vermont, habitat declines and trends reported from other states (MA drumming survey, NH regional biologist information, ME hunter reports) indicate a general decline in ruffed grouse populations across New England. The potential exists for some of these declines to be related to ruffed grouse "cycles", these cycles have been shown to be less prevalent in the NE than in midwest and northern/subarctic regions.

Early successional habitat components required by ruffed grouse are declining on both a statewide and regional basis. Losses of acceptable habitat continue due to conversion to non-forest use. On areas which remain forested, stand maturation due to reductions in active forest management have substantially reduced habitat quality and grouse population density has decreased significantly. Conservation efforts should focus on implementing a comprehensive program of habitat improvements on both public land and private land.

Distribution

Distributed statewide where acceptable habitat components are present.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Prefers mosaic of young and mid-aged hardwood and hardwood/conifer forests. Typically utilizes maturing forest habitats for nesting, sapling/pole stage hardwood forest habitats for breeding, and very young hardwood forest regeneration habitats for brood rearing. While species can utilize and survive within edge and other suboptimal habitats, larger patch sizes of required habitat components adjacent to one another usually result in greater productivity and survival.



Common Name: **Ruffed Grouse**
Scientific Name: **Bonasa umbellus**
Species Group: **Bird**

Habitat Types:

Upland Shores
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Hardwood Swamps
Seeps and Pools
Open Peatlands
Marshes and Sedge Meadows
Northern Hardwood
Floodplain Forests
Softwood Swamps
Wet Shores
Early Succession Boreal Conifers
Early Succession Boreal Hardwoods
Early Succession Northern Hardwoods
Early Succession Other Types
Shrub Swamps
Early Succession Spruce-Fir
Early Succession Pine and Hemlock
Early Succession Upland Oak
Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Fragmentation
Invasion by Exotic Species
Inadequate Disturbance Regime

Description of habitat threat(s): - permanent loss of forest habitat to non-forest uses; - regional forest maturation resulting in suboptimal brood survival due to lack of protective cover and resulting increased predation; - fragmentation of dense regeneration habitats by mature forest resulting in substantially declining species productivity and significant mortality in localized, and sometimes large areas. Coupled with habitat loss to conversion of non-forest habitats, some ruffed grouse sub-populations have been extirpated in areas of former population abundance.

Non-Habitat Threats:

Parasites



Common Name: **Ruffed Grouse**
 Scientific Name: **Bonasa umbellus**
 Species Group: **Bird**

Predation or Herbivory

Description of non-habitat threat(s): Herbivory by white-tailed deer can substantially limit necessary brood habitats in areas of high deer abundance, although this is essentially a manifestation of an early-succession habitat volume-related problem.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Information on optimal habitat component patch size in various landscapes and forest cover types would be helpful.
Research	Basic Life History	Low	
Research	Distribution and Abundance	Low	Institute a ruffed grouse drumming survey and small-game hunter survey to establish ruffed grouse breeding population trends and harvest levels.
Research	Threats and Their Significance	Medium	Information on mortality due to parasites or nest predation in northern hardwood forest would be helpful.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	Medium	Little monitoring has been done on either productivity or mortality. This has compromised efforts to adequately measure changes due to habitat loss and respond to these population changes with corrective actions.
Monitoring	Habitat Change	High	See above.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Ruffed Grouse**
 Scientific Name: **Bonasa umbellus**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Species Restoration	High	Reverse VT ruffed grouse population trend decline to support an annual average hunting harvest of 150,000 birds over 10 years through improvement of grouse breeding and rearing habitat Use CSWA habitat target of 82,000ha (Rosenberg 2004).	Population response to management, BBS surveys.	Ruffed Grouse Society, FWD	PR, EQUIP
Awareness Raising and Communications	Medium	Initiate public education campaigns to highlight the need for active, even age forest management on public and private lands to create and maintain seedling/sapling forest habitat complexes.	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs.		
Habitat Restoration	Medium	Determine appropriate old field habitat targets for state lands and restore and maintain old field habitats where needed to increase suitable ES songbird habitat.	Number of acres positively affected by management. Population response to management.	ANR, USFS, Audubon, Forest Products Association, VT Loggers	PR, EQUIP
Conservation Finance	Medium	Create a state-funded, private lands, early successional habitat improvement initiative. Fund for > \$50,000/yr with revenues from state lands forest management.	Level of funds raised.	FWD	ANR, PR

Bibliography

- Bump, G., R.W. Darrow, F.C. Edminster, and W.F. Crissey. 1947. The Ruffed Grouse: Life History, propagation, management. NY State Conserv. Dept., Albany.
- Dessecker, D.R. and D.G. McAuley. 2001. Importance of early successional habitat for game birds. Wildlife Society Bulletin 29: 456-465.
- Gullion, G.W. 1984. Managing northern forests for wildlife. Ruffed Grouse Society, Coraopolis, Pennsylvania, USA.
- Kibbe, D. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- Rosenberg, K. V. 2004. Partners in Flight continental priorities and objectives defined at the state and bird conservation region levels, Vermont. Cornell Lab of Ornithology, Ithaca, NY. 26 p.
- The Birds of North American, (A. Poole and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, D.C.: The American Ornithologists' Union.
- Weik, A. P. 2001. Ruffed Grouse Assessment Summary. Maine Department of Inland Fisheries and wildlife. Bangor, Maine.



Common Name: **Sora**
Scientific Name: **Porzana carolina**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S2S3B,S3N

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Currently listed as a species of special concern in Vermont. Known as a breeder in the state but distribution and abundance unknown. Although uncommon in the state, the species increased by 50% between the first and second breeding bird atlas, perhaps as a result of more concerted use of playbacks.

Distribution

In the second Vermont Breeding Bird Atlas, 60% of records were from the Champlain Valley, with the remainder scattered across the state, generally at lower elevations.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Unknown
Champlain Hills	Not Probable	Vermont Valley	Confident
Northern Green Mtns	Not Probable	Southern Green Mtns	Unknown
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Found primarily in seasonal or semi-permanent freshwater wetlands with shallow and intermediate water depths (~0.33 m), dominated by emergent vegetation, especially cattails (*Typha* spp.), sedges (*Carex* spp., *Cyperus* spp.), burreeds (*Sparganium* spp.) and bulrushes (*Scirpus* spp.; Melvin and Gibbs 2012). Some suggestion that the species is area sensitive, but data are inconclusive.

Habitat Types:

Marshes and Sedge Meadows

Current Threats

Habitat Threats:

Conversion of Habitat
Climate Change
Invasion by Exotic Species



Common Name: **Sora**
 Scientific Name: **Porzana carolina**
 Species Group: **Bird**

Description of habitat threat(s): Habitat Threats: Loss of wetlands will continue to limit the species. Invasion of wetlands by loosestrife and phragmites will degrade habitat quality. Greater frequency of storm events could lead to variation in water levels in wetlands, particularly along Lake Champlain, leading to decreased nesting success.

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Agricultural and urban runoff could reduce water quality and prey populations.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Determine habitat requirements specific to Vermont
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	Information about the distribution of Sora in VT is lacking. A state-wide (perhaps one-time) survey of all potential wetlands would also yield valuable information for other wetland-dependent species (VIRA, COGA, PBGR, AMBI, LEBI, BLTE). Marshbird monitoring programs are limited in their spatial extent in VT. An extensive initial survey would provide baseline data for a long-term monitoring program that would lay the foundation for a more representative marshbird monitoring program.
Research	Threats and Their Significance	Medium	Determine potential limiting factors
Monitoring	Population Change	High	Improving the standardization and spatial extent of marsh monitoring programs for wetland birds would greatly help our understanding of the species' distribution and population status.
Monitoring	Habitat Change	Medium	1) Most wetlands on which Sora are found are protected, but more information about wetland loss and degradation would be useful as loss of wetlands will continue to limit Sora. 2) Monitor habitat changes at known nesting locations
Monitoring	Monitor Threats	Low	

Bibliography

Kibbe, D. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Laughlin, S. B. and D.P. Kibbe, editors. The atlas of breeding birds of Vermont. University of New England Press. Hanover, New Hampshire, USA.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Lesser Yellowlegs**
Scientific Name: **Tringa flavipes**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S4S5N

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

There is limited information on the population size of Lesser Yellowlegs. The best estimate is 400,000 birds worldwide. The species has been experiencing long-term (> 30 year) population declines (Andres et al. 2012). Impacts to breeding areas (Canadian tundra, muskeg) include commercial development, extraction of earth resources, and oil and gas development. Wetlands along migration routes in VT may be jeopardized by development also. Pollutants and un-regulated hunting may be concerns on wintering grounds (S. America)

Distribution

Found in all biophysical regions of Vermont

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Unknown
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Nests in open forest and forest-tundra transitional habitat, the vast majority of which (over 90%) occurs in the boreal forest of Alaska and Canada. Typical nesting habitat is open to semi-open forest mixed with marshes, bogs, sedge meadows, and ponds (Clay et al. 2012), using shallow wetlands and muskeg areas with abundant aquatic invertebrates. Migratory habitats include lake shores, river banks, and wetlands near agricultural area and early successional forests and shrub patches.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Lesser Yellowlegs**
 Scientific Name: **Tringa flavipes**
 Species Group: **Bird**

Habitat Types:

Open Peatlands
 Marshes and Sedge Meadows
 Grasslands, Hedgerows, Old Field, Shrub, or Orchard
 Wet Shores
 Shrub Swamps
 Aquatic: Lower CT River
 Aquatic: Fluvial
 Aquatic: Large Lake Champlain Tribs Below Falls
 Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat

Description of habitat threat(s): Conversion of wetlands to agriculture and residential development along shorelines, oil and gas development, logging, and mining activities. As a boreal forest-breeding species it is susceptible to climate change as higher temperatures may lead to drought and habitat changes.

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Agrochemicals are used throughout the species' migration corridors and nonbreeding grounds, the effects of which are unknown. Hunting pressure has presumably declined, but unregulated hunting during fall migration is a concern in the Caribbean and the Guianas.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Water level management to attract shorebirds. Timing and duration of drawdowns is critical to providing stopover areas along migration routes in VT..
Research	Distribution and Abundance	Medium	More shorebird surveys are needed around Lake Champlain and during drawdowns at state wildlife management areas.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Policy & Regulations	Medium	Protect shorelines through ACT 250 and other regulatory processes.	Number of shoreline acres or feet impacted by development.	Audubon-VT, VT-DEC, USFWS	EPA, Nongame fund, Partners for Fish and Wildlife, EQIP



Common Name: **Lesser Yellowlegs**
Scientific Name: **Tringa flavipes**
Species Group: **Bird**

Bibliography

Andres, B.A., Smith, P.A., Morrison, R.I.G., Gratto-Trevor, C.L., Brown, S.C, and C. A. Friis. 2012. Population estimates of North American shorebirds, 2012. Wader Study Group Bulletin 119:178–194.

Clay, R.P., A.J. Lesterhuis, and S. Centrón. 2012. Conservation Plan for the Lesser Yellowlegs (*Tringa flavipes*). Version 1.0. Manomet Center for Conservation Sciences, Manomet, Massachusetts.

Tibbitts, T.L. and W. Moskoff. 1999. Lesser Yellowlegs (*Tringa flavipes*). In: The Birds of North America, No. 427.



Common Name: **Upland Sandpiper**
Scientific Name: **Bartramia longicauda**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2S3B,S3N

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Breeding population possibly extirpated in Vermont, no confirmed breeding records in at least 5 years. Currently listed as endangered in Vermont.

Upland Sandpiper numbers have dropped sharply in Vermont since the early 1990's, with population losses from 1966-2003 estimated at -9.6 per year (Sauer et al. 2014), and the breeding population has disappeared from the state since the first SWAP. The small numbers reported during the second Breeding Bird Atlas (2003-2007) in the Champlain Valley and at the Berlin airport are no longer present. Habitat loss due to direct loss and agricultural intensification possible causes, as well as range contraction in the northeastern US. Broadcast methods designed to encourage establishment of a breeding population have been unsuccessful.

Distribution

In recent years, only 1-2 individuals reported during breeding season, from same location in northern Franklin County.

Distribution by Biophysical Region:

Champlain Valley	Historic Records Only	Southern VT Piedmont	Unknown
Champlain Hills	Probable	Vermont Valley	Unknown
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Unknown
Northeastern Highlands	Unknown		

Distribution by Watershed:

Known Watersheds

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

A migrant that winters in South America, Upland Sandpiper prefers large grassland areas (50-100 acres) with a mosaic of grassland vegetation structures (short, medium, and taller grasses for foraging, nesting, and brood rearing, primarily pastures and hayfields in Vermont. Will use airports with suitable mowing regime.



Common Name: **Upland Sandpiper**
Scientific Name: **Bartramia longicauda**
Species Group: **Bird**

Habitat Types:

Lawns, Gardens, and Row Crops
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Other Cultural

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Habitat Fragmentation

Description of habitat threat(s): Loss of habitat due to development and succession of abandoned farmland, fragmentation of large agricultural grasslands, and agricultural intensification (conversion to row crops, early haying regimes).

Non-Habitat Threats:

Trampling or Direct Impacts

Description of non-habitat threat(s): Destruction of nest site from mowing.

Research and Monitoring Needs

Type	Need	Priority	Description
Monitoring	Population Change	Medium	Track, verify, and monitor any observations recorded during breeding period. Note habitat structure and landscape characteristics at any site where there is evidence of breeding.



Common Name: **Upland Sandpiper**
 Scientific Name: **Bartramia longicauda**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Compatible Resource Use	High	Implement Vermont grassland bird management and recovery plan (LaBarr et al. 2013)			
Technical Assistance, Training, Learning Networks	High	Educate agricultural community and general public about grassland birds and management options to protect habitat	Continue grassland bird outreach programs and initiation of landowner information-sharing network	VFWD, Audubon-VT, VCE, UVM	PR, SWG
Habitat Restoration	High	Maintain grassland habitat in focal areas through active management of woody vegetation.	Increase and maintain available habitat in focal habitat	VFWD, Audubon-VT, NRCS	SWG, EQIP
Conservation Payments/Financial Incentives	High	Protect potential breeding habitat in grassland bird focal areas (see LaBarr et al. 2013, Vermont Grassland Bird Management Plan) by focusing EQIP payments on private lands in these areas.	Protection of acreage through enrollment in EQIP	VFWD, Audubon-VT, NRCS	EQIP

Bibliography

Jones, A.L. and P.D. Vickery. 1997. Conserving grassland birds: managing large grasslands including conservation lands, airports, and landfills over 75 acres for grassland birds. Massachusetts Audubon Society, Lincoln, Mass. USA.

LaBarr, M. L., Strong, A., Renfrew, R., Buck, J. and S. Parren. 2013. The Vermont grassland bird management and recovery plan. Vermont Fish and Wildlife Dept. Montpelier, VT.

Sauer, J. R., J. E. Hines, J. E. Fallon, K. L. Pardieck, D. J. Ziolkowski, Jr., and W. A. Link. 2014. The North American Breeding Bird Survey, Results and Analysis 1966 - 2012. Version 02.19.2014 USGS Patuxent Wildlife Research Center, Laurel, MD.

Perlut, N. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.



Common Name: **American Woodcock**
Scientific Name: **Scolopax minor**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

PIF Tier IIA -- high regional concern. Annual singing-ground survey trends for the Eastern Region and Vermont for the period 1968-2004 = -2.1 (P<0.01) and -1.1%, respectively (Kelly 2004). Range wide declines in American woodcock have been tied to similar declines in habitat area and quality, and losses of these habitats appears to be accelerating. Existing, moist-soil early-successional hardwood habitat (especially alder and aspen-dominated sites), and open field components required by woodcock should be identified on both public and some conserved private land, and these habitats should be actively managed to prevent further losses or qualitative declines. Additional work should focus on identifying areas where active habitat management would re-establish quality regeneration and open field habitat components across all biophysical regions.

Distribution

American woodcock are present statewide where acceptable habitat exists. During migration, woodcock numbers increase and birds can often be found less optimal habitat. While distributed across all Vermont biophysical regions, the relative scarcity of critical habitat components w/in these polygons makes this distribution map somewhat a misleading gauge of habitat security.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconnic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Prefers young hardwood, hardwood/conifer and alder forests proximate to open-field habitats in moist soil areas. Typically utilizes dense alder or aspen regeneration forest habitats for nesting, brood rearing and adult feeding, open field or forest openings > 1 ac. for breeding and roosting. While species can utilize and survive within moist soil forest edge and other suboptimal habitats, larger patch sizes of required habitat components adjacent to one another usually result in greater productivity and survival.



Common Name: **American Woodcock**
Scientific Name: **Scolopax minor**
Species Group: **Bird**

Habitat Types:

Outcrops and Alpine
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Hardwood Swamps
Seeps and Pools
Northern Hardwood
Floodplain Forests
Open Peatlands
Wet Shores
Early Succession Boreal Hardwoods
Early Succession Pine and Hemlock
Early Succession Northern Hardwoods
Early Succession Other Types
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Shrub Swamps
Early Succession Upland Oak

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Inadequate Disturbance Regime
Habitat Fragmentation

Description of habitat threat(s): Losses of moist-soil regeneration/successional habitats has been implicated in 30-yr population decline in both eastern and central flyways. Additional concerns regarding conversion to agriculture and non-forest cover of overwintering habitat in southern U.S. Fragmentation of both field habitats by reforestation and feeding/brood cover by succession has likely increased brood mortality during post-hatch brood movements to adequate rearing habitat.

Non-Habitat Threats:

Pollution
Predation or Herbivory

Description of non-habitat threat(s): Some anecdotal and emerging scientific data concerns regarding soil contaminants (primarily heavy metals) inducing adult mortality and compromising



Common Name: **American Woodcock**
Scientific Name: **Scolopax minor**
Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Habitat/breeding success interrelationship information would help to formulate range-wide recovery strategies.
Research	Basic Life History	Medium	AOU recommends studies to investigate large-scale population dynamics as related to habitat.
Research	Distribution and Abundance	Low	AOU recommends studies to investigate potential range expansions, however these likely would not be warranted in Vermont or established , central portions of range.
Research	Threats and Their Significance	Medium	As related to Habitat requirements, IAFWA has conducted work on habitat v. hunting mortality. Additional work could expand on this type of comparative mortality assessment, including such elements as soil contamination, losses by domestic predators and potential breeding losses due to various statutory restrictions on vegetation management in riparian and other "buffer" areas.
Monitoring	Population Change	High	Ongoing singing ground surveys should be enhanced and updated to consider historic habitat changes and other factors such as degree of development, etc.
Monitoring	Habitat Change	High	Habitat losses have largely been "tracked" by USFS Forest statistics in terms of age class/cover type composition. A more focused approach to estimation of historic, current and projected "woodcock habitat" across the region is certainly warranted.
Monitoring	Range Shifts	Medium	See above research needs re: distribution.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Woodcock**
 Scientific Name: **Scolopax minor**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Species Restoration	High	Reverse declining woodcock breeding populations to BPOP of 3.0-3.2 males/singing ground route. Maintain population w/ breeding habitat enhancement and creation/maintenance of suitable migration/feeding habitat. Use CSWA habitat target (Rosenberg 2004)	Population response to management, BBS surveys.	Ruffed Grouse Society, FWD	PR, EQIP
Awareness Raising and Communications	Medium	Initiate public education campaigns to highlight the need for active, even age forest management on public and private lands to create and maintain seedling/sapling forest habitat complexes.	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs.		
Habitat Restoration	Medium	Determine appropriate old field habitat targets for state lands and restore and maintain old field habitats where needed to increase suitable ES songbird habitat.	Number of acres positively affected by management. Population response to management.	ANR, USFS, Audubon, Forest Products Association, VT Loggers	PR, EQIP
Conservation Finance	Medium	Create a state-funded, private lands, early successional habitat improvement initiative (modeled on NH's Small Landowner Grant program). Fund for > \$50,000/yr with revenues from state lands forest management. This could offset landowner EQIP obligations.	Level of funds raised.	FWD	EQIP
Species Restoration	Medium	Continue and increase efforts at singing ground survey participation and observer recruitment.			
Habitat Restoration	Medium	Increase the size and number of well distributed roosting/display field habitats in proximity to feeding and brood habitat on public land.	Number and distribution of roosting/display fields		



Common Name: **American Woodcock**
Scientific Name: **Scolopax minor**
Species Group: **Bird**

Bibliography

Dessecker, D.R. and D.G. McAuley. 2001. Importance of early successional habitat for game birds. Wildlife Society Bulletin 29: 456-465.

Association of Fish and Wildlife Agencies (IAFAWA), 1999. The Report of the Woodcock Task Force to the Migratory Shore & Upland Game Bird Subcommittee, Migratory Wildlife Committee. Declines in American Woodcock Populations: Probable Cause and Management Recommendations. March, 1999.

Kibbe, D. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Kelley, J.R., Jr. 2004. American woodcock population status, 2004. U.S. Fish and Wildlife Service, Laurel, Maryland. 15pp.

Keppie, D.M. and R.M. Whiting, Jr. 1994. American Woodcock (*Scolopax minor*). In: The Birds of North America, No. 100 (A. Poole and F. Gill, Eds.) Philadelphia: The Academy of Natural Sciences; Washington, D.C.: The American Ornithologists' Union.

Rosenberg, K. V. 2004. Partners in Flight continental priorities and objectives defined at the state and bird conservation region levels, Vermont. Cornell Lab of Ornithology, Ithaca, NY. 26 p.



Common Name: **Common Tern**
Scientific Name: **Sterna hirundo**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1S2B,S2N

State Trend: Stable

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

VT state endangered since 1988. Common Tern populations in Vermont declined during the 1970's and 80's from about 300-400 breeding pairs to approx. 50 breeding pairs in 1988. Since then numbers have increased steadily due to monitoring and management efforts and protection of nesting islands. Breeding numbers have recently exceeded the levels recommended for down-listing to Threatened in Vermont but continuing low productivity has prevented down-listing. Monitoring and management efforts will need to continue to avoid a population decline in the future.

Distribution

Nests only on 4-5 small rocky islands (<0.5 ha) in the NE arm of Lake Champlain. Has not been documented nesting elsewhere in VT. Popasquash and Rock island primary nesting sites. Both are small islands in the NE arm of Lake Champlain. Has nested in the past on 3 other island. A social attraction (decoys and sound) project in 2006-2008 was successful in bringing terns to Grammas Island but heavy owl predation at this site resulted in the discontinuation of this work. Can be observed throughout the northern part of the lake.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Unknown
Champlain Hills	Unknown	Vermont Valley	Unknown
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Unknown
Northeastern Highlands	Unknown		

Distribution by Watershed:

Known Watersheds

Lake Champlain

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☐

Nests on isolated islands, beaches, dredge spoils and human made structures in areas with little to no vegetation.

Habitat Types:

Aquatic: Lake Champlain



Common Name: **Common Tern**
Scientific Name: **Sterna hirundo**
Species Group: **Bird**

Current Threats

Habitat Threats:

Habitat Alteration

Description of habitat threat(s): Traditional nesting islands are currently protected for this species. No direct habitat problems documented although habitat degradation due to nesting and roosting Double-crested Cormorants possible. Current management actions have limited cormorant related degradation in recent years.

Non-Habitat Threats:

Competition

Predation or Herbivory

Trampling or Direct Impacts

Description of non-habitat threat(s): Predation by avian predators and ants, competition for nesting sites with Ring-billed gulls and human disturbance are primary problems to this species.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	Well documented.
Research	Basic Life History	Low	Well documented.
Research	Distribution and Abundance	Low	Well documented.
Research	Threats and Their Significance	Low	Well documented.
Research	Population Genetics	Low	Well documented.
Monitoring	Population Change	High	Annual monitoring needed to determine population size and reproductive success.
Monitoring	Habitat Change	Medium	Annual monitoring of impacts of Ring-billed Gull and Double-crested Cormorants and impacts on island vegetation.
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	High	Annual monitoring required to determine impacts of predation, nest-site competition, and human disturbance on breeding population

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Common Tern**
 Scientific Name: **Sterna hirundo**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Protected Area Management	High	Provide adequate nesting space by managing gull and cormorant populations as competition for nesting space on tern nesting islands by Ring-billed Gulls and Double-crested Cormorants may result in limited nesting space for terns	breeding success, number of nests per island.	VFWD, Audubon-VT	PR, private grants
Protected Area Management	High	Limit predation by owls, night herons and ants at nesting islands through active management (fencing, trapping owls, eradicating ants).	Sustained increase in reproductive success to 1 fledgling/pair	VFWD, Audubon-VT	PR, private grants
Species Restoration	High	Continue to prevent Double-crested Cormorants from nesting as nesting islands could be negatively impacted by Double-crested Cormorants resulting in alteration of current vegetative cover	Number of islands free of double-crested cormorants.	VFWD, Audubon-VT, Lake Champlain Land Trust	PR, private grants
Protected Area Management	High	Continue to restrict access to tern nesting islands during the breeding season.	No documented nest failure due to human disturbance	VFWD, Audubon-VT, VT State Police	PR, private grants

Bibliography

- Fowle M, M. LaBarr and K. Wohlfort. 2014. The 2014 Report For Common Tern, Peregrine Falcon, Bald Eagle & Other Species Of Greatest Conservation Need In Vermont. Unpubl. report. Audubon Vermont, Huntington, Vermont.
- LaBarr, M. 1996. The Vermont Common Tern Recovery Plan. Unpubl. report, Vermont Institute of Natural Science. Woodstock, VT.
- LaBarr, M. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Black Tern**
Scientific Name: **Chlidonias niger**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: S2B,S2N

State Trend: Stable

Extirpated in VT?

Regional SGCN? Yes

Assessment Narrative:

Listed as Endangered in Vermont. Black Tern populations have remained low and fluctuated during the past decade with a high count of 157 breeding pairs in 2010 and a low of 65 pairs in 2007. Totals for 2014 were 95 breeding pairs but this was considered an underestimate (J. Sefchick-Edwards, pers. com.). These numbers represent an overall increase from estimates between 1990 and 2005 (44-103 pairs). Missisquoi National Wildlife Refuge remains the only nesting location for Black Tern in Vermont. They are, however, found in 6 separate locations within the 6000+ acre refuge.

Distribution

Black Terns are currently only found at one location, Missisquoi National Wildlife Refuge on northern Lake Champlain. Nesting occurred regularly in Lake Memphremagog South Bay throughout the 1990's.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Unknown
Champlain Hills	Unknown	Vermont Valley	Unknown
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Probable	Taconic Mtns	Unknown
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Nests in wetlands consisting of both emergent (cattails) and shrub (buttonbush) vegetation with adequate floating vegetation to build nests on.

Habitat Types:

Marshes and Sedge Meadows

Shrub Swamps

Current Threats

Habitat Threats:

Unknown Habitat Threats



Common Name: **Black Tern**
 Scientific Name: **Chlidonias niger**
 Species Group: **Bird**

Description of habitat threat(s): Sufficient habitat seems available both at its known nesting site (MNWR) and at sites it has nested at in the past. High water levels on Lake Champlain seem to influence nesting numbers at MNWR by limiting available nesting habitat.

Non-Habitat Threats:

Unknown Non-Habitat Threats

Description of non-habitat threat(s): Direct problems to this species have been difficult to determine, however Vermont is on the periphery of this species range and declines in the core of its range may be causing peripheral populations to decline at a faster rate.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Determine if changes in habitat structure at nesting locations has resulted in abandonment of sites used in the past.
Research	Basic Life History	Low	Well documented
Research	Distribution and Abundance	Medium	Annually monitor current and past nesting locations
Research	Threats and Their Significance	High	Increased understanding of limiting factors to this species in Vermont
Research	Population Genetics	Medium	Relationship to core population
Research	Taxonomy	Low	
Research	Other Research	Low	
Monitoring	Population Change	High	Annually monitor breeding population at current and past nesting locations.
Monitoring	Habitat Change	Medium	Monitor changes in habitat structure, especially those due to invasive species.
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	Medium	Annually monitor known limiting factors
Monitoring	Other Monitoring Needs	Medium	Continue to manage protected wetlands to provide suitable habitat. Determine appropriate management actions (e.g. vegetation management, artificial nesting structures) that will enhance breeding success.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Black Tern**
Scientific Name: **Chlidonias niger**
Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Species Restoration	High	Determine appropriate management strategies which will increase population size and the number of breeding locations.	Increase in population size and number of colony sites at different geographic locations.	USFWS, VFWD, Audubon-VT	SWG, PR

Bibliography

- Kibbe, D.P. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- Laughlin, S.B. and D. P. Kibbe, editors. 1985. The Atlas of Breeding Birds of Vermont. University Press of New England, Hanover, New Hampshire, USA.
- SefchickEdwards, J. 2014, Vermont Black Tern Population Data, 2007-2014. Unpubl. report, Missisquoi National Wildlife Refuge, Swanton, VT.
- Shambaugh, Nathaniel. 2003. 2003 Black Tern Population Survey and Other Marshbird Monitoring Activities in Vermont. Unpubl. report. VT Fish and Wildlife Department, Waterbury. Vermont

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Black-billed Cuckoo**
Scientific Name: **Coccyzus erythrophthalmus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Bird Conservation Regions 13 & 14 High Regional Concern. Not on Audubon/ABC's Watch Lists. BBS shows a significant long-term (1966-2013) annual decline for Vermont (-4.62 %). Shorter-term (2003-2013) decline is higher (-6.28%), but not judged significant.

Distribution

Distributed statewide, although less common in northeastern quarter of the state. The Second Atlas of Breeding Birds of Vermont (Renfrew 2013) results show biggest losses from southern Green Mountains.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Brushy pastures, shrubby hedgerows at edges of fields, dry, open woods and groves (DeGraaf and Rudis 1986). Prefers groves of trees, forest edges, and thickets; frequently associated with water. In NE. U.S.; usually found in edges and clearings of young deciduous-coniferous woods; abandoned farmland...brushy hillsides and pastures hawthorn thickets " (Hughes 2001).



Common Name: **Black-billed Cuckoo**
Scientific Name: **Coccyzus erythrophthalmus**
Species Group: **Bird**

Habitat Types:

Early Succession Boreal Hardwoods
Early Succession Pine and Hemlock
Early Succession Northern Hardwoods
Lawns, Gardens, and Row Crops
Shrub Swamps
Early Succession Spruce-Fir
Early Succession Upland Oak
Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Fragmentation

Description of habitat threat(s): "Declines in some regions could be due in part to reversion of abandoned farmland to forests that are unsuitable (Erskine 1992). Other problems responsible for declines could be modification of habitat on wintering grounds, hazards during migration, and pesticide use." (Hughes 2001).

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Black-billed cuckoos may be highly vulnerable to pesticides used on insect outbreaks, perhaps especially on winter range in South America.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	The nature of breeding range habitat changes relatively well understood, however the magnitude of these changes should be documented.
Research	Distribution and Abundance	Medium	Continued monitoring of changes in distribution and abundance should be tied to tracking changes in habitat (succession of abandoned farmland).
Monitoring	Population Change	Medium	Continued monitoring of changes in distribution and abundance should be tied to tracking changes in habitat (succession of abandoned farmland).
Monitoring	Habitat Change	Medium	The nature of breeding range habitat changes relatively well understood, however the magnitude of these changes should be documented.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Black-billed Cuckoo**
 Scientific Name: **Coccyzus erythrophthalmus**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Policy & Regulations	Medium	Better control use of pesticides harming this species and its food sources.	Number of regulations restricting use of harmful pesticides. Reduction in contaminants present in cuckoo habitat.	ANR, Vt. Dept of Agriculture, USDA	USDA, FDA
Conservation Payments/Financial Incentives	Medium	Conservation of hedgerows could be incorporated into EQIP program goals.	Number of sites designated for hedgerow conservation and protection from development.	ANR, USFS, USFWS	SWG, PR, EQIP
Habitat Restoration	High	Early- successional habitat (shrubland) goals should be developed for public and private land to support 4,200 individuals (Rosenberg 2004).	Total area managed for ESH.	ANR, USFS, USFWS	SWG, PR, EQIP

Bibliography

Degraaf, R.M. and D.D. Rudis 1986. New England wildlife: habitat, natural history, and distribution. Gen. Tech. Rep. NE-108. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station; 1986. 491 p.

Hughes, J. M. 2001. Black-billed cuckoo (*Coccyzus erythrophthalmus*). In *The Birds of North America*, No. 587 (A. Poole and F. Gill, eds.). The Birds of North America, Inc., Philadelphia, PA.

Kibbe, D.P. 2013. *The Second Atlas of Breeding Birds of Vermont*. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Rosenberg, K. V. 2004. Partners in Flight continental priorities and objectives defined at the state and bird conservation region levels, Vermont. Cornell Lab of Ornithology, Ithaca, NY. 26 p.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Short-eared Owl**
Scientific Name: **Asio flammeus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S1B,S2N

State Trend: Unknown

Extirpated in VT? Unknown

Regional SGCN? Yes

Assessment Narrative:

Unknown breeding status in VT - regular winter sightings in Champlain Valley.

BBS data indicate significant long-term overall decline, although trend unknown for many areas. Decline likely attributed to habitat conversion of marshes, grasslands, and low-use pastures.

Distribution

Distribution info from VT Breeding Bird Atlas (Laughlin & Kibbe 1985). Two confirmed nestings found in Champlain Valley. Other sightings in Northern Green Mts, and Southern & Northern VT Piedmont - all single sightings. Significant wintering concentrations have been seen in the Champlain Valley.

Distribution by Biophysical Region:

Champlain Valley	Probable	Southern VT Piedmont	Probable
Champlain Hills		Vermont Valley	Unknown
Northern Green Mtns	Probable	Southern Green Mtns	Unknown
Northern VT Piedmont	Probable	Taconic Mtns	Unknown
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

Breeding: Open land (marshlands & grasslands preferred), but will also use agricultural land and other open habitat.

Wintering: same as above with little/no snow cover (DeGraaf & Yamasaki 2001).

Habitat Types:

Marshes and Sedge Meadows

Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat



Common Name: **Short-eared Owl**
 Scientific Name: **Asio flammeus**
 Species Group: **Bird**

Habitat Succession

Habitat Alteration

No Habitat Threats

Description of habitat threat(s): Loss of marshes and grasslands since 1930s has caused declines in population (DeGraaf & Yamasaki 2001).

Non-Habitat Threats:

Trampling or Direct Impacts

Description of non-habitat threat(s): Farming practices may impact nesting owls.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	Determine if there is a breeding population in VT
Research	Threats and Their Significance	Medium	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	Medium	
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	Medium	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Short-eared Owl**
 Scientific Name: **Asio flammeus**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Conservation Payments/Financial Incentives	High	Protect suitable grassland habitat from development and agricultural intensification by creating Grassland Bird Focus Areas to concentrate management efforts (see Vermont Grassland Bird Management Plan)	Development of Grassland bird focus Areas and increased protection of habitat through enrollment in EQIP and GRP	VFWD, Audubon VT, NRCS	USDA
Protected Area Management	High	Maintain nesting habitat throughout breeding season by developing site specific conservation plans which include restricting mowing after July 15 on publicly owned lands (WMAs and state airports).	Maintain and increase current acreage under management on state lands	VFWD, NRCS, VTrans	VFWD
Habitat Restoration	High	Maintain grassland habitat in suitable locations through active management of woody vegetation within Grassland Bird Focus Areas.	Increase and maintain available habitat in suitable locations	VFWD, Audubon VT, NRCS, USFWS	US government
Technical Assistance, Training, Learning Networks	High	Educate agricultural community and general public about grassland birds and management options to protect habitat	Develop a grassland bird outreach program	VFWD, Audubon VT, VCE, UVM	SWG, VFWD

Bibliography

DeGraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover, NH. 482 pp.

Laughlin, S. B., and D. P. Kibbe. 1985. The atlas of breeding birds of Vermont. University Press of New England, Hanover, NH. 456 pp.



Common Name: **Common Nighthawk**
Scientific Name: **Chordeiles minor**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The Common Nighthawk was listed as an endangered species in Vermont in 2012. In Vermont's first breeding bird atlas (1985) there were three confirmations in priority blocks and the bird occupied 35 blocks. During the second Breeding Bird Atlas (2013) in Vermont the bird was not confirmed in any survey blocks, and only occupied nine survey blocks. The species is declining due to lack of breeding habitat. Additional evidence that non-selective pest control for mosquitoes has resulted in declining food resource availability (largely moths).

Distribution

Confirmed breeding records are few, and decreased between the first and second Atlas surveys (1985-2013). Two probable breeding records were from southern Orange County and Brattleboro. The bird has been recorded via ebirders throughout the state over the past 10 years, but those sightings include birds that are migrating.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Prefers nesting/breeding habitats including dunes, beaches, logged or clearcut areas, open forests, rock outcrops, gravel outwashes and gravel on flat roofed buildings. Utilizes virtually all open habitats, above water and open woodlands, including urban and suburban areas, during crepuscular feeding and migration.



Common Name: **Common Nighthawk**
Scientific Name: **Chordeiles minor**
Species Group: **Bird**

Habitat Types:

Upland Shores
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Hardwood Swamps
Softwood Swamps
Northern Hardwood
Floodplain Forests
Marshes and Sedge Meadows
Wet Shores
Early Succession Boreal Conifers
Early Succession Boreal Hardwoods
Early Succession Pine and Hemlock
Early Succession Northern Hardwoods
Building or Structure
Shrub Swamps
Early Succession Spruce-Fir
Early Succession Upland Oak
Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Inadequate Disturbance Regime
Habitat Fragmentation

Description of habitat threat(s): Losses of upland openings, reductions of clearcut timber harvest, conversion of natural openings to non-suitable habitat (residential, etc.), conversion of flat, gravel-covered roofs to metal/rubberized coating/sheeting.

Non-Habitat Threats:

Predation or Herbivory
Loss of Prey Base

Description of non-habitat threat(s): Reductions of preferred prey due to non-selective pesticide use.
Loss of suitable wintering ground habitats due to changes in agricultural practices.



Common Name: **Common Nighthawk**
Scientific Name: **Chordeiles minor**
Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Factor accounting for decreased nesting on available gravel roofs and effects of deforestation; nest/roost site characteristics. Continue study on effectiveness of creating artificial gravel patches on rooftops
Research	Basic Life History	Medium	Longevity of breeders, reproductive output; male fidelity
Research	Distribution and Abundance	Low	Continued monitoring to determine if population is increasing or decreasing. Support for gathering data from citizen scientists, including Vermont ebird.
Research	Threats and Their Significance	High	Population status as related to pesticide use on breeding grounds, wintering grounds, and migration routes.
Monitoring	Population Change	High	Sustained monitoring is crucial to maintaining the existing population
Monitoring	Habitat Change	High	Changes of suitable nesting habitats, and use of rafts.
Monitoring	Monitor Threats	Medium	Monitoring of all limiting factors goes hand-in-hand with population monitoring and is critical to evaluate long-term viability and management needs of statewide population. death, including interference competition from others.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Common Nighthawk**
 Scientific Name: **Chordeiles minor**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Conduct complete vegetation removal on select areas of existing upland openings, and provide for even-age timber management on public lands to increase suitable common nighthawk nesting habitat.	Number of acres positively affected by management. Population response to management.	FWD, FPR, Audubon, Forest Products Association, VT Loggers	PR, EQIP
Habitat Restoration	High	Identify specific threats, and agricultural practices that negatively impact birds		FWD	PR
Awareness Raising and Communications	Medium	Initiate public education campaigns to highlight the need for active, even-age forest management and the need for mineral soil outcrops (gravel, ledge) on public and private lands to create suitable habitat complexes.	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs.	ANR, USFS, USFWS	SWG, PR
Conservation Finance	Medium	Create a state-funded, private lands, early successional habitat improvement initiative. Fund for > \$50,000/yr with revenues from state lands forest management. Allow the installation of gravel pads on flat roofed buildings as a conservation practice.	Level of funds raised.	FWD	PR
Habitat Restoration	Medium	Identify and enlarge/enhance suitable lakeshore and riparian gravel depositions to create additional nesting habitat.	Number of sites identified and positively affected by management. Population response to management.	VA, VCE	SWG

Bibliography

Brigham, R. M., Janet Ng, R. G. Poulin and S. D. Grindal. 2011. Common Nighthawk (Chordeiles minor), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/213>

Hunt, P. 2013. Common Nighthawk (Chordeiles minor). In: The Second Atlas of Breeding Birds of Vermont, (R. Renfrew, ed.). University Press of New England, Hanover and London

eBird. 2012. eBird: An online database of bird distribution and abundance [web application]. eBird, Cornell Lab of Ornithology, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: Date February 4, 2015).



Common Name: **Whip-poor-will**
 Scientific Name: **Caprimulgus vociferus**
 Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Whip-poor-will has been declining throughout much of its range, particularly in the northeastern US and eastern Canada, based on data from breeding bird atlases. In Vermont this species was found in 77% fewer blocks between the first (1979-85) and second (2003-2007) breeding bird atlases (Renfrew 2013). Based on expert knowledge, atlas data, and standardized whip-poor-will survey routes conducted since 2005, the statewide population was estimated at <200 breeding pairs in 2012 by the Bird-Scientific Advisory Group of the Agency of Natural Resources. Based on this estimate and declining trends, the species was listed as threatened in 2013. There are ongoing efforts to determine a more precise estimate of the state's population, and standardized monitoring continues. Causes of declines unknown, possibly succession of farmland, prey, and/or invasive plant and other understory growth in otherwise suitable forests.

Distribution

Statewide, but primarily in Champlain and Connecticut River valleys, rare and disjunct in northern and southern thirds of the state. Highest densities appear to be in West Haven area.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

In Vermont, dry open deciduous or open mixed lowland forest with little or no underbrush or ground cover, near or adjacent to open areas for foraging (Cink 2002), such as pastures. Occupied territories often consist of areas where forested mountainsides abut lowland open areas such as pastures, e.g. in the lower Champlain Valley (VFWD/VCE, unpubl. data). Open areas also may be forest clearings, meadows, water, or gravel pits. May occur along power line rights-of-way and pine-oak forests along major rivers.



Common Name: **Whip-poor-will**
Scientific Name: **Caprimulgus vociferus**
Species Group: **Bird**

Habitat Types:

Upland Shores
Outcrops and Alpine
Cliffs and Talus
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Hardwood Swamps
Seeps and Pools
Northern Hardwood
Floodplain Forests
Softwood Swamps
Open Peatlands
Marshes and Sedge Meadows
Wet Shores
Early Succession Boreal Hardwoods
Early Succession Pine and Hemlock
Early Succession Northern Hardwoods
Shrub Swamps
Early Succession Upland Oak
Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development
Habitat Succession
Habitat Alteration
Habitat Fragmentation
Invasion by Exotic Species
Inadequate Disturbance Regime
Impacts of Roads or Transportation Systems

Description of habitat threat(s): Habitat alteration due to forestry practices or conversion of grassland to crops. Habitat conversion due to rural/suburban development of fields and forest. Habitat succession via closure of forest openings and regrowth of understory. Roads present potential threat of direct impact on whip-poor-wills that forage along roads at night, particularly secondary and tertiary roads. Solar infrastructure can result in loss of open habitat, wind turbines present strike hazard. Late successional forest stands with little undergrowth needed for nesting. Exotic plants such as Buckthorn and Mountain Holly



Common Name: **Whip-poor-will**
Scientific Name: **Caprimulgus vociferus**
Species Group: **Bird**

compromise nesting habitat. .

Non-Habitat Threats:

Predation or Herbivory

Loss of Prey Base

Description of non-habitat threat(s): Pesticide/bio control for agricultural pests may have reduced prey base. Ground nests vulnerable to predators.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Habitat preference, ultimately to guide development of management recommendations. Seemingly suitable habitat is unoccupied, although this could be due to limitations other than breeding habitat.
Research	Basic Life History	High	Many unknowns in this species that are difficult to study, including: demography, including seasonal and year-round survival, site fidelity, inter- and intra-seasonal movements, and migratory connectivity.
Research	Threats and Their Significance	High	Reasons for population declines still not well understood. Whether food is limited still unknown. Vulnerability to wind turbines, especially during migration, not known for Vermont. Survival during non-breeding season, including post-fledgling survival, could be a limiting factor. Potential effects of climate change not known.
Monitoring	Population Change	High	Standardized survey to detect regional population changes inadequate for detecting population trends at state level. Need more intensive, standardized survey method to better monitor Vermont's population, to determine local detection probabilities, and to derive a more accurate population estimate.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Whip-poor-will**
 Scientific Name: **Caprimulgus vociferus**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Investigate and implement, where possible, understory prescribed burning to create open-forest habitat w/ little underbrush suitable for whip-poor-will nesting.	Number of sites identified and positively affected by management. Population response to management.		
Habitat Restoration	Medium	Determine appropriate old field habitat targets for state lands and restore and maintain old field habitats where needed to increase suitable ES songbird habitat.	Number of acres positively affected by management. Population response to management.	ANR, USFS, Audubon, Forest Products Association, VT Loggers	PR, EQUIP
Habitat Restoration	High	Stabilize or reverse declining population trend for whip-poor-wills to realize and maintain a population of >400 breeding pairs.	Standardized survey routes, occasional statewide counts with public participation	VFWD, VCE, Audubon	PR, EQUIP
Conservation Finance	Medium	Investigate and implement, where possible and where suitable habitat mosaic exists, understory prescribed burning to create open-forest habitat w/ little underbrush suitable for whip-poor-will nesting.	Number of acres identified and positively affected by management. Population response to management.		
Awareness Raising and Communications	Medium	Initiate public education campaigns on the need for active, even-age forest management on public and private lands and to eliminate undergrowth in prime potential forest habitat adjoining open habitat, especially removal of invasive plants (e.g. buckthorn)	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs, number of acres treated		

Bibliography

Cink, C.L. 2002. Whip-poor-will (*Caprimulgus vociferus*). In A. Poole and F. Gill, editors, *The Birds of North America*, No. 620. Academy of Natural Sciences, Philadelphia, Pa., and American Ornithologists' Union, Washington, DC. 20 pp.

Hunt, P. 2013. *The Second Atlas of Breeding Birds of Vermont*. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Rosenberg, K. V. 2004. Partners in Flight continental priorities and objectives defined at the state and bird conservation region levels, Vermont. Cornell Lab of Ornithology, Ithaca, NY. 26 p.



Common Name: **Chimney Swift**
Scientific Name: **Chaetura pelagica**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S4B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Vermont's second Breeding Bird Atlas found a 26% loss in the number of occupied blocks. These losses were primarily from the Champlain Valley, Northern Green Mountains, and Northern Vermont Piedmont. Declines for this species throughout its range have been shown with BBS surveys and Breeding Bird Atlas surveys. Once nested in old growth forests with scattered large hollow trees for roosting. Since settlement and industrialization, the species has shifted its habitat to mostly urban areas with large chimneys used for roosting and nesting, and agricultural areas where barns and silos provide similar roosting and nesting habitat. Reductions in sizes of chimneys due to change from coal burning to oil and electricity use have now further reduced this species population. Threats on migration pathways and wintering grounds are also a factor.

Distribution

Chimney Swifts breed in almost every region of Vermont; although the second Breeding Bird Atlas found a 26% loss in the number of occupied blocks. They travel miles to feed and are documented through ebird throughout Vermont. Chimney Swifts winter in the Amazon Basin.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Chimney Swifts formerly nested and roosted in caves and large dead trees. As European settlement and development increased, the birds nested mainly in large, tall chimneys in urban and suburban areas. The conversion from heating with coal to oil and electricity in residential homes and industrial plants has reduced the size of chimneys making them less suitable as nest sites for chimney swifts. Some experts believe the birds may still nest in large snags in rural areas. Chimney Swifts forage in a variety of habitats, but seem to prefer open areas over densely forested habitats.



Common Name: **Chimney Swift**
Scientific Name: **Chaetura pelagica**
Species Group: **Bird**

Habitat Types:

Building or Structure
Lawns, Gardens, and Row Crops
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat

Description of habitat threat(s): Originally, loss of old growth habitat with large trees used for roosting and nesting. After habitat shift, loss of large chimneys in urban areas.

Non-Habitat Threats:

Unknown Non-Habitat Threats

Description of non-habitat threat(s): Threats across their range include pesticides, collisions, catastrophic weather events, threats to wintering grounds

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Snag requirements for nesting. Determine if artificial nest structures can reverse declining population.
Research	Distribution and Abundance	Medium	Support of for gathering data from citizen scientists important, including Vermont ebird
Monitoring	Population Change	Medium	Continued monitoring needed to assess if declining trend in population is significant and warrants listing. Use of radio telemetry to determine specific habitat requirements.
Monitoring	Habitat Change	Medium	Assess forest succession conditions throughout state and determine if chimney swifts are re-occupying original forest habitats. Nest and roost trees identified and preserved.
Monitoring	Monitor Threats	Medium	Monitoring of all limiting factors goes hand-in-hand with population monitoring and is critical to evaluate long-term viability and management needs of statewide population.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Chimney Swift**
 Scientific Name: **Chaetura pelagica**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Restoration of old growth habitats should be monitored to determine if chimney swifts are re-occupying these areas.	Presence/absence of chimney swifts on breeding bird surveys, bird atlas; location of roost or nest trees.	USFS,USFWS,VFPR, private landowners.	PR, Nongame Fund, SWG
Habitat Restoration	High	Implement artificial nest structure program, especially on existing and new commercial buildings, to restore nesting opportunities for chimney swifts.	Number of artificial nest structures erected and utilized.	VFWD, VA, VCE, TWF, VLCT, VT Home Builders Association, Cons Comms	SWG

Bibliography

The Atlas of Breeding Birds of Vermont. 1985. S.B. Laughlin and D.P. Kibbe, eds. University Press of New England. 456 p.

Kibbe, D.P. 2013. Chimney Swift (*Chaetura pelagica*). In: The Second Atlas of Breeding Birds of Vermont, (R. Renfrew, ed.). University Press of New England, Hanover and London

eBird. 2012. eBird: An online database of bird distribution and abundance [web application]. eBird, Cornell Lab of Ornithology, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: Date February 4, 2015).

Steeves, Tanner K., Shannon B. Kearney-McGee, Margaret A. Rubega, Calvin L. Cink and Charles T. Collins. 2014. Chimney Swift (*Chaetura pelagica*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/646>



Common Name: **Black-backed Woodpecker**
 Scientific Name: **Picoides arcticus**
 Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S2B,S2N

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

Uncommon resident in northeastern Vermont. Thirty successful nesting pairs documented by Weinhausen (1998) in northeastern Vermont in 1996-97. Black-backed woodpeckers in Vermont are habitat specialists that depend on mature/over mature spruce-fir- tamarack forests for nesting and foraging. Trends in amount of required habitat are unknown. Conversion of substantial amount of potential habitat from intensively managed industrial forest land to public or private land w/easements over the past decade may lead to increased and/or stable amount of preferred habitats. Habitat losses may be caused by seasonal and/or permanent residential development, especially at shoreline sites, which may experience increase frequency in the coming decade in northeastern Vermont.

Distribution

Most nesting records concentrated in Northeast Highlands. Thirty successful nests were documented by Weinhausen (1998) in 1996-97. The Second Atlas of Breeding Birds of Vermont (Renfrew 2013) reports that 35% of the Northeast Highlands biophysical region was occupied by Black-backs and that 93% of all records of this species were found here as well.

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Probable	Southern Green Mtns	Probable
Northern VT Piedmont	Confident	Taconic Mtns	Not Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

The Black-backed woodpecker inhabits mature and decadent coniferous forests where it utilizes these older stands for both nesting and foraging habitat. Lowland spruce-fir forests and bogs and swamps supporting tamarack, red and black spruce, and balsam fir typify the natural community preferred by black-backed woodpeckers. Black-backed woodpeckers feed primarily on wood-boring beetle (cerambycids) larvae (Dixon and Saab 2000). In Vermont, Weinhausen (1998) found a mean stand diameter of 11.5 to 26.7 cm within a 900 meter radius of 62% of nest trees. Average dbh of nest trees was 27 cm (range 19-55). Natural mortality due to senescence or reduced tree vigor brought on by fire, windstorm, flooding, spruce budworm outbreak or residual stand damage from logging can all lead to cerambycid infestations followed by nesting black-backs (Dixon and Saab 2000).



Common Name: **Black-backed Woodpecker**
Scientific Name: **Picoides arcticus**
Species Group: **Bird**

Habitat Types:

Spruce Fir Northern Hardwood
Softwood Swamps
Open Peatlands

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Habitat Fragmentation

Description of habitat threat(s): Black-backed woodpeckers in Vermont are habitat specialists that depend on mature/over-mature spruce-fir-tamarack forests for nesting and foraging. Trends in amount of required habitat are unknown. Conversion of substantial amount of potential habitat from intensively managed industrial forest land to public or private land w/easements over the past decade may lead to increased and/or stable amount of preferred habitats. Habitat losses may be caused by seasonal and/or permanent residential development, however, especially at shoreline sites. Some predictions are that lakeshore development may experience increased frequency in the coming decade in northeastern Vermont.

Description of non-habitat threat(s):

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Minimum and optimum patch size and degree of home range overlap should be determined. Effects of selective or shelterwood silvicultural practices on nesting success should be investigated
Research	Distribution and Abundance	Medium	Periodic surveys of known woodpecker occupancy throughout the Northeast Highland biophysical region to determine extent of distribution and abundance.
Research	Threats and Their Significance	Medium	Known nest sites should be monitored periodically for limiting factors.
Monitoring	Population Change	Medium	Periodically monitor occupancy of known & potential nesting habitats
Monitoring	Habitat Change	Medium	Known nest sites should be monitored periodically for limiting factors.



Common Name: **Black-backed Woodpecker**
 Scientific Name: **Picoides arcticus**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Increase rotation age in some managed forests.	Number of sites and total area with increased rotation ages.	ANR	SWG, PR
Publically-Owned Protected Areas	Medium	Identify potential areas on public lands for designation as reserve or no-cut status (including some areas where wildfires and insect outbreaks would not be controlled).	Number of sites and total area of designated reserves.	ANR, UVM, TNC	SWG, PR
Protected Area Management	High	Area of suitable breeding habitat (mature spruce-fir patches > 100 ha) should be mapped. A subset of this data that is in reserve or no-cut status should be determined. Use research findings to inform whether additional acreage is necessary.	Number of sites inventoried. Number of these sites occupied by breeding black-backed woodpeckers.	ANR, UVM	SWG, PR

Bibliography

- Alexander, C. 2013. Second Atlas of Breeding Birds of Vermont. Rosalind Renfrew, ed. 547 pp. University Press of New England.
- Cadman, M.D., D.A. Southerland, G.G. Beck, D. Lepage, and A.R. Couturier. 2007. Atlas of the Breeding Birds of Ontario., 2001-2005. Bird Studies Canada, Toronto, ON.
- Dixon, R.D. and V.A. Saab. 2000. Black-backed woodpecker (*Picoides arcticus*). In The birds of North America, No. 509. (A. Poole and F. Gill, eds). The Birds of North America. Inc. Philadelphia, PA.
- McGowan, K.J., and K. Corwin, editors. 2008. The Second Atlas of Breeding Birds in New York State. Cornell University Press, Ithaca, NY.
- Weinhagen, A. C. 1998. Nest-site selection by the Black-backed Woodpecker in northeastern Vermont. Master's thesis, Univ. of Vermont, Burlington.

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Olive-sided Flycatcher**
Scientific Name: **Contopus cooperi**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: S4B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Strong NS negative trends on VT BBS routes (Sauer et al. 2004).

Population declines throughout North America (Sauer et al. 2004). VT populations widespread but local (Fichtel 1985).

Distribution

Based on Fichtel (1985).

Distribution by Biophysical Region:

Champlain Valley	Probable	Southern VT Piedmont	Confident
Champlain Hills		Vermont Valley	Confident
Northern Green Mtns	Probable	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Montane and northern coniferous forests. Frequently associated with forest openings, forest edges near natural openings, or open to semi-open forest stands (Altman and Sallabanks 2000)].

Habitat Types:

Spruce Fir Northern Hardwood
Other Cultural

Current Threats

Habitat Threats:

Conversion of Habitat
Inadequate Disturbance Regime

Description of habitat threat(s): Species seems to require disturbances in coniferous forests. Beaver ponds, burns, clearcuts, or wind throws all appear to be appropriate. The species has a large territory size (10-20 ha), therefore there seems to be wide spacing between territories. Other problems are conversion to



Common Name: Olive-sided Flycatcher
Scientific Name: *Contopus cooperi*
Species Group: Bird

nonforest habitat, loss of wintering habitat, and decrease in prey species (summarized from Altman and Sallabanks 2000).

Description of non-habitat threat(s):

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Range-wide, a better understanding of habitat associations is a research priority.
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	Habitat-specific demographic data would be useful to understanding population ecology.
Research	Threats and Their Significance	High	The relative importance of breeding versus wintering habitat loss and degradation would help target effective conservation strategies.
Monitoring	Population Change	High	Distribution in Vermont and areas of high abundance to target habitat management activities.
Monitoring	Habitat Change	High	Determine effects of disturbance regimes on habitat quality.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Create more early successional habitat, particularly in the northeast kingdom of VT.	Number of acres positively affected by management. Population response to management.	ANR, USFWS	PR, SWG

Bibliography

- Altman, B., and R. Sallabanks. 2000. Olive-sided Flycatcher (*Contopus cooperi*). In *The birds of North America*, No. 502 (A. Poole and F. Gill, eds.). The birds of North America, Inc., Philadelphia, Pennsylvania, USA.
- DeGraaf, R. M., and M. Yamasaki. 2001. *New England wildlife: habitat, natural history, and distribution*. University of New England Press, Hanover, New Hampshire, USA, and London, UK.
- Fichtel, C. 1985. Olive-sided Flycatcher. Pages 170-171 in S. B. Laughlin and D. P. Kibbe, eds. *The atlas of breeding birds of Vermont*. University Presses of New England, Hanover, New Hampshire, USA.
- Sauer, J. R., J. E. Hines, and J. Fallon. 2004. *The North American Breeding Bird Survey, Results and Analysis 1966-2003*. Version 2004.1. USGS Patuxent Wildlife Research Center, Laurel, Maryland, USA.



Common Name: **Purple Martin**
Scientific Name: **Progne subis**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S3S4B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

Populations have shown a nation-wide decline, especially since 1980. The Purple Martin once lived along riparian, lake shoreline, and swamp edges. It is now almost completely found inhabiting areas close to human settlement. It originally nested in tree cavities but almost all nesting now occurs in bird houses erected by humans, except in western U.S. where some natural nesting still occurs.

Distribution

Breeding in Vermont is almost exclusively the Champlain Valley, although the bird is recorded from other places within the state. In 2012 the Bird Scientific Advisory Group and many volunteers surveyed the state to try to record all Purple Martins breeding. This is possible with this species, as they are completely dependent upon human-provided nest sites. At that time 421 nesting pairs were counted and documented on ebird. Purple Martins migrate outside of the US to central south America. Recent research suggests that some fly across the Gulf of Mexico.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Historically found along riparian areas, lake and pond shorelines, and edges of forest openings. Currently found almost exclusively around human settlements. All confirmed breeding records in the second Breeding Bird Atlas were in the Champlain Valley, with the majority of records less than 10 kilometers from the Lake. Colonies were clustered in Grand Isle County and Addison County. In Vermont a survey was completed in 2012 to find all Purple Martin nests. Birds will nest as one pair or in larger colonies. The 2012 survey ranged from 1 occupied compartment to 53 occupied compartments. Some limitations may be due to competition with non-native species such as the House Sparrow and the European Starling.



Common Name: **Purple Martin**
Scientific Name: **Progne subis**
Species Group: **Bird**

Habitat Types:

Wet Shores
Building or Structure
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops
Aquatic: Fluvial
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Habitat Alteration
Climate Change
Invasion by Exotic Species
No Habitat Threats

Description of habitat threat(s): Change in rainfall resulting wet springs impact food supplies for adults returning from wintering grounds and adults feeding young. Competes for nest sites with introduced House Sparrow and European starling. Habitat requires maintenance of nest boxes by humans. Inadequate number of nest boxes erected to attract martins and support population viability.

Non-Habitat Threats:

Competition

Description of non-habitat threat(s): Competes with introduced house sparrow and European starling for nest sites. Inadequate number of nest boxes erected to attract martins and support population viability. Nest boxes may not be maintained adequately by owners.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Distribution and Abundance	Medium	Support for gathering data from citizen scientists important, including Vermont ebird. Support education to ensure new martin houses are built and maintained.
Research	Threats and Their Significance	Medium	Maintain existing boxes in a manner to attract martins, reduce exotic species, and prevent disease.
Monitoring	Population Change	Medium	Continued monitoring needed to assess if declining trend in population is significant and warrants listing. Use of radio telemetry to determine specific habitat requirements.
Monitoring	Habitat Change	Low	Support citizens if needed, to ensure martin houses are constructed and maintained.
Monitoring	Monitor Threats	Medium	Monitoring of all limiting factors goes hand-in-hand with population monitoring and is critical to evaluate long-term viability and management needs of statewide population.

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Purple Martin**
Scientific Name: **Progne subis**
Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Species Restoration	High	Establish artificial nest box program on public and private lands.	Presence/absence of breeding martins, number of young fledged, number of nests abandoned due to house sparrows and starlings.	Audubon-VT, VCE, BOVM, Cornell Lab of Ornithology, USFWS, NRCS	Nongame Fund, SWG, Neotropical Bird Conservation Act grants, National Fish and Wildlife Foundation grants, USFWS, NRCS
Awareness Raising and Communications	Medium	Educate public about nest box program, nest box maintenance, and Purple Martin Society	Number of cooperating homeowners who erect martin boxes	VFWD, VA, VCE	SWG, PR

Bibliography

Hunt, P., D.P. 2013. Purple Martin (*Progne subis*). In: The Second Atlas of Breeding Birds of Vermont, (R. Renfrew, ed.). University Press of New England, Hanover and London

eBird. 2012. eBird: An online database of bird distribution and abundance [web application]. eBird, Cornell Lab of Ornithology, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: Date February 4, 2015).

National Audubon Society. 2014. Audubon's Birds and Climate Change Report: A Primer for Practitioners. National Audubon Society, New York. Contributors: Gary Langham, Justin Schuetz, Candan Soykan, Chad Wilsey, Tom Auer, Geoff LeBaron, Connie Sanchez, Trish Distler. Version 1.2.

Strong, A. and E. Talmage, 2012, (personal communication).

Tarof, Scott and Charles R. Brown. 2013. Purple Martin (*Progne subis*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/287>



Common Name: **Gray Jay**
Scientific Name: **Perisoreus canadensis**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1S2B,S1S2N

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

Partners In Flight (PIF) reports a continental population estimate of 20,000 birds based on Breeding Bird Survey (BBS) estimates. The Canadian estimate of 17,000 individuals is categorized as of medium quality while the American estimate of 3.4 million (sic) birds is categorized as high quality. Breeding Bird Survey (BBS) data for Vermont is nonexistent but survey data for Quebec report increases in population size (<1.5% annual increases since 1966). However, the same indices suggest a greater than 1.5% annual decrease for grey jay in the Maritime Provinces and northern New England states. The Second Breeding Bird Atlas (2003-07) reports a 14% decrease in breeding blocks (7 blocks to 6) since the first atlas (1982). Increased softwood harvest in northeastern Vermont between 1978 and 1984 heightened concerns for the Gray Jay's continued existence. The Scientific Advisory Group on Birds (SAG) proposed it be listed as threatened, but the proposal was rejected. Public lands and conservation easements acquired over the last 2 decades in prime gray jay range, along with higher than expected densities found by Barnard in his ongoing field studies (begun in 1991) have reduced this concern. However, concerns for the species are again increased due to projected effects of long-term climate change influences potentially decreasing the abundances and distribution of the already limited boreal habitat (southern edge of range). For this reason SAG recommends this species be reclassified from medium to a high priority assessment.

Distribution

Distribution limited to largest boreal forest patches of northeastern Vermont. Strongholds include the Victory, Nulhegan and Coaticook River Basins. Also occurs at high elevation coniferous forest.

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Coniferous forests and nearby deciduous or mixed woodlands.



Common Name: **Gray Jay**
Scientific Name: **Perisoreus canadensis**
Species Group: **Bird**

Habitat Types:

Spruce Fir Northern Hardwood
Softwood Swamps
Early Succession Boreal Conifers
Early Succession Spruce-Fir

Current Threats

Habitat Threats:

Habitat Alteration

Habitat Fragmentation

Description of habitat threat(s): Gray jays utilize all seral stages of coniferous forests, readily occupy managed landscapes, and readily visit bird feeders or take handouts directly from humans. Conversion of coniferous forests in Vermont strongholds unlikely, however, smaller occupied patches (if they exist) could be degraded by residential or commercial development.

Non-Habitat Threats:

Harvest or Collection

Description of non-habitat threat(s): Accidental capture by trappers has been reported in Ontario. Susceptibility to land trap losses depends on types of baits used (Strickland and Ouellet 1993).

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	1) Investigate minimum patch size required for successful breeding and map all potential breeding habitat. 2) Determine total minimum area of suitable coniferous forest patches necessary to support 500 breeding pairs of gray jays in northeastern Vermont.
Monitoring	Population Change	Medium	Monitor population trends via surveys in targeted habitats and track the number of sites inventoried for breeding gray jays.
Monitoring	Habitat Change	Medium	

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks	Low	Investigate the occurrence of gray jays accidentally caught in furbearer traps in Vermont and, if needed, educate trappers on baiting techniques to minimize losses.	Number of accidental losses.	ANR, VT Trappers Association	SWG, PR



Common Name: **Gray Jay**
Scientific Name: ***Perisoreus canadensis***
Species Group: **Bird**

Bibliography

Barnard, W. 2013. Second Atlas of Breeding Birds of Vermont. 2013. Rosalind Renfrew, ed. 547 pp. University Press of New England

Stirckland, D. and H. Ouellet. 1993. Gray Jay. In The Birds of North America, No. 40 (A. Poole, P. Stettenheim, and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union.



Common Name: **Sedge Wren**
Scientific Name: **Cistothorus platensis**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1B

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Rangewide BBS population trends have been stable long-term and in the last ten years, although northeast populations have shown non-significant declines in recent years (Sauer et al. 2014), and the species' persistence as a breeding species in the Northeast is considered to be in jeopardy. In nearly all the states where it occurs, it is state-listed as endangered or threatened. Populations have evidently undergone dramatic declines in the latter part of 20th century after a northern expansion in range which was probably due to the clearing of forests in the 1800s. Land conversion in recent decades from old fields and pasturelands to forests or development has reduced nesting habitat. However, it appears that sedge wren populations today remain well below the level that available habitats could support, and the species is rare and local in Vermont (Renfrew 2013). Breeding pairs that are present in one year are not necessarily present in subsequent years (A. Strong, pers. obs.).

Distribution

Occasionally appear in June somewhere in Champlain Valley, location not consistent. Records during the second Breeding Bird Atlas were limited to the southern half of the Champlain Valley (Renfrew 2013).

Distribution by Biophysical Region:

Champlain Valley	Probable	Southern VT Piedmont	Unknown
Champlain Hills	Unknown	Vermont Valley	Historic Records Only
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Unknown
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Dense, tall growths of sedges and grasses in wet meadows, hayfields, retired croplands, upland margins of ponds and marshes, coastal marshes, and sphagnum bogs. Avoids short, sparse, or open vegetative cover, flooded areas, and wetlands dominated by cattails (Herkert et al. 2001). In Vermont, Sedge Wrens have been observed in June on older hayfields (both well-drained and wet), in a ditch, and on ungrazed pasture (Perlut 2013).



Common Name: **Sedge Wren**
Scientific Name: **Cistothorus platensis**
Species Group: **Bird**

Habitat Types:

Marshes and Sedge Meadows

Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Succession

Habitat Alteration

Description of habitat threat(s): Early haying of grasslands perhaps not as critical as SEWR in VT may be second nesting attempts of birds breeding further north earlier in summer. Consequently, drainage ditches in wet hayfields and meadows may reduce habitat availability. Succession of grassland habitats and conversion of agricultural habitats to urban/suburban developments problematic. Habitat conversion is likely not as severe a problem as for other grassland species, as SEWR requires wet meadow habitat and at least to some degree protected through regulatory measures.

Description of non-habitat threat(s):

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Determine habitat requirements, including preferred soil moisture regimes, vegetation height/density/composition, specific nesting cover requirements, minimum effective habitat area.
Research	Basic Life History	High	Determine where VT birds are coming from.
Research	Other Research	High	Habitat Requirements: Define relationships between habitat use, invertebrate prey abundance and soil moisture, rainfall, wetland proximity, grassland type. There is potential to evaluate through landsat imagery, although substantial ground-truthing would be necessary.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Sedge Wren**
 Scientific Name: **Cistothorus platensis**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	High				
Technical Assistance, Training, Learning Networks	High	Educate agricultural community and general public about grassland birds and management options to protect habitat	Continue grassland bird outreach and landowner networking programs	VFWD, Audubon-VT, VCE, UVM	SWG, PR
Protected Area Management	High	Maintain nesting habitat throughout breeding season by developing site specific conservation plans which include restricting mowing after July 15 on publicly owned lands (WMAs and state airports).	Maintain and increase current acreage under management on state lands	VFWD, NRCS, VTrans	SWG, PR
Compatible Resource Use	Medium	Maintain consistency of timing of cutting for potential habitat, as early season mowing of potential habitat is a problem in dry springs.		UVM, NRCS	NRCS (EQIP, GRP), USDA.
Habitat Restoration	Medium	Late season mowing of potential habitat would reduce problem of succession.	Long-term maintenance of wet meadow habitat.	UVM, NRCS.	NRCS (EQIP, GRP), USDA.

Bibliography

Heckert, J. R., D. E. Kroodsma, and J. P. Gibbs. 2001. Sedge Wren (*Cistothorus platensis*). In *The birds of North America*, No. 582 (A. Poole and F. Gill, eds.). The birds of North America, Inc., Philadelphia, Pennsylvania, USA.

LaBarr, M. L., Strong, A., Renfrew, R., Buck, J. and S. Parren. 2013. The Vermont grassland bird management and recovery plan. Vermont Fish and Wildlife Department. Montpelier, VT.

Perlut, N. P. 2013. Sedge Sren (*Cistothaurus platensis*). Pp. 312-313 In Renfrew, R.B., ed. *The second atlas of breeding birds of Vermont*. University Press of New England, Hanover, WI.

Sauer, J. R., J. E. Hines, and J. Fallon. 2014. *The North American Breeding Bird Survey, Results and Analysis 1966-2003*. Version 2004.1. USGS Patuxent Wildlife Research Center, Laurel, Maryland, USA.

Kibbe, D.P. 2013. *The Second Atlas of Breeding Birds of Vermont*. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.



Common Name: **Bicknell's Thrush**
Scientific Name: **Catharus bicknelli**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: S3B

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Species extensively studied by VINS since 1992, but no baseline data before that, so population trends still poorly known. Recent data from Mountain Birdwatch project indicate that species experienced an annual regionwide decline of 9.1% from 2001-2004. A habitat specialist whose overall population trends and abundance are not well known. Conservation limiting factors on both breeding and wintering grounds, combined with rarity and occupancy of naturally fragmented habitats, place the species at conservation risk. A high priority for attention in VT.

Distribution

Distributed throughout high elevation montane forests of VT

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Montane forests dominated by balsam fir with lesser amounts of red spruce, heart-leaved white birch, and mountain ash. Often associated with recently disturbed areas undergoing vigorous succession, characterized by standing dead conifers and dense regrowth of balsam fir. Highest densities typically found in chronically disturbed stands of dense, stunted fir on exposed ridgelines or along human-created openings (e.g. ski trails) or in regenerating fir waves.

Habitat Types:

Spruce Fir Northern Hardwood
Early Succession Boreal Conifers
Early Succession Spruce-Fir



Common Name: **Bicknell's Thrush**
Scientific Name: **Catharus bicknelli**
Species Group: **Bird**

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Habitat Fragmentation
Impacts of Roads or Transportation Systems
Climate Change

Description of habitat threat(s): The primary problems in Vermont and the Northeast (there are other, likely more significant problems on the species' Caribbean wintering grounds) are degradation and fragmentation of montane forests. Atmospheric pollution may be affecting forest health, and climate change could profoundly impact long-term viability of montane balsam fir forests. Immediate problems include loss and fragmentation of habitat from ski area development, communications tower development, and wind turbine development.

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Mercury contamination may be a problem, as research has shown significantly higher atmospheric deposition rates in montane forests than in surrounding low elevation habitats. Recent research has indicated that Hg levels in adult BITH increase with age.



Common Name: **Bicknell's Thrush**
Scientific Name: **Catharus bicknelli**
Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	Habitat requirements are reasonably well known, though responses to human-created developments need study.
Research	Basic Life History	Medium	Much basic life history is known, but some aspects of ecology and demography need further study.
Research	Distribution and Abundance	Medium	Distribution is well-documented, but better information is needed on abundance and population ecology in different subhabitat types.
Research	Threats and Their Significance	Medium	1) Expand studies of mercury contamination in BITH and synergistic effects with other environmental stressors (such as calcium depletion); evaluate effects on reproductive success, behavior, survivorship. Design and implement expanded research program on multiple peaks (at least Stratton, Mansfield, and East Mt) 2) It is not known how developments affect local breeding populations.
Research	Population Genetics	High	Knowledge of natal dispersal and migratory connectivity could help elucidate population structure and guide conservation planning.
Research	Taxonomy	Low	Taxonomy, while still debatable to some, has been resolved and is not crucial to conservation.
Research	Other Research	High	1) Conduct research that will enable robust predictions of breeding densities in different sub-habitat types, which can be extrapolated across VT and entire breeding range to derive population estimates. Design specific studies that will quantify BITH breeding densities, correlate density measures with GIS habitat data to estimate overall population numbers in different montane forest subhabitat types. Use data to generate overall population estimates. 2) There are needs for additional research on the species' wintering grounds.
Monitoring	Population Change	High	Mountain Birdwatch is a critical, ongoing program to monitor population trends.
Monitoring	Habitat Change	High	Mountain Birdwatch will collect habitat information with bird population data, as a means to evaluate local changes that may be occurring. Landscape level monitoring of montane forest habitats is essential.
Monitoring	Range Shifts	High	Ongoing monitoring through Mountain Birdwatch will help assess distributional changes, as will programs underway in Quebec and Canadian Maritime provinces, where the species also breeds.
Monitoring	Monitor Threats	High	Landscape level studies of the impacts of development on montane forest species will be the onnly means to document changes that occur. The current strategy of reacting to site-specific projects (e.g. East Mountain wind farm) is unlikely to provide rigor rigorous information that can be applied across the species' range.
Monitoring	Other Monitoring Needs	High	Monitoring must be continued on the species' winter range.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Bicknell's Thrush**
 Scientific Name: **Catharus bicknelli**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks	High	Identify the 15 core monitoring sites, ensure that funding is available to monitor them annually and indefinitely	Coordinate with Mountain Birdwatch program to ensure annual coverage of sites	VCE, VFWD, GMC, USFS	SWG, USFWS, GMNF
Compatible Resource Use	High	Evaluate impacts of human development (ski area expansion/construction, wind power, telecommunications facility) on Bicknell's Thrush and montane forest habitat, and use results to guide future development	Evaluate published results of past studies, implement new studies as appropriate to investigate development impacts, develop and periodically revise as necessary guidelines to minimize and mitigate impacts, monitor post-construction response of BITH	ANR, USFS, TNC, VCE	SWG, GMNF, USFWS
Habitat Restoration	Medium	Implement experimental habitat manipulation measures to evaluate the possibility of creating suitable habitat for BITH through artificial disturbance	Conduct controlled habitat manipulations of montane forest to mimic natural disturbance events (e.g., fir waves, catastrophic storm events); carefully monitor BITH and vegetation responses over time	VFWD, VFPR, GMNF, VCE	SWG, GMNF, NFWF
Planning & Zoning	High	Develop a planning process whereby explicit mitigation and management guidelines are specified. Further develop a means to ensure that these are followed and monitored, both in short- and long-term.	Formalize as policy existing recommendations for ski areas and develop new recommendations as research findings warrant. Establish accountability by land owners/managers to adopt specified measures.	ANR, USFS, VCE	SWG, PR
Publically-Owned Protected Areas	High	Identify top 15 breeding sites (those with largest habitat blocks and/or largest known breeding concentrations), specify these as highest priority for long-term protection/ conservation/monitoring. Ensure minimal or no further habitat loss at these sites	Use GIS to identify 15 largest montane forest habitat patches, review current protected status of each, assess further needs for long-term protection, develop site-specific plans for each site.	VFWD, USFS, TNC, VCE	State Wildlife Grants, GMNF, USFWS



Common Name: **Bicknell's Thrush**
Scientific Name: **Catharus bicknelli**
Species Group: **Bird**

Bibliography

- Atwood, J.L., C.C. Rimmer, K.P. McFarland, S.H. Tsai, and L.R. Nagy. 1996. Distribution of Bicknell's Thrush in New England and New York. *Wilson Bull.* 108:650-661.
- Goetz, J.E., K.P. McFarland, and C.C. Rimmer. 2003. Multiple paternity and multiple male feeders in Bicknell's Thrush. *Auk* 120:1044-1053.
- Lambert, J.D., K.P. McFarland, C.C. Rimmer, S.D. Faccio, and J.L. Atwood. 2005. A practical model of Bicknell's Thrush habitat in the Northeastern United States. *Wilson Bulletin*. In press.
- Laughlin, S. B., and D. P. Kibbe. 1985. *The Atlas of Breeding Birds of Vermont*. University Press of New England, Hanover, NH. 456 pp.
- Rimmer, C.C., K.P. McFarland, W.G. Ellison, and J.E. Goetz. 2001. Bicknell's Thrush (*Catharus bicknelli*). In *The Birds of North America*, No. 592 (A. Poole & F. Gill, eds.). The Birds of North America, Inc., Philadelphia, PA.
- Rimmer, C.C., K.P. McFarland, D.C. Evers, E.K. Miller, Y. Aubry, D. Busby, and R.J. Taylor. 2005. Mercury levels in Bicknell's Thrush and other insectivorous passerine birds in montane forests of northeastern North America. *Ecotoxicology* 14:223-240.
- Rimmer, C.C. and K.P. McFarland. 2000. Migrant stopover and postfledging dispersal at a montane forest site in Vermont. *Wilson Bull.* 112:124-136.
- Rimmer, C.C., J.L. Atwood, K.P. McFarland, and L.R. Nagy. 1996. Population density, vocal behavior and recommended survey methods for Bicknell's Thrush. *Wilson Bull.* 108:639-649.
- Rimmer, C.C. 2013. *The Second Atlas of Breeding Birds of Vermont*. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- Strong, A.M., C.C. Rimmer, and K.P. McFarland. 2004. Effect of prey biomass on the reproductive success and mating system of a polygynandrous songbird, the Bicknell's Thrush (*Catharus bicknelli*). *Auk* 121:446-451.
- Strong, A.M., C.C. Rimmer, K.P. McFarland, and K. Hagen. 2002. Effects of mountain resorts on wildlife. *Vermont Law Review* 26:689-716.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Wood Thrush**
Scientific Name: **Hylocichla mustelina**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The USGS Breeding Bird Survey (BBS) reports for the period 1966-2012 the Vermont Wood Thrush population experienced an estimated 3.5% annual decline. The same BBS reported annual Wood Thrush declines of 4.8% in New Hampshire and 2.8% in the New England/Atlantic Coast Region. Partners In Flight (PIF) estimates the continental population of 11,000,000 birds. The species is declining at similar rates in NH, ME, MA, and NY and throughout USFWS Region 5. A common but declining breeding species of northern hardwoods forests, in many ways an "umbrella" species that merits focused conservation attention in VT. Its decline may be due in part to maturation of northern hardwoods forests and the fragmentation of existing forests for human development. Certain forest management practices that encourage younger aged stands and conservation of forest blocks may help stabilize Wood Thrush declines. Factors on the species' Central American wintering grounds may also be involved in its declining populations. As a forest health umbrella species the Wood Thrush's re-categorization to High Priority represents concerns for the future of the northern hardwood forest in the face of ever expanding human development.

Distribution

Distributed widely throughout state in n. hardwood forests

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Upland, mesic northern hardwood forests, Interior and edges of deciduous and mixed forests, especially well-developed, upland, mesic sites. Key elements for suitability include: trees > 16 in height, high variety of deciduous tree species, moderate subcanopy and shrub density, shade, fairly open forest floor, moist soil, and decaying leaf litter

Habitat Types:

Northern Hardwood



Common Name: **Wood Thrush**
 Scientific Name: **Hylocichla mustelina**
 Species Group: **Bird**

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Fragmentation

Description of habitat threat(s): Forest fragmentation may introduce nest predators and cowbirds that lower reproductive success, especially in smaller, isolated patches

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Acidification of northern hardwoods forests and consequent calcium depletion may affect population ecology of this species

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Habitat-specific data needed in relation to assess area sensitivity in VT
Research	Basic Life History	Medium	Age- and sex-specific survivorship and reproductive success in different forest subhabitats and patch sizes needed.
Research	Distribution and Abundance	Medium	Age- and sex-specific partitioning (relative abundance and density) in different forest subhabitats and patch sizes needed
Research	Threats and Their Significance	High	1) Need to understand relative importance of differing fragmentation effects on demography, productivity, and site persistence; also how species responds to different forestry practices. Conduct field studies in different forest types (successional stage, patch size and configuration, proximity to edge habitat), use results to guide conservation planning that incorporates forestry and sustainable development. 2) Investigate environmental stressors like mercury and calcium depletion in Wood Thrush, as a means to understand their synergistic role in avian population and forest health; use results to guide regulatory planning for Hg and acidic ion emissions. Conduct studies to measure levels of Hg and Ca in WOTH and in ecosystem food chain; correlate measures to WOTH demographics and reproductive success.
Monitoring	Population Change	High	Need continued, habitat-specific monitoring in core no. hardwoods habitats, both managed and unmanaged landscapes
Monitoring	Habitat Change	High	Need to document and understand impacts of landscape-level forestry practices and atmospheric pollution on species' population biology
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	Medium	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Wood Thrush**
 Scientific Name: **Hylocichla mustelina**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks	Medium	Educate landowners, private foresters, local conservation and planning commissions about habitat conservation needs of Wood Thrush, as a means to guide sustainable land use practices and local regulations	Develop educational materials based on known information about Wood Thrush ecology, habitat needs and conservation in VT. Provide planning expertise to local planners, landowners, foresters.	VFWD, VFPR, USFS, TNC	Municipal planning grants, NFWF
Standards	High	Evaluate and refine current forestry practices as a means to promote optimal habitat suitability for this species, and to reverse population declines	Synthesize management studies from other parts of species' breeding range and evaluate applications to VT; conduct focused studies to assess species' response to differing forestry regimes	VFPR, VFWD, USFS, VT Assoc Loggers, private foresters, VWA, Coverts	NFWF, USFWS, SWG

Bibliography

Roth, R.R., M.S. Johnson, and T.J. Underwood. 1006. Wood Thrush (*Hylocichla mustelina*). In *The Birds of North America*, No. 246 (A. Poole and F. Gill, eds.). The Birds of North America, Philadelphia, PA.

Schwenk, S.. 2013. *Second Atlas of Breeding Birds of Vermont*. Rosalind Renfrew, ed. 547 pp. University Press of New England.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Brown Thrasher**
Scientific Name: **Toxostoma rufum**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

BBS trend 1966-2012 for VT was -4.5 and for 2002-2012 was -3.77 ($p = 0.25$). The Second Vermont Breeding Bird Atlas showed a 47% decrease in breeding block occupation from the first atlas. These trends suggest recent declines remain similar to long-term declines and as such Brown Thrasher has been moved from Medium to High Priority. Species declining along with shrub dominated and successional habitats throughout the East as forests mature and suitable habitat is converted to non-forest/non-habitat use.

Distribution

Statewide

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Probable	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Thicket/shrub complexes; hedgerows and early successional habitat w/ high stem densities coupled with low (10%-30%) canopy coverage.



Common Name: **Brown Thrasher**
Scientific Name: **Toxostoma rufum**
Species Group: **Bird**

Habitat Types:

Early Succession Boreal Hardwoods
Early Succession Pine and Hemlock
Early Succession Other Types
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops
Shrub Swamps
Early Succession Northern Hardwoods
Early Succession Upland Oak

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Inadequate Disturbance Regime
Impacts of Roads or Transportation Systems
Habitat Fragmentation
Incompatible Recreation

Description of habitat threat(s): Losses of shrub dominated and early successional woody regeneration habitats due to conversion and forest maturation.

Non-Habitat Threats:

Pollution
Trampling or Direct Impacts

Description of non-habitat threat(s): Evidence exists that nest discovery/disturbance evokes high rate of abandonment; evidence exists that pesticide use in feeding areas resulted in substantial declines.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Effects of land management practices on nest site selection and productivity/mortality.
Research	Basic Life History	Medium	More comprehensive breeding ecology information.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Brown Thrasher**
 Scientific Name: **Toxostoma rufum**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Determine appropriate old field habitat targets for state and private lands and restore and maintain old field habitats where needed to increase suitable ES songbird habitat.	Number of acres positively affected by management. Population response to management.	ANR, USFS, Audubon, Forest Products Association, Power companies, VT Loggers Association	PR, EQUIP, private grants
Awareness Raising and Communications	High	Initiate public education campaigns to highlight the need for active, management on public and private lands to create and maintain early successional shrubland habitat for the suite of shrubland birds.	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs.	VFWD, Audubon, VCE, power companies	PR, EQUIP, private grants

Bibliography

Cavitt, J. F. and C.A. Haas. 2000. Brown Thrasher (*Toxostoma rufum*). In *The Birds of North America*, No. 557 (A. Poole and F. Gill, eds.). The Birds of North America, Inc. Phila. Pa.

Laughlin, S. B. and D. P. Kibbe, editors. *The atlas of breeding birds of Vermont*. University of New England Press. Hanover, New Hampshire, USA.

Hunt, P. 2013. *The Second Atlas of Breeding Birds of Vermont*. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.



Common Name: **Blue-winged Warbler**
Scientific Name: **Vermivora pinus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2S3B

State Trend: Increasing

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

National Breeding Bird Survey data indicate a 2.44% long term population decline in the New England/Mid-Atlantic region for Blue-winged Warbler (Sauer 2011). But, results of the Second Vermont Breeding Bird Atlas (Renfrew 2013) indicate a 125% increase in Blue-winged Warblers between this publication and the First Vermont Breeding Bird Atlas (Laughlin and Kibbe 1985). Species distribution has shifted since the early 1980s, and appears to be moving north. PA and MD observed declines in their recent atlases, while Ontario and NY had increases (Renfrew 2013). Species is listed as Special Concern due to small population size and declining habitat.

Distribution

The Second Vermont Breeding Bird Atlas (Renfrew 2013) reports Blue-winged Warbler distribution to be largely in the western biophysical regions that border New York State where 81% of the species records are found. The Northern Green Mountain and Southern Vermont Piedmont regions hold the remaining 5% and 14%, of the distribution respectively. The species likely occurs in other regions of the state during migration.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Unknown
Northern Green Mtns	Probable	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Confident
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Nests in brushy growth near the borders of swamps or streams, forest edges, abandoned fields and pastures, thickets, and second-growth woods. Prefers brushy old pastures and old fields with saplings < 3 m tall" (DeGraaf and Yamasaki 2001).



Common Name: **Blue-winged Warbler**
Scientific Name: **Vermivora pinus**
Species Group: **Bird**

Habitat Types:

Early Succession Pine and Hemlock
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Shrub Swamps
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration

Description of habitat threat(s): Succession of old field to forest, loss of habitat to development, and fragmentation are major problems.

Non-Habitat Threats:

Genetics
Predation or Herbivory

Description of non-habitat threat(s): Hybridization with Golden-winged Warbler. Although hybridizes extensively with Golden-winged Warbler, introgressive hybridization appears to be asymmetric with Blue-winged Warbler gene pool remaining largely "pure" (Gill et al. 2001). Competition between Blue-winged Warbler and Golden-winged Warbler appears to be leading to continual northward shift in the range of Golden-winged Warbler. Colonization of Golden-winged Warbler breeding sites by Blue-winged Warbler may have negative impacts on Golden-winged Warbler populations. Brown-headed Cowbird parasitism is also a significant problem,



Common Name: **Blue-winged Warbler**
Scientific Name: **Vermivora pinus**
Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Develop a better understanding of the characteristics of high quality habitat would be valuable, as are optimal management activities. Audubon VT and NRCS are working together to answer these questions.
Research	Distribution and Abundance	High	Likely more common in Vermont than Breeding Bird Atlas data suggest. Because habitat is transitory and song is not necessarily a good indicator of presence (because of hybridization) there is a need to characterize the species' distribution in the State. Audubon VT is currently doing some of this work.
Research	Threats and Their Significance	High	Understanding effects of Brown-headed Cowbirds, Golden-winged Warbler, development, and succession are necessary to manage and conserve the species in VT.
Monitoring	Population Change	Medium	Knowing how long a patch remains suitable would be useful, as is knowing how long after management Blue-winged Warbler will return to a site. Additionally, understanding how species interacts with Golden-winged Warbler presence and Brown-headed Cowbird parasitism would also be helpful.
Monitoring	Habitat Change	High	Quantify the relative importance of succession and development.
Monitoring	Range Shifts	Medium	Determine if the species is moving northward in VT and if there are areas that are "pure" Blue-winged Warbler sites.
Monitoring	Monitor Threats	Medium	Get better information about current limiting factors to habitat (development versus habitat succession). Implement periodic assessment (5 year?) of grassland/shrubland acreage in Vermont, likely through GIS analysis.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Blue-winged Warbler**
 Scientific Name: **Vermivora pinus**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications	Medium	Initiate education campaigns highlighting the need for active, even age forest management on public and private lands to create and maintain seedling/sapling forest habitat complexes. Use demonstration sites to educate public and professionals about BMPs	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs.	Audubon VT, NRCS, VFWD, Towns, land trusts, etc.	Private grants
Easements	High	Create at least one large (>1000 ha) management area dedicated to early successional species.	Acres of land purchase or conservation easements with dedicated management plan acquired.	UVM, NRCS, VFWD.	NRCS, USDA.
Habitat Restoration	High	Determine appropriate old field habitat targets for state lands and restore and maintain old field habitats where needed to increase suitable ES songbird habitat.	Number of acres positively affected by management. Population response to management.	ANR, USFS, Audubon VT, Forest Products Association, VT Loggers	PR, EQIP

Bibliography

- Clark, D. B. 1985. Blue-winged Warbler. Pages 274-275 in S. B. Laughlin and D. P. Kibbe, eds. The atlas of breeding birds of Vermont. University Presses of New England, Hanover, New Hampshire, USA.
- DeGraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University of New England Press, Hanover, New Hampshire, USA, and London, UK.
- Gill, F. G., R. A. Canterbury, and J. L. Confer. 2001. Blue-winged Warbler (*Vermivora pinus*). In The Birds of North America, No. 584 (A. Poole and F. Gill, Eds.). The Birds of North America, Inc., Philadelphia, PA.
- Kibbe, D.P. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- Laughlin, S. B., and D. P. Kibbe. 1985. The Atlas of Breeding Birds of Vermont. University Press of New England, Hanover, NH. 456 pp.
- Sauer, J. R., J. E. Hines, J.E. Fallon, K.L. Pardieck, D.J. Ziolkowski, Jr., and W.A. Link. 2011. The North American breeding bird survey, results and analysis 1966-2009. Version 3.23.2011. USGS Patuxent Wildlife Research Center, Laurel, MD.



Common Name: **Golden-winged Warbler**
Scientific Name: **Vermivora chrysoptera**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: S2S3B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

National Breeding Bird Survey data show a long term Golden-winged Warbler population decline of 10.8% in the Northeast/Mid-Atlantic regions (Sauer 2011). Results of the Second Vermont Breeding Bird Atlas (Renfrew 2013) indicate a 7% decline between this effort and the First Vermont Atlas (Laughlin 1985). It is suspected the species has been negatively affected by Brown-headed Cowbird parasitism and hybridization with Blue-winged Warblers (Confer 1992), as well as forest succession (Renfrew 2013). Audubon Vermont surveys conducted in 2013 and 2014 found approximately 60 Golden-winged Warblers in the southern Champlain Valley, which is significantly more than the previously estimated population of 20 pairs (Audubon VT, unpubl. data). Partners in Flight report Golden-winged Warblers to be a Tier 2 species (Rosenberg and Wells 2005). It is likely that forest succession and development of abandoned agricultural habitat will continue to limit this species.

Distribution

The Second Vermont Breeding Bird Atlas indicates 100% of the confirmed nesting distribution of the species is found among the Champlain Valley, Vermont Valley, and Taconic Mountain biophysical regions (Renfrew 2013). The species may occur in other regions as migrant.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Confident
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Preferred habitat in Vermont is a mosaic of shrub thickets, forbs, grasses, scattered trees or saplings, and forest edges (Confer 1992, and C. Smalling, pers. comm.). Also uses power line rights-of-way (ROWS) (Golden-winged Warbler Working Group 2013). The Golden-winged Warbler Working group (2013) recommends the following for optimal Golden-winged Warbler habitat, as long as the area is next to a primarily deciduous forest patch and non-active agriculture (such as row crops):

- 30-70% tall shrubs and saplings (3–13 ft.) distributed unevenly
- Herbaceous openings (mostly forbs and some grasses)
- Deciduous trees (5–15/acre), creating 10–30% canopy cover



Common Name: **Golden-winged Warbler**
Scientific Name: **Vermivora chrysoptera**
Species Group: **Bird**

Habitat Types:

Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration

Description of habitat threat(s): Loss of habitat through development and succession. Habitat quality may decline with presence of Brown-headed Cowbird and Blue-winged Warbler.

Non-Habitat Threats:

Genetics

Description of non-habitat threat(s): Hybridization with Blue-winged Warbler may be causing range shifts and contribute to overall decline of the Golden-winged Warbler.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	A better understanding of the characteristics of high quality habitat would be valuable, as are optimal management activities. Audubon VT and NRCS are working together to answer these questions.
Research	Distribution and Abundance	High	Because habitat is transitory and song is not necessarily a good indicator of presence (because of hybridization) there is a need to characterize the species' distribution in the State. Audubon VT is currently doing some of this work.
Research	Threats and Their Significance	High	Understanding effects of cowbirds, BWWA, development, and succession are necessary to manage and conserve the species in VT.
Research	Population Genetics	Medium	Quantifying the genetic "purity" of the VT population would be valuable, especially understanding N-S variation. Hybridization likely a factor.
Monitoring	Population Change	High	Knowing how long a patch remains suitable would be useful, as is knowing how long after management GWWA will return to a site. Additionally, understanding how species reacts to BWWA presence and BHCO parasitism would also be helpful.
Monitoring	Habitat Change	Medium	Quantify the relative importance of succession and development.
Monitoring	Range Shifts	Medium	Determine if the species being pushed northward in VT as a result of hybridization and genetic swamping by Blue-winged Warbler
Monitoring	Monitor Threats	Medium	Monitor the effects of Blue-winged Warbler, development, succession, and Brown-headed Cowbird. In particular better information about current limiting factors to habitat (development versus succession). Periodic assessment (5 year?) of early successional acreage in Vermont, possibly through GIS analysis.



Common Name: **Golden-winged Warbler**
 Scientific Name: **Vermivora chrysoptera**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Stabilize or reverse declining population trend for GWWA through habitat management via NRCS, USFWS funds, and private and corporate landowners	Population response to management, targeted surveys.	VFWD, NRCS, Audubon VT, VELCO, private & corporate landowners	NRCS, private and other public funding sources (NFWF)
Research	Medium	Determine if there are any habitats that are used solely by GWWA and prioritize for conservation. Research genetics of VT GWWA/BWWA population to determine the level of genetic swamping.	Stable population of GWWA in the presence of BWWA.	UVM, Audubon VT, VFWD.	NFWF. PR, SWG
Awareness Raising and Communications	Medium	Initiate education campaigns highlighting the need for active, even-age forest management on public and private lands to create and maintain seedling/sapling forest habitat complexes. Use demonstration sites to educate public and professionals about BMPs.	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs	Audubon VT, NRCS, VFWD, Towns, land trusts	Private foundations
Habitat Restoration	High	Establish at least one large (>1000 ha) management area dedicated to early successional species.	Land purchase or conservation easements and dedicated management plan.	VFWD, NRCS, TNC.	SWG, NRCS

Bibliography

- Clark, D. B. 1985. Golden-winged Warbler. Pages 276-277 in S. B. Laughlin and D. P. Kibbe, eds. The atlas of breeding birds of Vermont. University Presses of New England, Hanover, New Hampshire, USA.
- Confer, J. L. 1992. Golden-winged Warbler (*Vermivora chrysoptera*). In The Birds of North America, No. 20 (A. Poole and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, D. C.; The American Ornithologists' Union.
- DeGraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University of New England Press, Hanover, New Hampshire, USA, and London, UK.
- Golden-winged Warbler Working Group. 2013. Best management practices for Golden-winged Warbler habitats in the Great Lakes region. www.gwwa.org.
- Kibbe, D.P. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- Laughlin, S. B., and D. P. Kibbe. 1985. The Atlas of Breeding Birds of Vermont. University Press of New England, Hanover, NH. 456 pp.
- Sauer, J. R., J. E. Hines, J.E. Fallon, K.L. Pardieck, D.J. Ziolkowski, Jr., and W.A. Link. 2011. The North American breeding bird survey, results and analysis 1966-2009. Version 3.23.2011. USGS Patuxent Wildlife Research Center, Laurel, MD.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Chestnut-sided Warbler**
Scientific Name: **Dendroica pensylvanica**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

This species is slowly declining in Vermont according to Breeding Bird Survey data. However, it was found widely distributed throughout the state in both breeding bird atlases. Most gains from the first atlas occurred in the Champlain Valley, including the Lake Champlain islands (Renfrew2013). Shrub habitats are a necessary component for nesting either through succession of agricultural fields or periodic active forest management.

Distribution

Found in all regions of the state with increases noted mainly in the northern part of the Champlain Valley. Distribution has remained consistent between the two breeding bird atlases.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Forested or shrubby habitats during migration. The species requires shrubs for nesting mostly in early successional hardwood forests that are 1-3m in height (DeGraaf et. al. 2006). Highly specialized habitat for breeding, confined to early-successional deciduous forest ranging from wet to dry sites. This habitat can be provided using even-aged management, or uneven-aged forest management using group selection tree cutting that promotes shrub development. Not known to use coniferous forest habitats.



Common Name: **Chestnut-sided Warbler**
Scientific Name: **Dendroica pensylvanica**
Species Group: **Bird**

Habitat Types:

Early Succession Boreal Hardwoods
Early Succession Pine and Hemlock
Early Succession Northern Hardwoods
Shrub Swamps
Early Succession Upland Oak
Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Inadequate Disturbance Regime
Habitat Fragmentation

Description of habitat threat(s): Loss of successional habitats due to conversion to non-forested uses and suppression of natural disturbance (flood, fire), and reductions in active forest management.

Non-Habitat Threats:

Pollution
Loss of Prey Base

Description of non-habitat threat(s): Some indications that Lepidoptera larvae reductions due to pesticides and biological controls can decrease productivity.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Basic Life History	Medium	Basic research on breeding activities is needed, particularly to determine trends in re-nesting and lifetime broods/reproductive success

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Chestnut-sided Warbler**
 Scientific Name: **Dendroica pensylvanica**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Conservation Finance	Medium	Create a state-funded, private lands, early successional habitat improvement initiative (modeled on NH's Small Landowner Grant program). Fund for > \$50,000/yr with revenues from state lands forest management. This could offset landowner EQIP obligations.	Level of funds raised.	FWD	SWG
Awareness Raising and Communications	Medium	Initiate public education campaigns to highlight the need for active, even age forest management on public and private lands to create and maintain seedling/sapling forest habitat complexes.	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs.	VFWD, USFS, USFWS	SWG, PR
Habitat Restoration	High	Stabilize or reverse declining population trend for Chestnut-sided warblers to realize and maintain a survey value of 14-15 per BBS route or between 120,000 to 180,000 individuals (Rosenberg 2004).	Population response to management, BBS surveys.	Audubon, VFWD	PR, EQIP
Habitat Restoration	Medium	Determine appropriate old field habitat targets for state lands and restore and maintain old field habitats where needed to increase suitable ES songbird habitat.	Number of acres positively affected by management. Population response to management.	ANR, USFS, Audubon, Forest Products Association, VT Loggers	PR, EQIP

Bibliography

- DeGraaf, R.M., M. Yamasaki, W.B. Leak, and A.M. Lester. 2006. Technical Guide to Forest Wildlife Management in New England. University Press of New England, Lebanon, NH. 305pp.
- Kibbe, D. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- Richardson, M., and D.W. Brauning. 1995. Chestnut-sided warbler (*Dendroica pensylvanica*). In: Birds of North America, No. 190. (A. Poole and F. Gill, eds.) . The Academy of Natural Sciences, Philadelphia, and the American Ornithologists' Union, Washington, D.C.
- Rosenberg, K. V. 2004. Partners in Flight continental priorities and objectives defined at the state and bird conservation region levels, Vermont. Cornell Lab of Ornithology, Ithaca, NY. 26 p.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Black-throated Blue Warbler**
Scientific Name: **Dendroica caerulescens**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Stable

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

BBS survey trends variable by time period. Survey-wide, significant positive long- (1966-2012, +2.26%) and short-term trends (2002-2012; +3.77%). Vermont shows non-significant increases in both long- and short-term periods. Slight (3%) increase in number of blocks with breeding evidence between first and second breeding bird atlases. Population likely secure. Primary breeding ground limiting factors result from consequences of fragmentation (by permanent land use changes) on reproductive success. Because winter range is restricted to the Caribbean, there is some concern over the effects of deforestation on these islands, particularly the Greater Antilles.

Distribution

From Kibbe (1985).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Large, continuous tracts of undisturbed deciduous or mixed deciduous/coniferous forests, often in hilly or mountainous terrain (Holmes 1994). " Uneven aged timber harvest methods such as selection or group cuts can effectively mimic the natural disturbance regime and seem to be compatible with Black-throated Blue Warbler conservation" (Burdett and Niemi 2003).

Habitat Types:

Northern Hardwood

Current Threats

Habitat Threats:

Conversion of Habitat



Common Name: **Black-throated Blue Warbler**
 Scientific Name: **Dendroica caerulescens**
 Species Group: **Bird**

Habitat Alteration

Habitat Fragmentation

Climate Change

Description of habitat threat(s): Productivity and density decrease in forest fragments. Susceptible to parasitism by Brown-headed Cowbirds in these forests when located in predominately agricultural landscapes. Habitat conversion through development or habitat alteration from clearcutting can limit species. Conversely, as forest regenerates from abandonment of agricultural lands, additional habitat will become available through succession.

Species susceptible to decreased overwinter survival rates with El Niño cycle droughts in the Caribbean (Silllett et al. 2000). Some evidence that ENSO cycles are becoming more extreme. Simulations of range shift in response to climate change suggests extirpation from Vermont by 2080 (National Audubon Society 2014).

Description of non-habitat threat(s):

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	Medium	1) Compare changes in population trends across BBS routes in areas with different land use patterns. 2) More information is needed on distribution and overwinter survival in disturbed habitats on wintering grounds.
Research	Threats and Their Significance	High	Effects of fragmentation on breeding season productivity, particularly the effects of predation and cowbird parasitism should be further researched. Landscapes dominated by contiguous forests have not consistently shown increased predation trends associated with relatively temporary disturbances, such as timber harvests; "It is critical to better comprehend the complex relationships that exist between nest predation, habitat fragmentation, and landscape context." (Burdett and Niemi 2003).
Monitoring	Population Change	Medium	A better assessment of population trends across a variety of landscape types is needed.
Monitoring	Habitat Change	High	1) Better information on land use change in Vermont would help concentrate development in areas that would be less likely to affect forest interior species. Trends in rate of forest loss and fragmentation across range should be investigated. 2) Continue to monitor populations at Hubbard Brook Experimental Forest (unfragmented), uneven-aged-managed forests, and forest fragments in Vermont to better assess the effect of patch size and management on population trends using a source-sink framework

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Black-throated Blue Warbler**
 Scientific Name: **Dendroica caerulescens**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Compatible Resource Use	Medium	Where timber resource extraction and/or other habitat management goals requiring timber cutting are desired, uneven aged management, using selection harvests, should be employed on a portion of public lands	Amount of public forests designated for unevenaged management.	ANR, USFS, USFWS	SWG, PR
Habitat Restoration	Medium	Identify contiguous forests blocks w/mature components & encourage their conservation via easements or other financial incentives on private lands. Conserve contiguous forest blocks on public lands via appropriate long-range management plan designations.	Number and distribution of core forest blocks conserved on private and public lands.		

Bibliography

Burdett, C.L. and G.J. Niemi. 2003. Conservation assessment for Black-throated Blue Warbler (*Dendroica caerulescens*) USDA Forest Service, Eastern Region. Natural Resources Research Institute, Univ. of Minnesota, Duluth, Minnesota, USA.

Kibbe, D. P.. 1985. Black-throated Blue Warbler. Pages 292-293 in S. B. Laughlin and D. P. Kibbe, eds. The atlas of breeding birds of Vermont. University Presses of New England, Hanover, New Hampshire, USA.

Holmes, R. T. 1994. Black-throated Blue Warbler (*Dendroica caerulescens*). In The Birds of North America, No. 87 (A. Poole and F. Gill, Eds.). Philadelphia: The Academy of Natural Sciences; Washington, D. C.; The American Ornithologists' Union.

National Audubon Society. 2014. National Audubon Society Climate Report. Available online at: <http://climate.audubon.org/birds/btbwar/black-throated-blue-warbler>

Robbins, C. S., D. K. Dawson, and B. A. Dowell. 1989. Habitat area requirements of breeding forest birds of the Middle Atlantic States. Wildlife Monographs 103:1-34.

Sauer, J. R., J. E. Hines, and J. Fallon. 2004. The North American Breeding Bird Survey, Results and Analysis 1966-2003. Version 2004.1. USGS Patuxent Wildlife Research Center, Laurel, Maryland, USA.

Sillett, T. S., R. T. Holmes, and T. W. Sherry. 2000. Impacts of a global climate cycle on population dynamics of a migratory songbird. Science 288:2040-2042.

Strong, A. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Prairie Warbler**
Scientific Name: **Dendroica discolor**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S3B

State Trend: Stable

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

BBS trend for eastern region, 1966-2012 = -2.04 and 2002-2012 = -1.07. Significant declines in the US per BBS data 1966 -2012; less significant decline 2002-2012. Distribution during first Vermont Bird Atlas was limited to the eastern foothills but shifted in large part to the southern Champlain Valley during the second Vermont Breeding Bird Atlas. Second Breeding Bird Atlas showed a 66% increase in priority block occupancy (4 blocks to 7).

Distribution

Rare breeder in southern Champlain Valley and northern CT River drainage.

Distribution by Biophysical Region:

Champlain Valley	Probable	Southern VT Piedmont	Probable
Champlain Hills	Probable	Vermont Valley	Unknown
Northern Green Mtns	Unknown	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Breeding habitat of early successional hardwood forest regeneration, old field, shrub/dune, upland shrub habitats; prefers open canopy (however uses closed canopy palustrine forest in Mid-Atlantic breeding areas). Utilizes, powerline corridors, Christmas tree farms and gravel pit/mine shrub habitats.



Common Name: **Prairie Warbler**
Scientific Name: **Dendroica discolor**
Species Group: **Bird**

Habitat Types:

Upland Shores
Outcrops and Alpine
Cliffs and Talus
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Hardwood Swamps
Softwood Swamps
Seeps and Pools
Northern Hardwood
Floodplain Forests
Early Succession Boreal Hardwoods
Early Succession Pine and Hemlock
Early Succession Northern Hardwoods
Early Succession Other Types
Early Succession Spruce-Fir
Early Succession Upland Oak
Mine
Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Inadequate Disturbance Regime
Habitat Fragmentation

Description of habitat threat(s): Succession of old field habitats and forest maturation have caused habitat decline in parts of Vermont.

Non-Habitat Threats:

Parasites

Description of non-habitat threat(s): Parasitized by brown-headed cowbird.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Prairie Warbler**
 Scientific Name: **Dendroica discolor**
 Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Effects of land management practices on nest site selection and productivity/mortality
Research	Basic Life History	Medium	More comprehensive breeding ecology information.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Determine appropriate old field habitat targets for state and private lands and restore and maintain old field habitats where needed to increase suitable ES songbird habitat.	Number of acres positively affected by management. Population response to management.	Audubon, VFWD, NRCS, Ruffed Grouse Society, Wild Turkey Federation	PR, EQUIP, private grants, power companies
Awareness Raising and Communications	Medium	Initiate public education campaigns to highlight the need for active, management on public and private lands to create and maintain early successional shrubland habitat for the suite of shrubland birds.	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs.	VFWD, Audubon, NRCS, VCE, power companies	PR, EQUIP, Private grants, Power companies

Bibliography

University Press of New England, Hanover N.H., USA and London, U.K.

Nolan, V., Jr., E.D. Ketterson, and C.A. Buerkle. 1999. Prairie Warbler (*Dendroica discolor*). In *The Birds of North America*, No. 455 (A. Poole and F. Gill, eds.). The Birds of North America, Inc. Phila. Pa.

Laughlin, S. B. and D. P. Kibbe, editors. *The atlas of breeding birds of Vermont*. University of New England Press. Hanover, New Hampshire, USA.

Hunt, P. 2013. *The Second Atlas of Breeding Birds of Vermont*. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.



Common Name: **Bay-breasted Warbler**
 Scientific Name: **Dendroica castanea**
 Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S1B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Breeding evidence documented in three priority blocks during the first breeding bird atlas and four blocks during the second atlas, none of which were the same between atlases. Only three confirmed breeding records for the state. Breeding Bird Survey results are not available for Vermont. The species' trends are variable across regions. However, analyses of population trends from Canada (which supports 98% of the breeding population) suggests 3-5%/year declines depending on the time period analyzed (Venier et al. 2011).

Distribution

First breeding confirmation was near Sable Mountain, Granby (1980). Subsequently confirmed nesting at Wenlock Wildlife Management Area, Ferdinand (1987) and Brighton State Park, Brighton (1995) and four blocks during the second atlas.

Distribution by Biophysical Region:

Champlain Valley	Unknown	Southern VT Piedmont	Unknown
Champlain Hills	Unknown	Vermont Valley	Unknown
Northern Green Mtns	Confident	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Unknown
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Mainly breeds in dense, boreal forests of mature spruce and fir; also inhabits old mixed-wood stands and prefers moist, swampy areas to dry, upland locations. (Mayasich and Niemi 2002). Dramatic increases reported in response to outbreaks of spruce budworm (*Choristoneura fumiferana*). "Foraging microhabitat preferences are the inner portions of mid-level branches that are among the large lichen-covered (scant foliage) limbs of conifers; also the inner portions of conifer branches among the dead limbs at lower heights." (Mayasich and Niemi 2002).



Common Name: **Bay-breasted Warbler**
 Scientific Name: **Dendroica castanea**
 Species Group: **Bird**

Habitat Types:

Spruce Fir Northern Hardwood
 Softwood Swamps
 Early Succession Boreal Conifers
 Early Succession Spruce-Fir

Current Threats

Habitat Threats:

Conversion of Habitat
 Habitat Fragmentation

Description of habitat threat(s): Some suggestion that Bay-breasted Warblers are susceptible to habitat conversion of wintering ground habitat. Additionally, conversion of natural forests to plantations of black spruce (*Picea mariana*) and jack pine (*Pinus banksiana*) plantations decrease habitat quality as these species are less susceptible to spruce-budworm (a key food resources for the Bay-breasted Warbler) outbreaks. Loss and fragmentation of late successional, lowland spruce-fir forest; management practices that favor short-cutting cycles preventing establishment of late successional habitat.

Description of non-habitat threat(s): Possible impacts from aerial spraying for spruce budworm (declines following application of organophosphate insecticides documented in New Brunswick in 1970s)

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Specific habitat association in VT needs better documentation
Research	Basic Life History	Medium	
Research	Distribution and Abundance	High	Not well documented in VT. A targeted survey of this and other late successional, lowland boreal forest birds is needed. Virtually no information exists now.
Research	Threats and Their Significance	High	Need to understand landscape level limiting factors (primarily via timber harvesting) to persistence of late successional, lowland boreal forests and population biology of this species
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	High	Traditional monitoring methods (BBS, VT Forest Bird Monitoring Program) do not adequately cover this species. Need to document trends in VT via a targeted survey of lowland boreal forests.
Monitoring	Habitat Change	High	Need to understand landscape level changes and limiting factors (primarily via poorly-planned timber harvesting) to persistence of late successional, lowland boreal forests. Quantify extent of current suitable breeding habitat.
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	Medium	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Bay-breasted Warbler**
 Scientific Name: **Dendroica castanea**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Publically-Owned Protected Areas	Medium	Identify potential areas on public lands for designation as reserve or no-cut status (including some areas where wildfires and insect outbreaks would not be controlled).	Number of sites and total area of designated reserves.	ANR, USFS, USFWS, UVM, TNC	SWG, PR
Habitat Restoration	Medium	Apply increased rotation ages to some managed spruce-fir forests on public lands.	Number of sites and total area with increased rotation ages.	ANR, USFS, USFWS	SWG, PR
Species Restoration	Medium	Determine current management regimes, and ensure that overall management is compatible with goal of maintaining or increasing current population levels. No PIF target was set for VT as "Population numbers are unavailable at this time" (Rosenberg 2004).			

Bibliography

Kibbe, D. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.

Mayasich, J. M. and G. J. Niemi 2002. Conservation assessment for Bay-breasted warbler (*Dendroica castanea*). USDA Forest Service, Eastern Region, Milwaukee, WI.

Rosenberg, K. V. Partners In Flight continental priorities and objectives defined at the state and bird conservation region levels -- Vermont. Cornell Lab of Ornithology, Ithaca, NY.

Tumosa, J. and S. Lemieux. 2002. Bay-breasted warbler excerpts from USFS GMNF reviews. www.natureserve.org.

Venier, L., S. Holmes, and J. M. Williams. 2011. Bay-breasted Warbler (*Setophaga castanea*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/206doi:10.2173/bna.206>

Williams, J.M. 1996. Bay-breasted Warbler. In The Birds of North America, No. 206 (A. Poole and F. Gill, eds.). The Birds of North America, Philadelphia, PA.



Common Name: **Blackpoll Warbler**
Scientific Name: **Setophaga striata**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S4S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

The breeding distribution for Blackpoll Warbler appears to not have changed much in last 25 years. However, data indicates significant changes within the distribution have occurred. This is also the case in Vermont. The species occupied 33 priority 1 blocks during the first Vermont atlas (1982) and 28 priority 1 blocks during the second atlas (2013). But during that interval, priority block gains numbered 8 while losses numbered 13. Blackpolls are found largely among the high elevation spruce-fir forests of the Green Mountains. Elevations greater than 2800 feet most often provide the greatest area of montane forest, the blackpoll's preferred habitat. Although the population appears to have been static during the last 30 years threats such as habitat loss to human development and accumulated environmental toxins such as mercury are likely an important factor in the species' long-term status. Current climate change models project the warbler's preferred montane forest habitat will decrease in size as it recedes northward. This suggests a seriously imperiled future for the species.

Distribution

This species is a high elevation species that breeds above 2800 feet in Vermont. It is detected throughout Vermont during spring and fall migration. Recent work by VCE biologists and biologists in Nova Scotia used geotrackers to show these birds fly non-stop over the Atlantic Ocean an average of 2540 km to the Greater Antilles or the northeastern coast of South America.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

In Vermont Blackpoll Warblers breed in dense thickets within montane forests dominated by balsam fir, with lesser amounts of spruce, white birch, and mountain ash.



Common Name: **Blackpoll Warbler**
Scientific Name: **Setophaga striata**
Species Group: **Bird**

Habitat Types:

Spruce Fir Northern Hardwood
Spruce Fir Northern Hardwood
Early Succession Spruce-Fir
Early Succession Spruce-Fir

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development
Habitat Alteration
Habitat Fragmentation
Climate Change

Description of habitat threat(s): Audubon's climate model suggest that this species will be effected by climate change as the spruce-fir habitat is shifting upslope, requiring the and the species breeding range to contract or shift north. Other potential problems include loss and fragmentation of montane forests from ski area, wind power and telecommunications development. Collision mortalities have been documented with wind energy facilities and telecommunication towers.

Non-Habitat Threats:

Pollution
Trampling or Direct Impacts

Description of non-habitat threat(s): Atmospheric pollution, including airborne mercury, could impact the species directly, as well as damage its habitat. Collision mortalities have been documented with buildings.



Common Name: **Blackpoll Warbler**
Scientific Name: **Setophaga striata**
Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Habitat needs reasonably well known, although population structure in different sub habitat types (krummholz, regenerating chronically disturbed forests, taller stature and more open forests, transitional spruce-fir-birch forests) not well known.
Research	Basic Life History	High	Demographics and breeding success need more study, especially in different sub habitat types.
Research	Distribution and Abundance	Medium	Fairly well known, although relative abundance in different sub habitat types needs to be better quantified. Conduct research that will enable robust predictions of breeding densities in different sub-habitat types, which can be extrapolated across breeding range to derive population estimates. Documenting shift range may be necessary if climate change has a big impact on Vermont's high elevation habitat.
Research	Threats and Their Significance	High	1) Species' susceptibility and response to habitat fragmentation and conversion from development (ski area, wind turbines, telecommunications facilities) needs to be better understood. Evaluate impacts of human development (ski area expansion/construction, wind power, telecommunications facility) on montane forest habitat, and use results to guide future development. 2) Impacts of atmospheric pollutants (e.g. mercury) and possible role of calcium depletion should be studied.
Research	Population Genetics	Low	Genetic structure of breeding populations in Northeast, and relation to core breeding populations in Canada interesting, but probably not crucial for conservation.
Research	Taxonomy	Low	Genetic structure of breeding populations in Northeast, and relation to core breeding populations in Canada interesting, but probably not crucial for conservation.
Monitoring	Population Change	High	Species poorly monitored by traditional methods like BBS. VCE Mountain Birdwatch program monitors adequately, but must be maintained for long-term. Very important to monitor this species as an avian indicator of montane forests. Continue long-term monitoring at a minimum of 15-20 sites in VT to document population trends. Support for gathering data from citizen scientists important.
Monitoring	Habitat Change	High	Important to document habitat changes in concert with population changes.
Monitoring	Range Shifts	Medium	This should be covered by a regional monitoring program (i.e. Mountain Birdwatch).
Monitoring	Monitor Threats	Medium	Important to monitor limiting factors like development, atmospheric pollution, mercury burdens, climate change, impacts of collisions.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Blackpoll Warbler**
 Scientific Name: **Setophaga striata**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Planning & Zoning	Medium	Develop a planning process whereby explicit mitigation and management guidelines are specified. Further develop a means to ensure that these are followed, and results monitored, both in short- and long-term.			
Research	High	Monitor development, atmospheric pollution, mercury burdens, climate change, impacts of collisions.		VCE, Audubon, BOVM, FWS	SWG
Habitat Restoration	High	Identify 10-15 core breeding sites and ensure that a long-term protection plan exists for each.	The number of sites protected		

Bibliography

DeLuca, William, Rebecca Holberton, Pamela D. Hunt and Bonita C. Eliason. 2013. Blackpoll Warbler (*Setophaga striata*), The Birds of North America Online (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/431>

DeLuca, W., Bradley K. Woodworth, Christopher C. Rimmer, Peter P. Marra, Philip D. Taylor, Kent P. McFarland, Stuart A. Mackenzie, D. Ryan Norris. 2015. Transoceanic migration by a 12 g songbird, *Biology Letters*. 11 20141045; DOI: 10.1098/rsbl.2014.1045.

eBird. 2012. eBird: An online database of bird distribution and abundance [web application]. eBird, Cornell Lab of Ornithology, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: Date February 4, 2015).

Laughlin, S. B. and D. P. Kibbe. 1985. The Atlas of Breeding Birds in Vermont. University Press of New England, Hanover, N.H.

National Audubon Society. 2014. Audubon's Birds and Climate Change Report: A Primer for Practitioners. National Audubon Society, New York. Contributors: Gary Langham, Justin Schuetz, Candan Soykan, Chad Wilsey, Tom Auer, Geoff LeBaron, Connie Sanchez, Trish Distler. Version 1.2.

Rimmer, C. and K. P. McFarland. 2013. Blackpoll Warbler (*Dendroica striata*). In: The Second Atlas of Breeding Birds of Vermont, (R. Renfrew, ed.). University Press of New England, Hanover and London.



Common Name: **Cerulean Warbler**
Scientific Name: **Dendroica cerulea**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G4

Global Trend:

State Rank: S1B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Strong significant long-term decline (-4.2%) based on survey-wide BBS data. No data from Vermont.

Historic population at the mouth of the Lamoille River appears to be extirpated. In 2002-2004, singing males have been located in Niquette Bay State Park, Highgate, and near Colchester Pond. More populations may be discovered as Vermont's forests continue to mature (and the second breeding bird atlas is completed), however declines throughout its range suggest that this species will never be common in Vermont. Preliminary survey data from the wintering range suggest that it can be found in a diversity of forest types in Columbia, Venezuela, Ecuador, and Peru (Hamel 2000).

Distribution

Data from Ellison (1985).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Unknown
Champlain Hills		Vermont Valley	Unknown
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Unknown
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

As summarized by Hamel (2000), this species requires a closed canopy, presence of scattered tall, old growth canopy trees, and distinct layering of foliage from ground cover to canopy. Area sensitivity varies by area with minimum patch size 20-30 ha in Ohio to 700 ha in Middle Atlantic States to 1,600 ha in the Mississippi Alluvial Valley, but breeding occurs in 10 ha patches in Ontario (summarized by Hamel 2000).

Habitat Types:

Oak-Pine Northern Hardwood
Northern Hardwood
Floodplain Forests



Common Name: **Cerulean Warbler**
Scientific Name: **Dendroica cerulea**
Species Group: **Bird**

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Fragmentation

Description of habitat threat(s): Area sensitive in parts of its range, suggesting fragmentation a problem to population. Development or harvest of mature upland forests will decrease available habitat.

Description of non-habitat threat(s):

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Species has been relatively well-studied on its breeding range, including recent studies from Ontario. However, better summaries of this information may lead to directed searches for new populations in Vermont. Better information about habitat requireme
Research	Basic Life History	Medium	Also relatively well-studied, however more information about non-breeding social system, particularly as to whether or not they are an obligate flock follower.
Research	Distribution and Abundance	High	1) Directed surveys in Vermont are necessary to better understand their present status in the state. 2) Better information on distribution in Vermont will be critical to conserving the species and predicting future distribution. Intensively monitor (as le
Research	Threats and Their Significance	Low	Presumably habitat quality in Vermont will increase as forests mature. However, some information on minimum patch size would help in understanding the effects of development.
Monitoring	Population Change	High	Population trends in Vermont will be difficult to assess without more information on distribution. But all known local populations should be carefully monitored.
Monitoring	Habitat Change	Low	As forest regenerates from abandonment of agricultural lands, habitat will become available through succession. Population response of CERW will be difficult to assess. Forest growth models might be useful in helping to predict future occurrences in the s

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Cerulean Warbler**
 Scientific Name: **Dendroica cerulea**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Identify contiguous forests blocks w/mature components & encourage their conservation via easements or other financial incentives on private lands. Conserve contiguous forest blocks on public lands via appropriate long-range management plan designations.	Number and distribution of core forest blocks conserved on private and public lands	ANR, USFS, USFWS, VHCB, VLT, TNC	SWG, PR, VHCB
Easements	Medium	Maintain of large forest tracts, particularly in Champlain Valley and Taconic regions.	Maintenance of large forest tracts, particularly in Champlain Valley and Taconic regions.	TNC, VFWD, Forest Legacy program	TNC

Bibliography

Ellison, W.G. 1985. Cerulean Warbler. Pages 308-309 in S. B. Laughlin and D. P. Kibbe, eds. The atlas of breeding birds of Vermont. University Presses of New England, Hanover, New Hampshire, USA.

Hamel, P. B. 2000. Cerulean Warbler (Dendroica cerulea). In The Birds of North America, No. 511 (A. Poole and F. Gill, Eds.). Philadelphia: The Birds of North America, Philadelphia, PA.

Sauer, J. R., J. E. Hines, and J. Fallon. 2004. The North American Breeding Bird Survey, Results and Analysis 1966-2003. Version 2004.1. USGS Patuxent Wildlife Research Center, Laurel, Maryland, USA.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Canada Warbler**
Scientific Name: **Cardellina canadensis**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S4B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

In Vermont the results from the second Breeding Bird Atlas indicate this bird is declining in Vermont. Block occupancy dropped 31% between the Atlases. Regional atlases and breeding bird surveys have also demonstrated a decline throughout the region over the past 30 years. As the climate and the forests change this species may be at a greater risk, not only on its breeding grounds, but also on its wintering grounds.

Distribution

Canada Warblers are found throughout Vermont. The Canada Warbler is considered a neotropical migrant and migrates from its breeding grounds to northern South America

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

A wide range of coniferous and deciduous forests, and mixed forests at all elevations, but especially mid-slopes in Green Mountains. Uses both mature and regenerating forest. Seem to prefer a dense understory with moss, and an uneven forest floor. Hummocks, roots, and debris are used to hide the nest and fledglings. Clearcuts and shelterwood cuts received more use than mature forest in northern New Hampshire. First appear in clearcuts 5 years after harvest, become common after 15 years and remain abundant until the next cutting cycle.



Common Name: **Canada Warbler**
Scientific Name: **Cardellina canadensis**
Species Group: **Bird**

Habitat Types:

Spruce Fir Northern Hardwood
Northern Hardwood
Softwood Swamps
Early Succession Boreal Conifers
Early Succession Spruce-Fir

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Climate Change
Habitat Fragmentation

Description of habitat threat(s): Forest succession, loss of forested wetlands, and development all may influence suitable nesting sites. Climate change may alter the plant structure increasing the likelihood that the birds move further north (and out of Vermont to breed). Significant problems may occur on South American wintering grounds (mid-slope of Andes Mts)

Non-Habitat Threats:

Pollution
Trampling or Direct Impacts

Description of non-habitat threat(s): Atmospheric pollution, including airborne mercury, could impact the species directly, as well as damage its habitat. Although not well documented, collisions with glass buildings and wind towers can be a source of mortality for migrating, especially with birds that migrate at night

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Canada Warbler**
Scientific Name: **Cardellina canadensis**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S4B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

In Vermont the results from the second Breeding Bird Atlas indicate this bird is declining in Vermont. Block occupancy dropped 31% between the Atlases. Regional atlases and breeding bird surveys have also demonstrated a decline throughout the region over the past 30 years. As the climate and the forests change this species may be at a greater risk, not only on its breeding grounds, but also on its wintering grounds.

Distribution

Canada Warblers are found throughout Vermont. The Canada Warbler is considered a neotropical migrant and migrates from its breeding grounds to northern South America

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

A wide range of coniferous and deciduous forests, and mixed forests at all elevations, but especially mid-slopes in Green Mountains. Uses both mature and regenerating forest. Seem to prefer a dense understory with moss, and an uneven forest floor. Hummocks, roots, and debris are used to hide the nest and fledglings. Clearcuts and shelterwood cuts received more use than mature forest in northern New Hampshire. First appear in clearcuts 5 years after harvest, become common after 15 years and remain abundant until the next cutting cycle.



Common Name: **Canada Warbler**
Scientific Name: **Cardellina canadensis**
Species Group: **Bird**

Habitat Types:

Spruce Fir Northern Hardwood
Northern Hardwood
Softwood Swamps
Early Succession Boreal Conifers
Early Succession Spruce-Fir

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Climate Change
Habitat Fragmentation

Description of habitat threat(s): Forest succession, loss of forested wetlands, and development all may influence suitable nesting sites. Climate change may alter the plant structure increasing the likelihood that the birds move further north (and out of Vermont to breed). Significant problems may occur on South American wintering grounds (mid-slope of Andes Mts)

Non-Habitat Threats:

Pollution
Trampling or Direct Impacts

Description of non-habitat threat(s): Atmospheric pollution, including airborne mercury, could impact the species directly, as well as damage its habitat. Although not well documented, collisions with glass buildings and wind towers can be a source of mortality for migrating, especially with birds that migrate at night

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Canada Warbler**
 Scientific Name: **Cardellina canadensis**
 Species Group: **Bird**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	These are reasonably well known overall, but important to understand ecological and demographic differences in core populations that inhabit in prime habitats vs. smaller, more peripheral populations in patchy, secondary habitats
Research	Basic Life History	High	Nest success and productivity are poorly understood, as is age structure of populations in different habitat types. Need to understand demographics in secondary habitats (i.e. small patches) vs. those in core habitats
Research	Distribution and Abundance	Low	
Research	Threats and Their Significance	High	Need continued research on effects of forestry practices on populations in both prime and secondary habitats. Research on effects of climate change.
Research	Taxonomy	Low	Taxonomic research led to the recent name change.
Monitoring	Population Change	High	Need to ensure a long-term monitoring program that adequately samples this species, to clearly document declines or increases. Support of Vermont ebird for gathering data from citizen scientists important.
Monitoring	Habitat Change	High	Important to know how species responds to both natural and human-caused habitat changes
Monitoring	Range Shifts	Medium	Ability to shift range may be necessary if climate change has a big impact on Vermont forests
Monitoring	Monitor Threats	medium	

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Design and implement forest management strategies to enhance habitat suitability.	Area of potential habitat with long-range management plans which provide for beneficial forms of active forest management.	VFWD, USFS, USFWS	SWG, PR
Planning & Zoning	Medium	Conserve large tracts of core breeding habitats (mid-slope mixed forests, cedar swamps, red maple-conifer swamps).	Number of large forest tracts conserved via public ownership, easements, or town planning/zoning	ANR, USFS, USFWS, Town and RPCs	SWG, PR



Common Name: **Canada Warbler**
Scientific Name: **Cardellina canadensis**
Species Group: **Bird**

Bibliography

Conway, C.J. 1999. Canada Warbler. In *The Birds of North America*, No. 421 (A. Poole and F. Gill, eds.). The Birds of North America, Inc., Philadelphia, PA

Degraaf, R. M. and Yamasaki, M. 2001. *New England Wildlife: Habitat, Natural History, and Distribution*. University Press of New England, Hanover, NH, USA and London, UK.

eBird. 2012. eBird: An online database of bird distribution and abundance [web application]. eBird, Cornell Lab of Ornithology, Ithaca, New York. Available: <http://www.ebird.org>. (Accessed: Date April 2, 2015).

Faccio, S.D., C.C. Rimmer, and K.P. McFarland. 1997. Monitoring forest bird populations in Vermont: results of the Vermont forest bird monitoring program, 1989-1996. Report of the Vermont Institute of Natural Science, Woodstock, VT.

Lambert, J.D. and L. Reitsma. 2013. Canada Warbler (*Cardellina canadensis*). In: *The Second Atlas of Breeding Birds of Vermont*, (R. Renfrew, ed.). University Press of New England, Hanover and London

Miller, N.A. 1999. Landscape and habitat predictors of Canada Warbler (*Wilsonia canadensis*) and Northern waterthrush (*Seiurus noveboracensis*) occurrence in Rhode Island swamps. Masters thesis. University of Rhode Island, Kingston, R.I.

National Audubon Society. 2014. *Audubon's Birds and Climate Change Report: A Primer for Practitioners*. National Audubon Society, New York. Contributors: Gary Langham, Justin Schuetz, Candan Soykan, Chad Wilsey, Tom Auer, Geoff LeBaron, Connie Sanchez, Trish Distler. Version 1.2.

Reitsma, Len, Marissa Goodnow, Michael T. Hallworth and Courtney J. Conway. 2010. Canada Warbler (*Cardellina canadensis*). *The Birds of North America Online* (A. Poole, Ed.). Ithaca: Cornell Lab of Ornithology; Retrieved from the Birds of North America Online: <http://bna.birds.cornell.edu/bna/species/421>

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Eastern Towhee**
Scientific Name: **Pipilo erythrophthalmus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Species declining across region due to conversion of necessary early successional/shrub dominated habitats to either non-forest condition or via maturation of forest cover to an unsuitable forest age structure.

Distribution

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills		Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Early-successional/shrub/edge habitats, both mesic and xeric, characterized by dense shrub-small tree cover near ground and well-developed litter layer. Cover may be continuous or discontinuous patches interspersed w/in more open ground. Overstory trees may or may not be present, however open-canopied woodlands are favored over closed canopy coverage.

Habitat Types:

Early Succession Boreal Hardwoods
Early Succession Pine and Hemlock
Early Succession Northern Hardwoods
Shrub Swamps
Early Succession Upland Oak
Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Towhee**
 Scientific Name: **Pipilo erythrophthalmus**
 Species Group: **Bird**

Current Threats

Habitat Threats:

Conversion of Habitat
 Habitat Succession
 Habitat Alteration
 Inadequate Disturbance Regime
 Habitat Fragmentation

Description of habitat threat(s):

Description of non-habitat threat(s): Possible nest parasitism by cowbirds.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Basic Life History	High	Studies on reproductive success and demography especially desirable in northeastern U.S. To acquire baseline data via marked birds.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Conservation Finance	Medium	Create a state-funded, private lands, early successional habitat improvement initiative (modeled on NH's Small Landowner Grant program). Fund for > \$50,000/yr with revenues from state lands forest management. This could offset landowner EQIP obligations.	Level of funds raised.	FWD	SWG, PR
Awareness Raising and Communications	Medium	Initiate public education campaigns to highlight the need for active, even age forest management on public and private lands to create and maintain seedling/sapling forest habitat complexes.	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs.	VCE, VA, USFS	SWG, PR
Habitat Restoration	Medium	Determine appropriate old field habitat targets for state lands and restore and maintain old field habitats where needed to increase suitable ES songbird habitat.	Number of acres positively affected by management. Population response to management.	ANR, USFS, Audubon, Forest Products Association, VT Loggers	PR, EQIP



Common Name: **Eastern Towhee**
Scientific Name: **Pipilo erythrophthalmus**
Species Group: **Bird**

Bibliography

Greenlaw, J.S. 1996. Eastern Towhee (*Pipilo erythrophthalmus*). In: The Birds of North America, No. 262 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, D.C.

Dettmers, R.D. and K. V. Rosenberg. 2000. Partners in Flight Landbird Conservation Plan: Physiographic Area 9: Southern New England. Version 1.0.

Hodgman, T.P. and K. V. Rosenberg. 2000. Partners in Flight Landbird Conservation Plan: Physiographic Area 27: Northern New England. Version 1.0.

Rosenberg, K. V. 2004. Partners in Flight continental priorities and objectives defined at the state and bird conservation region levels, Vermont. Cornell Lab of Ornithology, Ithaca, NY. 26 p.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Field Sparrow**
Scientific Name: **Spizella pusilla**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Significant long-term population declines in Vermont of 5.3 percent annually 1966-2003, and 4.3 percent annually in the ten years following the first SWAP, 2004-2013 (Sauer et al. 2014). Atlas block occupancy declined by 39% between the first (1979-85) and second (2003-07) Vermont breeding bird atlases (Renfrew 2013).

Distribution

Widely distributed in eastern and western Vermont on either side of the Green Mountains, except in the Northeastern Highlands (Renfrew 2013).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Grasslands with scattered, shrubby vegetation with elevated perches. Habitat declines as woody encroachment progresses. Can be found in orchards and Christmas tree farms (Carey et al. 1994). Areas close to suburban development are avoided (Carey et al. 2008). In Vermont, often found in overgrown meadows dominated by juniper (Renfrew 2013).

Habitat Types:

Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession



Common Name: **Field Sparrow**
Scientific Name: **Spizella pusilla**
Species Group: **Bird**

Description of habitat threat(s): Primary problems to the species are likely due to succession of old fields and conversion of agricultural habitat to urban/suburban development.

Non-Habitat Threats:

Predation or Herbivory

Description of non-habitat threat(s): Parasitism is a possible limiting factor, more information needed.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	A better understanding of optimal stem densities and mowing rotations would inform specific management strategies. Better habitat-specific demographics would enable a more thorough understanding of when and why habitat decreases in quality for FISP.
Research	Threats and Their Significance	Medium	Species could be heavily parasitized by cowbirds. More intensive demographic data would elucidate BHCO limiting factor.
Monitoring	Population Change	High	Improved monitoring would elucidate population distribution and trends. A BBS-type survey route for early successional species could help monitor FISP, BWWA, GWWA, BRTH, PRAW, etc.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Field Sparrow**
 Scientific Name: **Spizella pusilla**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Conservation Finance	Medium	Create a state-funded, private lands, early successional habitat improvement initiative (modeled on NH's Small Landowner Grant program). Fund for > \$50,000/yr with revenues from state lands forest management. This could offset landowner WHIP obligations.	Level of funds raised.	VFWD	PR
Habitat Restoration	High	Stabilize declining population trend for Field Sparrows.	Population response to management, BBS surveys.	VFWD, NRCS, TNC.	NRCS
Awareness Raising and Communications	Medium	Initiate public education campaigns to highlight the need for active, even age forest management on public and private lands to create and maintain seedling/sapling forest habitat complexes	Number of media outlets reached, number of audiences reached, number of media products developed, number of participants in programs.		
Habitat Restoration	High	Determine appropriate old field habitat targets for state lands and restore and maintain old field habitats where needed to increase suitable ES songbird habitat.	Number of acres positively affected by management. Population response to management.	ANR, USFS, Audubon-VT, Forest Products Association, VT Loggers	PR, EQIP

Bibliography

- Carey, M., M. Carey, D. E. Burhans, and D. A. Nelson. 2008. Field Sparrow (*Spizella pusilla*). In *The birds of North America online* (A. Poole, Ed.). Cornell Lab of Ornithology, Ithaca, NY. Retrieved from <http://bna.birds.cornell.edu/bna/species/103>.
- Corwin, K. *The Second Atlas of Breeding Birds of Vermont*. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- LaBarr, M. L., Strong, A., Renfrew, R., Buck, J. and S. Parren. 2013. *The Vermont Grassland Bird Management and Recovery Plan*. Vermont Fish and Wildlife Department, Montpelier, Vt.
- Sauer, J. R., J. E. Hines, and J. Fallon. 2014. *The North American Breeding Bird Survey, Results and Analysis 1966-2003*. Version 2004.1. USGS Patuxent Wildlife Research Center, Laurel, Maryland, USA.



Common Name: **Vesper Sparrow**
Scientific Name: **Pooecetes gramineus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S3B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

In Vermont a 10.9 percent per year decline in 1966-2003, and losses of 7.2 percent annually in the decade since the first SWAP, 2004-2013 (Sauer et al. 2014). Vermont atlas block occupancy declined from 35 to 11 (69%) from the first (1979-85) to second (2003-07) atlas (Renfrew 2013). Also long-term decline survey-wide (Sauer et al. 2014). The generally small size of farming operations in VT seem as though they should create sufficient habitat to support a larger population in the state. The relative rarity of this species suggests that their habitat requirements may be somewhat more specialized than currently understood.

Distribution

An uncommon breeder in Vermont that is sparse and widely distributed. Most records are from the southwestern Champlain Valley (Renfrew 2013).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Unknown
Northern Green Mtns	Probable	Southern Green Mtns	Unknown
Northern VT Piedmont	Confident	Taconic Mtns	Unknown
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Breeds in dry, open habitats with short, sparse, and patchy herbaceous vegetation; some bare ground; and low to moderate shrub or tall forb cover. In the East, suitable habitats include reclaimed surface mines, crop and haylands, weedy roadsides, natural meadows, and grasslands (Jones and Cornely 2002). In Vermont, suitable habitat generally occurs in agricultural and other human-modified landscapes such as airports, and should be at least 20 hectares (Renfrew 2013).



Common Name: **Vesper Sparrow**
Scientific Name: **Pooecetes gramineus**
Species Group: **Bird**

Habitat Types:

Open Peatlands
Marshes and Sedge Meadows
Wet Shores
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops
Shrub Swamps

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Succession

Description of habitat threat(s): Early hay harvest and more intensive management of other row crops substantially reduces nesting success. Conversion of agricultural habitats to urban/suburban development also a problem. Old field succession and farm abandonment also decreasing habitat availability.

Description of non-habitat threat(s):

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Better information about precise habitat requirements, in particular nest site selection would be helpful for ascertaining potential habitat and developing management recommendations.
Research	Distribution and Abundance	High	Conduct focused surveys, including in areas where they were found during the second atlas, to obtain better information about population distribution in VT.
Monitoring	Population Change	High	Population monitoring, particularly in response to changing agricultural and development practices.
Monitoring	Habitat Change	Medium	Understanding habitat-specific demographic parameters would help us assess management options.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Vesper Sparrow**
 Scientific Name: **Pooecetes gramineus**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks	High	Educate agricultural community and general public about grassland birds and management options to protect habitat.	Develop a grassland bird outreach program	VFWD, Audubon-VT, VCE, UVM	SWG, PR
Market Forces	Medium	Enroll land into EQIP, GRP as well as FRPP programs to reduce the Impact of development on this species.		NRCS, VHCB	NRCS
Conservation Payments/Financial Incentives	Medium	Conserve grassland/shrubland habitats on private lands.	Number and total area of sites conserved.	USDA, USFWS, VHCB	FSA, SWG, VHCB

Bibliography

- DeGraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University of New England Press, Hanover, New Hampshire, USA, and London, UK.
- Jones, S. L., and J. E. Cornely. 2002. Vesper Sparrow (*Pooecetes gramineus*). In *The birds of North America*, No. 624 (A. Poole and F. Gill, eds.). The birds of North America, Inc., Philadelphia, Pennsylvania, USA.
- LaBarr, M. L., Strong, A., Renfrew, R., Buck, J. and S. Parren. 2013. The Vermont grassland bird management and recovery plan. Vermont Fish and Wildlife Department. Montpelier, Vt.
- Sauer, J. R., J. E. Hines, J. E. Fallon, K. L. Pardieck, D. J. Ziolkowski, Jr., and W. A. Link. 2014. The North American Breeding Bird Survey, Results and Analysis 1966 - 2012. Version 02.19.2014 USGS Patuxent Wildlife Research Center, Laurel, MD.
- Schwenk S. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- Vickery, P. D., M. L. Hunter, Jr., and S. M. Melvin. 1994. Effects of habitat area on the distribution of grassland birds of Maine. *Conservation Biology* 8:1087-1097.



Common Name: **Grasshopper Sparrow**
Scientific Name: ***Ammodramus savannarum***
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Currently listed as Threatened in Vermont. Species has declined throughout region due primarily to loss of grassland habitat and agricultural intensification (early mowing regimes). BBS data show a significant long-term (1966-2012; -2.86%/year) and short-term (2002-2012; -2.79%). The species is too rare to assess trends in Vermont; data from the second breeding bird atlas showed a 75% decline in number of blocks with breeding evidence, but the sample size is small (4 blocks to 1 block; Renfrew 2013). Only two or three locations in Vermont consistently support more than a few breeding pairs.

Distribution

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Unknown
Northern Green Mtns	Probable	Southern Green Mtns	Confident
Northern VT Piedmont	Unknown	Taconic Mtns	Unknown
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Grasslands, pastures, old fields and airports with minimal grass and litter cover and patches of bare ground. Specific habitat use patterns vary geographically (Vickery 1996). In most locations the species is area-sensitive, with occupancy significantly reduced in patches less than 30 ha (Vickery et al. 1994).

Habitat Types:

Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Other Cultural

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession



Common Name: **Grasshopper Sparrow**
Scientific Name: ***Ammodramus savannarum***
Species Group: **Bird**

Habitat Alteration

Habitat Fragmentation

Description of habitat threat(s): Direct loss of nesting habitat due to habitat conversion and agricultural intensification (mowing regimes)

Non-Habitat Threats:

Trampling or Direct Impacts

Description of non-habitat threat(s): Early and frequent mowing regimes directly impact nesting and reproductive success. Insufficient information on statewide population size.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Determine habitat requirements specific to Vermont
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Accurately determine population size and location of breeding pairs statewide
Research	Threats and Their Significance	High	Determine impacts of habitat loss and agricultural practices on distribution and nesting success.
Research	Population Genetics	Low	
Monitoring	Population Change	High	Accurately determine population size and trend information throughout the state and particularly at known nesting locations (airports).
Monitoring	Habitat Change	High	Determine statewide changes in grassland habitats and agricultural practices. Identify habitat changes at known nesting locations (airports)
Monitoring	Monitor Threats	High	Monitor limiting factors at current nesting locations (airports) including habitat loss due to development of the site and mowing practices.



Common Name: **Grasshopper Sparrow**
Scientific Name: ***Ammodramus savannarum***
Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks	High	Educate agricultural community and individuals with grasslands about grassland birds and management options.	Development of a grassland bird outreach program	VFWD, Audubon-VT, VINS, UVM	
Protected Area Management	High	Maintain nesting habitat throughout breeding season by developing site-specific conservation plans which include restricting field mowing until after July 15th on publicly owned lands (WMAs and state airports)	Maintain and increase current acreage under management on state lands	VFWD, Audubon-VT, NRCS, Vtrans	VFWD, Vtrans
Conservation Payments/Financial Incentives	High	Maintain nesting habitat throughout breeding season by restricting field mowing until after July 15th	Increase protection of available habitat through enrollment in EQIP and GRP	VFWD, Audubon-VT, NRCS	
Habitat Restoration	High	Maintain grassland habitat in suitable locations through active management of woody vegetation within Grassland Bird Focus Areas.	Increase and maintain available habitat in suitable locations	VFWD, private landowners	
Conservation Payments/Financial Incentives	High	Protect privately owned known nesting sites and suitable grassland habitat from development and agricultural intensification by creating Grassland Bird Focus Areas to concentrate management efforts (see the Vermont Grassland Bird Management Plan).	Development of Grassland Bird Focus Areas and increase protection of available habitat through enrollment in EQIP and GRP.	VFWD, Audubon-VT, NRCS, private landowners	USFWS
Conservation Payments/Financial Incentives	High	Maintain large tracts (> 100 acres) of suitable grassland habitat for entire suite of grassland bird species.	Increase protection of available habitat through enrollment in WHIP and GRP	VFWD, Audubon-VT, NRCS	

Bibliography

Vickery, P. D., M. L. Hunter, Jr., and S. M. Melvin. 1994. Effect of habitat area on the distribution of grassland birds in Maine. *Conservation Biology* 8:1087-1097.

Vickery, P.D. 1996 Grasshopper Sparrow. In A. Poole and F. Gill Editors. *The birds of North America*, no. 580 Academy of natural Sciences, Philadelphia PA, and American Ornithologists' Union, Washington, D.C., USA.

LaBarr, M. L., Strong, A., Renfrew, R., Buck, J. and S. Parren. 2013. The Vermont grassland bird management and recovery plan. Vermont Fish and Wildlife Department. Montpelier, Vt



Common Name: **Bobolink**
Scientific Name: **Dolichonyx oryzivorus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Significant long-term population declines both in VT and survey-wide. Although Vermont atlas block occupancy changed little (Renfrew 2013), abundance declined by 2.6 percent annually from 1966 to 2003, and 2.4 percent per year since the first SWAP, 2004-2013 (Sauer et al. 2014). Much of VT hayed grasslands are likely population sinks. Main threats are loss and degradation of quality habitat, including fragmentation, due to field succession and conversion to development after farms are lost, and intensive management of hay fields (more frequent mowing).

Distribution

Distributed throughout Vermont, most abundant in lowlands of Champlain Valley, less so in other areas of the state in open landscapes that include fields not under intensive rowcrop production.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Grasslands, primarily managed for hay or to a lesser extent, low-intensity grazing. Generally avoids alfalfa, row crops, and grass habitats with standing water. More common in larger (> 5ha), more blocky (as opposed to linear) fields, and in relatively less forested landscapes with large expanses of grassland habitat. Social attraction also plays a role in habitat selection.

Habitat Types:

Marshes and Sedge Meadows

Grasslands, Hedgerows, Old Field, Shrub, or Orchard



Common Name: **Bobolink**
Scientific Name: **Dolichonyx oryzivorus**
Species Group: **Bird**

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development
Habitat Succession
Habitat Alteration
Habitat Fragmentation
Invasion by Exotic Species
Climate Change

Description of habitat threat(s): Greatest problems are the frequency and timing of hayfield mowing, the succession of agricultural land, and the conversion of agricultural land to development. Fragmentation of grasslands limits patch size and openness of landscape, which is important for nesting Bobolinks. Bobolink distribution expected to shift northward due to climate change, although models need to be refined. Takeover of hayfields and other grasslands by exotic plants such as parsnip and chervil renders habitat unsuitable

Non-Habitat Threats:

Pollution
Predation or Herbivory
Trampling or Direct Impacts

Description of non-habitat threat(s): Pesticides on migration and wintering grounds, predation of eggs and nestlings, direct mortality from mowing

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Threats and Their Significance	Medium	1) Demographic model of climate change impacts to predict future distribution. 2) Determine relative contribution of seasonal survival of juveniles and adults, and immigration/emigration, to improve assessments of relative importance of productivity and survival in determining population size
Research	Other Research	High	Determine most effective use of resources to maximize acreage of quality nesting habitat using combination of approaches for different types of landowners and interests.
Monitoring	Population Change	High	Determine if the Champlain Valley is a source or sink for Bobolink.
Monitoring	Habitat Change	High	Better information is necessary regarding the timing of hay mowing in landscapes with various proportions of agriculture throughout VT.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Bobolink**
 Scientific Name: **Dolichonyx oryzivorus**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Easements	Medium	Reduce the amount of grassland habitat being lost to development through strategic acquisition of grassland easements.	Grassland acreage enrolled in easements	NRCS	NRCS, USDA.
Conservation Payments/Financial Incentives	High	Decrease nest losses due to early mowing regimes on fields used for animal forage via EQIP conservation payments. Continue outreach to landowners about incentive programs	Increase in proportion and total area of grasslands in which hay cutting is delayed.	NRCS, UVM, VCE, Audubon-VT	NRCS, USDA
Species Restoration	High	Implement the Vermont grassland bird management and recovery plan (LaBarr et al. 2013)		VFWD, UVM, TNC, NRCS.	SWG, PR, NRCS
Technical Assistance, Training, Learning Networks	High	Improve outreach to (and exchange of information among) landowners with flexibility (e.g., those primarily interested in preventing succession).	Number of acres under a late-mowing management regime, number of landowners contacted	UVM, NRCS, Audubon-VT, VCE	NRCS, USDA

Bibliography

- Martin, S. G., and T. A. Gavin. 1995. Bobolink (*Dolichonyx oryzivorus*). In *The birds of North America*, No. 176 (A. Poole and F. Gill, eds.). The birds of North America, Inc., Philadelphia, Pennsylvania, USA.
- Sauer, J. R., J. E. Hines, J. E. Fallon, K. L. Pardieck, D. J. Ziolkowski, Jr., and W. A. Link. 2014. *The North American Breeding Bird Survey, Results and Analysis 1966 - 2012*. Version 02.19.2014 USGS Patuxent Wildlife Research Center, Laurel, MD.
- Renfrew, R.. 2013. *The Second Atlas of Breeding Birds of Vermont*. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.
- LaBarr, M. L., Strong, A., Renfrew, R., Buck, J. and S. Parren. 2013. *The Vermont Grassland Bird Management and Recovery Plan*. Vermont Fish and Wildlife Department, Montpelier, Vt.



Common Name: **Eastern Meadowlark**
Scientific Name: **Sturnella magna**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S5B

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Upgraded from medium to high priority. One of the most severely declining population trends of grassland bird species throughout its range. In Vermont Eastern Meadowlark populations have declined by 9.6 percent annually from 1966 to 2003 (Sauer et al. 2014), and since the first SWAP, meadowlarks have disappeared from much of Vermont except in the Champlain Valley, which supports most of the remaining population. Between the first and second Vermont Breeding Bird atlas, the species was lost from 63 of 155 (55%) blocks (Renfrew 2013). Regrowth of abandoned farmlands and agricultural intensification resulting in grassland habitat loss, fragmentation, and degradation are the primary causes of declines.

Distribution

Sparsely distributed in relatively large open agricultural (or airfield) areas throughout much of the state, except in the Champlain and Vermont valleys, where it is fairly common in open, agricultural areas with suitable nesting habitat. Nearly absent from southeastern Vermont (Renfrew 2013).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

Meadows, old fields, hayfields with thick layer of dead grass. Requires large, open landscapes, large patches of grasslands (>10ha). Can occur at airports with compatible mowing program.

Habitat Types:

Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Other Cultural



Common Name: **Eastern Meadowlark**
Scientific Name: **Sturnella magna**
Species Group: **Bird**

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development
Habitat Succession
Habitat Alteration
Habitat Fragmentation
Invasion by Exotic Species

Description of habitat threat(s): Loss and degradation of habitat due to frequent mowing of hayfields, habitat loss due to succession of farmland to forest, conversion of grassland habitat to development and potentially, solar panel arrays. Takeover of hayfields and other grasslands by exotic plants such as parsnip and chervil renders habitat unsuitable.

Non-Habitat Threats:

Trampling or Direct Impacts

Description of non-habitat threat(s): Direct mortality due to mowing.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Other Research	Medium	Determine whether Vermont habitats in the Champlain Valley are sources or sinks
Monitoring	Population Change	High	Species no longer tracked well with BBS methods. Carry out more intensive, standardized monitoring scheme to track population status in the state and determine important breeding areas and compatible management practices.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Meadowlark**
 Scientific Name: **Sturnella magna**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Conservation Payments/Financial Incentives	High	Maintain nesting habitat by delaying mowing until after July 15th	Increased protection of habitat through enrollment in EQIP and GRP	VFWD, Audubon VT, NRCS	
Technical Assistance, Training, Learning Networks	High	Educate agricultural community, landowners, and general public about grassland birds and management options to protect habitat	Continue grassland bird outreach and networking programs	VFWD, Audubon, VCE, UVM	SWG, PR
Protected Area Management	High	Maintain nesting habitat throughout the breeding season by developing site specific conservation plans which include restricting mowing until after July 15 on publicly owned lands (WMAs, state airports).	Maintain and increase current acreage under management on state and federal lands	VFWD, Audubon-VT, VCE, USFWS NRCS Vtrans	SWG, NRCS
Habitat Restoration	High	Maintain grassland habitat in suitable locations through active management of woody vegetation within focal grassland areas.	Increase and maintain available habitat in suitable locations	VFWD, Audubon-VT, VCE, NRCS, USFWS	USFWS, NRCS
Conservation Finance	High	Decrease nest losses due to early mowing regimes on fields used for animal forage via EQIP conservation payments. Continue outreach to landowners about incentive programs.	Increase in acreage in which hay cutting is delayed.	NRCS, UVM, VCE, Audubon-VT	NRCS
Conservation Finance	High	Focus efforts on relatively large fields (>50 acres) of suitable grassland habitat in open landscapes	Strategic enrollment in EQIP	NRCS	NRCS

Bibliography

LaBarr, M. L., Strong, A., Renfrew, R., Buck, J. and S. Parren. 2013. The Vermont Grassland Bird Management and Recovery Plan. Vermont Fish and Wildlife Department, Montpelier, Vt.

Puryear, K. 2004. Landscape-level Grassland Bird conservation in the southern Champlain Valley, Vermont. Masters Project. University of Vermont, Burlington Vermont, USA.

Sauer, J. R., J. E. Hines, J. E. Fallon, K. L. Pardieck, D. J. Ziolkowski, Jr., and W. A. Link. 2014. The North American Breeding Bird Survey, Results and Analysis 1966 - 2012. Version 02.19.2014 USGS Patuxent Wildlife Research Center, Laurel, MD.

Renfrew, R. 2013. The Second Atlas of Breeding Birds of Vermont. R. Renfrew, ed. University Press of New England, Hanover and London. 548pp.



Common Name: **Rusty Blackbird**
Scientific Name: **Euphagus carolinus**
Species Group: **Bird**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S3B

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The continental Rusty Blackbird population has undergone a precipitous decline since the 1966 beginning of the Breeding Bird Survey (BBS) and most significantly in the last 20 years. In eastern North America its decline has been most significant at the southern edge of its breeding range (Maine, New Hampshire, and Vermont). From BBS estimates, Partners in Flight (PIF) estimates there are 5,000,000 Rusty Blackbirds in North America with 20% of them residing in the US. PIF categorizes this estimate as a medium quality. This is likely due to the species' preference for remote boreal coniferous forests near water's edge. The Second Atlas of Breeding Birds of Vermont (2002-07) reported a 26% decline in occupied blocks. The majority of the decline occurred within the Northern Vermont Piedmont biophysical region. In addition to the significant loss of population there is concern among biologists that climate change projections estimate a decline in boreal forest that includes the blackbird's New England habitat. In light of these foreboding estimates Vermont listed the species as endangered in 2014.

Distribution

"Rusty Blackbirds are local and uncommon summer residents of the Northeast Highlands, the North Central region, and the Green Mountains." (Nichols 1985)

Distribution by Biophysical Region:

Champlain Valley	Probable	Southern VT Piedmont	Probable
Champlain Hills	Probable	Vermont Valley	Probable
Northern Green Mtns	Probable	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Wooded swamps, tree-bordered marshes, beaver ponds, boreal bogs and stream borders with alder and willow thickets (DeGraff and Rudis 1986). "Disturbance can be favorable to this species; e.g., nests found in modest openings regenerating from clearcuts (Ellison 1990)" (Avery 1995).

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Rusty Blackbird**
Scientific Name: **Euphagus carolinus**
Species Group: **Bird**

Habitat Types:

Hardwood Swamps
Softwood Swamps
Early Succession Boreal Conifers
Early Succession Northern Hardwoods
Shrub Swamps
Early Succession Spruce-Fir

Current Threats

Habitat Threats:

Conversion of Habitat

Description of habitat threat(s): Permanent residence and/or vacation home development on lakeshores/pondshores may reduce available habitat.

Non-Habitat Threats:

Harvest or Collection

Description of non-habitat threat(s): "Substantial mortality to local populations may occur when Rusty Blackbirds are in mixed-species winter roosts subjected to blackbird control in the s. U.S.(Stickley et al. 1986)" (Avery 1995).

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Basic Life History	Low	Some evidence of colonial nesting, however nesting by widely-separated individuals seems to prevail in Vermont. Factors governing how habitat might influence whether Rusty Blackbirds nest singly or colonially should be investigated (Avery 1995).
Research	Distribution and Abundance	Medium	More complete surveys of the distribution of breeding Rusty Blackbirds in Vermont are warranted to obtain a better estimate of its true status.
Research	Threats and Their Significance	Low	
Monitoring	Population Change	Medium	
Monitoring	Habitat Change	Low	
Monitoring	Monitor Threats	Medium	Shoreline development in the Rusty Blackbird strongholds in Vermont should be monitored.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Rusty Blackbird**
 Scientific Name: **Euphagus carolinus**
 Species Group: **Bird**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Easements	Medium	Known nesting habitats should be monitored over time to track impacts from development. Easements should be considered to protect important breeding habitats from development. PIF Vermont target population is 226 breeding individuals.	Number of sites identified and conserved.	VFWD, VCE, VA, VHCB	SWG, VHCB
Policy & Regulations	Low	Assist PIF with efforts to reduce mortality from pesticides used on wintering grounds.	Reduction in mortalities due to pesticides	PIF, USFWS	USFWS, USDA, SWG

Bibliography

Avery, M. L. 1995. Rusty Blackbird (*Euphagus carolinus*). In *The Birds of North America*, No. 200 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, and The American Ornithologist's Union, Washington, D.C.

Degraaf, R.M. and D.D. Rudis 1986. New England wildlife: habitat, natural history, and distribution. Gen. Tech. Rep. NE-108. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station; 1986. 491 p.

Ellison, W.G. 1990. The status and habitat of the Rusty Blackbird in Caledonia and Essex counties. Vermont Fish Wildl. Dep., Waterbury, VT.

Nichols, W. D. 1985. Rusty Blackbird. Pp. 364-365 in *The atlas of breeding birds of Vermont* (S.B. Laughlin and D.P. Kibbe, eds.). Univ. Press of New England, Hanover, NH.

Fisher, C. and L.L. Powell. 2013. *Second Atlas of Breeding Birds of Vermont*. 2013. Rosalind Renfrew, ed. 547 pp. University Press of New England.

Vermont's Wildlife Action Plan

Fish SGCN Conservation Reports

Appendix A3

Draft 10/1/2015

Species	Page
Northern Brook Lamprey	1
Silver Lamprey	6
American Brook Lamprey	11
Mottled Sculpin	16
Lake Sturgeon.....	19
American Eel (Lake Champlain population)	24
American Eel (CT River population)	28
Blueback Herring (CT River population).....	33
American Shad	36
Mooneye	41
Cisco or Lake Herring	44
Lake Whitefish	48
Round Whitefish.....	53
Atlantic Salmon (naturally reproducing populations in Lakes Champlain & Memphremagog).....	58
Brook Trout (naturally reproducing populations).....	63
Lake Trout (naturally reproducing populations)	66
Redfin Pickerel	71
Bridle Shiner	76
Blackchin Shiner	81
Blacknose Shiner	85
Silver Redhorse	89
Shorthead Redhorse	94
Greater Redhorse	99
Stonecat.....	104
Redbreast Sunfish	109
Eastern Sand Darter	113
Channel Darter.....	118
Sauger	122

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Northern Brook Lamprey**
Scientific Name: **Ichthyomyzon fossor**
Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend: Unknown

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Only two sub-populations of Northern Brook Lamprey are known to occur in Vermont, i.e. the Malletts Creek-Indian Brook watershed (Langdon 2014). On a range-wide scale no major threats are known, but the species is vulnerable to local extirpation through indiscriminant use of fish toxicants (COSEWIC 2007; NatureServe 2014). Li et al. (2014) investigated the status of native lamprey populations in the Great Lakes drainage basin where recent declines have been observed and attributed to habitat loss and degradation, anthropogenic stresses, and stream treatment with lampricides. They report that of six species historically found in Pennsylvania streams, current surveys confirmed the presence of four species and absence of two species, including Northern Brook Lamprey, in streams previously known to support them possibly indicating extirpation. The use of lampricides to control Sea Lamprey in the Lake Champlain basin probably represents the greatest threat to Vermont populations. As of yet neither population has been exposed to chemical treatment; however, the U. S. Fish and Wildlife Service recently made it known that it wishes to treat Malletts Creek (Langdon 2015). Northern Brook Lamprey is currently listed as endangered in Vermont, Pennsylvania, Ohio, Indiana and Illinois; and special concern in Minnesota, Ontario and Quebec. Recent research into the genetic relationship between Northern Brook Lamprey and Silver Lamprey seems to indicate that the two species may be ecotypes of one species (Docker et al. 2012), even though each is considered a distinct species in all taxonomic accounts (COSEWIC 2011) and that this convention should remain in place until such time that there is "strong evidence [that] rejects the hypothesis that parasitic and nonparasitic members of a paired species represents distinct species" (Renaud et al. 2009 cited in COSEWIC 2011). Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

Northern brook lamprey occur in the Mississippi and Great Lakes drainages from western New York, Quebec, Ontario, Michigan and eastern Wisconsin; from the north shore of Lake Superior to northern Indiana and Ohio (Scott and Crossman 1973). Only one northern brook lamprey population is known in Vermont. This population is limited to one watershed consisting of Indian Brook and Malletts Creek, both tributaries of Malletts Bay on Lake Champlain.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lamoille River



Common Name: **Northern Brook Lamprey**
Scientific Name: **Ichthyomyzon fossor**
Species Group: **Fish**

Lake Champlain
Missiquoi River
Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Clear streams and small to moderate sized rivers (Scott and Crossman 1973; Smith 1985; Langdon et al, In press; Morman 1979). The non-parasitic northern brook lamprey spends its entire life in streams. No known populations reside on Lake Champlain's deltas, unlike sea lamprey and American brook lamprey. It has a 4 to 6 year life span and spends most of it buried in the substrate, where it grows to about 150mm, metamorphoses, and becomes sexually mature. Adults generally migrate upstream to spawn. Larval habitat includes lotic depositional and estuarine areas of streams with organic matter for feeding and fairly stable substrate in order to maintain burrow. Stream riffle areas with sand and pea gravel up to approximately 15 mm diameter are used for spawning (W. Bouffard, USFWS, personal communication). Spawning water depth is typically <450 mm deep (Scott and Crossman 1998).

Habitat Types:

Aquatic: Lacustrine
Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Habitat Alteration
Sedimentation
Habitat Fragmentation

Description of habitat threat(s): Culverts at road crossings may fragment habitat by creating barriers to the upstream migration of spawning adult northern brook lamprey. Sedimentation of stream bottoms reduces the quality of spawning habitat. The lower reaches of Malletts Creek and Indian Brook, which converge to form a large wetland complex, may reach the lethal temperature limit of 30.5 °C thereby limiting population distribution in those streams.

Non-Habitat Threats:

Genetics
Competition
Pollution

Description of non-habitat threat(s): Due to the small population size, distance from other regional northern brook lamprey populations, and absence of gene flow between populations, there is a potential for inbreeding depression to occur naturally within Vermont's population. No diagnostic characteristics exist for differentiating between northern brook and silver lamprey accomocoetes. These two species were not readily distinguishable using mitochondrial sequence and mitochondrial restriction fragment length polymorphism analyses, raising the question whether they are different species or different ecomorphotypes



Common Name: Northern Brook Lamprey
Scientific Name: *Ichthyomyzon fossor*
Species Group: Fish

(Mandrak et al. 2004). This presents difficulties in understanding current distributions and population trends for northern brook lamprey in the Lake Champlain basin. Competition for larval habitat with sea lamprey and silver lamprey could reduce survival and/or fitness of northern brook lamprey larvae. Small population sizes and extremely limited geographic distribution increase the vulnerability of the population to a potential pollution event. The Lake Champlain Fish and Wildlife Management Cooperative is currently involved in a sea lamprey control program that includes the use of lampricides to kill stream-resident sea lamprey larvae. These lampricides are toxic to all species of lamprey. Currently, sea lamprey are being controlled in streams with northern brook lamprey through the use of traps, eliminating the need for lampricides. The repeated use of lampricides has been shown to adversely affect populations of northern brook lamprey in Lake Superior tributaries (Schuldt and Goold 1980).

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Given difficulties in identifying Northern Brook Lamprey from other <i>Ichthyomyzon</i> spp. and apparent low population densities, efforts to survey other streams having suitable habitat to locate potential new populations and better define current distribution
Research	Threats and Their Significance	High	Threats and their significance are poorly understood for this species.
Research	Population Genetics	Medium	Genetic similarities between the disjunct Vermont population and other nearest populations within the region have not been investigated. Investigate gene flow within and between Northern Brook Lamprey populations in the Lake Champlain basin.
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	Current estimates are needed to ascertain status of known population.
Monitoring	Habitat Change	High	Habitat within species' limited distribution in state is not being monitored.
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Northern Brook Lamprey**
 Scientific Name: **Ichthyomyzon fossor**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Natural Processes Restoration	Medium	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, VTrans, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners, VFWD (SWG), USFWS, VDEC, VTrans
Invasive Species Control & Prevention	High	Manage potential non-target impacts of the Lake Champlain Sea Lamprey control program on Northern Brook Lamprey populations.	Number of existing populations of Northern Brook Lamprey protected and sustained.	LCFWMC, USFWS, VDEC	VFWD (DJ, SWG)
Research	Low	Assess, monitor and manage as appropriate the potential for Sea Lamprey competition with Northern Brook Lamprey for spawning and juvenile habitats.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM
Research	High	Current research indicates Northern Brook Lamprey and Silver Lamprey may be ecotypes of a single species. Both ecotypes should be conserved.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM
Natural Processes Restoration	Medium	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)



Common Name: **Northern Brook Lamprey**
Scientific Name: **Ichthyomyzon fossor**
Species Group: **Fish**

Bibliography

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2007. COSEWIC assessment and update status report on the northern brook lamprey *Ichthyomyzon fossor* (Great Lakes – Upper St. Lawrence populations and Saskatchewan – Nelson population) in Canada. COSEWIC, Ottawa. Available at: www.sararegistry.gc.ca/status/status_e.cfm. (December 2014).

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2011. COSEWIC assessment and status report on the silver lamprey *Ichthyomyzon unicuspis* (Great Lakes – Upper St. Lawrence populations and Saskatchewan – Nelson Rivers populations) in Canada. COSEWIC, Ottawa. Available at: www.sararegistry.gc.ca/status/status_e.cfm. (December 2014).

Docker, M. F., N. E. Mandrak, and D. D. Heath. 2012. Contemporary gene flow between “paired” silver (*Ichthyomyzon unicuspis*) and northern brook (*I. fossor*) lampreys: implications for conservation. *Conservation Genetics* 13:823-835.

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.

Li S., K-M. Werner, and J. R. Stauffer, Jr. 2014. An examination of Petromyzontidae in Pennsylvania: current distribution and habitat preferences of lampreys. *Northeastern Naturalist* 21(4):606-618.

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).

Renaud, C. B., M. F. Docker, and N. E. Mandrak. 2009. Taxonomy, distribution, and conservation of lampreys in Canada. Pages 293-309 in L. R. Brown et al., editors. *American Fisheries Society, Symposium 72*, Bethesda, Maryland.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Silver Lamprey**
Scientific Name: **Ichthyomyzon unicuspis**
Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S2?

State Trend: Stable

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

In Vermont, Silver Lamprey is widely distributed in Lake Champlain and its tributaries. The use of lampricides to control Sea Lamprey in the Lake Champlain basin probably represents the greatest threat to Vermont populations. Other threats to the species include construction of barriers that prevent access to spawning habitats, water pollution, habitat alteration, siltation, water level fluctuation and competition with introduced species (COSEWIC 2011). In conjunction with the program to control Sea Lamprey abundance in Lake Champlain the USFWS conducts quantitative assessment sampling (QAS) typically every four years to estimate Sea Lamprey and Silver Lamprey abundance in seven Vermont treatment rivers: Missisquoi River, Stonebridge Brook, Lamoille River, Winooski River, Lewis Creek, Poultney River, and Hubbardton River. Abundance trends based on QAS (raw data provided by Allaire 2015) indicate declining populations in the Winooski and Poultney rivers; and no discernible trends (decreasing or increasing) in the other streams where multiple treatments have been conducted. However, needs to be pointed out that QAS estimates typically have very large confidence intervals (A. Allaire, USFWS, personal communication), so these results must be interpreted with caution. Recent research into the genetic relationship between Silver Lamprey and Northern Brook Lamprey seems to indicate that the two species may be ecotypes of one species (Docker et al. 2012), even though each is considered a distinct species in all taxonomic accounts (COSEWIC 2011) and that this convention should remain in place until such time that there is “strong evidence [that] rejects the hypothesis that parasitic and nonparasitic members of a paired species represents distinct species” (Renaud et al. 2009 cited in COSEWIC 2011). Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

Silver lamprey are restricted to eastern North America, from the St. Lawrence River as far down as Montmagny, Quebec, west through the Great Lakes, through the upper Mississippi valley from Wisconsin, to eastern Manitoba; from Manitoba tributaries of Hudson Bay in the north to the Ohio River basin as far south as Kentucky (Scott and Crossman 1973). In Vermont, this species is at the eastern edge of its North American range. Vermont populations are limited to Lake Champlain and the lower reaches of several tributaries up to the fall line.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Mettawee River

Probable Watersheds

Otter Creek



Common Name: **Silver Lamprey**
Scientific Name: **Ichthyomyzon unicuspis**
Species Group: **Fish**

Lake Champlain
Lamoille River
Missiquoi River
Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Generally occurs in large streams and lakes where host fish are present and can be parasitized. Larval habitat includes lotic depositional and estuarine areas of streams with organic matter for feeding and fairly stable substrate required to maintain burrows. Spawning occurs in riffle areas with sand and gravel up to approximately 30 mm diameter (W. Bouffard, U. S. Fish and Wildlife Service, personal communication). In Vermont, the silver lamprey spawns in the lower sections of several tributaries of Lake Champlain (Langdon et al. In press).

Habitat Types:

Aquatic: Lacustrine
Aquatic: Fluvial
Aquatic: Lower CT River
Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Habitat Alteration
Sedimentation
Habitat Fragmentation

Description of habitat threat(s): Culverts and dams may constitute barriers to the upstream migration of spawning adults. Sedimentation of stream bottoms reduces the quality of spawning habitat.

Description of non-habitat threat(s): The Lake Champlain Fish and Wildlife Management Cooperative is currently involved in a sea lamprey control program that includes the use of lampricides to kill stream-resident larvae. These lampricides are toxic to all species of lamprey.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Silver Lamprey**
Scientific Name: **Ichthyomyzon unicuspis**
Species Group: **Fish**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	
Research	Threats and Their Significance	Medium	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	Population monitoring to determine current status and changes.
Monitoring	Habitat Change	Medium	Habitat assessment and monitoring to assess habitat change and identify limiting factors.
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Silver Lamprey**
 Scientific Name: **Ichthyomyzon unicuspis**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Invasive Species Control & Prevention	High	Manage potential non-target impacts of the Lake Champlain Sea Lamprey control program on Silver Lamprey populations.	Number of existing populations of Silver Lamprey protected and sustained.	LCFWMC, USFWS, VDEC	VFWD (DJ, SWG)
Natural Processes Restoration	Medium	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners
Research	High	Current research indicates Northern Brook Lamprey and Silver Lamprey may be ecotypes of a single species. Both ecotypes should be conserved.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM
Natural Processes Restoration	Medium	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, VTrans, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners, VFWD (SWG), USFWS, VDEC, VTrans
Invasive Species Control & Prevention	Low	Assess, monitor and manage as appropriate the potential for Sea Lamprey competition with Silver Lamprey for spawning and juvenile habitats.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Silver Lamprey**
 Scientific Name: **Ichthyomyzon unicuspis**
 Species Group: **Fish**

Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

Bibliography

Allaire, A. 2015. Personal communication. U.S. Fish and Wildlife Service, Essex.

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2011. COSEWIC assessment and status report on the silver lamprey *Ichthyomyzon unicuspis* (Great Lakes – Upper St. Lawrence populations and Saskatchewan – Nelson Rivers populations) in Canada. COSEWIC, Ottawa. Available at: www.sararegistry.gc.ca/status/status_e.cfm. (December 2014).

Docker, M. F., N. E. Mandrak, and D. D. Heath. 2012. Contemporary gene flow between “paired” silver (*Ichthyomyzon unicuspis*) and northern brook (*I. fossor*) lampreys: implications for conservation. *Conservation Genetics* 13:823-835.

Renaud, C. B., M. F. Docker, and N. E. Mandrak. 2009. Taxonomy, distribution, and conservation of lampreys in Canada. Pages 293-309 in L. R. Brown et al., editors. *American Fisheries Society, Symposium 72*, Bethesda, Maryland.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Brook Lamprey**
Scientific Name: **Lethenteron appendix**
Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend: Unknown

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Populations of American Brook Lamprey are currently known to occur in seven Vermont streams (Langdon 2014), all located in the Lake Champlain Valley biophysical region. NatureServe (2014) identifies no known major threats to the species at the range-wide scale. Nonetheless, all lamprey species are susceptible to mortality caused by the use of lampricides; as such, chemical control of Sea Lamprey in the Lake Champlain basin probably represents the greatest threat to the state's American Brook Lamprey populations. Two brook lamprey populations that have been exposed to lampricides are Trout Brook in Milton (treated in 1995) and Winooski River in Winooski (treated in 2004). Brook lamprey mortalities were confirmed to have occurred as a result of treatments in both streams (B. Chipman, Vermont Fish & Wildlife Department, personal communication). The coastal population in New Hampshire is listed as endangered and those in Massachusetts and Maryland are threatened. Threats to these populations are generally identified as sedimentation, water temperature increases, pollutants, extreme water level changes, and habitat fragmentation often associated with land development activities (MDNR; MDFW 2005; NHFGD 2005). Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

American brook lamprey is restricted to east-central North America: on the eastern seaboard from southeastern New Hampshire south to Maryland; and in the St. Lawrence River and tributaries from Montmagny southwest through the Great Lakes, west to southeastern Minnesota, south into the Mississippi River drainage to Tennessee and Missouri (Scarola 1973; Scott and Crossman 1973). In Vermont, it is known to occur in the Missisquoi River and tributaries, Hungerford and Kelly brooks; Winooski River. and its tributary, Sunderland Brook; Youngman Brook; and Trout Brook.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain
Missisquoi River
Winooski River



Common Name: **American Brook Lamprey**
Scientific Name: **Lethenteron appendix**
Species Group: **Fish**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Clear, cool streams and small rivers characterized by a large proportion of groundwater inflow (Scott and Crossman 1973; Smith 1985; Langdon et al, in press; Morman 1979). The non-parasitic American brook lamprey spends its entire life in streams and infrequently migrates to lakes, except to reside on some Lake Champlain river deltas particularly in New York, i.e. Ausable and Salmon rivers (Langdon et al. in press). It has a 4 to 6 year life span and spends most of it buried in the substrate, where it grows to about 200mm, metamorphoses, and becomes sexually mature. Adults generally migrate upstream to spawn. American brook lamprey spawn the earliest and initiate spawning in the coolest water (~7°C)(Hardisty and Potter 1971) compared to other species of lamprey in the Lake Champlain Basin. Larval habitat consists of lotic depositional, estuarine, and lentic areas of streams, with organic matter for feeding and fairly stable substrate in order to maintain burrow. Spawning occurs in riffle areas with sand and pea gravel up to approximately 20 mm diameter (Manion and Hanson 1980; W. Bouffard, USFWS, personal communication). Adults require certain physical factors for successful spawning such as suitable substrate (pea gravel), water velocities (0.3-0.5 m/s), and temperatures (mean 14°C, range 6.7-20.6°C) (Manion and Hanson 1980; Hardisty and Potter 1971; Morman 1979).

Current Threats

Habitat Threats:

Habitat Alteration

Sedimentation

Habitat Fragmentation

Description of habitat threat(s): Culverts at road crossings and dams may fragment habitat by creating barriers to upstream spawning migration of adult brook lamprey. Sedimentation of stream bottoms reduces the quality of spawning habitat.

Non-Habitat Threats:

Competition

Description of non-habitat threat(s): Competition for larval habitat with sea lamprey and silver lamprey may reduce survival and/or fitness of American brook lamprey larvae (Scott and Crossman 1998). The Lake Champlain Fish and Wildlife Management Cooperative is currently involved in a sea lamprey control program that includes the use of lampricides to kill stream-resident larvae. These lampricides are toxic to all species of lamprey. In most streams where American brook lamprey and sea lamprey co-occur, traps are used to remove adult sea lamprey before they spawn, eliminating the need to use lampricides. The repeated use of lampricides has been shown to adversely affect populations of American brook lamprey in Lake Superior tributaries (Schuldt and Goold 1980).

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Brook Lamprey**
Scientific Name: **Lethenteron appendix**
Species Group: **Fish**

Research and Monitoring Needs

<i>Type</i>	<i>Need</i>	<i>Priority</i>	<i>Description</i>
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	
Research	Threats and Their Significance	High	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	Population monitoring to determine current status and changes.
Monitoring	Habitat Change	High	Habitat assessment and monitoring to assess habitat change and identify limiting factors.
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	Monitor known limiting factors to populations and their habitats.
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Brook Lamprey**
 Scientific Name: **Lethenteron appendix**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Natural Processes Restoration	Medium	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners
Invasive Species Control & Prevention	High	Manage potential non-target impacts of the Lake Champlain Sea Lamprey control program on American Brook Lamprey populations.	Number of existing populations of American Brook Lamprey protected and sustained.	LCFWMC, USFWS, VDEC	VFWD (DJ, SWG)
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Brook Lamprey**
 Scientific Name: **Lethenteron appendix**
 Species Group: **Fish**

Natural Processes Restoration	Medium	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, VTrans, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners, VFWD (SWG), USFWS, VDEC, VTrans

Bibliography

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.

Maryland Department of Natural Resources (MDNR). American brook lamprey (Lampetra appendix). MDNR, Wildlife and Heritage Service, Annapolis. Available at: http://www.dnr.state.md.us/wildlife/Plants_Wildlife/rte/rteamericanbrooklamprey.asp. (January 2015).

Massachusetts Division of Fish & Wildlife (MDFW). 2005. Commonwealth of Massachusetts comprehensive wildlife conservation strategy. MDFW, Natural Heritage & Endangered Species Program, Boston.

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).

New Hampshire Fish and Game Department (NHFGD). 2005. New Hampshire wildlife action plan. NHFGD, Concord



Common Name: **Mottled Sculpin**
Scientific Name: **Cottus bairdi**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S2

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

At present eight populations distributed among the same number of streams are known to occur in Vermont (Langdon 2014). All are in the Lake Champlain Valley biophysical region and are peripheral to the species' continental range. Very little is known of the biology, abundance of these populations, or population threats. On a range-wide scale no major threats are known (NatureServe 2014).

Distribution

The mottled sculpin has a wide but discontinuous distribution in North America with the more eastern range encompassing northern Quebec and Labrador, and the drainages of the Great Lakes, Hudson and James bays. Western populations are largely limited to the Columbia River drainage (Scott and Crossman 1973). In Vermont, seven populations of mottled sculpin have been identified, all in tributaries to northern Lake Champlain (Allen Brook, Colchester; Bartlett Brook, South Burlington; Englesbee Brook, Burlington; Lamoille River, Milton; Mill River, Georgia; Stonebridge Brook, Milton; Trout Brook, Milton. It is also suspected to inhabit shoreline areas of Lake Champlain.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Mottled sculpin occur in cool streams, small rivers and in lakes. It prefers a bottom of gravel or rubble, rarely being found over sand. In streams it is a riffle dweller. In lakes it inhabits rocky shorelines. Although most reports label mottled sculpin as a cold or cool water species, Vermont specimens have been recorded in warmer streams below the fall line (Langdon et al. In press). Mottled sculpin are found in areas with substrates composed of clean gravel and rubble/cobble. They are intolerant of habitat degradation (siltation and turbidity) and populations have been reduced in some parts of its range. Spawning takes place in cavities beneath rocks, ledges, or logs generally in May when water temperatures reaches 10 C (Smith 1985; Trautman 1981).



Common Name: **Mottled Sculpin**
Scientific Name: **Cottus bairdi**
Species Group: **Fish**

Habitat Types:

Aquatic: Lacustrine

Aquatic: Fluvial

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Habitat Alteration

Sedimentation

Description of habitat threat(s): Mottled sculpin are intolerant of habitat degradation due to sedimentation, siltation and turbidity. This is believed to be one cause for the reduction or extirpation of populations in some parts of its range.

Non-Habitat Threats:

Unknown Non-Habitat Threats

Description of non-habitat threat(s):

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Very little is known about the distribution and abundance
Research	Threats and Their Significance	Medium	Evaluate and monitor the impacts of sedimentation and strategies to reduce sedimentation
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	Medium	Monitoring population status needed.
Monitoring	Habitat Change	Medium	Monitor known habitat to ascertain current status and future changes.
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	Medium	Monitor the impacts of sedimentation
Monitoring	Other Monitoring Needs	N/A	



Common Name: **Mottled Sculpin**
Scientific Name: **Cottus bairdi**
Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Low	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

Bibliography

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.



Common Name: **Lake Sturgeon**
Scientific Name: **Acipenser fulvescens**
Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G3G4

Global Trend: Unknown

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Lake Champlain and the lower reaches of three major tributaries (Winooski, Lamoille and Missisquoi rivers, all directly accessible from the lake) are habitats for the only Lake Sturgeon population occurring in Vermont. This population is on the eastern edge of its continental range. Prior to closure of the sturgeon fishery in 1967 commercial landings indicated a rapid decline in sturgeon abundance during the 1940s. In 1975 Lake Sturgeon was designated endangered in the state. Occasional encounters with Lake Sturgeon during VFWD fisheries assessments and incidental catches by anglers persist. Current data has found adult sturgeon ascend the Lamoille and Winooski rivers during the spring spawning season, and spawning has been confirmed by egg trap and drift net sampling done in those rivers as well as the Missisquoi River (MacKenzie 2014). Abundance, age class structure and distribution of sturgeon in the lake are unknown. Sturgeon ascending Missisquoi and Lamoille rivers to spawn are limited to limited suitable habitat located downstream of Swanton Dam and Peterson Dam, respectively, which likely have significantly reduced their historic range within these rivers. Threats to sturgeon populations generally include overexploitation (including poaching), dams (direct and indirect effects), contaminants, habitat degradation, and introduced species (COSEWIC 2006). Because Lake Sturgeon is a slow growing, late maturing, intermittently spawning species, depleted populations, even when protected, may take many years to recover, if at all (COSEWIC 2006). The species is classified as threatened in New York State (NYSDEC 2005) as well as in the Great Lakes – Upper St. Lawrence unit of Canada (COSEWIC 2006). Selected as an Regional-SGCN by the 13 Northeastern states in 2014

Distribution

Lake sturgeon has a wide distribution occurring from the St. Lawrence River to Hudson Bay, west to the Saskatchewan River in Alberta, south through Lake Champlain, the Mississippi River to the Tennessee River in Alabama and in northern Mississippi; from lakes Winnipeg and Manitoba south through eastern North and South Dakota, northeastern Nebraska and Kansas to eastern Missouri and Arkansas (Scott and Crossman 1973). In Vermont, it inhabits only Lake Champlain with small spawning runs recently documented to occur in the Missisquoi, Lamoille and Winooski rivers (C. MacKenzie, Vermont Fish and Wildlife Department, personal communication). The Vermont population is on the eastern edge of the species North American range.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds



Common Name: **Lake Sturgeon**
Scientific Name: **Acipenser fulvescens**
Species Group: **Fish**

Lake Champlain
Lamoille River
Missiquoi River
Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Lake sturgeon is a demersal species inhabiting lakes and large rivers, usually at 5-9 m depth, over mud, sand, and gravel (Page and Burr 1991). Sturgeon prefer spawning in fast, shallow water with rocky substrate (Harkness and Dymond 1961). Chiotti et al. (2008) located spawning sites in the Big Manistee River of Michigan at water depths in the range of 1.5 to 3 m and average water velocities in the range of 0.34 to 1.32 m/s. Lahaye et al. (1992) collected sturgeon eggs on spawning grounds in the Des Prairies and L'Assomption rivers (Quebec) at minimum and maximum water velocities of 0.02 and 1.39 m/s. Age-0 sturgeon prefer shallow (<2 m), riverine areas with substrate of coarse sand or pea-sized gravel, low current velocity (<0.60 m/s), and an absence of rooted vegetation (Kempinger 1996).

Habitat Types:

Aquatic: Lacustrine
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Habitat Alteration
Sedimentation
Habitat Fragmentation

Description of habitat threat(s): Lake sturgeon eggs require clean river bottoms for survival and the species has declined in areas where siltation has been high. Dams fragment habitat and create barriers to upstream migrating fish during the spawning season.

Non-Habitat Threats:

Genetics
Pollution
Harvest or Collection
Reproductive Traits

Description of non-habitat threat(s): Lake sturgeon are long-lived. Maturity is attained at 14-20 years, and thereafter reproduction occurs approximately every four years. These life history characteristics make lake sturgeon extremely vulnerable to harvest and other disturbances. Lake sturgeon populations exhibit



Common Name: **Lake Sturgeon**
 Scientific Name: **Acipenser fulvescens**
 Species Group: **Fish**

long recovery times because of delayed maturation and the number of years between spawning events. Low population size could lead to inbreeding depression. Over-harvest by sport and commercial fishermen prior to closure of the Vermont fishery in 1967 may have contributed to the decline of the species in Lake Champlain.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Little is known about the current distribution and abundance of Lake Sturgeon in Lake Champlain and spawning tributaries.
Research	Threats and Their Significance	High	
Research	Population Genetics	Medium	Genetic testing of tissue samples from Lake Sturgeon show s that the population is most closely related to that occurring in the St. Lawrence River. Genetic diversity appears to be relative consistent among populations and heterozygosity is consistent with that observed for most other freshwater fish species. Even though the Lake Champlain population has declined significantly, genetic diversity within the population is not depressed nor is there signs of inbreeding at this time.
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	Estimates of lake population size and age structure are lacking.
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Lake Sturgeon**
 Scientific Name: **Acipenser fulvescens**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Invasive Species Control & Prevention	Medium	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)
Invasive Species Control & Prevention	Medium	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)
Natural Processes Restoration	Medium	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, dam owners	Dam owners
Legislation	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP	
Habitat Restoration	High	Enforce and monitor compliance with applicable laws & regulations. Monitor habitat conditions & stressors effects on habitats; restore critical habitats or ameliorate threats as opportunities arise to [CLIPPED]	Increase and/or maintain available habitat in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, NRCS, USFWS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG, WPAG, VBBRG, 604b), NRCS (EQIP)
Natural Processes Restoration	High	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, dam owners	Dam owners
Natural Processes Restoration	Medium	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, dam owners	Dam owners

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Lake Sturgeon
Scientific Name: *Acipenser fulvescens*
Species Group: Fish

Awareness Raising and Communications	High	Enhance public awareness of SGCN and threats to VT populations; a greater understanding of the effects of their own actions on SGCN and steps they can take to restore the lake's population; develop public and [CLIPPED]	Number of outreach efforts made to better inform the public .	VFWD, USFWS, VDEC, TNC, Echo Center, LCBP, LCI	VFWD, USFWS, VDEC, TNC, Echo Center, LCBP, LCI
Species Restoration	High	Develop and implement a plan for restoring Lake Sturgeon to Lake Champlain and historic spawning tributaries.	Components of the restoration plan that are implemented within the timeframe of the current WAP.	VFWD, USFWS	SWG, DJ, USFWS
Invasive Species Control & Prevention	High	Reduce in Sea Lamprey wounding rates observed on Lake Sturgeon.	Sea lamprey wounding rates.	VFWD, USFWS, NYDEC	USFWS, VFWD, NYDEC, GLFC
Compliance & Enforcement	Medium	Protect Lake Sturgeon from directed and incidental harvest by anglers through law enforcement and enhanced public outreach.	Documented annual number of illegal harvest cases and number of fish harvested.	VFWD	VFWD

Bibliography

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2006. COSEWIC assessment and update status report on the lake Sturgeon *Acipenser fulvescens* in Canada. COSEWIC, Ottawa. Available at: www.sararegistry.gc.ca/status/status_e.cfm. (November 2014).

MacKenzie, C. 2014. Vermont management plan for lake sturgeon in Lake Champlain (draft). Vermont Department of Fish and Wildlife, Montpelier.

NYSDEC (New York State Department of Environmental Conservation). 2005. A strategy for conserving New York's fish and wildlife resources. Final Submission Draft, Albany. Available at: www.dec.ny.gov/docs/wildlife_pdf/cwcs2005.pdf. (November 2014).

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Eel (Lake Champlain population)**
Scientific Name: **Anguilla rostrata**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G4

Global Trend: Unknown

State Rank: S2

State Trend: Increasing

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

American eel use to be an abundant species in both Connecticut and Lake Champlain drainages. As recently as the early 1980s, it was harvested commercially in Lake Champlain. For at least 150 years the Richelieu River, which flows out of Lake Champlain, supported a significant commercial eel fishery. Canadian fishery scientist estimated the average annual eel harvest between 1920 and 1980 was 35 metric tons; however, since 1981 landings have decreased from 72.9 tons to only 4.7 tons (Dumont et al. 2004). Today in Vermont, it is rarely encountered. The construction of large dams on rivers once ascended by eels have blocked juvenile fish access to critical rearing habitats. Dams used for hydropower generation have also been implicated in causing high levels of mortality as eels are subjected to turbine losses during their outmigration to sea. High commercial harvest of juvenile eels in coastal waters has further contributed to depressed U. S. stocks. Global climate change has also been identified as a potential limiting factor. The current status of American eel on a national level has given support to the species being recommended for federal endangered species designation. In 2000, the Atlantic States Marine Fisheries Commission adopted an "Interstate Fishery Management Plan for American eel."

Distribution

The American eel occurs from Greenland throughout much of eastern Canada, south through most of eastern United States to the Gulf coast, along the eastern seaboard of Mexico to the Yucatan Peninsula, the West Indies and Bermuda to the Gulf of Mexico, Panama and the West Indies and Bermuda (Scott and Crossman 1973). Spawning grounds are in the Sargasso Sea. In Vermont, eel historically were found through much of Vermont (Lake Champlain and Connecticut River drainages).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Probable
Northern VT Piedmont	Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Probable		

Distribution by Watershed:

Known Watersheds

Probable Watersheds

Lake Champlain



Common Name: **American Eel (Lake Champlain population)**
Scientific Name: **Anguilla rostrata**
Species Group: **Fish**

Lamoille River
Missiquoi River
Otter Creek

Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

American eel is a catadromous species and is reported to only spawn in the Sargasso Sea of the Atlantic Ocean. Larvae are carried by ocean currents to coastal areas where they transform into glass eels, then elvers and begin a long upstream migration to inland waters where they can live in excess of 20 years (as immature yellow eels) before returning to the sea to spawn (silver eels). The only life stages occurring in Vermont is the immature yellow phase and at the beginning of their downstream migrating the silver phase. Yellow-phase American eels have occurred in both the Connecticut River and Lake Champlain drainages of Vermont, where they can live in a wide variety of habitats including ponds, lakes, rivers and streams. They often occupy areas where they can find cover (rocks, snags, weeds) during daylight hours.

Habitat Types:

Aquatic: Lower CT River
Aquatic: Lacustrine
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lake Champlain
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Habitat Fragmentation
Climate Change

Description of habitat threat(s): American eels migrate into Vermont from the Atlantic Ocean. Therefore, obstructions, such as dams, in the migration route can fragment habitat and limit access. Recent studies by Knights (2003) and Wirth and Bernatchez (2003) suggest that climate change may be affecting ocean temperatures and currents such that food and dispersal of eel larvae have been negatively impacted, thereby reducing survival and recruitment.

Non-Habitat Threats:

Pollution



Common Name: **American Eel (Lake Champlain population)**
Scientific Name: **Anguilla rostrata**
Species Group: **Fish**

Harvest or Collection

Description of non-habitat threat(s): For decades, eels from Lake Champlain have been harvested in Canada as they migrate out along the Richelieu and St. Lawrence rivers. This has also occurred elsewhere throughout the range of the species in North America. In addition, eels were harvested for commercial sale in Vermont waters of Lake Champlain by electrofishing for a few years in the 1980s. The commercial harvest of eels in Lake Champlain was made illegal in 2002. In-migrating glass eels and elvers have also been harvested in coastal areas and tributaries of North America for many years and shipped to Asia to supply aquacultural operations. Commercial exploitation has contributed to over-harvested eel stocks. Given the long length of time American eels can reside in freshwater and being piscivores pollution may contribute to bio-accumulation of contaminants in their tissues. Being a catadromous species, American eel are subjected to a wide variety of problems associated with extensive migrations and residency in both marine and freshwater environments. During outmigration from freshwater rearing waters to saltwater silver eels can be exposed to high mortality having to pass through turbines at hydroelectric power dams.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	Contribution of eels in northern regions to overall stock is unknown.
Research	Threats and Their Significance	High	1) Pollution and contaminants: analyze tissue samples for contaminants. 2) Passage: determine passage efficiency of eelways on Richilieu R., assess impacts of turbine mortality on outmigrating eels, and investigate providing eelways at other critical dams
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	Monitor populations and maintain an eel database.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Eel (Lake Champlain population)**
 Scientific Name: **Anguilla rostrata**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Natural Processes Restoration	Medium	Improve flow regimes below hydroelectric generation and flood control projects.	Number of projects which operate under flow regimes that provide suitable habitat for American Eel.	VDEC, VFWD, USFWS, dam owners	Dam owners
Research	Low	Investigate the significance of eel harvest in Lake Champlain and whether harvest protection measures are warranted.	Issue was reviewed with recommendations.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM
Natural Processes Restoration	High	Restore fish passage at dams to allow upstream migrants access to spawning and juvenile habitats and expedite outmigrants (post-spawned adults, juveniles) to sea. Operate and maintain existing fishways for peak passage performance.	Number of dams or other obstructions removed or mitigated to restore fish access to critical habitat; and number of miles of habitat that access was restored.	VDEC, VFWD, USFWS, dam owners	Dam owners
Invasive Species Control & Prevention	Low	Monitor health of American Eels populating Lake Champlain Basin. Enforce fish importation regulations and disease management protocols.	Enforcement of importation regulations.	VFWD, USFWS, NYDEC	VFWD (SWG, DJ), USFWS
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **American Eel (CT River population)**
Scientific Name: **Anguilla rostrata (CT R)**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G4

Global Trend: Unknown

State Rank: S2

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Historically, American Eels were common in Vermont and found in watersheds lying east and west of the Green Mountains (Thompson 1853), but since colonial settlement of the state eels have been negatively affected by artificial barriers to their migrations (dams) and habitat loss and alteration (NatureServe 2014). Other identified threats to eel populations are hydro turbine mortality, oceanic conditions, overfishing, parasitism, predation and pollution (NatureServe 2014). MacMartin (1962) reported of his statewide survey of Vermont streams (1952-1960) that eels were found in only one tributary to the Connecticut River, i.e. the West River, where eels were encountered far upstream in the mainstem and in headwater streams. From the late 1980s through the 1990s eels were observed with some regularity in the lower West River (i.e. downstream of Ball Mountain Dam) and in certain tributary streams. However, since that time, eel sightings have become much less frequent in occurrence (K. Cox, Vermont Fish & Wildlife Department, personal communication). In recent years American eel populations have been in decline. The ASMFC (2012) reported “[a]ccording to the 2012 benchmark stock assessment, American Eel population is depleted in U.S. waters. The stock is at or near historically low levels due to a combination of historical overfishing, habitat loss, food web alterations, predation, turbine mortality, environmental changes, toxins and contaminants, and disease.” Restoration of American Eel in Vermont tributaries to the Connecticut River is largely contingent upon appropriate management actions implemented by coastal state fishery agencies and the federal government. In 2010 the U.S. Fish & Wildlife Service (USFWS) was petitioned to list American Eel as an endangered species (ASMFC 2014). The USFWS concluded the petition may be warranted and undertook a status review. The USFWS is expected to publish its proposed rule in 2015. American eel was selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

The American eel occurs from Greenland throughout much of eastern Canada, south through most of eastern United States to the Gulf coast, along the eastern seaboard of Mexico to the Yucatan Peninsula, the West Indies and Bermuda to the Gulf of Mexico, Panama and the West Indies and Bermuda (Scott and Crossman 1973). Spawning grounds are in the Sargasso Sea. In Vermont, eel historically were found through much of Vermont (Lake Champlain and Connecticut River drainages).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Probable
Northern VT Piedmont	Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Probable		

Distribution by Watershed:

Known Watersheds

Probable Watersheds



Common Name: **American Eel (CT River population)**
Scientific Name: **Anguilla rostrata (CT R)**
Species Group: **Fish**

Middle Connecticut	Waits
West	Upper Connecticut - Mascoma
Black - Ottauquechee	Deerfield

Passumpsic
Upper Connecticut
White

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

American eel is a catadromous species and is reported to only spawn in the Sargasso Sea of the Atlantic Ocean. Larvae are carried by ocean currents to coastal areas where they transform into glass eels, then elvers and begin a long upstream migration to inland waters where they can live in excess of 20 years (as immature yellow eels) before returning to the sea to spawn (silver eels). The only life stages occurring in Vermont is the immature yellow phase and at the beginning of their downstream migrating the silver phase. Yellow-phase American eels have occurred in both the Connecticut River and Lake Champlain drainages of Vermont, where they can live in a wide variety of habitats including ponds, lakes, rivers and streams. They often occupy areas where they can find cover (rocks, snags, weeds) during daylight hours.

Habitat Types:

Aquatic: Lacustrine

Aquatic: Fluvial

Aquatic: Large Lake Champlain Tribs Below Falls

Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Habitat Fragmentation

Climate Change

Description of habitat threat(s): American eels migrate into Vermont from the Atlantic Ocean. Therefore, obstructions, such as dams, in the migration route can fragment habitat and limit access. Recent studies by Knights (2003) and Wirth and Bernatchez (2003) suggest that climate change may be affecting ocean temperatures and currents such that food and dispersal of eel larvae have been negatively impacted, thereby reducing survival and recruitment.



Common Name: **American Eel (CT River population)**
Scientific Name: **Anguilla rostrata (CT R)**
Species Group: **Fish**

Non-Habitat Threats:

Pollution

Harvest or Collection

Description of non-habitat threat(s): For decades, eels from Lake Champlain have been harvested in Canada as they migrate out along the Richelieu and St. Lawrence rivers. This has also occurred elsewhere throughout the range of the species in North America. In addition, eels were harvested for commercial sale in Vermont waters of Lake Champlain by electrofishing for a few years in the 1980s. The commercial harvest of eels in Lake Champlain was made illegal in 2002. In-migrating glass eels and elvers have also been harvested in coastal areas and tributaries of North America for many years and shipped to Asia to supply aquacultural operations. Commercial exploitation has contributed to over-harvested eel stocks. Given the long length of time American eels can reside in freshwater and being piscivores pollution may contribute to bio-accumulation of contaminants in their tissues. Being a catadromous species, American eel are subjected to a wide variety of problems associated with extensive migrations and residency in both marine and freshwater environments. During outmigration from freshwater rearing waters to saltwater silver eels can be exposed to high mortality having to pass through turbines at hydroelectric power dams.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	
Research	Threats and Their Significance	High	1) Pollution and contaminants: analyze tissue samples for contaminants. 2) Passage: determine passage efficiency of fishways on the Connecticut River and tributaries, assess impacts of turbine mortality on outmigrating eels, and investigate need for eelways at critical dams
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	Monitor populations and maintain an eel database.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Eel (CT River population)**
Scientific Name: **Anguilla rostrata (CT R)**
Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Natural Processes Restoration	High	Restore fish passage at dams to allow upstream migrants access to spawning and juvenile habitats and expedite outmigrants (post-spawned adults, juveniles) to sea. Operate and maintain existing fishways for peak passage performance.	Number of dams or other obstructions removed or mitigated to restore fish access to critical habitat; and number of miles of habitat that access was restored.	VDEC, VFWD, NHDES, NHFGD, CRASC, USFWS, CRWC, dam owners	Dam owners
Invasive Species Control & Prevention	Low	Monitor health of American Eels populating the Connecticut River Basin. Enforce fish importation regulations and disease management protocols.	Enforcement of importation regulations.	VFWD, NHFGD, CRASC, USFWS	VFWD, NHFGD, CRASC, USFWS
Policy & Regulations	Low	Support and cooperate with the inter-agency program for the restoration of anadromous fishes to the Connecticut River basin (e.g., CRASC).		VFWD, NHFGD, CRASC, USFWS	VFWD, NHFGD, CRASC, USFWS
Natural Processes Restoration	Medium	Improve flow regimes below hydroelectric generation and flood control projects.	Number of projects which operate under flow regimes that provide suitable habitat for American Eel.	VDEC, VFWD, NHDES, NHFGD, CRASC, USFWS, CRWC, dam owners	Dam owners
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, NHDES, NHFGD, CRASC, CRWC, NRCS, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Eel (CT River population)**
Scientific Name: **Anguilla rostrata (CT R)**
Species Group: **Fish**

Bibliography

ASMFC (Atlantic States Marine Fisheries Commission). 2012. American eel benchmark stock assessment. ASMFC Stock Assessment Report No. 12-01, Washington, D.C. Available at: http://www.asmfc.org/uploads/file/AmericanEelBenchmarkStockAssessmentReport_May2012.pdf. (December 2014).

ASMFC (Atlantic States Marine Fisheries Management Commission). 2014. American eel. Available at: <http://www.asmfc.org/species/American-eel>. (December 2014).

Chipman, B. 2014. Personal communication. Vermont Fish and Wildlife Department, Essex.

FTC (Fisheries Technical Committee). 2009. Strategic plan for Lake Champlain fisheries. Lake Champlain Fish and Wildlife Management Cooperative, U. S. Fish and Wildlife Service, Essex Junction, Vermont.

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).

Staats, N. Not dated. American eel sampling in Lake Champlain 2014 progress report, draft. U. S. Fish and Wildlife Service, Essex Junction, Vermont.

Thompson, Z. 1853. Natural history of Vermont, 1972 edition. Charles E. Tuttle Company, Rutland, Vermont



Common Name: **Blueback Herring (CT River only)**
Scientific Name: **Alosa aestivalis**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: SU

State Trend: Historic

Extirpated in VT? Yes

Regional SGCN? Yes

Assessment Narrative:

Blueback herring were once common in the Connecticut River mainstem as far upriver as the natural barrier (falls) at Bellows Falls, Vermont and North Walpole, New Hampshire, a distance of 174 river-miles from the river mouth on Long Island Sound. Construction of the first dam on the river in 1798 near present-day Turners Falls, Massachusetts extirpated the species from the upper river (Gephard and McMenemy 2004). With restoration of fish passage at Turners Falls (MA) and Vernon (VT) dams during the early 1980s Blueback Herring as well as other anadromous fishes once again have access to critical habitats in the upper river. During the first 10 years that the Vernon fish ladder was operated (1981-1990) an average of 35 bluebacks were observed annually passing into the Vernon headpond which extends upriver nearly to Bellows Falls. In 1991 the Vernon Dam herring run peaked at 383 fish which was followed over the next nine years (1992-2000) with a downward trend (average annual passage count 22 fish). Since then, Blueback Herring have not been observed above Vernon Dam. A similar trend has occurred in the lower Connecticut River, represented by passage counts made at the Holyoke (MA) Dam fishlift, i.e. an increasing trend from the 1970s through mid-1980s followed decreasing numbers thereafter. Such declines have been noted in other U.S. river stocks ranging from Maine to Delaware. Stock assessments for Blueback Herring and Alewife, collectively reported as river herring, made by the Atlantic State Marine Fisheries Commission (ASMFC) concluded that 23 (including the Connecticut River) of the 52 stocks assessed are depleted relative to historic levels (ASMFC 2012). NatureServe (2013) lists Blueback Herring as vulnerable based on "drastic declines in abundance...in many areas in recent decades." Causes for stock declines are not fully understood at present; however, restoration of Blueback Herring to the VT-NH shared section of the Connecticut River is contingent on appropriate management actions implemented by coastal state fishery agencies and the federal government. Freshwater threats to Blueback Herring stocks include obstruction of migration routes by dams and other barriers, entrainment and impingement mortality, habitat degradation, overfishing, and predators e.g. striped bass (NatureServe 2013; Savoy and Crecco 2004). Selected as an Regional-SGCN by the 13 Northeastern states in 2014

Distribution

The blueback herring reaches its northern limit in Canadian waters and occurs along the eastern North American seaboard from Cape Breton, Nova Scotia south to northern Florida (Scott and Crossman 1973). Landlocked populations are also known to occur, including Lake Champlain. Anadromous blueback herring occur naturally in the Connecticut River basin, although their historic occurrence in the Vermont-New Hampshire shared section of the Connecticut River is reported to have been similar to that of American shad. The upriver limit to their distribution extended to the natural barrier on the river between Bellows Falls, Vermont and North Walpole, New Hampshire. Although blueback herring are also found in Lake Champlain (since 1979), the species is not indigenous to that waterbody and is believed to have gained access to the lake by the Hudson Barge Canal connection to the Hudson River. The last year blueback herring (2 individuals) have been observed in Vernon fish ladder was 2000.

Distribution by Biophysical Region:

Champlain Valley Not Probable

Southern VT Piedmont Probable

Champlain Hills Not Probable

Vermont Valley Not Probable



Common Name: **Blueback Herring (CT River only)**
 Scientific Name: **Alosa aestivalis**
 Species Group: **Fish**

Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Probable Watersheds

Middle Connecticut

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Blueback herring is an anadromous species spending portions of its life in both marine and freshwater environments. In freshwater, it migrates up rivers to spawn. Generally the upstream migration of blueback herring does not extend as far as for other clupeid species (e.g., alewife); spawning may occur in both brackish and fresh waters (Scott and Crossman 1973). Spawning occurs in a diversity of habitats, including large rivers, small streams, ponds and large lakes over a range of substrates, such as gravel, sand, detritus, and submersed vegetation and other structures. Swift flowing waters are used as spawning sites. Blueback herring spawn in rivers releasing their eggs into the water column whereupon the eggs settle and adhere to the substrate, including stones, gravel and sticks (Scott and Crossman 1973). Larval and juvenile blueback herring reside in the river during their first summer before out-migrating the following fall. Spawning success and survival of adults and juveniles are dependent on successful passage past manmade obstacles, such as dams and hydroelectric generation plants.

Current Threats

Habitat Threats:

Habitat Fragmentation

Description of habitat threat(s): Spawning success and survival of adults and juveniles are limited by the existence and development of dams and hydroelectric generation plants which may impede access to spawning and nursery habitats, as well as cause mortality to outmigrating fish. River flow regimes as manipulated by hydroelectric power generation activities can affect blueback herring habitat.

Non-Habitat Threats:

Predation or Herbivory

Description of non-habitat threat(s): Migrating adult and juvenile blueback herring are susceptible to predation, especially fish predators. Improvements in stock strength of striped bass and bluefish are believed to have increased predatory pressure on migrating clupeid populations, including blueback herring and American shad. Being an anadromous species, blueback herring are subjected to a wide variety of problems associated with extensive migrations and residency in both marine and freshwater environments. Outmigrating adult and juvenile herring may be exposed to turbine mortality at power dams resulting in high mortality.



Common Name: **Blueback Herring (CT River only)**
Scientific Name: **Alosa aestivalis**
Species Group: **Fish**

Bibliography

ASMFC (Atlantic States Marine Fisheries Commission). 2012. ASMFC stock assessment overview: river herring. ASMFC, Washington, D.C. Available at: http://www.asmfc.org/uploads/file/riverHerringStockAssessmentOverview_May2012.pdf. (October 2014).

Gephard, S., and J. McMenemy. 2004. An overview of the program to restore Atlantic salmon and other diadromous fishes to the Connecticut River with notes on the current status of these species in the river. Pages 287-317 in P. M. Jacobson et al., editors. The Connecticut River ecological study (1965-1973) revisited: ecology of the lower Connecticut River 1973-2003. American Fisheries Society Monograph 9, Bethesda, Maryland.

NatureServe. 2013. *Alosa aestivalis*. IN: The IUCN Red List of Threatened Species. Version 2014.2. Available at: <http://www.iucnredlist.org/details/full/201946/0>. (October 2014).

Savoy, T. F., and V. A. Crecco. 2004. Factors affecting the recent decline of blueback herring and American shad in the Connecticut River. Pages 361-377 in P. M. Jacobson et al., editors. The Connecticut River ecological study (1965-1973) revisited: ecology of the lower Connecticut River 1973-2003. American Fisheries Monograph 9, Bethesda, Maryland.



Common Name: **American Shad**
 Scientific Name: **Alosa sapidissima**
 Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S4

State Trend: Increasing

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

American Shad were once common in the Connecticut River mainstem as far upriver as the natural barrier at Bellows Falls, Vermont and North Walpole, New Hampshire, a distance of 174 river-miles from the river mouth on Long Island Sound. Construction of the first dam on the river in 1798 near present-day Turners Falls, Massachusetts extirpated the species from the upper river (Gephart and McMenemy 2004). With restoration of fish passage at Turners Falls (MA) and Vernon (VT) dams during the early 1980s shad as well as other anadromous fishes are once again provided access to critical habitats in the upper river. During the first eleven years that upstream fish passage was provided at Vernon Dam the number of adult shad passed above the dam followed an increasing trend from 97 shad counted in 1981 to 37,197 shad in 1991. Over the following 20 years the shad passage above the dam declined from 31,155 fish in 1992 to 290 in 2011. Since then annual shad passage counts at the dam have been in the range of 10,175 and 27,706 fish. Factors affecting shad abundance above Turners Falls Dam are complicated but illustrate the some of the threats that beset the species and challenge restoration of the species to the upper Connecticut River. Specifically, after 1990 commercial shad landings along the Atlantic seaboard and stock abundance in several rivers declined significantly. In the Connecticut River, 1.6 million shad were estimated to have returned to the river in 1992; by 1995 returns had declined to 304,500 fish (ASFC 1999). Striped Bass predation on adult and juvenile shad has also been suggested (Savoy and Crecco 2004). It was also hypothesized that the thermal discharge from Vermont Yankee Nuclear Power into the Vernon headpond may have had negative effects on adult and juvenile shad migrations and survival. And lastly, structural and operational factors at both Turners Falls and Vernon fish ladders affecting fish passage were identified and corrected which appears to explain increased passage at Vernon observed in 2012-2014. Threats to American Shad habitats include the following: barriers to migration, water withdrawals, toxic and thermal wastewater discharge, channelization, dredging and instream construction, inappropriate land uses, atmospheric deposition, climate change, competition and predation by invasive and managed species, fisheries activities, and instream flow regulation (ASMFC 2010). Selected as an Regional-SGCN by the 13 Northeastern states in 2014

Distribution

American shad inhabit waters along the Atlantic Coast from Newfoundland south to Florida; it has also been introduced to the Pacific Coast of North America (Scott and Crossman 1973). In Vermont, it is restricted to the Connecticut River from the Massachusetts state line upstream to at least Bellows Falls dam. Low numbers in some years are passed above Bellows Falls via the fish ladder there. The last year shad were observed in the ladder was 2005. Shad have also been observed in the lower West River.

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Confident
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		



Common Name: **American Shad**
Scientific Name: **Alosa sapidissima**
Species Group: **Fish**

Distribution by Watershed:

Known Watersheds

Middle Connecticut
West

Probable Watersheds

Black - Ottauquechee

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

As an anadromous fish, the American shad divides its life between marine and freshwater environments. Adults ascend rivers and streams to spawn. Once the eggs hatch, larval and juvenile shad inhabit riverine habitats, including setbacks, through their first summer of life before out-migrating to the ocean in the fall. Maturity is attained at sea. Stier and Crance (1985) review the habitat requirements of American shad. Adults utilize well oxygenated (≥ 5 ppm), flowing water, although they do not appear to have specific preferences for spawning locations other than broad flats and shallow water. Spawning may occur over a variety of substrate types providing water velocity is sufficient enough to keep sedimentation minimal. Spawning generally occurs at water temperatures of 8-26 C with peak activity occurring within the range of 14-21 C. Temperatures for maximum egg hatch and survival is 15.5-26 C. Temperatures at or near 11 C are minimal for egg incubation, and temperatures in excess of 80.1 F are unsuitable. Juvenile shad are found at water temperatures of 10-31 C. Temperatures less than 10 C cannot be tolerated. Juvenile outmigration begins when the water temperature goes below 15.5 C.

Current Threats

Habitat Threats:

Habitat Fragmentation

Description of habitat threat(s): Spawning success and survival of adults and juveniles are limited by the existence and development of dams and power generation plants which may impede access to spawning and nursery habitats, as well as impose artificial flow regimes associated which in turn alter and degrade habitat for shad. Fishway design deficiencies at ladders on the Connecticut River at Turners Falls, Massachusetts are being examined as likely causes for the significant reduction in shad run abundance into Vermont and New Hampshire. This exacerbates the habitat fragmentation problem.

Non-Habitat Threats:

Harvest or Collection

Predation or Herbivory

Description of non-habitat threat(s): Migrating adult and juvenile shad are susceptible to predation, especially fish predators. Improvements in stock strength of striped bass and bluefish are believed to have increased predatory pressure on migrating clupeid populations, including shad. Being an anadromous species, shad are subjected to a wide variety of problems associated with extensive migrations and residency in both marine and freshwater environments. Excessive commercial harvest off the Atlantic seaboard and within the estuaries represents a problem for shad stocks. Outmigrating adult and juvenile shad may be exposed to turbine mortality at power dams resulting in high mortality.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Shad**
Scientific Name: **Alosa sapidissima**
Species Group: **Fish**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Low	
Research	Threats and Their Significance	High	Data are needed to design fish ladders for improved fish passage performance.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	High	Research is being conducted by other agencies to which Vermont is a cooperator.
Monitoring	Population Change	High	
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	
Monitoring	Other Monitoring Needs	N/A	

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **American Shad**
Scientific Name: **Alosa sapidissima**
Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Natural Processes Restoration	High	Improve flow regimes below hydroelectric generation and flood control projects.	Number of projects which operate under flow regimes that provide suitable habitat for American Shad.	VDEC, VFWD, NHDES, NHFGD, CRASC, USFWS, CRWC, dam owners	Dam owners
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, NHDES, NHFGD, CRASC, CRWC, NRCS, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, NHDES, NHFGD, CRASC, CRWC, NRCS, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Legislation	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, NHDES, NHFGD, CRASC, USFWS, CRWC	
Natural Processes Restoration	High	Restore fish passage at dams to allow upstream migrants access to spawning and juvenile habitats and expedite outmigrants (post-spawned adults, juveniles) to sea. Operate and maintain existing fishways for peak passage performance.	Number of dams or other obstructions removed or mitigated to restore fish access to critical habitat; and number of miles of habitat that access was restored.	VDEC, VFWD, NHDES, NHFGD, CRASC, USFWS, CRWC, dam owners	Dam owners

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Shad**
 Scientific Name: **Alosa sapidissima**
 Species Group: **Fish**

Species Restoration	High	Provide technical support to regional restoration plans and efforts as necessary.	ASMFC Interstate Fishery Management Plan for Shad & River Herring	CRASC, USFWS, NMFS, ASFMC, CRWC	CRASC, USFWS, NMFS, ASFMC, CRWC
Research	Medium	Striped Bass & other predator threats to American Shad stocks.	ASMFC Interstate Fishery Management Plan for Shad & River Herring	CRASC, NMFS, ASMFC	CRASC, NMFS, ASMFC
Policy & Regulations	High	Support and cooperate with the inter-agency program for the restoration of anadromous fishes to the Connecticut River basin (e.g., CRASC).	Implement Connecticut River Basin Management Plan for Blueback Herring. Percent of tasks implemented as prescribed in the plan.	VFWD, NHFGD, CRASC, USFWS, ASMFC	VFWD, NHFGD, CRASC, USFWS

Bibliography

ASMFC (Atlantic States Marine Fisheries Commission). 2010. Amendment 3 to the interstate fishery management plan for shad and river herring (American shad management). ASMFC, Washington, D.C. Available at: http://www.asmfc.org/uploads/file/Amendment3_FINALshad.pdf. (January 2015).

ASMFC. 1999. Amendment 1 to the interstate fishery management plan for shad & river herring. Fishery management report No. 35, Washington, D.C. Available at: <http://www.asmfc.org/uploads/file/shadam1.pdf>. (January 2015).

Gephard, S., and J. McMenemy. 2004. An overview of the program to restore Atlantic salmon and other diadromous fishes to the Connecticut River with notes on the current status of these species in the river. Pages 287-317 in P. M. Jacobson et al., editors. The Connecticut River ecological study (1965-1973) revisited: ecology of the lower Connecticut River 1973-2003. American Fisheries Society Monograph 9, Bethesda, Maryland.

Savoy, T. F., and V. A. Crecco. 2004. Factors affecting the recent decline of blueback herring and American shad in the Connecticut River. Pages 361-377 in P. M. Jacobson et al., editors. The Connecticut River ecological study (1965-1973) revisited: ecology of the lower Connecticut River 1973-2003. American Fisheries Monograph 9, Bethesda, Maryland

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Mooneye**
Scientific Name: **Hiodon tergisus**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: SU

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

In Vermont, Mooneye is confined to Lake Champlain, where it is on the eastern edge of its continental range. Very little is known of its biology, distribution and habitat use, and past and present population abundance in the lake. On a range-wide scale no major threats are known (NatureServe 2014); however in New York, it is listed as threatened (NYSDEC 2014). The cause(s) for declines in New York populations are not as yet known, but increased siltation in part is suspected (NYSDEC 2014). Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

Mooneye is a North American species with a range extending from James Bay, Ottawa River to the Lake Champlain and St Lawrence watershed, southwest of the Appalachian Mountains through western New York and Pennsylvania to Arkansas and Oklahoma, north through eastern Kansas to North Dakota, southeastern Saskatchewan and southern Manitoba (Scott and Crossman 1973). In Vermont, it is a peripheral species at the eastern most extent of its range and has been recorded only from Lake Champlain, including the southern part of the lake, and near the mouths of the Missisquoi and Lamoille rivers and Otter Creek.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain

Probable Watersheds

Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Mooneye are found in shallow areas of large lakes and deep pools of clear rivers where the bottom is relatively free of silt (Langdon et al. In press). Although it is generally found in non-flowing waters, it may also occur in swift waters, such as below dams (Trautman 1957). Mooneye has rarely been taken with collection gear at depths greater than 10.7 m (Scott and Crossman 1973). The mooneye is a warm water species, preferring water temperatures in the range of 27.5-29.0 C, and migrate up rivers to spawn when the water temperatures reach 19.4 C (Langdon et al. In press).



Common Name: **Mooneye**
Scientific Name: **Hiodon tergisus**
Species Group: **Fish**

Habitat Types:

Aquatic: Lower CT River

Aquatic: Lacustrine

Aquatic: Fluvial

Current Threats

Habitat Threats:

Sedimentation

Habitat Fragmentation

Description of habitat threat(s): Mooneye is not tolerant of silted habitats or turbidity (Scott and Crossman 1973).

Non-Habitat Threats:

Unknown Non-Habitat Threats

Description of non-habitat threat(s):

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Assess habitat requirements and use by mooneye in Lake Champlain.
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Very little is known about the distribution and abundance of mooneye in Lake Champlain.
Research	Threats and Their Significance	Medium	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	Monitoring population status needed.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	Medium	
Monitoring	Other Monitoring Needs	N/A	

***Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report***



Common Name: **Mooneye**
Scientific Name: **Hiodon tergisus**
Species Group: **Fish**

Bibliography

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).

NYSDEC (New York State Department of Environmental Conservation). 2014. Mooneye fact sheet. Available at: <http://dec.ny.gov/animals/26032.html>. (November 2014).



Common Name: **Cisco or Lake Herring**
Scientific Name: **Coregonus artedii**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S4

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

At present eight populations distributed among the same number of streams are known to occur in Vermont (Langdon 2014). All are in the Lake Champlain Valley biophysical region and are peripheral to the species' continental range. Very little is known of the biology, abundance of these populations, or population threats. On a range-wide scale no major threats are known (NatureServe 2014).

Distribution

This species is the most widely distributed of the 14 species of the genus *Coregonus* occurring in North America. It is found from eastern Quebec to the Hudson Bay, through the Great Lakes system, Ontario, Manitoba, Saskatchewan and Alberta, north into the Northwest Territories, and in the Mackenzie River system north to Great Bear Lake (Scott and Crossman 1973). In Vermont, where it occurs on the eastern edge of its range, cisco are restricted to Lake Champlain.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

In Vermont, cisco inhabit the cold, deep areas of Lake Champlain. This is a schooling, open-water species inhabiting cool mid-lake areas during the summer, shifting to shallower waters inshore from fall to spring. It cannot tolerate water temperatures of over 26.1 C. Cisco spawn in late fall, just prior to ice formation, at depths of about .9-3 m, over gravel to rubble bottom (Langdon et al. In press).



Common Name: **Cisco or Lake Herring**
Scientific Name: **Coregonus artedii**
Species Group: **Fish**

Habitat Types:

Aquatic: Lower CT River

Aquatic: Lacustrine

Aquatic: Fluvial

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Invasion by Exotic Species

Climate Change

Description of habitat threat(s): The appearance of alewife in Lake Champlain is of great concern due to their potential impact through competition with other pelagic planktivores, such as cisco. The species also require deep, cold water. If climate change has a significant impact on the thermal structure of Lake Champlain, this could affect the cisco population.

Non-Habitat Threats:

Competition

Parasites

Loss of Prey Base

Description of non-habitat threat(s): Sea lamprey have negatively impacted cisco in other bodies of water (Smith 1985, Bronte et al. 2003). As soft-scaled members of the salmon family, cisco are susceptible to parasitism/predation by sea lamprey. Lamprey predation was identified as a contributing factor to the decline of cisco in Lake Superior (Bronte et al. 2003), and 80% of cisco in Oneida Lake, NY that died off during high summer temperatures had lamprey scars (Smith 1985). So far there are no direct reports of lamprey impacts on cisco in Lake Champlain. Cisco are primarily pelagic (open-water) feeders on zooplankton, and therefore might compete with other species with similar feeding habits, such as smelt. The appearance of alewife in Lake Champlain could pose a problem to both of these native open-water planktivores.



Common Name: **Cisco or Lake Herring**
Scientific Name: **Coregonus artedii**
Species Group: **Fish**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	
Research	Basic Life History	Medium	
Research	Distribution and Abundance	High	Develop population indices through forage base monitoring in Lake Champlain.
Research	Threats and Their Significance	Medium	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	Medium	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Cisco or Lake Herring**
 Scientific Name: **Coregonus artedii**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Invasive Species Control & Prevention	High	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitats occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)
Research	Medium	Currently under VFWD fishing regulations Cisco is an unregulated activity. Whether or not harvest poses a threat to Cisco population(s) should be reviewed.	Review was conducted and recommendations were considered.	VFWD, UVM	VFWD (SWG, DJ), UVM
Research	Medium	Investigate whether predation on Cisco presents significant threat to the species.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM
Legislation	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, NRCS, USFWS, LCBP	
Research	Medium	Investigage whether Sea Lamprey parasitism is a significant threat to Cisco.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM

Bibliography

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2005. COSEWIC assessment and update status report on the lake whitefish (Lake Simcoe population) *Coregonus clupeaformis* in Canada. COSEWIC, Ottawa. Available at: www.sararegistry.gc.ca/status/status_e.cfm. (November 2014).

Halnion, L. C. 1963. Historical survey of Lake Champlain's fishery. Vermont Fish and Game Department Federal Aid in Fish Restoration Job Completion Report F-1-R-10, Job 6, Montpelier.

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).

Staats, N. 2015. Personal communication. U. S. Fish and Wildlife Service, Essex.

Staats, N. R., and B. Pientka. 2014. Lake Champlain fisheries restoration and management – forage fish monitoring. Vermont Fish and Wildlife Department Federal Aid Job Performance Report F-35-R-16, Montpelier.



Common Name: **Lake Whitefish**
Scientific Name: **Coregonus clupeaformis**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S4?

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Seven populations have been identified in Vermont (Langdon 2014). On a range-wide scale no major threats are known (NatureServe 2014); however, interspecies hybridization has been reported (*Esox americanus* X *E. niger* and *E. americanus* X *E. lucius*) which could pose a threat to populations where the Redfin Pickerel is the only resident esocid in the fish community and whether by intention or accident another esocid species might be introduced. Redfin Pickerel is listed as endangered in Maine (MDIFW 2013).

Distribution

Lake whitefish is a widely distributed North American species. Its range includes virtually all of Canada and Alaska, the Great Lakes, the St. Lawrence River including northern New York, Vermont, New Hampshire and Maine (Scott and Crossman 1973). In Vermont, it is confined to Lake Champlain. However, in 1878 lake whitefish were introduced into five other Vermont lakes, such as Lake Dunmore in Addison County and Lake St. Catherine in Rutland County, but none of these introductions were successful (Langdon et al. In press).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Lake whitefish inhabit cold lakes which are deep and clear. Lake whitefish spawn during November and December at water temperatures below 7.8 C; spawning occurs near the surface in water less than 7.6 m deep, and the adhesive eggs sink to the bottom onto a usually gravel or rubble and occasionally sand substrate (Langdon et al. In press).



Common Name: **Lake Whitefish**
Scientific Name: **Coregonus clupeaformis**
Species Group: **Fish**

Habitat Types:

Aquatic: Lower CT River

Aquatic: Lacustrine

Aquatic: Fluvial

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Climate Change

Description of habitat threat(s): Lake whitefish also require deep, cold water. If climate change has a significant impact on the thermal structure of Lake Champlain, this could affect the population of lake whitefish.

Non-Habitat Threats:

Competition

Parasites

Description of non-habitat threat(s): The appearance of alewife in Lake Champlain is of great concern due to their potential impact through competition with other pelagic planktivores, such as lake whitefish. Sea lamprey have negatively impacted lake whitefish in Lake Superior (Bronte et al. 2003). As soft-scaled members of the salmon family, lake whitefish are susceptible to parasitism/predation by sea lamprey. Lamprey predation was identified as a contributing factor to the decline of lake whitefish in Lake Superior (Bronte et al. 2003), and whitefish populations rebounded after implementation of lamprey control.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Lake Whitefish**
Scientific Name: **Coregonus clupeaformis**
Species Group: **Fish**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	
Research	Basic Life History	Medium	
Research	Distribution and Abundance	High	Develop population indices through forage base monitoring in Lake Champlain.
Research	Threats and Their Significance	Medium	
Research	Population Genetics	Medium	University of Vermont (E. Marsden) is currently engaged in a genetics study of sub-populations in Lake Champlain,.
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	Medium	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Lake Whitefish**
 Scientific Name: **Coregonus clupeaformis**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Natural Processes Restoration	Medium	UVM is studying the effects of physical isolation/reduced connectivity on Lake Champlain bays and changed water movement, nutrient retention, and potential population sub-structuring of Lake Whitefish.	Completion of study and recommendations that may come from it.	VDEC, VFWD, USFWS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Invasive Species Control & Prevention	High	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)
Legislation	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, NRCS, USFWS, LCBP	
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Invasive Species Control & Prevention	High	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)
Medium	Investigage whether Sea Lamprey parasitism is a significant threat to Lake Whitefish.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Lake Whitefish**
 Scientific Name: **Coregonus clupeaformis**
 Species Group: **Fish**

Medium	Currently under VFWD fishing regulations Lake Whitefish is an unregulated activity. Whether or not harvest poses a threat to Lake Whitefish population(s) should be reviewed.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM

Bibliography

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2005. COSEWIC assessment and update status report on the lake whitefish (Lake Simcoe population) *Coregonus clupeaformis* in Canada. COSEWIC, Ottawa. Available at: www.sararegistry.gc.ca/status/status_e.cfm. (November 2014).

Halnon, L. C. 1963. Historical survey of Lake Champlain's fishery. Vermont Fish and Game Department Federal Aid in Fish Restoration Job Completion Report F-1-R-10, Job 6, Montpelier.

Herbst, S. J., J. E. Marsden, and S. J. Smith. 2011. Lake whitefish in Lake Champlain after commercial fishery closure and ecosystem changes. *North American Journal of Fisheries Management* 31:1106-1115.

Herbst, S. J., J. E. Marsden, and B. J. Lantry. 2013. Lake whitefish diet, condition, and energy density in Lake Champlain and the lower four Great lakes following dreissenid invasions. *Transactions of the American Fisheries Society* 142:388-398.

Marsden, J. E., and R. W. Langdon. 2012. The history and future of Lake Champlain's fishes and fisheries. *Journal of Great Lakes Research* 38:19-34.

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).



Common Name: **Round Whitefish**
Scientific Name: **Prosopium cylindraceum**
Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S1

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Until recently Round Whitefish populations were reported from four lakes, all in northeastern Vermont and the St. Francis River drainage (Willoughby Lake, Lake Seymour, Holland Pond, Beaver Pond), and the upper Connecticut River. There is historic reference (Titcomb and Bailey 1896) to a whitefish species distinct from Lake Whitefish (*Coregonus clupeaformis*) in Lake Memphremagog, but it is ambiguous which coregonine species was intended; the persistence of Round Whitefish upstream in the watershed is evidence that Round Whitefish was indeed the so-called "shad" (i.e. whitefish) noted for Lake Memphremagog (Gerardi 2015). Sampling conducted by electrofishing, experimental gill nets and beach seines between 2006 and 2012 failed to collect whitefish in Seymour Lake, Holland Pond and Beaver Pond suggesting these populations may be extirpated (Kratzer 2011). Likewise, extensive sampling by McGill University staff and students at Lake Memphremagog in the early 1970s provided no documentation of Round Whitefish remaining in the lake (Nakashima and Leggett 1975; Gascon and Leggett 1977). Several possible causes of population elimination are Smallmouth Bass introduction and predation in Lake Seymour, competition with the resident White Sucker population in Beaver Pond, Chain Pickerel (*Esox niger*) predation and/or anoxia in the hypolimnion in Holland Pond (Kratzer 2011), along with major fish community changes in Lake Memphremagog brought about by introduction of many species not native to the lake (Gerardi 2015). As recently as 2011 sampling of the upper Connecticut River by New Hampshire Fish & Game Department captured Round Whitefish, which population is characterized as being abundant (Kratzer 2011). Historically, Round Whitefish were reported to be in Lake Champlain as early as 1894 (Evermann and Kendall 1902) but does not appear to have been encountered during the 1929 biological survey of the lake (Greeley 1930). Current reports of Round Whitefish in the lake appear to be lacking (Marsden 2014; Parrish 2014; Pientka 2014). Potential threats to the species have been identified by NatureServe (2014) to include habitat loss through either siltation of spawning streams or lake shores and decreased water quality resulting from poor forest management practices and land development; road, bridge and in-stream construction projects with negative effects on spawning habitat or obstruction of whitefish access to spawning habitat; piscicides; and exotic fish introductions. Round Whitefish is classified as an endangered species in New York in 1983. Historically, the species was found in over 80 lakes in the state but has now declined to populations numbering fewer than eight (NYSDEC 2015a). Possible reasons for the decline of the species in New York are predation by invading Smallmouth Bass and Yellow Perch, competition with Lake Whitefish, overfishing, loss of spawning habitat, siltation, and lake acidification (NYSDEC 2015b). Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

In Vermont, round whitefish is found in lakes Seymour, Willoughby and in Beaver Pond in Holland. A historic record of "lake whitefish" in Holland Pond (located near Beaver Pond in Holland) is believed actually to have been a round whitefish.

Distribution by Biophysical Region:

Champlain Valley	Historic Record(s) Only	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Round Whitefish**
Scientific Name: **Prosopium cylindraceum**
Species Group: **Fish**

Northern VT Piedmont Not Probable

Taconic Mtns Not Probable

Northeastern Highlands Confident

Distribution by Watershed:

Known Watersheds

St. Francois River

Upper Connecticut

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The round whitefish is an inhabitant of cold, clear lakes and rivers. In rivers it occurs over rocky bottoms. In the Great Lakes it is commonly found inshore at depths of less than 36.6 m. It sometimes occurs in brackish waters. The round whitefish is a benthic insectivore, feeding on benthic invertebrates and occasionally fishes and fish eggs. Round whitefish require gravel for spawning. Since eggs incubate overwinter with no parental care, a silt-free spawning substrate probably is essential for successful recruitment as for other salmonids.

Habitat Types:

Aquatic: Fluvial

Current Threats

Habitat Threats:

Habitat Alteration

Climate Change

Sedimentation

Invasion by Exotic Species

Description of habitat threat(s): This species was once common to many of New York's Adirondack lakes. Recent surveys have shown its distribution there significantly reduced. The New York DEC has speculated on the possible causes: "predation by invading yellow perch on whitefish eggs and fry; predation by smallmouth bass; competition with lake whitefish; over fishing; loss of spawning sites; siltation; and lake acidification". Given the current and past locations in Vermont, only the following causes are possible for Vermont lakes: predation by yellow perch on whitefish eggs and fry (in lakes Seymour and Willoughby); predation by smallmouth bass (Lake Seymour only); loss of spawning sites and siltation (lakes Seymour and Willoughby). Beaver Pond is a small remote pond with no human structures along the shore and only seasonal human use within the watershed. With a low alkalinity (about 4mg/l) lake acidification, however, exists as a potential problem to that population. pH values for beaver Pond have been observed to drop well below 6.0 during spring runoff. Since this species is a benthic insectivore it relies entirely on invertebrates on the bottom as a food source. Loss or reduction of this food through sedimentation would limit its existence. Sedimentation may also limit egg survival by reducing oxygen exchange with surrounding water. Eggs incubate for months unguarded and unmaintained making a low sedimentation rate necessary for egg survival. Loss of spawning sites through siltation is also a threat. Climate change could result in the warming of the coldwater thermal regime that round whitefish require.



Common Name: **Round Whitefish**
 Scientific Name: **Prosopium cylindraceum**
 Species Group: **Fish**

Non-Habitat Threats:

Competition
 Pollution
 Predation or Herbivory
 Loss of Prey Base

Description of non-habitat threat(s): Beaver Pond is a small remote pond with no human structures along the shore and only seasonal human use within the watershed. With a low alkalinity (about 4mg/l) lake acidification is a potential threat to that population. Acidification would first impact newly hatched eggs, since this stage is normally the most vulnerable to acidity. Multiple year class failures would result in reductions or possible extermination of the species. Predation on whitefish eggs and fry by yellow perch in lakes Seymour and Willoughby and by smallmouth bass in Lake Seymour is a threat to these populations. Lake whitefish may be a competitor with round whitefish.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	
Research	Threats and Their Significance	High	Identify potential limiting factors.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Round Whitefish**
Scientific Name: **Prosopium cylindraceum**
Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Legislation	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, USFWS, NRCS, TNC, CRWC	
Invasive Species Control & Prevention	High	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, VFWD	VDEC (ANCG)
Policy & Regulations	High	Currently under VFWD fishing regulations Round Whitefish is an unregulated activity. Whether or not harvest poses a threat to Round Whitefish population(s) should be reviewed.	Review was conducted and recommendations were considered.	VFWD	VFWD (SWG, DJ)
Species Restoration	High	Develop and implement a plan for restoring Round Whitefish to additional waters that have the habitat and fish community makeup to sustain whitefish populations.	Components of the restoration plan that are implemented within the timeframe of the current WAP.	VFWD, USFWS	SWG, DJ, USFWS
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, NHDES, NHFGD, CRWC, NRCS, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Research	High	Investigate whether predation on Round Whitefish presents significant threat to the species.	Review was conducted and recommendations were considered.	VFWD, UVM	VFWD (SWG, DJ), UVM

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Round Whitefish**
Scientific Name: ***Prosopium cylindraceum***
Species Group: **Fish**

Bibliography

Evermann and Kendall. 1896. An annotated list of the fishes known from the State of Vermont. Appendix 8 Report of the U.S. Commissioner of Fish and Fisheries for 1894. Government Printing Office, Washington, D.C.

Gascon, D., and W. C. Leggett. 1977. Distribution, abundance, and resource utilization of littoral zone fishes in response to a nutrient/production gradient in lake Memphremagog. *Journal of the Fisheries Research Board of Canada* 34(8):1105-1117.

Gerardi, L. 2015. Personal communication. Vermont Fish and Wildlife Department, St. Johnsbury.

Greeley, J. R. 1930. Fishes of the Lake Champlain watershed. A Biological Survey of the Champlain watershed. State of New York Conservation Department. J.B. Lyons Company, Albany, pp. 44-87.

Kratzer, J. 2011. Round whitefish survey and inventory. Vermont Fish and Wildlife Department, State Wildlife Grant Final Report T-1-7 Job 2.37.

Marsden, J. E. 2014. Personal communication. Wildlife and Fisheries Biology Program, University of Vermont, Burlington.

Nakashima, B. S., and W. C. Leggett. 1975. Yellow perch (*Perca flavescens*) biomass responses to different levels of phytoplankton and benthic biomass in Lake Memphremagog, Quebec-Vermont. *Journal of the Fisheries Research Board of Canada* 32(10):1785-1797.

NYSDEC (New York State Department of Environmental Conservation). 2015a. Round whitefish restoration. NYSDEC, Albany. Available at: <http://www.dec.ny.gov/animals/26044.html>. (January 2015).

NYSDEC. 2015b. Round whitefish fact sheet. NYSDEC, Albany. Available at: <http://www.dec.ny.gov/animals/26013.html>. (January 2015).

Parrish, D. 2014. Personal communication. Vermont Fish and Wildlife Cooperative Research Unit, University of Vermont, Burlington.

Pientka, B. 2014. Personal communication. Vermont Fish and Wildlife Department, Essex.

Titcomb, J. W., and H. W. Bailey. 1896. Thirteenth Biennial Report of the Commissioners of Fisheries and Game of the State of Vermont, Argus and Patriot Company, Montpelier.



Common Name: **Atlantic Salmon** (naturally reproducing populations in Lakes Champlain & Memphremagog)
Scientific Name: **Salmo salar**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S4

State Trend: Historic

Extirpated in VT? Yes

Regional SGCN? Yes

Assessment Narrative:

Whether or not landlocked Atlantic Salmon were endemic to Vermont is not at all clear. Thompson (1853) described salmon spawning runs in the state as follows: "They came up the Connecticut river [sic] about the 25th of April, and proceeded to the highest branches. Shortly after they appeared [emphasis added] in lake [sic] Champlain and the large streams which fall into it." In that salmon are described as "appearing" in the lake seems to suggest that adult fish are not present year round but rather seasonally as is characteristic of anadromous salmon. Thompson (1842) mentions salmon as being in Lake Champlain but does not make any distinction between the population as being anadromous and/or landlocked ecotypes. However, the State of Vermont Fish Commissioners (VFC 1892) state of landlocked salmon that it "is not known to be indigenous to any lakes in Vermont" and that "Atlantic salmon...entered Lake Champlain by way of the St. Lawrence and Richelieu rivers during their migratory period for the purpose of spawning..." As a consequence of the construction of dams and other obstructions anadromous salmon were prevented access to the lake, and subsequently landlocked salmon were introduced (VFC 1892). Greeley (1930) identified landlocked salmon in Lake Champlain as an introduced species. On the other hand, Behnke (2007) offers the opinion that landlocked salmon were native to Lake Champlain. Likewise, by all accounts, landlocked salmon were not indigenous to Lake Memphremagog but were introduced from Maine (Grand Lake, East Grand lake or Sebago strains) in the late 1800s (Gerardi 2015). As a result of stocking, landlocked salmon became naturalized in the Clyde River and established a vibrant fishery, based at least in part on natural reproduction until it collapsed during the 1940s as a result of expanded hydroelectric generation on the river (Gerardi 2015). Although not substantiated, anadromous Atlantic Salmon may have had access to Lake Memphremagog and its tributaries prior to European colonization and damming of the St. Francis River at Drummondville, Quebec (Gerardi 2015). Without regard to which ecotype was native to Lake Champlain and Lake Memphremagog threats to future restored populations are similar to both: dams and water pollution, stream acidification, sedimentation of spawning and holding habitats, and possible changes in water temperature have been attributed to declines of some populations; landlocked populations have also suffered from northern pike and/or *Esox* hybrid predation (NatureServe 2014). With the appearance of Alewife (*Alosa pseudoharengus*) in Lake Champlain salmon have become more reliant on this species as prey. This may result in thiamine deficiency and consequently lead to salmon reproductive failure otherwise known as early mortality syndrome of salmon fry (Chipman 2015).

Distribution

Landlocked populations of Atlantic salmon occur principally in Newfoundland, Labrador and Quebec (Scott and Crossman 1973). Natural populations were also located in Maine and presumably in Vermont, i.e. lakes Champlain and Memphremagog. Scarola (1973) reports all landlocked salmon populations in New Hampshire are introduced. Landlocked populations have been stocked extensively to supplement natural populations, as well as to establish new fisheries. Even though salmon still occur in Lake Champlain, this is principally the result of a stocking program designed to restore a naturally reproducing population to the watershed, albeit there is little evidence of significant natural reproduction occurring there at this time. The Lake Memphremagog population does reproduce naturally with spawning occurring in the Clyde River.

Distribution by Biophysical Region:

Champlain Valley Historic Record(s) Only

Southern VT Piedmont Not Probable

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Atlantic Salmon** ((naturally reproducing populations in Lakes Champlain & Memphremagog)
Scientific Name: **Salmo salar**
Species Group: **Fish**

Champlain Hills	Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

St. Francois River

Probable Watersheds

Lake Champlain

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

The landlocked salmon is a fish of oligotrophic waters, that is deep, well oxygenated, relatively infertile lakes. As the spawning season sets in fish leave the depths of the lake to suitable spawning habitat located in shallow lake shore areas or to ascend tributary streams or use lake outlets (Scarola 1973). Habitat studies conducted in Lake Winnepesaukee, New Hampshire found the salmon have a preference for depths below the thermocline during the summer period (12-21 m) (Scarola 1973). Water temperatures at this time of year are in the mid 10s (C). Salmon can survive water temperatures into the 21s for brief periods of time but such temperatures are usually avoided if cooler water is available. Longer exposure to these temperatures can cause the fish physiological stress. Temperatures into the 27s can be lethal. Spawning habitat preferences are similar to those for anadromous Atlantic salmon.

Habitat Types:

Aquatic: Fluvial

Aquatic: Lower CT River

Current Threats

Habitat Threats:

Habitat Alteration

Climate Change

Sedimentation

Habitat Fragmentation

Invasion by Exotic Species

Description of habitat threat(s): Salmon require cool streams for spawning and juvenile rearing. Many of the historical salmon streams in Vermont have barriers that impede their access to these habitats (e.g., Missisquoi, Lamoille, Winooski, Clyde rivers and Otter Creek). Dams and other impassable barriers, such as culverts, also fragment habitats that are needed to access spawning and rearing habitats but also for seasonal movements (e.g., summer feeding, temperature refugia, overwintering habitats). Water quality degradation, such as in mean annual temperatures, have occurred due to losses in mature riparian canopies, impacts from land use practices, urban stormwater runoff, and water retention in impoundments. Stream habitats have been degraded and habitat complexity decreased from channelization and removal and

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Atlantic Salmon** (naturally reproducing populations in Lakes Champlain & Memphremagog)
Scientific Name: **Salmo salar**
Species Group: **Fish**

reduced recruitment of large woody debris to rivers. Habitat degradation from a variety of stressors has reduced the quality of spawning and juvenile rearing habitats in Vermont streams. The appearance of alewife in Lake Champlain is of great concern due to their potential impact through competition with other pelagic planktivores, such as landlocked Atlantic salmon.

Non-Habitat Threats:

Genetics

Competition

Parasites

Description of non-habitat threat(s): The native stocks of landlocked salmon in Lake Champlain were extirpated by the mid 1800s. It is questionable whether past and current restoration efforts are using genetically appropriate strains for restoration given the habitat conditions present in the lake. Competition and displacement by introduced salmonids, such as rainbow trout, may impact juvenile rearing stages of salmon. Sea lamprey parasitism has been found to be a significant source of mortality affecting salmon restoration.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Low	
Research	Threats and Their Significance	High	Impacts of Alewife consumption on thiamine levels in salmon and reproduction. Medium priority: inter-specific non-native salmonid competition.
Research	Population Genetics	High	Evaluation of strains being stocked in Lake Champlain for restoration purposes.
Research	Taxonomy	Low	
Research	Other Research	High	Accelerate the development of Lake Champlain salmon strain(s) which are more appropriate to the environmental conditions of those waters.
Monitoring	Population Change	High	
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Atlantic Salmon** (naturally reproducing populations in Lakes Champlain & Memphremagog)
 Scientific Name: **Salmo salar**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Invasive Species Control & Prevention	Medium	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP, TU	VDEC (ANCG)
Natural Processes Restoration	High	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, TU, dam owners	Dam owners
Natural Processes Restoration	High	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, TU, dam owners	Dam owners
Species Restoration	Medium	Evaluate strains of salmon to be used for species restoration with focus on identifying ones best adapted to the environments where restoration is to occur and have desirable survival and growth characteristics.	Assess strains.	VFWD, USFWS, NYDEC	USFWS, VFWD, NYDEC, GLFC
Invasive Species Control & Prevention	Medium	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP, TU	VDEC (ANCG)
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, TU, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Atlantic Salmon** (naturally reproducing populations in Lakes Champlain & Memphremagog)
Scientific Name: **Salmo salar**
Species Group: **Fish**

Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, TU, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Species Restoration	High	Implement salmon restoration plan in Lake Champlain.	Components of the restoration plan that are implemented within the timeframe of the current WAP.	VFWD, USFWS	SWG, DJ, USFWS
Legislation	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, TU	
Invasive Species Control & Prevention	Medium	Adopt/implement appropriate actions that minimize the potential for new invasive species (including pathogens) introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	Enforcement of importation regulations.	VFWD	VFWD (DJ)
Species Restoration	High	Adopt/implement appropriate actions that minimize the potential for new invasive species (including pathogens) introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail. Reduce Sea Lamprey woun	Enforcement of importation regulations. Reduction of Sea lamprey wounding rates.	VFWD, USFWS, NYDEC	USFWS, VFWD, NYDEC, GLFC

Bibliography

- Behnke, R. J. 2007. About trout: the best of Robert J. Behnke from Trout Magazine. The Lyons Press, Guilford, Connecticut. pp. 109-116.
- Chipman, B. 2015. Personal communication. Vermont Fish and Wildlife Department, Essex.
- Gerardi, L. 2015. Personal communication. Vermont Fish and Wildlife Department, St. Johnsbury.
- Greeley, J. R. 1930. Fishes of the Lake Champlain watershed. A Biological Survey of the Champlain watershed. State of New York Conservation Department. J.B. Lyons Company, Albany, pp. 44-87.
- NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).
- Thompson, Z. 1853. Natural history of Vermont. Charles E. Tuttle Company, Rutland, Vermont.
- Vermont Fish Commissioners. 1892. Eleventh biennial report of the fish commissioners of the State of Vermont. Argus and Patriot Book and Job printing, Montpelier.



Common Name: **Brook Trout (naturally reproducing populations)**
Scientific Name: **Salvelinus fontinalis**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S5

State Trend: Stable

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Brook Trout is the most widely distributed and abundant salmonid species indigenous to Vermont. Today, wild, stocked and mixed populations occur throughout the state wherever suitable habitat exist. Brook Trout is sensitive to pollution and habitat degradation. Deforestation, water development projects, pollution, habitat degradation, and competition and/or predation from introduced fishes have all taken a toll on wild populations throughout its range. Beginning around the mid-1800s and continuing up to present time Brook Trout have been cultured in hatcheries and stocked to restore fishable populations to many waters throughout the state. Fish culture programs have relied upon brook trout strains originating from multiple wild genetic sources most of which are populations located outside of Vermont. Strains may very well have been selected for characteristics other than their adaptability and survival in Vermont environments but rather for how they perform in hatcheries (e.g. growth and disease resistance) and sport fishing qualities. The ability of hatchery trout to survive in Vermont waters beyond the first season after release has been evaluated with results indicating multi-season survival is generally poor. In many respects our established self-sustaining wild Brook Trout populations may have certain advantages over traditional hatchery strains in certain situations, such as greater adaptability to Vermont environments. In recent years other states within the natural range of the species (e.g. New York) have been investigating wild populations to identify heritage (native) brook trout strains. Research into the existence of so-called heritage populations has not been pursued in Vermont. Extensive stocking of hatchery strains on top of wild populations throughout the Vermont over the past 150 or so years raises doubts as to whether it is possible to identify within acceptable certainty truly native populations. Therefore, the fallback position taken by the Vermont Fish & Wildlife Department is conservation of all wild naturally reproducing Brook Trout population in the state. Threats to Brook Trout populations include loss and degradation of habitats resulting from channel alterations, artificial flow regimes, water pollution, habitat fragmentation by dams and other obstructions (e.g. culverts), reduction of riparian vegetation resulting in stream water temperature increases and loss of instream fish cover (e.g. large wood). Spawning habitat and trout egg and fry survival are negatively affected by sedimentation. Climate change is predicted to increase water temperatures at the expense of populations at state as well as continental range levels. Climatic warming will likely decrease thermally suitable summer habitat for lotic brook trout populations Meisner (1990). Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

The brook trout is native to most of eastern Canada from Newfoundland to west of Hudson Bay; south in the Atlantic, Great lakes and Mississippi River basins to Minnesota and through the Appalachian Mountains to Georgia (Page and Burr 1991). It has been introduced widely outside of its natural range. In Vermont, the species is distributed throughout the state where suitable habitat is available and competing species are absent or low in abundance.

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Certain
Northern Green Mtns	Confident	Southern Green Mtns	Confident

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Brook Trout (naturally reproducing populations)**
Scientific Name: **Salvelinus fontinalis**
Species Group: **Fish**

Northern VT Piedmont Confident

Taconic Mtns Certain

Northeastern Highlands Confident

Distribution by Watershed:

Known Watersheds

West

Waits

Upper Connecticut-Mascoma

Black-Ottawquechee

Deerfield

Hudson-Hoosic

Lamoille River

Missisquoi River

Otter Creek

Passumpsic

St. Francois River

Upper Connecticut

White

Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Brook trout occur in headwater streams, small to medium rivers, and lakes and ponds where waters are clear, cool and well-oxygenated. Raleigh (1982) gives a comprehensive review of brook trout habitat requirements. Riverine brook trout habitat is characterized by being influenced by cold-springs, having silt-free rocky substrate in riffles and runs, well vegetated stream banks, abundant instream cover, and relatively stable stream flows, temperatures and stream banks. Lacustrine habitats are typically oligotrophic in character. The temperature range for brook trout is 0-24 C, with optimal temperatures for growth and survival in the range of 11-16 C. Warm water temperatures appears to be the single most critical factor influencing brook trout survival and production. Brook trout normally require high dissolved oxygen concentrations, optimally near saturation or ≥ 7 mg/L at temperatures ≤ 15 C and ≥ 9 mg/L at temperatures ≥ 15 C. Instream and riparian cover is recognized as an important component of brook trout habitat. Brook trout tends to be more tolerant of low pH water than other salmonid species. Most spawning occurs in stream habitat, although brook trout may spawn directly in lakes and ponds where there are upwellings.



Common Name: **Brook Trout (naturally reproducing populations)**
Scientific Name: **Salvelinus fontinalis**
Species Group: **Fish**

Habitat Types:

Aquatic: Fluvial

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Habitat Alteration

Climate Change

Sedimentation

Habitat Fragmentation

Description of habitat threat(s): Brook trout habitat has been degraded by alterations of natural stream channel morphology and flow regimes; water pollution; fragmentation (e.g., dams and culverts); reduction of riparian vegetation resulting in stream water temperature increases and loss of instream cover (e.g., large woody debris) Spawning habitat and trout egg and fry survival are negatively affected by sedimentation. Climate change could potentially degrade temperature regimes required by brook trout throughout its distribution.

Non-Habitat Threats:

Genetics

Competition

Disease

Pollution

Predation or Herbivory

Description of non-habitat threat(s): Brook trout compete poorly with introduced salmonids, such as brown and rainbow trout, as well as warmwater species (e.g., centrachids, percids and esocids). Additionally, these competitors may prey upon brook trout. Stocking of non-native brook trout strains on wild populations may result in inbreeding, loss of genetic characteristics necessary for species survival, and intra-specific competition. Stocking also puts heritage populations at risk of introducing disease causing pathogens, such as whirling disease.

Bibliography

Kirn, R. 2015. Personal communication. Vermont Fish and Wildlife Department, Roxbury.

Meisner, J. D. 1990. Potential loss of thermal habitat for brook trout, due to climatic warming, in two southern Ontario streams. Transactions of the American Fisheries Society 119:282-291.



Common Name: **Lake Trout (naturally reproducing populations)**
Scientific Name: **Salvelinus namaycush**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S4

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Eighteen lakes in Vermont are currently being managed for Lake Trout fisheries. With a few exceptions more than half of these populations exhibit little to no natural reproduction and therefore stocking hatchery-reared fish is necessary to maintain fishing quality. Presently, eight lakes (up from six reported in the 2005 WAP), all located in the northeastern part of the state, have populations that exhibit significant natural reproduction: Big and Little Averill lakes, Averill; Maidstone Lake, Maidstone; Echo Lake, Charlestown; Caspian lake, Greensboro; Willoughby Lake, Westmore; Seymour Lake Morgan; Crystal Lake, Barton; two other populations, Eligo Lake in Greensboro-Craftsbury and Shadow Lake in Glover, also have been demonstrated support wild reproduction but currently considered too low to maintain Lake Trout fisheries without stocking (Gerardi 2015). Despite natural reproduction occurring in Willoughby and Seymore lakes, these populations are supplemented with stocked Lake Trout, although this contribution is minor in comparison to the total population at large, estimated to be in the range of 5 to 30% (Gerardi 2015). Maidstone Lake has populations representing two distinct morphs (i.e. common and lunge) (Baille et als. In press). Historically, Lake Champlain supported a wild population, which subsequently became extirpated. A restoration program is in progress and although Lake Trout reproduction has been documented (Ellrott and Marsden 2004) a fully self-sustaining population without stocking has yet to be established. Restored naturally reproducing and self-sustaining Lake Trout populations within the species natural range warrant special conservation attention. Lake Trout populations are susceptible to habitat alteration or degradation including lake eutrophication, oxygen depletion in the hypolimnion, spawning shoal sedimentation, and lake level manipulation. Climate change is also a concern in that it may alter the temperature regimes of habitats adversely affecting wild Lake Trout populations as well as favoring other species (e.g. Smallmouth Bass) that may compete and/or prey upon trout. Year-class strengths of Lake Trout populations are projected to weaken, and that of Smallmouth Bass to strengthen as a consequence of water temperature increases during each species respective spawning season (Casselman 2002). Introduced species, such Alewife, are known to contribute to reproductive failure by disruption of thiamine metabolism in Lake Trout. Sea Lamprey parasitism on Lake Trout in the Great Lakes and Lake Champlain has been responsible for major population declines.

Distribution

The natural occurrence of lake trout is limited to North America. Its natural range closely aligns with the limits of the Pleistocene glaciation. In Vermont, the species has been stocked extensively throughout the state; however, with perhaps very few exceptions these populations are not self-sustainable and are completely dependent on continued stocking for the populations to exist. The few self-sustaining populations in Vermont are all located in the Northeast Kingdom: Big and Little Averill ponds, Averill; Maid stone Lake, Maidenstone; Echo Lake, Charleston; Caspian Lake, Greensboro; and Crystal Lake, Barton (L. Gerardi, Vermont Fish and Wildlife Department, personal communication). Natural reproduction has been documented in Lake Champlain, but that population is not at the present time sustainable without stocking.

Distribution by Biophysical Region:

Champlain Valley Historic Record(s) Only

Southern VT Piedmont Not Probable

Champlain Hills Not Probable

Vermont Valley Not Probable

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Lake Trout (naturally reproducing populations)**
Scientific Name: **Salvelinus namaycush**
Species Group: **Fish**

Northern Green Mtns	Certain	Southern Green Mtns	Not Probable
Northern VT Piedmont	Confident	Taconic Mtns	Not Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Known Watersheds

St. Francois River

Upper Connecticut

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

Lake trout is a species of oligotrophic lakes. In the southern part of their range the species inhabits deep, cold lakes whereas at more northern latitudes they are also found in shallow and deep waters. This species is intolerant of waters with low oxygen content, and prefers cold water, seeking areas with temperatures below 16 C (Langdon et al. In press). Lake trout spawn over rocky shoals and along wave-swept shorelines.

Spawning in riverine habitat is rare. Lake trout can spawn in depths of water from .3-61 m over gravel that measures 3.8-10.2 cm in diameter, typically aggregating in the fall over clean substrate, with deep interstitial spaces (Langdon et al. In press). Lake trout prefer eating small crustaceans, insects, and fish. Young lake trout eat plankton, insects, and small aquatic invertebrates.

Habitat Types:

Aquatic: Lacustrine

Aquatic: Fluvial

Aquatic: Lower CT River

Current Threats

Habitat Threats:

Habitat Alteration

Climate Change

Sedimentation

Habitat Fragmentation

Invasion by Exotic Species

Description of habitat threat(s): Lake trout are susceptible to habitat degradation, such as lake eutrophication, hypolimnion oxygen depletion, spawning shoal sedimentation, lake level manipulations. Introduced species, such as alewives, are known to contribute to reproductive failure by disruption of thiamine metabolism in lake trout. Sea lamprey in Lake Champlain predate on lake trout and may be a significant factor in population declines there. Dams constructed at lake outlets raise lake levels that can erode shoreline soils, increasing sedimentation of spawning shoals and decreasing reproductive success. Water level fluctuations, associated with water storage for hydropower, may result in dewatered lake trout



Common Name: **Lake Trout (naturally reproducing populations)**
Scientific Name: **Salvelinus namaycush**
Species Group: **Fish**

egg and embryos, also negatively impacting reproductive success.

Non-Habitat Threats:

Genetics

Parasites

Harvest or Collection

Description of non-habitat threat(s): Lake Maidstone contains a population of lake trout (a.k.a. locally as lunge) that is morphologically distinct from lake trout in all other Vermont lakes. It is possible that this population is genetically distinct and represents a unique population that is native to Vermont. Because it is unknown whether this population is genetically distinct, no strategies exist to protect this population. Lake trout are highly valued by anglers due to the large size that this species can attain. Overfishing could result in the loss of self-sustaining, native lake trout populations (Ellrott and Marsden 2004).

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Low	
Research	Threats and Their Significance	Low	
Research	Population Genetics	Low	Investigated: see Baille et al. In press. Deciphering hatchery stock influences on wild populations of Vermont Lake Trout <i>Salvelinus namaycush</i> . Transactions of the American Fisheries Society, Bethesda, Maryland.
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	
Monitoring	Habitat Change	High	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Lake Trout (naturally reproducing populations)**
Scientific Name: **Salvelinus namaycush**
Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Invasive Species Control & Prevention		Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)
Invasive Species Control & Prevention		Adopt/implement appropriate actions that minimize the potential for new invasive species (including pathogens) introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	Enforcement of importation regulations.	VDEC, USFWS, LCBP	VDEC (ANCG)
Natural Processes Restoration		Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, dam owners	Dam owners
Habitat Restoration		Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, TU, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Research		Evaluate strains of Lake Trout to be used for species restoration with focus on identifying ones best adapted to the environments where restoration is to occur and have desirable survival and growth characteristics.	Assess strains.	VFWD, USFWS, NYDEC	USFWS, VFWD, NYDEC, GLFC
Natural Processes Restoration		Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, dam owners	Dam owners

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Lake Trout (naturally reproducing populations)
Scientific Name: *Salvelinus namaycush*
Species Group: Fish

Invasive Species Control & Prevention		Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN. Reduction in Sea Lamprey wounding rates observed on Lake Sturgeon.	VFWD, VDEC, USFWS, TU	VDEC (ANCG)
Legislation		Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP	
Species Restoration	High	Implement Lake Trout restoration plans in Lake Champlain. Undertake appropriate management actions as needed to conserve and enhance populations in other lakes in Vermont.	Components of the restoration plan that are implemented within the timeframe of the current WAP.	VFWD, NYDEC, USFWS	SWG, DJ, USFWS
Invasive Species Control & Prevention	High	CLIPPED	Enforcement of importation regulations. Sea lamprey wounding rates.	VFWD, VDEC, USFWS, NYDEC	USFWS, VFWD, NYDEC, GLFC

Bibliography

Baille, S. M. et als. In press. Deciphering hatchery stock influences on wild populations of Vermont lake trout *Salvelinus namaycush*. Transactions of the American Fisheries Society.

Casselman, J. M. 2002. Effects of temperature, global extremes, and climate change on year-class production of warmwater, coolwater, and coldwater fishes in the Great Lakes basin. Pages 39-60 in N. A. McGinn, editor. Fisheries in a changing climate. American Fisheries Society Symposium 32, Bethesda, Maryland.

Ellrott, B. J., and J. E. Marsden. 2004. Lake trout reproduction in Lake Champlain. Transactions of the American Fisheries Society 133:252-264.

Gerardi, L. 2015. Personal communication. Vermont Fish and Wildlife Department, St. Johnsbury.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Redfin Pickerel**
Scientific Name: **Esox americanus**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S4

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

Seven populations have been identified in Vermont (Langdon 2014). On a range-wide scale no major threats are known (NatureServe 2014); however, interspecies hybridization has been reported (*Esox americanus* X *E. niger* and *E. americanus* X *E. lucius*) which could pose a threat to populations where the Redfin Pickerel is the only resident esocid in the fish community and whether by intention or accident another esocid species might be introduced. Redfin Pickerel is listed as endangered in Maine (MDIFW 2013).

Distribution

The redfin pickerel (*Esox americanus americanus*) is the subspecies restricted to eastern United States; the grass pickerel (*E. a. vermiculatus*) is the western subspecies. Redfin pickerel is a fish primarily associated with the eastern coastal plain. It is distributed from the St. Lawrence River (Lac St. Pierre) south through the Richelieu-Champlain-Hudson system into New York, east through southern Vermont and New Hampshire to Long Island and south along the coastal plain to Georgia (Scott and Crossman 1973). In Vermont, redfin pickerel appears to be limited to the Poultney-Castleton rivers drainage, South Fork of East Creek in Orwell, and Pond Brook in Monkton.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Confident
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The redfin pickerel occurs in weedy areas of lakes, ponds and slow rivers. The redfin is often found in tea-colored, acidic waters with pH values as low as 4.3. The redfin pickerel spawns during April and May when water temperatures reach 10 C (Langdon et al. In press). Adults congregate to spawn and adhesive eggs are broadcast randomly in heavily vegetated, shallow areas along lakeshores or streambanks. Young redfins first consume zooplankton, snails and crustaceans, switching to fish as they grow older. Adults feed on fish and crayfish, but may supplement these food items with small crustaceans and insects (Jenkins and Burkhead 1993). This species may play a significant role in fish community structure because of its preference to predate



Common Name: **Redfin Pickerel**
Scientific Name: **Esox americanus**
Species Group: **Fish**

on fish.

Habitat Types:

Aquatic: Lacustrine

Aquatic: Fluvial

Current Threats

Habitat Threats:

Habitat Alteration

Description of habitat threat(s): Loss or alteration (i.e. through flow alteration, dewatering, sedimentation) of vegetated, shallow areas along lakeshores or streambanks would pose a problem to the reproductive success of the species.

Non-Habitat Threats:

Genetics

Description of non-habitat threat(s): This species has a limited distribution in Vermont, located in 3 drainages. If any of these populations undergo a reduction in population size (i.e., population bottleneck), then it is possible that genetic variation will be lost, forfeiting the evolutionary potential of the species. Natural selection can only act in the presence of genetic variation, and, therefore, the higher the genetic variability in a population, the higher the likelihood for population persistence. If gene flow between the 3 populations is limited, then the genetic variability of each population could decrease over time. Also, redfin pickerel have been reported to hybridized with northern pike and chain pickerel (Jenkins, 1993), which poses further genetic concerns. The introduction of northern pike or chain pickerel to waters populated by redfin pickerel poses the risk of hybridization, as well as introduce inter-specific predation and competition for habitat and forage. Decreased genetic variation in redfin pickerel would hinder the ability of the populations to adapt to changing conditions over time.

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Redfin Pickerel**
Scientific Name: **Esox americanus**
Species Group: **Fish**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Identify number of redfin pickerel populations in Vermont and those which are not co-habitants with other esocid species.
Research	Threats and Their Significance	Medium	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	Medium	
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	The introduction of northern pike and chain pickerel to redfin pickerel waters limits the species by hybridization and should be monitored.
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Redfin Pickerel**
 Scientific Name: **Esox americanus**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Invasive Species Control & Prevention	High	Identified populations which are not sympatric with other Esocid species should be managed to prevent to potential for interspecific competition and hybridization.	Number (percentage) of redfin pickerel waters remaining free of competing esocid species.	VDEC, VFWD, lake associations	VDEC (ANCG), VFWD (SWG, DJ)
Natural Processes Restoration	Medium	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, dam owners, lake associations	Dam owners, lake associations
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Invasive Species Control & Prevention	Medium	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, VFWD, lake associations	VDEC (ANCG), VFWD (SWG, DJ)
Invasive Species Control & Prevention	High	Protect habitats currently supporting Redfin Pickerel populations from the introduction of other esocid species (e.g., Chain Pickerel, Northern Pike) which may compete for available habitat and/or potentially hybridize with Redfin Pickerel.	Number (percentage) of redfin pickerel waters remaining free of competing esocid species.	VDEC, VFWD, lake associations	VDEC (ANCG), VFWD (SWG, DJ)
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

***Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report***



Common Name: **Redfin Pickerel**
Scientific Name: **Esox americanus**
Species Group: **Fish**

Bibliography

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.

MDIFW (Maine Department of Inland Fisheries and Wildlife). 2013. State list of endangered species. Available at: http://www.maine.gov/ifw/wildlife/endangered/listed_species_me.htm. (November 2014).

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).



Common Name: **Bridle Shiner**
 Scientific Name: **Notropis bifrenatus**
 Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G3

Global Trend: Unknown

State Rank: S1?

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Seven populations are known to occur in Vermont (Langdon 2014), although two or more of these are outdated reports (Hospital Creek, 1935; Lake Memphremagog, 1975). Bridle shiners were captured in the Canadian portion of Lake Memphremagog in 1999 (COSEWIC 2013). The species was also collected in the Connecticut River in the vicinity of Hanover, New Hampshire in 2008 (Yoder et al. 2010). Scarola (1987) states that bridle shiner in New Hampshire only occurs in the Merrimack and Coastal drainages, therefore the origin of the species in the upper Connecticut River is unknown but is presumed to be native. The species was frequently encountered in the Richelieu River watershed between 1965 and 1970; in 1990 six were collected in Missisquoi Bay near its mouth but were not captured during sampling efforts done in 2003 and 2012 (COSEWIC 2013). Threats to populations have been identified as reduction or removal of aquatic vegetation and increased turbidity, including agricultural pollution, as well as predation from introduced fish species (NatureServe 2014). Bridle Shiner is also potentially at risk to climate change (COSEWIC 2013). While population status and specific threats acting on Vermont populations are unknown, until water quality issues identified for Lake Champlain including Missisquoi Bay (LCBP 2012) are corrected, efforts for the population to recover to a more secure status may not be possible. Bridle shiner is listed as endangered in New Hampshire and Pennsylvania; special concern in Massachusetts, Connecticut, Ontario and Quebec; and extirpated in Maryland. Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

This North American species occurs in the Atlantic drainage from southern Maine to Virginia, west through Lake Champlain to New York and the Lake Ontario basin (Scott and Crossman 1973). In Vermont, bridle shiner are found in Lake Champlain, several other locations within that watershed, and has been reported to occur in Lake Memphremagog.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Confident
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Historic Record(s) Only	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain

Probable Watersheds

Missisquoi River



Common Name: **Bridle Shiner**
Scientific Name: **Notropis bifrenatus**
Species Group: **Fish**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

The bridle shiner is a species of quiet streams, lakes and ponds (Scott and Crossman 1973, Page and Burr 1991). Honeyfield and Ross (2004) describe its habitat as slow, warm backwater eddies of low gradient streams and ponds with dense vegetation and substrate of mud, sand or gravel. Spawning occurs in areas of calm water, at a depth of about two feet, and in openings within stands of dense emergent aquatic vegetation (Cornell web site). Holms et al. (1999) suggest bridle shiner require open water above aquatic plant stands to spawn. The species has a strong preference for clear water necessary for this daytime, sight-feeder to forage on prey organisms (Honeyfield and Ross 2004). It is tolerant of brackish water but is not acid tolerant (Holm et al. 1999).

Habitat Types:

Aquatic: Lacustrine

Aquatic: Fluvial

Aquatic: Lower CT River

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Habitat Alteration

Sedimentation

Habitat Fragmentation

Invasion by Exotic Species

Description of habitat threat(s): Threats include stream channelization, erosion, point-source discharges, loss of riparian vegetation, and large-scale development all have been identified as practices that have been identified as practices that have increased turbidity and altered bridle shiner habitat where the species have become extirpated (Sabo 2000). Sedimentation and siltation are suspected as being major factors for the decline of bridle shiner in the Delaware River Basin (Honeyfield and Ross 2004). They also suggest the species may be vulnerable to highway construction activities which alter bridle shiner habitat including streamflow regimes, channel structure, water quality, and aquatic plant abundance. Being a sight-feeder turbid water conditions interfere with bridle shiner feeding and suppresses the growth of aquatic vegetation on which the fish is dependent for feeding, reproduction and cover (Holm et al. 1999). They also identify the spread of Eurasian milfoil as a potential problem to the species. This plant can alter the composition of the plant community by replacing native vegetation and invading the entire water column thereby eliminating clear water areas above the plants that are necessary for spawning (Sabo 2000).

Non-Habitat Threats:

Genetics

Pollution

Predation or Herbivory



Common Name: **Bridle Shiner**
 Scientific Name: **Notropis bifrenatus**
 Species Group: **Fish**

Description of non-habitat threat(s): Scott and Crossman (1973) state bridle shiner are an important forage species for pickerel as well as other piscivorous fishes (yellow perch, smallmouth bass, crappie) where these species co-exist. In some New England lakes and ponds, where significant reduction or removal of submersed aquatic plant stands has occurred, fish predators (e.g., bass) have decimated bridle shiner populations (Sabo 2000). Honeyfield and Ross (2004) state, that within the species range populations appear to be highly fragmented and declining with separation distances between known populations exceeding 200 km. They suggest this may have resulted in genetic divergence among populations, although this has not been investigated. This species is sensitive to sedimentation and chemical runoff from agricultural lands (Ontario's Biodiversity website). The bridle shiner is not tolerant of acidic water making it vulnerable to atmospheric deposition (Holm et al. 1999).

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	
Research	Basic Life History	Medium	
Research	Distribution and Abundance	High	Little quantitative data exists for Vermont populations; better distributional data is needed.
Research	Threats and Their Significance	High	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	Monitoring program is recommended to assess changes in species abundance and distribution.
Monitoring	Habitat Change	High	Known populations, particularly those exposed to aquatic plant control activities, are in need of monitoring.
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	Known populations, particularly those exposed to aquatic plant control activities, are in need of monitoring.
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Bridle Shiner**
 Scientific Name: **Notropis bifrenatus**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VFWD, VDEC, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Invasive Species Control & Prevention	High	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)
Natural Processes Restoration	High	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, VTrans, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners, VFWD (SWG), USFWS, VDEC, VTrans
Natural Processes Restoration	High	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners
Legislation	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Bridle Shiner**
 Scientific Name: **Notropis bifrenatus**
 Species Group: **Fish**

Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VFWD, VDEC, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

Bibliography

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2005. COSEWIC assessment and status report on the bridle shiner *Notropis bifrenatus* in Canada. COSEWIC, Ottawa. Available at: www.sararegistry.gc.ca/status/status_e.cfm. (November 2014).

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.

LCBP (Lake Champlain Basin Program). 2012. State of the lake and ecosystems indicators report. LCBP, Grand Isle, Vermont. Available at: http://sol.lcbp.org/introduction_what-is-state-of-lake-report.html. (December 2014).

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).

Scarola, J. F. 1987. Freshwater fishes of New Hampshire. New Hampshire Fish and Game department, Concord.

Yoder, C. O., L. E. Hersha, and B. R. Apell. 2010. Fish assemblage and habitat assessment of the upper Connecticut River: a preliminary report and presentation of data. Midwest Biodiversity Institute, MBI Technical Report MBI/2009-8-3, Columbus, Ohio.



Common Name: **Blackchin Shiner**
 Scientific Name: **Notropis heterodon**
 Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

Seven populations are known to occur in Vermont (Langdon 2014). Very little is known of its biology, distribution and habitat use, and past and present abundance of these populations. On a range-wide scale no major threats are known (NatureServe 2014). Blackchin Shiner has all but disappeared from southern New York watersheds; possible threats identified include fluctuating water levels and habitat loss due to siltation; species also has a low tolerance to salt (NYNHP 2013).

Distribution

This North American species occurs only in the Great Lakes basin, the upper Mississippi River drainage, and downstream through the St. Lawrence River drainage into western Quebec (Scott and Crossman 1973). In Vermont, the species on the eastern edge of its distribution and is limited to a few locations within the Lake Champlain watershed.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Confident
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Probable Watersheds

Lake Champlain
 Metawee River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Trautman (1957) describes the blackchin shiner as a fish of glacial lakes and streams characterized by having very clear water, substrate of clean sand, gravel or organic debris, and the presence of dense beds of submersed vegetation. Scott and Crossman (1973) also note the species' preference for quiet pools in streams and weedy inshore areas of lakes. This shiner is also found in inlet and outlet streams of lakes (Becker 1983). Scott and Crossman (1973) state "clear, clean, weedy waters are essential for the survival of the blackchin shiner," but little information has been reported on spawning site preference and behavior. This species is an indicator of good water quality.



Common Name: **Blackchin Shiner**
Scientific Name: **Notropis heterodon**
Species Group: **Fish**

Habitat Types:

Aquatic: Lacustrine

Aquatic: Fluvial

Aquatic: Lower CT River

Aquatic: Large Lake Champlain Tribs Below Falls

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Sedimentation

Invasion by Exotic Species

Description of habitat threat(s): The blackchin shiner is intolerant of silt. Trautman (1957) notes that it has disappeared from Ohio waters once they became turbid, silt covered the bottom, and the aquatic vegetation disappeared. The continuing expansion of the submersed exotic weed Eurasian milfoil in Vermont's lakes and ponds threatens blackchin shiner populations. Milfoil may displace native plant communities and alter the composition and fish habitat value of the aquatic plant community. Additionally the propensity for this invasive plant to establish dense beds impairing the use of some lakes for boating, swimming and fishing has resulted in the increased use of herbicides. Significant reduction in aquatic plant abundance could degrade habitat necessary for this species. One such herbicide in common use in Vermont is fluridone (Sonar). The Michigan Environmental Science Board has concluded that this broad spectrum herbicide will not only control Eurasian milfoil but also significantly impact native aquatic plant species when applied at the labeled rate (Premo et al. 1999). Significant reduction in milfoil beds can subject blackchin shiner populations to increased predation pressure before native aquatic plants become reestablished restoring cover habitat for the shiner.

Non-Habitat Threats:

Predation or Herbivory

Description of non-habitat threat(s): The blackchin shiner does not appear to be particularly abundant in any of the waters it is known to occur in Vermont. Aggressive aquatic plant control activities in these waters could significantly reduce this important refuge habitat and subject the shiner populations to increased predation.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Blackchin Shiner**
 Scientific Name: **Notropis heterodon**
 Species Group: **Fish**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	A greater understanding of the habitat requirements of this species is needed, especially the association with aquatic vegetation.
Research	Basic Life History	Medium	
Research	Distribution and Abundance	High	
Research	Threats and Their Significance	High	Impacts of aquatic plant control (e.g., herbicide and mechanical treatments) on Blackchin Shiner habitat, biology, and aquatic community structure and function (e.g., species interactions, increased vulnerability to predation). Investigate and monitor these effects.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	Known populations, particularly those exposed to aquatic plant control activities, are in need of monitoring.
Monitoring	Habitat Change	High	Known populations, particularly those exposed to aquatic plant control activities, are in need of monitoring.
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	Known populations, particularly those exposed to aquatic plant control activities, are in need of monitoring.
Monitoring	Other Monitoring Needs	N/A	

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Blackchin Shiner**
Scientific Name: **Notropis heterodon**
Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Invasive Species Control & Prevention	High	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VFWD, VDEC, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VFWD, VDEC, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

Bibliography

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).

New York Natural Heritage Program (NYNHP). 2013. Blackchin shiner (Notropis heterodon). NYNHP Conservation Guide, Albany. Available at: <http://www.acris.nynhp.org/animals.php>. (January 2015).



Common Name: **Blacknose Shiner**
Scientific Name: **Notropis heterolepis**
Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend: Unknown

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Six populations are known to occur in Vermont (Langdon 2014). Very little is known of its biology, distribution and habitat use, and past and present abundance of these populations. On a range-wide scale no major threats are known (NatureServe 2014). Blacknose Shiners require very clear water and moderate amounts of vegetation; the species quickly disappears from habitat that becomes turbid and substrate that is degraded by sedimentation (ODNR 2012).

Distribution

The blacknose shiner is a widely distributed species occurring from the Hudson Bay drainage to the New England states west to Iowa (Scott and Crossman 1973). In Vermont, it appear to be most frequently encountered in the Lake Champlain watershed, albeit there are a few occurrences from other locations within the state.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Confident	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain
Missiquoi River
Winooski River

Probable Watersheds

Black-Ottawquechee

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The blacknose shiner is a species of clear, shallow water habitats in glacial lakes and small streams (Scott and Crossman 1973). Most recorded occurrences in Vermont (11 out of 13 records) have come from riverine habitats (R. Langdon, Vermont Department of Environmental Conservation, personal communication). The species does not appear to have any particular preference for substrate type, although Trautman (1957) states it has been found in waters with bottoms comprised of sand, gravel, muck, peat or organic debris. The presence of submersed vegetation is an important component of blacknose shiner habitat (Page and Burr 1991,



Common Name: **Blacknose Shiner**
Scientific Name: **Notropis heterolepis**
Species Group: **Fish**

Trautman 1957). Backlund (1995) reports the fish requires cool well-oxygenated water. This species is intolerant of sedimentation and turbid water (Backlund 1995, Eddy and Underhill 1974, Trautman 1957). This species is an important indicator of pristine, high quality waters (Backlund 1995). The biology and detailed habitat requirements of the blacknose shiner apparently have either been not thoroughly investigated or reported. Backlund (1995) states that the blacknose shiner is a host fish for the freshwater mussel cylindrical papershell *Anodontoidea ferussacianus*, a state listed endangered species in Vermont.

Habitat Types:

Aquatic: Lacustrine

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Sedimentation

Invasion by Exotic Species

Description of habitat threat(s): Backlund (1995) states the blacknose shiner is intolerant of turbid, polluted waters and that in South Dakota it has disappeared from many streams due to sedimentation, loss of aquatic vegetation and food, water temperature increases, and lowered dissolved oxygen. Aquatic plant control also poses a problem to blacknose shiner populations. In Minnesota the removal of aquatic vegetation along lake shorelines and increase sedimentation and turbidity levels have reduced this species (Eddy and Underhill 1974).

Non-Habitat Threats:

Pollution

Predation or Herbivory

Description of non-habitat threat(s): The blacknose shiner is reported to be intolerant of water pollution (Backlund 1995). Elimination of aquatic plant beds can expose this species to increased predation.



Common Name: **Blacknose Shiner**
 Scientific Name: **Notropis heterolepis**
 Species Group: **Fish**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	A greater understanding of the habitat requirements of this species is needed, especially the association with aquatic vegetation.
Research	Basic Life History	Medium	
Research	Distribution and Abundance	High	Little quantitative data exists for Vermont populations; better distributional data is needed.
Research	Threats and Their Significance	High	Investigate and monitor the effects of aquatic vegetation control programs (e.g., Eurasian milfoil) on Blacknose Shiner populations.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	Known populations, particularly those exposed to aquatic plant control activities, are in need of monitoring.
Monitoring	Habitat Change	High	Known populations, particularly those exposed to aquatic plant control activities, are in need of monitoring.
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	Known populations, particularly those exposed to aquatic plant control activities, are in need of monitoring.
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Blacknose Shiner**
 Scientific Name: **Notropis heterolepis**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VFWD, VDEC, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Invasive Species Control & Prevention	High	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VFWD, VDEC, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

Bibliography

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).

ODNR (Ohio Department of Natural Resources). 2012. Blacknose shiner – Notropis heterolepis. ODNR Division of Wildlife, Columbus. Available at: <http://wildlife.ohiodnr.gov/species-and-habitats/species-guide-index/fish/blacknose-shiner>. (January 2015).

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Silver Redhorse**
Scientific Name: **Moxostoma anisurum**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S2

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

In Vermont, Silver Redhorse is confined to Lake Champlain and the lower accessible reaches of larger tributary rivers to the lake, e.g. Missisquoi River, Winooski River, Otter Creek and Poultney River (Ferguson 2014). The population is on the eastern edge of its continental range. Little is known of its distribution and population status; however, based on anecdotal observations it is but appears to be less abundant than Shorthead Redhorse (*M. macrolepidotum*) but more abundant than Greater Redhorse (*M. valenciennesi*) (Ferguson 2014; Langdon 2014). Threats include poor water quality resulting from agricultural and urban pollution, artificial flow regimes, and habitat fragmentation (Cook et al. 2005). Eaton and Scheller (1996) identify Silver Redhorse as potentially at risk from climate change. NatureServe (2014) identifies no known major threats to the species at the range-wide scale, but acknowledges that localized threats may exist. In Vermont, little is known of this population or threats to it. Because populations are not being monitored, trends influenced by threats may not be detected.

Distribution

A North American species which occurs east from the St. Lawrence River, south through central New York, southwest through the western half of the coastal states to northern Alabama, northwest through eastern Arkansas and Missouri, north through North Dakota, and into Saskatchewan and Alberta (Scott and Crossman 1973). In Vermont, the silver redhorse is on the eastern edge of its North American range and occurs only in Lake Champlain and up to the fall line in the Missisquoi and Poultney rivers.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain
Winooski River

Probable Watersheds

Lamoille River
Missisquoi River
Otter Creek



Common Name: **Silver Redhorse**
Scientific Name: **Moxostoma anisurum**
Species Group: **Fish**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The silver redhorse is found in small to moderately-sized rivers and occasionally lakes. In rivers it prefers deep pools with some current (Meyer 1962). During their first year of life silver redhorses sometimes remain in small streams where they were hatched. This species avoids silty bottoms and may also be intolerant to general environmental degradation (Langdon et al. In press). Adult silver redhorse perform annual migrations to spawn. Seasonal movement patterns may prove important for successful spawning. Silver redhorse are early spawners and in their southern range breed from April through early May in water temperatures of 11-15 C. Spawning usually occurs in shallow riffles over gravel and cobble (Jenkins and Burkhead 1993). Silver redhorse feed on insect larvae, microcrustaceans, mollusks, algae, detritus, crayfishes, and the fry of shiners (Jenkins and Burkhead 1993).

Habitat Types:

Aquatic: Lacustrine

Aquatic: Fluvial

Aquatic: Lower CT River

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Habitat Alteration

Sedimentation

Habitat Fragmentation

Description of habitat threat(s): Flow alteration, temperature alteration, or decreased habitat diversity (i.e. loss of deep pool habitat) will most likely have negative effects for different life stages of silver redhorse. Fragmentation of silver redhorse habitat may disrupt the seasonal movement patterns of this species. For example, these movement patterns may prove critical for successful reproduction, and therefore the completion of the species life cycle. Disruption to the spawning efforts of this species poses a problem to population viability (i.e. weak year classes over time compound negative influences and population declines). If the quantity or quality of silver redhorse habitat is limited in a system, then interconnected river reaches will prove necessary for this species to find and occupy optimal or suitable habitat. Loss of riparian vegetation, general construction activity, road maintenance activities (ditching, sanding), bridge and culvert construction, agriculture, timber harvest, dam failure, rapid drawdown of dam impoundments, streambank erosion, and shifts in channel form or location are sources of sediment for silver redhorse habitat. Controlling sediment input into streams may be crucial to prevent detrimental effects to silver redhorse, because sedimentation decreases the quality and quantity of optimal habitat (i.e. spawning, feeding) for this species. Sedimentation eliminates interstitial spaces which could be critical for egg deposition and development and for production of benthic organisms, such as aquatic insects, a source of food for silver redhorse. Sedimentation has been shown to cause loss or reduction in fish populations, and disrupt the feeding and reproductive activities of fishes (Berkman and Rabeni, 1987).

Non-Habitat Threats:

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Silver Redhorse**
Scientific Name: **Moxostoma anisurum**
Species Group: **Fish**

Pollution

Description of non-habitat threat(s): Water pollution may indirectly influence silver redhorse through negative impacts to its prey base. Depletion of food items will negatively affect species growth and survival.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Distribution, abundance and dynamics of Silver Redhorse populations in Vermont are poorly understood.
Research	Threats and Their Significance	Medium	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	Medium	
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	Medium	Sediment and pollution
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Silver Redhorse**
 Scientific Name: **Moxostoma anisurum**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Legislation	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP	
Research	Medium	Currently under VFWD fishing regulations Moxostoma species are "cull" fishes and as such their harvest and collection is essentially an unregulated activity. Whether or not Moxostoma are threatened by unrestricted harvest should be reviewed.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Natural Processes Restoration	Medium	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, dam owners	Dam owners
Natural Processes Restoration	Medium	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, dam owners	Dam owners



Common Name: **Silver Redhorse**
Scientific Name: **Moxostoma anisurum**
Species Group: **Fish**

Bibliography

Cooke, S. J. et al. 2005. Threats, conservation strategies, and prognosis for suckers (Catostomidae) in North America: insights from regional case studies of a diverse family of non-game fishes. *Biological Conservation* 121:317-331.

Eaton, J. G., and R. M. Scheller. 1996. Effects of climate warming on fish thermal habitats in streams of the United States. *Limnology and Oceanography* 41(5):1109-1115.

Ferguson, M. 2014. Personal communication. Vermont Fish and Wildlife Department, Montpelier.

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).



Common Name: **Shorthead Redhorse**
Scientific Name: **Moxostoma macrolepidotum**
Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S2

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

Shorthead Redhorse in Vermont is confined to Lake Champlain and the lower accessible reaches of larger tributary rivers to the lake (Langdon et al. 2006). It appears to be a widespread species within its Vermont range but based on anecdotal observations appears to be fairly abundant where found (Ferguson 2014; Langdon 2014). Threats include poor water quality resulting from agricultural and urban pollution, artificial flow regimes, and habitat fragmentation (Cook et al. 2005). NatureServe (2014) identifies no known major threats to the species at the range-wide scale, but acknowledges that localized threats may exist. In Vermont, little is known of this population or threats to it. Because populations are not being monitored, trends influenced by threats may not be detected.

Distribution

The shorthead redhorse is the most widely distributed of the redhorse species in North America. It occurs from the upper St. Lawrence River, south into the Lake Champlain drainage to the coast in New York, east of the Appalachian Mountains to South Carolina, west through Pennsylvania and Ohio, southwest into Indiana and Arkansas, the Tennessee River drainage in Alabama, west to Texas, northwest through eastern Colorado and Montana, north to central Alberta, east to southern Hudson Bay and the east shore of James Bay (Scott and Crossman 1973). In Vermont, the species is on the eastern edge of its North American range and is confined to Lake Champlain and several large tributaries up to the fall line.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Winooski River

Missiquoi River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

This species prefers the clear water of small to large rivers and sometimes lakes. Most individuals have been observed holding or feeding in deep pools. It is found over clean sand, gravel, and cobble substrate, and is



Common Name: Shorthead Redhorse
Scientific Name: *Moxostoma macrolepidotum*
Species Group: Fish

tolerant of water temperatures up to 37 C. It requires a silt-free habitat and is thought to be susceptible to many forms of water pollution (Langdon et al. In press). It is common to find this species living in the same areas as other redhorse species. In Vermont, the shorthead redhorse is restricted to the larger tributaries of Lake Champlain (Langdon et al. In press). The spawning period for shorthead redhorse occurs in spring from early April to early July, as influenced by local regional conditions (i.e. climate). Spawning water temperature is 11-21 C. Spawning occurs in slow and moderate runs and pools over large gravel (Jenkins and Burkhead 1993). Shorthead redhorse may perform spawning migrations to find optimal spawning habitat. Spawning groups of this species have been observed in streams where adults are normally not found except during breeding time (Jenkins and Burkhead 1993). Seasonal movement patterns may prove important for successful spawning. Shorthead redhorse is specialized to benthically feed on aquatic insects, small crustaceans, mollusks, algae, and detritus (Jenkins and Burkhead 1993). Highly silted or embedded substrate may preclude this species from consuming its preferred food items. Studies have shown that the principal, insect food items of redhorses are chironomids, ephemeropterans, and trichopterans (Meyer 1962).

Habitat Types:

Aquatic: Fluvial

Aquatic: Lower CT River

Current Threats

Habitat Threats:

Habitat Alteration

Sedimentation

Habitat Fragmentation

Description of habitat threat(s): Flow alteration, temperature alteration, or decreased habitat diversity (i.e. loss of deep pool habitat) will most likely pose negative effects for different life stages of shorthead redhorse. For instance, shallow, channel margin habitats that are indicative of slower velocities are important for young redhorses. Anthropogenic flow alteration has been shown to alter and limit this habitat, affecting juvenile life stages (Scheidegger and Bain, 1995). Fragmentation of shorthead redhorse habitat may disrupt the seasonal movement patterns of this species. For example, these movement patterns may prove critical for successful reproduction, and therefore the completion of the species life cycle. Disruption to the spawning efforts of this species poses a problem to population viability (i.e. weak year classes over time compound negative influences and population declines). If the quantity or quality of shorthead redhorse habitat is limited in a system, then interconnected river reaches will prove necessary for this species to find and occupy optimal or suitable habitat. Loss of riparian vegetation, general construction activity, road maintenance activities (ditching, sanding), bridge and culvert construction, agriculture, timber harvest, dam failure, rapid drawdown of dam impoundments, streambank erosion, and shifts in channel form or location are sources of sediment for shorthead redhorse habitat. Controlling sediment input into streams may be crucial to prevent detrimental effects to shorthead redhorse, because sedimentation decreases the quality and quantity of optimal habitat (i.e. spawning, feeding) for this species. Sedimentation eliminates interstitial spaces which could be critical for egg deposition and development and for production of benthic organisms, such as aquatic insects, a source of food for shorthead redhorse. Sedimentation has been shown to cause loss or reduction in fish populations, and disrupt the feeding and reproductive activities of fishes (Berkman and Rabeni, 1987).

Non-Habitat Threats:

Pollution



Common Name: **Shorthead Redhorse**
 Scientific Name: **Moxostoma macrolepidotum**
 Species Group: **Fish**

Description of non-habitat threat(s): Water pollution may indirectly influence shorthead redhorse through negative impacts to its prey base. Depletion of food items will negatively affect species growth and survival.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Distribution, abundance and dynamics of Shorthead Redhorse populations in Vermont are poorly understood.
Research	Threats and Their Significance	Medium	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	Medium	
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	Medium	Sedimentation and pollution
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Shorthead Redhorse**
Scientific Name: **Moxostoma macrolepidotum**
Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Natural Processes Restoration	Medium	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, dam owners	Dam owners
Legislation	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP	
Natural Processes Restoration	Medium	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, dam owners	Dam owners
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Policy & Regulations	Medium	Currently under VFWD fishing regulations Moxostoma species are "cull" fishes and as such their harvest and collection is essentially an unregulated activity. Whether or not Moxostoma are threatened by unrestricted harvest should be reviewed.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)



Common Name: **Shorthead Redhorse**
Scientific Name: **Moxostoma macrolepidotum**
Species Group: **Fish**

Bibliography

Berkman, H. E., and C. F. Rabeni. 1987. Effect of siltation on stream fish communities. *Environmental Biology of Fishes* 18(4):285-294.

Jenkins, B., and N. Burkhead. 1993. *Freshwater fishes of Virginia*. American Fisheries Society, Bethesda, Maryland.

Langdon, R. W., M. T. Ferguson, and K. M. Cox. In press. *Fishes of Vermont*. Vermont Fish and Wildlife Department, Waterbury.

Meyer, W. H. 1962. Life history of three species of redhorse (*Moxostoma*) in the Des Moines River, Iowa. *Transactions of the American Fisheries Society* 91(4):412-419.

Scott, W. B., and E. J. Crossman. 1973. *Freshwater fishes of Canada*. Fisheries Research Board of Canada Bulletin 184, Ottawa.

Sheidegger, K. J., and M. B. Bain. 1995. Larval fish distribution and microhabitat use in free-flowing and regulated rivers. *Copeia* 1995(1):125-135.



Common Name: **Greater Redhorse**
Scientific Name: **Moxostoma valenciennesi**
Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend: Unknown

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? No

Assessment Narrative:

Based on current survey information, which is scant, Greater Redhorse appears to have the most restricted distribution of the three Moxostoma species occurring in Vermont, i.e. limited to northern Lake Champlain and the lower accessible reach of the Missisquoi River (Langdon et al. 2006). In 2009, it was documented to occur in Lake Memphremagog (Kratzer 2015). Whether Greater Redhorse has long been resident of the lake but has evaded detection until recently is no known. It is reported to be not difficult to find in the Missisquoi River during the spring season (Ferguson 2014). Threats include poor water quality resulting from agricultural and urban pollution, artificial flow regimes, and habitat fragmentation (Cook et al. 2005). However, in Vermont, little is known of this population or threats to it. Because populations are not being monitored, trends influenced by threats may not be detected.

Distribution

The greater redhorse is generally distributed in central and eastern North America primarily in the upper Mississippi and Great Lakes-St. Lawrence systems (Scott and Crossman 1979). This species has been recorded from northern Lake Champlain and the lower reach of the Missisquoi River (Langdon et al., In press). In Vermont, the species is on the eastern edge of its North American range.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain
Missisquoi River
St. Francois River

Probable Watersheds

Lamoille River
Otter Creek
Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The greater redhorse inhabits medium to large rivers, large lakes and impoundments. It prefers coarse substrate, such as gravel, cobble, and boulders in clean water. In rivers, this species is frequently found in



Common Name: Greater Redhorse
Scientific Name: *Moxostoma valenciennesi*
Species Group: Fish

moderate to swift current in run or riffle habitat and may also be found in large river pools. Generally, it is not found in silty areas and is believed to be intolerant of silt and turbidity (Jenkins and Burkhead 1993). Spawning occurs in spring or summer in high velocity riffle habitat over gravel or cobble substrate that is silt free. Spawning has been found to occur in moderate stream velocities (3.8-116.9 cm/s) and at shallow depths (10-100 cm) (Healy 2002). Greater redhorse may perform annual migrations upstream to spawn and downstream after spawning. In an Ontario river, this species was observed dispersing up to 15 km downstream from its spawning habitat (Healy 2002). This species demonstrates important seasonal movement patterns. Different life stages have specific habitat preferences. Age-0 fish were found in shallow (20 cm), slow velocity pools (21 cm/s). Juvenile fish (greater than age-0 but not sexually mature) were found in slightly deeper pools (60-149 cm) and higher velocities (37 cm/s) (Healy 2002). The greater redhorse is a specialized benthic feeder such that highly silted or embedded substrate may preclude this species from consuming its preferred food items.

Habitat Types:

Aquatic: Lacustrine

Aquatic: Fluvial

Aquatic: Lower CT River

Current Threats

Habitat Threats:

Habitat Alteration

Sedimentation

Habitat Fragmentation

Unknown Habitat Threats

Description of habitat threat(s): Flow alteration, temperature alteration, or low habitat diversity (i.e., loss of deep pool habitat, shallow riffles, or large woody debris due to human-induced change) will most likely pose negative effects for different life stages of greater redhorse. For instance, shallow channel margin habitats that are indicative of slower velocities are important for young redhorses. Flow alteration has been shown to alter and limit this habitat for juvenile life stages (Scheidegger and Bain 1995). Fragmentation of greater redhorse habitat may disrupt the seasonal movement patterns of this species. For example, these movement patterns may prove critical for successful reproduction, and therefore the completion of the species life cycle. Viability of greater redhorse populations most likely depends on optimal habitat availability (i.e., optimal or suitable depths, velocities, substrate, temperature, and flow regimes). Some evidence suggests that greater redhorse presence and abundance are correlated with longer contiguous river reaches (Healy 2002). If the quantity and quality of greater redhorse habitat is limited in a system, then interconnected river reaches will prove necessary for this species to find and occupy optimal or suitable habitat. Loss of riparian vegetation, general construction activities, road maintenance activities (ditching, sanding), bridge and culvert construction, agriculture, timber harvest, dam failure, rapid drawdown of dam impoundments, streambank erosion, and shifts in channel form or location are sources of sediment into greater redhorse habitat. Controlling sediment input into streams may be crucial to prevent detrimental effects to greater redhorse, because sedimentation decreases the quality and quantity of optimal habitat (i.e., spawning, feeding) for this species. Sedimentation eliminates interstitial spaces which could be critical for egg deposition and development and for production of benthic organisms, a primary food source for greater redhorse. Specialized benthic feeders, such as greater redhorse, represent a very ecologically vulnerable group to increased sedimentation, because they are unable to modify their feeding



Common Name: **Greater Redhorse**
 Scientific Name: **Moxostoma valenciennesi**
 Species Group: **Fish**

habits. Sedimentation has been shown to cause loss or reduction in fish populations, and disrupt the feeding and reproductive activities of fish (Berkman and Rabeni 1987). The greater redhorse seems to be rare over the majority of its range (Healy 2002). Determining the primary mechanism behind this trend is a challenge. Unknown habitat problems may exist.

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): The reproductive strategy of the greater redhorse is a crucial aspect to its conservation. It becomes sexually mature at a late age, is highly fecund, and spawns seasonally. Disruption to the spawning efforts of this species poses a problem to population viability (i.e., week year classes over time compound negative influences and population declines). Water pollution may indirectly influence greater redhorse through negative impacts on its prey base. Depletion of food items will negatively affect species growth and survival.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Distribution, abundance and dynamics of Greater Redhorse populations in Vermont are poorly understood.
Research	Threats and Their Significance	High	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	Medium	Sediment and pollution
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Greater Redhorse**
 Scientific Name: **Moxostoma valenciennesi**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Natural Processes Restoration	Medium	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, dam owners	Dam owners
Natural Processes Restoration	Medium	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, dam owners	Dam owners
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Policy & Regulations	Medium	Currently under VFWD fishing regulations Moxostoma species are "cull" fishes and as such their harvest and collection is essentially an unregulated activity. Whether or not Moxostoma are threatened by unrestricted harvest should be reviewed.	Review was conducted and recommendations were considered.	VFWD, USFWS, NYDEC, UVM	VFWD (SWG, DJ), USFWS, UVM
Legislation	High	Support efforts, such as state, federal, regional and international Climate Change Action Plans to reduce greenhouse gas emissions in the Northeast and climate change risks to SGCN.	Adopt appropriate legislation & policies developed to reduce greenhouse emissions & reduce climate change risks to SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP	



Common Name: **Greater Redhorse**
Scientific Name: **Moxostoma valenciennesi**
Species Group: **Fish**

Bibliography

Cooke, S. J. et al. 2005. Threats, conservation strategies, and prognosis for suckers (Catostomidae) in North America: insights from regional case studies of a diverse family of non-game fishes. *Biological Conservation* 121:317-331.

Ferguson, M. 2014. Personal communication. Vermont Fish and Wildlife Department, Montpelier.

Kratzer, J. 2015. Personal communication. Vermont Fish and Wildlife Department, St. Johnsbury.

Langdon, R. W., M. T. Ferguson, and K. M. Cox. 2006. *Fishes of Vermont*. Vermont Department of Fish and Wildlife, Waterbury.

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).



Common Name: **Stonecat**
Scientific Name: **Noturus flavus**
Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Two populations are known to occur in Vermont: one in the Missisquoi River and the other in the LaPlatte River. In the Missisquoi River sub-populations occur immediately below Swanton Dam and the other from that dam upriver to the Highgate Falls Dam. In the LaPlatte River sub-populations occur down- and upstream of Shelburne Falls. The Vermont Cooperative Fish and Wildlife Research Unit at the University of Vermont is currently conducting research of populations in both rivers including estimating abundance, age structure, distribution and habitat use. The project is partially funded by a Vermont Fish and Wildlife Department State Wildlife Grant. On a wide-range scale no major threats are known; however, localized threats to populations may exist such as siltation, pollution and impoundment of habitat (NatureServe 2013). Low winter flows effects on overwinter survival are believed to be the most likely factor influencing stonecat abundance in the Milk River in Alberta (ASRD 2004). During the summer of 2012, stonecat stranding and mortality was observed immediately below Swanton Dam (Pientka 201X; Puchala 201X). The lack of adequate spillage over the dam to keep stonecat habitat under water as a consequence of unusually low base river flows and dam leakage are believed to have been the cause. Chemical control of parasitic Sea Lamprey also has the potential to cause non-target mortality. Large numbers of Stonecats are typically killed when the Great Chazy River in New York is treated with lampricides. Incidental mortality has also occurred in the lower Missisquoi River when lampricide has been applied to habitat holding stonecats. Recently, the U.S. Fish & Wildlife Service announced an interest in treating the LaPlatte River with lampricide to control Sea Lamprey ammocoetes numbers there (Langdon 2014).

Distribution

This is a North American species with a distribution described by Scott and Crossman (1973) as the St. Lawrence River and tributaries in Quebec, south in the Hudson, Allegheny and Mohawk systems in New York, west to the Appalachian Mountains, to western North Carolina and northern Alabama (Tennessee River), north through central Tennessee, west through northern Missouri, Kansas and northeastern Colorado, Wyoming to Alberta, east through North Dakota into Manitoba, southeast through the tip of Lake Superior to central Michigan, and into southern Ontario and Quebec (Scott and Crossman 1973). Only two populations of stonecat is known to occur in Vermont. One population is located within a relatively short reach of the LaPlatte River immediately above and below the fall line; and the second is in Hungerford Brook, a tributary of the Missisquoi River. In Vermont, the species is on the eastern edge of its North American range.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:



Common Name: **Stonecat**
Scientific Name: **Noturus flavus**
Species Group: **Fish**

Known Watersheds

Lake Champlain

Missiquoi River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

The stonecat prefers moderate currents of medium to large rocky-bottomed streams. It is absent, however, from high gradient streams with fast currents. It is also found in lakes near gravel shoals where the current is produced by wave action. The stonecat appears to require a current to prosper, since it has been eliminated from streams where flows have been slowed by the construction of dams. It appears to be intolerant to siltation and general habitat degradation. The stonecat is a state listed endangered species in Vermont with one known population in the state. This population is in a very short section of the river encompassing habitat immediately below and above the fall line. Population monitoring suggests the population has been declining due to unidentified causes. It appears from the literature and Vermont data from the LaPlatte River that this species requires moderate current and a low silt, coarse substrate. Stonecat prefer to use large cobble and boulders for hiding. The combination of habitat requirements of low silt, moderate current, and large substrate represent a somewhat restrictive combination within the Champlain Valley biophysical region.

Habitat Types:

Aquatic: Lacustrine

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Habitat Alteration

Sedimentation

Description of habitat threat(s): It has been reported that this species is sensitive to siltation but the exact mechanism of impact is not known. It may be that siltation covers the developing eggs; however, this may not be a problem, since parents are cavity nesters, preparing the nest and providing care for the young. Or, siltation may embed coarse substrate materials eliminating cover habitat and nesting sites. Since in Vermont the stonecat is only found in the LaPlatte River and Hungerford Brook, a primary conservation consideration is the limiting of upstream land use activities that increase siltation in moderate gradient habitats.

Non-Habitat Threats:

Genetics

Loss of Prey Base

Description of non-habitat threat(s): Stonecat is a benthic insectivore, specializing in aquatic insects. Excess sedimentation can impact aquatic insects populations and reduce this species' food base. This species would have difficulty shifting to non-benthic foods. Because stonecat has one of the most restricted distributions of any other fish species in Vermont, reductions in population size causing a bottleneck which could possibly result in a loss of genetic variation forfeiting the evolutionary potential of the species.



Common Name: **Stonecat**
Scientific Name: **Noturus flavus**
Species Group: **Fish**

Natural selection can only act in the presence of genetic variation, and therefore, the higher the genetic diversity in a population, the higher the likelihood for population persistence. If gene flow is limited to within one population of stonecat (estimated number probably much less than 100 individuals), the species is not prepared to adapt to environmental changes of time.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Comparative studies of habitat occupied by the more abundant New York populations to LaPlatte River habitat.
Research	Basic Life History	High	
Research	Distribution and Abundance	High	
Research	Threats and Their Significance	High	Investigate genetic characteristics of the LaPlatte River Stonecat population and genetic similarity to populations in New York.
Research	Population Genetics	Medium	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Stonecat**
 Scientific Name: **Noturus flavus**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VFWD, VDEC, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Invasive Species Control & Prevention	High	Manage potential non-target impacts of the Lake Champlain Sea Lamprey control program on Stonecat populations.	Number of existing populations of Stonecat protected and sustained.	VFWD, USFWS	USFWS, VFWD
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VFWD, VDEC, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Natural Processes Restoration	High	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners
Natural Processes Restoration	High	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, VTrans, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners, VFWD (SWG), USFWS, VDEC, VTrans



Common Name: **Stonecat**
Scientific Name: **Noturus flavus**
Species Group: **Fish**

Bibliography

ARSD (Alberta Sustainable Resource Development). 2004. Status of the stonecat (*Noturus flavus*) in Alberta. ARSD, Fish and Wildlife Division, and Alberta Conservation Association, Wildlife Status Report No. 52, Edmonton.

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.

NatureServe. 2013. *Noturus flavus*. IN: The IUCN Red List of Threatened Species. Version 2014.2. Available at: www.iucnredlist.org. (November 2014).

Pientka, B. 201X. Personal communication. Vermont Fish and Wildlife Department, Essex.

Puchala, E. 201X. Personal communication. Vermont Fish and Wildlife Cooperative Research Unit, University of Vermont, Burlington.



Common Name: **Redbreast Sunfish**
 Scientific Name: **Lepomis auritus**
 Species Group: **Fish**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S4

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Scarola (1973) reported Redbreast Sunfish to be in the New Hampshire portion of the Connecticut River watershed. More recently (2008), Yoder et al. (2010) during a fish assemblage assessment of the Connecticut River extending from Lake Francis in NH downriver to Turners Falls Dam in MA collected two Redbreast Sunfish from separate locations in the river situated between Bellows Falls and Wilder dams. NAI (2004) fish community monitoring conducted in the vicinity of Vermont Yankee Nuclear Power Station (Vernon, VT and Hinsdale, NH) captured a total of 14 Redbreast Sunfish from the Connecticut River during the years 1991-1999. Reports of the species occurring in Vermont “inland” waters, as far as known, appear to be limited to Lake Morey and Lake Fairlee. However, in more recent years the species has not been encountered in either lake (Kim 2015). The disappearance of Redbreast Sunfish in Lake Morey seems to coincide with the appearance of bluegill (Kim 2015). Perhaps this indicates bluegill have outcompeted and replaced Redbreast Sunfish in the lake. Nonetheless, this strongly suggests the species is uncommon in both the Connecticut River and rare or even extirpated from the two “inland” lakes where it was previously reported in Vermont. NatureServe (2014) states there are no known threats to the species; however, dams and pollution have caused local declines. COSEWIC (2008) reports there is little information in the literature which addresses limiting factors affecting Redbreast Sunfish, although the species appears adaptable to a wide range of environmental conditions. COSEWIC (2008) identifies the shoreline development including seasonal and year-round homes, forestry and agricultural activities, and stream alterations to be the most obvious threat to the species in New Brunswick, although waterbody-specific assessments conducted in that province failed to demonstrate correlations between shore development and water quality parameters to Redbreast Sunfish abundance. Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

The redbreast sunfish occurs along the Atlantic Slope from New Brunswick to Florida. It is also found in Gulf Coast drainages in Georgia and Florida, north into Kentucky and Arkansas. In Vermont, redbreast sunfish are found in the Connecticut River and lakes Morey and Fairlee (Orange County) and the Black River (Windsor County). Historically, the species has been recorded from New York waters within the Lake Champlain drainage (e.g., Lake George and its outlet, the Mettawee River and Little Chazy River) (Greeley 1930); however, no records, past or present, are known from Lake Champlain.

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Confident
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:



Common Name: **Redbreast Sunfish**
Scientific Name: **Lepomis auritus**
Species Group: **Fish**

Known Watersheds

Middle Connecticut

West

Upper Connecticut-Mascoma

Black-Ottawquechee

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

This species inhabits the shores of lakes and ponds, and pools of clear streams with little current, but unlike other Vermont sunfishes it is more of a stream-adapted species. Aho and Terrell (1986) report cover, current velocity, and variables correlated with velocity (e.g., gradient, riffle/pool ratios) to be major factors determining the distribution and abundance of redbreast sunfish in riverine systems. Both juveniles and adults are usually found in shallow water near cover, although fish may occupy deeper habitats under warmwater summer conditions and during winter. Important cover include fallen trees, stumps and aquatic vegetation. These hard structures appear to be important habitat components for spawning site selection. Additionally, hard structures have been attributed to being the substrate producing more than 60% of the food organisms consumed by sunfish species, including redbreast sunfish. Scarola (1973) states redbreast sunfish can be found over gravelly bottoms with or without vegetation; however, unlike the pumpkinseed, it does not rely as much on there being aquatic vegetation present. Aho and Terrell (1986) quantify variables critical to habitat suitability models in both lotic and lentic environments for redbreast sunfish. Water temperatures regarded as suitable for growth and survival of adult and juvenile fish are assumed to be in the range of 15-35 C; for spawning and incubation the optimal range is assumed to be 21.1-27.2 C. Nests are generally constructed at depths less than 1.5 m. Water velocities at nest sites are less than 0.06 m/s with an average of 0.02 m/s. Based on available information for other sunfish species, 25-70% hard structure cover is estimated to be most productive for redbreast sunfish. This species appears to require a mixture of coarse sand and gravel substrate at spawning sites to be successful.

Habitat Types:

Aquatic: Fluvial

Current Threats

Habitat Threats:

Habitat Alteration

Description of habitat threat(s): Abundance of hard structures for cover are critical components of redbreast sunfish habitat. Removal of such cover or inadequate structure being recruited into lakes and streams (e.g., from forested riparian areas) may negatively affect the suitability of habitat for this species. It may be sensitive to acidity (i.e., long term pH values <4.0), but is tolerant of high temperatures (<35 C) (Aho and Terrell 1987). Rapid reductions in water level of more than 0.9 m during the spawning season may adversely affect embryo development and survival (Aho and Terrell 1987).

Non-Habitat Threats:

Pollution



Common Name: **Redbreast Sunfish**
Scientific Name: **Lepomis auritus**
Species Group: **Fish**

Description of non-habitat threat(s): Aho and Terrell (1987) identify several potential threats to this species. Low to moderate turbidity levels are suitable to this species; however, excessive levels may impact fish growth and abundance. Pesticide contamination of waters supporting redbreast sunfish has been a suspected cause for the observed decline of some populations.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	1) Determine its distribution in Vermont waters. It may be present in more streams of suitable habitat in the Connecticut Valley than is presently known. 2) The spatial extent of its presence in the Connecticut River and its larger tributaries should also
Research	Threats and Their Significance	Low	Evaluate and monitor pesticide levels in known populations of Redbreast Sunfish.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	Medium	Monitor known populations.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	Medium	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Redbreast Sunfish**
 Scientific Name: **Lepomis auritus**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, CRWC, TNC	VDEC (ERG, VWG), VFWD (DJ, SWG)
Habitat Restoration	Medium	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, CRWC, TNC	VDEC (ERG, VWG), VFWD (DJ, SWG)
Natural Processes Restoration	Medium	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, CRWC, TNC, lake associations, dam owners	Lake associations, dam owners

Bibliography

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2008. COSEWIC assessment and update status report on the redbreast sunfish (*Lepomis auritus*) in Canada. COSEWIC, Ottawa. Available at: www.sararegistry.gc.ca/status/status_e.cfm. (December 2014).

Kirn, R. 2015. Personal communication. Vermont Fish and Wildlife Department, Roxbury.

NAI (Normandeau Associates, Inc. (NAI). 2004. §316(a) demonstration in support of a request for increased discharge temperature limits at Vermont Yankee Nuclear Power Station during May through October. Report of NAI to Entergy Nuclear Vermont Yankee, Vernon, Vermont.

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).

Scarola, J. F. 1987. Freshwater fishes of New Hampshire. New Hampshire Fish Department, Concord.

Yoder, C. O., L. E. Hersha, and B. R. Apell. 2010. Fish assemblage and habitat assessment of the upper Connecticut River: a preliminary report and presentation of data. Midwest Biodiversity Institute, MBI Technical Report MBI/2009-8-3, Columbus, Ohio.



Common Name: **Eastern Sand Darter**
Scientific Name: **Ammocrypta pellucida**
Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G3

Global Trend: Unknown

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The number of populations known to occur in Vermont stands at four located in the Missisquoi, Lamoille, Winooski and Poultney rivers (Grandmaison et al. 2004; Langdon 2014), or five if the Winooski River is considered to have two populations separated by the natural fall line (present day Winooski One dam). Vermont populations are disjunct from the species' main distribution which encompasses the Midwestern states. Although populations in the Vermont are presumed to be stable, systematic and temporal assessments have not been conducted to quantify abundance trends. However, there is general agreement that the abundance of the species is declining throughout much of its continental range (Kuehne and Barbour 1983 cited in Grandmaison et al. 2004). Various sources summarized by Grandmaison et al. (2004) speculate that Vermont populations at present appear to be abundant enough to be viable; however, historical data and current monitoring activities are inadequate or lacking from which confident abundance trends can be derived and such conclusions are at best speculative. Grandmaison et al. (2004) present a summary of potential threats to Eastern Sand Darters in nine states within the species range. The most cited threats to the species are habitat destruction or degradation resulting from impoundment, channelization, channel dredging and siltation. Threats identified for Vermont populations are sedimentation resulting from bank erosion and storm water discharges; water quality degradation from livestock manure runoff, and chemicals and other catastrophic spills. The potential for sedimentation to impair critical habitat necessary for eastern sand darters is a persistent problem within its Vermont range given most of the populations lie within drainages with high agricultural and land development activity. Sea Lamprey control measures employing lampricides (e.g. TFM) has also been a concern in Vermont. Bioassay testing on sand darters has determined the maximum no-effect concentration is 1.3 to 1.4 times the minimum lethal concentration of TFM needed to control Sea Lamprey. The range of Eastern Sand Darter encompasses nine states and two Canadian provinces. Of these, it is a species of special concern in two (Indiana, Ohio), threatened in five (Illinois, Michigan, New York, Vermont, Canada) and endangered in one (Pennsylvania) (Grandmaison et al. 2004). Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

Eastern sand darters range from the St. Lawrence River drainage, southern Quebec, Vermont and New York; through the Great Lakes and Ohio River basins from western New York to eastern Illinois; and south to Kentucky (Page and Burr 1991). In Vermont, populations are known to occur below the fall line in the Missisquoi, Lamoille, Winooski, and Poultney rivers. There is one recent occurrence of the species being collected from above the fall line on the Winooski River. One individual has also been collected in Lake Champlain at the mouth of the Lamoille River in Malletts Bay. In Vermont, this species is on the eastern edge of its range.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Sand Darter**
Scientific Name: **Ammocrypta pellucida**
Species Group: **Fish**

Northeastern Highlands Not Probable

Distribution by Watershed:

Known Watersheds

Lake Champlain

Lamoille River

Missiquoi River

Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The eastern sand darter shows a strong preference for sandy areas of rivers and streams with slow to moderate currents, where it spends most of its time burrowed into the sand with only its eyes or head protruding. It has also been reported from sandy shoals in Lake Erie, but has not been reported in Lake Champlain, except for one individual at the mouth of the Lamoille River in Malletts Bay. The eastern sand darter requires medium to fine sand, so water velocity and sedimentation are important factors in habitat suitability. Habitat use and preference studies indicate that the fish use areas with a large percentage of sand particles 0.23 to 1 mm in size. It is quite sensitive to sedimentation and poor water quality.

Current Threats

Habitat Threats:

Habitat Alteration

Sedimentation

Description of habitat threat(s): The range of the eastern sand darter is believed to be shrinking due to the loss of clean sand habitat caused by increased siltation from soil erosion and agricultural activities. Hydroelectric power generation should be regulated to maintain suitable flows and habitat.

Description of non-habitat threat(s): The Lake Champlain Fish and Wildlife Management Cooperative is currently involved in a sea lamprey control program that includes the use of lampricides to kill stream resident lamprey larvae. Bioassay LCD levels on eastern sand darter have determined.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Sand Darter**
Scientific Name: ***Ammocrypta pellucida***
Species Group: **Fish**

Research and Monitoring Needs

<i>Type</i>	<i>Need</i>	<i>Priority</i>	<i>Description</i>
Research	Habitat Requirements	High	Determination of optimal microhabitat requirements (e.g., depth, velocity and substrate).
Research	Basic Life History	Medium	
Research	Distribution and Abundance	High	Increase sampling efforts in rivers with known populations, including sampling beyond known areas of occurrence.
Research	Threats and Their Significance	High	Effects of limiting factors (e.g., hydrogeneration) on habitat, and possible long term effects of lampricide treatment on populations.
Research	Population Genetics	Medium	How closely are Vermont populations linked genetically to one another and to other populations located outside of the state.
Research	Taxonomy	Low	
Research	Other Research	Low	Diet studies.
Monitoring	Population Change	High	
Monitoring	Habitat Change	Medium	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Sand Darter**
 Scientific Name: **Ammocrypta pellucida**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Invasive Species Control & Prevention	Medium	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)
Natural Processes Restoration	Medium	Restore/maintain connectivity within aquatic systems supporting sustainable SGCN population(s); provide for safe & efficient up- and downstream SGCN passage at dams & other obstructions.	Based on historic distribution of the SGCN, number of miles of habitat to which access to critical habitat has been restored or maintained.	VDEC, VFWD, USFWS, dam owners	Dam owners
Natural Processes Restoration	High	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, dam owners	Dam owners
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VDEC, VFWD, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Invasive Species Control & Prevention	High	Manage potential non-target impacts of the Lake Champlain Sea Lamprey control program on Eastern Sand Darter populations.	Number of existing populations of Eastern Sand Darter protected and sustained.	VFWD, USFWS	USFWS, VFWD

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Sand Darter**
Scientific Name: ***Ammocrypta pellucida***
Species Group: **Fish**

Invasive Species Control & Prevention	High	Adopt/implement appropriate actions that minimize the potential for new invasive species introductions of potential threat to SGCN; control in-state invasive species populations when/where opportunities avail.	No increase in numbers of invasive organisms in habitat occupied by the SGCN.	VDEC, USFWS, LCBP	VDEC (ANCG)

Bibliography

Grandmaison, D., J. Mayasich, and D. Etnier. 2004. Eastern sand darter assessment. Report of Natural Resources Research Institute to U. S. Fish and Wildlife Service. NRRI Technical report No. NRRI/TR-2003/40, Fort Snelling, Minnesota.

Langdon, R. 2014. Personal communication: DEC fish distribution database. Vermont Department of Environmental Conservation, Montpelier.

Kuehne, R. A., and R. W. Barbour. 1983. The American darters. The University Press of Kentucky, Lexington



Common Name: **Channel Darter**
 Scientific Name: **Percina copelandi**
 Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend: Unknown

State Rank: S1

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Four populations are currently known in Vermont. These are located in the Lamoille, Winooski, LaPlatte and Poultney rivers. Little, if anything, is known of the size, structure, and trends of these populations. Channel Darter has very specific habitat requirements and as such populations tend to be restricted in size and distribution (COSEWIC 2002). NatureServe (2014) ranks the overall threat to populations to be high with reductions of Channel Darter populations occurring throughout its continental range. Threats to populations include habitat loss and degradation due to siltation, pollution, flow modification, and impoundments; fragmented populations have a reduced likelihood of recovering (COSEWIC 2002; NatureServe 2014). Potential causes of declines in Lake Erie include eutrophication, shoreline modifications from development, and invasive Round Goby (NatureServe 2014). Population declines and extirpation of some populations have been reported from Lake Erie, Ohio, Michigan, Ontario and Quebec (NatureServe 2014). Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

This is a wide ranging species but is highly localized in the St. Lawrence, Great Lakes and Mississippi River drainages from southern Quebec and Vermont, south to northern Louisiana; along the Gulf Slope in Mobile, Pascagoula and Pearl River drainages (Page and Burr 1991). In Vermont, the species is on the eastern edge of its range with populations known to occur below the fall line in the Winooski, LaPlatte and Poultney rivers. There is a historic record from Lake Champlain on the New York side (Greeley 1930); however, no occurrences have been made within the Vermont portion of the lake.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Mettawee River
 Lake Champlain
 Winooski River
 Lamoille River



Common Name: **Channel Darter**
Scientific Name: **Percina copelandi**
Species Group: **Fish**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The channel darter is a bottom dweller of gravelly or sandy shoals of warm lakes and rivers. In rivers, it is found in areas with coarse sand and gravel substrate. These areas have low to moderate current, but enough water velocity to prevent silt deposition. Channel darters are found in areas with substrates composed of gravel and sand. Preferred habitat is low in sediments and turbidity. Some studies of spawning in rivers and aquaria indicate that channel darters require swift currents (0.03-0.04 m/sec) presumably with gravel substrate.

Habitat Types:

Aquatic: Lacustrine

Aquatic: Lake Champlain

Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Habitat Alteration

Sedimentation

Description of habitat threat(s): Channel darters are limited by the loss of clean gravel substrate resulting from increased siltation and turbidity from soil erosion and agricultural activities. Alteration of river flow regimes from hydroelectric power generation may also degrade habitat quality.

Description of non-habitat threat(s): The Lake Champlain Fish and Wildlife Management Cooperative is currently involved in a sea lamprey control program that includes the use of lampricides to kill stream resident lamprey larvae.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Channel Darter**
Scientific Name: **Percina copelandi**
Species Group: **Fish**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Increase sampling efforts in rivers with known populations, including sampling beyond known areas of occurrence.
Research	Threats and Their Significance	High	
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	
Monitoring	Habitat Change	High	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	High	Monitor impacts of sea lamprey control in the Lake Champlain watershed on channel darter populations.
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Channel Darter**
 Scientific Name: **Percina copelandi**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VFWD, VDEC, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)
Natural Processes Restoration	High	Restore flow regimes and/or water levels that support sustainable SGCN population(s) & at targeted abundance levels.	Number of miles of SGCN habitat improved or restored.	VDEC, VFWD, USFWS, TU, dam owners, watershed associations, town & regional planning & Cons Comms	Dam owners
Invasive Species Control & Prevention	High	Manage potential non-target impacts of the Lake Champlain Sea Lamprey control program on Channel Darter populations.	Number of existing populations of Channel Darter protected and sustained.	VFWD, USFWS	USFWS, VFWD
Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.	VFWD, VDEC, USFWS, NRCS, LCBP, watershed associations, town & regional planning & Cons Comms	VDEC (ERG, VWG), VFWD (DJ, SWG)

Bibliography

COSEWIC (Committee on the Status of Endangered Wildlife in Canada). 2002. COSEWIC assessment and status report on the channel darter *Percina copelandi* in Canada. COSEWIC, Ottawa. Available at: www.sararegistry.gc.ca/status/status_e.cfm. (November 2014).

NatureServe. 2014. NatureServe Explorer: an online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available at: <http://explorer.natureserve.org>. (November 2014).

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Sauger**
Scientific Name: **Sander canadense**
Species Group: **Fish**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend: Unknown

State Rank: S4S5

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Lake Champlain supports the only Sauger population in Vermont, where the species is on the eastern edge of its continental range. In Lake Champlain, historically they were more abundant in the southern than northern portion of the lake (Halnon 1952, 1956). Once described as common in Lake Champlain (Anderson 1978), saugers have been rarely seen during the last 20 years (MacKenzie 2014) and has apparently declined to the point that it eluded detection by New York and Vermont fisheries biologists from 1994 (Nettles et al. 2005) until 2010 (NYDEC 2013). The species was once widely distributed in New York but is now extirpated from much of its historic range with exception of Lake Champlain. Sauger has declined in abundance and distribution across its range (Rawson and Scholl 1978; Hesse 1994; Pegg et al. 1997). Threats to the species include angler harvest, channelization, water flow fluctuations, migration barriers, loss of spawning and rearing habitat, and environmental degradation (Rawson and Scholl 1978; Hesse 1994; Pegg et al. 1997). Selected as an Regional-SGCN by the 13 Northeastern states in 2014.

Distribution

The distribution of sauger in North America is from the St. Lawrence-Lake Champlain system south, west of the Appalachian Mountains to Tennessee River in Alabama, southwest to northern Louisiana, northwest through eastern Oklahoma to central Montana and central Alberta east below James Bay to Quebec (Scott and Crossman 1998). The distribution of sauger in Vermont is limited to Lake Champlain, where it may have been more numerous in the southern portion of the lake. Anderson (1978) reported sauger to be present in all sections of the lake except for the Main Lake.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Not Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Lake Champlain



Common Name: **Sauger**
Scientific Name: **Sander canadense**
Species Group: **Fish**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Many of the general habitat requirements are similar between sauger and walleye; however, sauger habitat preferences are for large, shallow sections of lakes which are turbid with colloidal clay suspension, as well as large, turbid, slow flowing rivers (Anderson 1978; Scott and Crossman 1978). Scott and Crossman (1978) considered sauger "less adaptable" than walleye because of these preferences. Walleye and sauger may utilize the same shoals or gravel to rubble in large turbid lakes for spawning (Scott and Crossman 1978). Preferred spawning habitats are shallow shoreline and shoals of lakes and riffles in rivers, including areas immediately below dams providing there is rocky substrate and good water circulation from wave action and river currents (McMahon et al. 1984). Sauger have been found to be highly selective for spawning sites and in some parts of their range have been shown to be reliant on access to a few discrete areas in large tributaries (Nelson 1968; Gardner and Steward 1987; Penkal 1992; Jaeger 2004). Sauger fry must reach their initial feeding grounds (habitat type?) within 3-5 days before yolk-sac absorption or they will perish from lack of food (McMahon et al. 1984).

Habitat Types:

Aquatic: Fluvial

Aquatic: Lower CT River

Current Threats

Habitat Threats:

Habitat Alteration

Habitat Fragmentation

Invasion by Exotic Species

Description of habitat threat(s): Sauger are considered to be the most migratory percid and are heavily dependent throughout their life history on unimpeded access to a wide diversity of physical habitats (Collette 1977; Jaeger 2004). The historic spawning grounds of sauger in Lake Champlain are not well known. Undoubtedly, dams have decreased their accessibility to many of the historical spawning grounds in the basin. For example, recent lake sturgeon and walleye habitat assessments conducted on the Missisquoi River indicate most of the quality spawning habitat occurs above Swanton Dam (Madeline Lyttle, U. S. Fish and Wildlife Service, personal communication). Sauger also appear to be sensitive to changes in water quality. Sauger may be more dominant than walleye under very turbid water conditions where they co-occur; however, dominance may shift with changing water quality (Scott and Crossman 1998). Improvements in Lake Champlain water quality may explain the perceived reduction in sauger abundance, but this needs to be investigated.

Non-Habitat Threats:

Competition

Description of non-habitat threat(s): The sauger population was once abundant in portions of Lake Champlain and were captured in considerable numbers as recently as the 1980s. Recent surveys of the South Bay, where sauger were formerly abundant, failed to produce even in a single capture. Predation by native species, such as smallmouth bass (Johnson and Hale 1977) have been found to influence recruitment of walleye, a close relative to sauger, in natural systems (as referenced in Quist et al. 2003). Others have

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Sauger**
 Scientific Name: **Sander canadense**
 Species Group: **Fish**

speculated that native piscivorous predators, such northern pike, smallmouth bass, lake trout, burbot and Atlantic salmon, can be a major source of mortality for age-0 walleye in Lake Champlain (Frater 2002). We would expect these interactions to be as important, if not more so, for sauger. For example, the introduction of black crappie in Black Lake (New York) was believed to have caused successive walleye year-class failures (Schiavone 1983). While black crappie are believed to be native to Lake Champlain, its cogenitor the white crappie is not. It too has been found to be a significant walleye fry predator in some systems (Quist et al. 2003). White crappie are known to occur in large numbers in areas where sauger were historically abundant, e.g. South Bay (David Nettles, U. S. Fish and Wildlife Service, personal communication). Another exotic in lake Champlain, the white perch, has been found to be an important predator of walleye eggs (Roseman et al. 1996; Schaeffer and Margraf 1987). White perch have become or are becoming one of the most dominant species in the fish assemblage in some areas of the lake, e.g. Missisquoi Bay (Pierre Bilodeau, Quebec Parks and Wildlife, personnel communication).

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Determine the current population status of Sauger in Lake Champlain (Vermont and New York sections), and identify critical spawning and juvenile habitats.
Research	Threats and Their Significance	High	1) Determine the effect of recent invasions of non-indigenous species (e.g., White Crappie, White Perch, Zebra Mussel) on Sauger in Lake Champlain. 2) Determine the effect, if any, changing water quality may have on the Sauger population.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	N/A	
Monitoring	Population Change	High	
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	N/A	
Monitoring	Monitor Threats	Medium	
Monitoring	Other Monitoring Needs	N/A	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Sauger**
 Scientific Name: **Sander canadense**
 Species Group: **Fish**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Sander canadense	Harvest or Collection	Compliance & Enforcement	High	Close Lake Champlain fishery to the Sauger harvest.	Enforcement of
Sander canadense	Habitat Alteration/Degradation	Habitat Restoration	High	Monitor habitat conditions & effects of stressors on habitats; restore critical habitats or ameliorate threats when/where opportunities arise to secure/restore numbers of SGCN populations & targeted abundance levels.	Increase and/or maintain available habitat (river-miles or surface acres) in terms of quantity and quality required for all life stages of the SGCN.

Bibliography

Hesse, L. W. 1994. The status of Nebraska fishes in the Missouri River. 6. Sauger (Percidae: Stizostedion canadense). Transactions of the Nebraska Academy of Sciences 21:109-121.

Mackenzie, C. 2014. Sauger regulation proposal, 2014. Vermont Fish and Wildlife Department, Montpelier.

Nettles, D. C., C. D. Martin, and N. R. Staats. 2005. South Bay, lake Champlain walleye and sauger assessments, 2003-2004. Draft report. U. S. Fish & Wildlife Service, Ray Brook (DEC office), New York.

NYDEC (New York Department of Environmental Conservation). 2013. Conservation management plan for sauger in New York State, 2013-2020. New York Division of Fish, Wildlife and Marine Resources, Bureau of Fisheries, Albany.

Pegg, M. A., P. W. Bettoli, and J. B. Layzer. 1997. Movement of saugers in the Tennessee River determined by radio telemetry, and implications for management. North American Journal of Fisheries Management 17:763-768

Vermont's Wildlife Action Plan

Invertebrate SGCN Conservation Reports

Appendix A4

Draft 10/1/2015

Species	Page
Ant Group	1
Bumble Bee Group	5
Beetles-Carabid Group	9
Beetles-Tiger Beetle Group.....	21
Butterflies-Grassland Group.....	26
Butterflies-Hardwood Forest Group.....	30
Butterflies-Wetland Group	34
Moths Group	38
Mayflies/Stoneflies/Caddisflies Group	45
Odonates-Bog/Fen/Swamp/Marshy Pond Group	48
Odonates-Lakes/Ponds Group.....	54
Odonates-River/Stream Group.....	59
Crustaceans Group.....	64
Freshwater Mussels Group	68
Freshwater Snails Group	80

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Ant Group**
Scientific Name: **Ant Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Unknown

Extirpated in VT? No

Regional SGCN?

Assessment Narrative:

This group consists of the following two species:

-- *Temnothorax pilagens*: This species is known from only three sites worldwide and is rarely observed. The only VT site is in a state park, which could provide some protection; however, the ant is not a conservation target for the park at this time. More survey work is needed to determine the extent of its range. First collected in 1986 in VT, it hasn't been found here since then despite 14 years of subsequent collecting effort at the park. Recent efforts to collect it again in NY have been unsuccessful.

-- *Myrmica lobifrons*: This species is not considered rare, but is a habitat specialist. At least in New England, it has only been collected in or very close to ombrotrophic bogs. The potential that it could be found in fens, marshes, or other bodies of water with more flow and higher calcium and nutrient levels is uncertain.

Distribution

-- *Temnothorax pilagens*: Known VT location is in Niquette Bay State Park, Colchester. Described as a Nearctic species, it is found in northeastern parts of the United States and possibly southeastern Canada. Only known from three sites, all in the northern US: Niquette Bay State Park, VT (1986); E.N. Huyck Preserve, Rensselaerville, NY (2002 and 2003); and Sleeping Bear National Lakeshore, Empire, MI (2011 and 2013).

-- *Myrmica lobifrons*: May occur throughout the state where ombrotrophic bogs occur.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Probable
Champlain Hills	Probable	Vermont Valley	Probable
Northern Green Mtns	Probable	Southern Green Mtns	Probable
Northern VT Piedmont	Probable	Taconic Mtns	Probable
Northeastern Highlands	Probable		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

Temnothorax pilagens: Occurs in forests, woodlands, and parks; preferentially wooded sites with little understory, and a high density of suitable nest sites. The Vermont site is described in literature as second-growth temperate deciduous forest that has been protected from logging since the 1930's. The forest is dominated by oaks, while hemlock, birch, and pine contribute strongly to the canopy. The site is flat, possibly situated on a floodplain at about 30m elevation. Nests occur in preformed cavities in acorns, hickory nuts or sticks. This species utilizes a social "parasite" strategy, enslaving workers of two congeneric species (T.



Common Name: **Ant Group**
Scientific Name: **Ant Group**
Species Group: **Invert**

longispinosus and T. ambiguous) to perform tasks of the nest. The slave-makers and their hosts live in nest sites in the litter and surface soil layers. In all three known populations, T. pilagens was enslaving both T. longispinosus and T. ambiguus; most often, nests contained slaves of both host species. Nests contain on average four T. pilagens workers (ranging from 0 to 16) and 13 slaves (ranging from 2 to 50 workers). One exceptional example contained 27 T. pilagens and 55 slave workers.

--Myrmica lobifrons: A habitat specialist. At least in New England, it has only been collected in or very close to ombrotrophic bogs. The potential that it could be found in fens, marshes, or other bodies of water with more flow and higher calcium and nutrient levels is uncertain.

Habitat Types:

Oak-Pine Northern Hardwood

Open Peatlands

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Alteration

Habitat Fragmentation

Climate Change

Description of habitat threat(s): --T. pilagens: Oaks are necessary, as they supply acorns used for habitation; hickory nuts are also used. Loss/reduction of oaks would impact habitat availability. Habitat loss may occur in areas that have not been surveyed, but may support this species.

--M. lobifrons: occurs in bog habitat that may be vulnerable to climate change.

Non-Habitat Threats:

Competition

Pollution

Loss of Relationship with Other Species

Unknown Non-Habitat Threats

Description of non-habitat threat(s): Pesticide use could negatively impact the ground-dwelling T. pilagens. Abundance has been declining at the state park for many species of ants, including the hosts of T. pilagens. This could result in loss of colony functions and could reduce survival. The other slave-making ant known from the T. pilagens site is more abundant and utilizes the same host ant species and habitat. It does not appear to yet be impacted by declining host numbers. There may be significant competition for nesting cavities. Low abundance of T. pilagens makes this ant vulnerable to fluctuations in population density. Recolonization or rebound following large population drops may not be possible.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Ant Group**
 Scientific Name: **Ant Group**
 Species Group: **Invert**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Determine whether <i>M. lobifrons</i> habitat requirements are broader than ombrotrophic bogs.
Research	Distribution and Abundance	High	Use existing habitat description to help identify other sites in the state where <i>T. pilagens</i> may exist. An intensive survey of these potential sites will be needed.
Research	Threats and Their Significance	High	Potential limiting factors to populations of <i>T. pilagens</i> need to be evaluated. This should particularly focus on factors at the state park site.
Monitoring	Population Change	High	Original site where <i>T. pilagens</i> was reported should continue to be intensively surveyed to determine if this ant is still present and, if so, at what level of abundance.
Monitoring	Monitor Threats	High	Limiting factors identified as significant at sites of <i>T. pilagens</i> occurrence need to be monitored.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	High	Use existing habitat description to help identify other sites in the state where <i>T. pilagens</i> may exist. Intensive surveys of these potential sites should be conducted. Niquet State Park needs to be intensively resurveyed and level of abundance determined.	Number of new potential sites surveyed.	FWD, FPR, USFS, UVM, VT Entomological Society	SWG, FPR
Easements	High	Acquisition/easement of high priority SGCN ant sites	Number of unprotected sites that become protected	FWD, FPR, USFS, VLT, TNC, other land trusts	SWG, FPR, USFS, VHCB
Protected Area Management	High	Ensure that bog sites are protected for <i>M. lobifrons</i> . Inform landowners and managers of its presence.	Number of landowners/managers provided with technical assistance specific to <i>M. lobifrons</i>	FPR, FWD, USFS, land trusts, UVM, TNC, other landowners and land managers	SWG, FPR, USFS
Protected Area Management	High	Work with FPR to address conservation and enhancement of <i>T. pilagens</i> in management plan for Niquet State Park.	Inclusion of <i>T. pilagens</i> in management plan, with monitoring plan.	FPR, FWD	SWG, FPR
Compatible Resource Use	High	Evaluate potential threats for <i>T. pilagens</i> populations. This should particularly focus on factors at the state park site.	Completion of threat assessment with recommendations.	FWD, FPR, USFWS, UVM, VT Entomological Society	SWG, FPR



Common Name: **Ant Group**
Scientific Name: **Ant Group**
Species Group: **Invert**

Bibliography

Ellison, A.M., N.J. Gotelli, E.J. Farnsworth, G.D. Alpert. 2012. A Field Guide to the Ants of New England. Yale University Press. 256 pp.

Seifert B, I. Kleeberg, B. Feldmeyer, T. Pamminger, E. Jongepier, and S. Foitzik. 2014. *Temnothorax pilagens* sp. n. A new slave-making species of the tribe Formicoxenini from North America (Hymenoptera, Formicidae). *ZooKeys* 368: 65–77.



Common Name: **Bumble Bee Group**
Scientific Name: **Bumble Bee Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend:

Extirpated in VT?

Regional SGCN?

Assessment Narrative:

This group consists of nine species. Research indicates that declines in North American bumble bees have been associated with increased levels of pathogen infections and reduced genetic diversity. Also, habitat loss/degradation, pesticides, climate change, and competition with honey bees may have contributed to the range-wide decline of several species.

- B. affinis: A previously widespread species in Vermont, with no records since 1999.
- B. ashtoni: nest parasite of only B. affinis and B. terricola, two species which are in severe decline.
- B. citrinus: Despite being a parasite of very common Bombus species, it is reported to have declined precipitously in much of the range, including Vermont.
- B. fernaldae: Widely scattered records across the eastern range with only one known for Vermont, despite being in the central part of the eastern range.
- B. fervidus: Much more prevalent in historic VT collections than presently found. Above ground nesting in thatch in grassy areas and gardens may make it vulnerable.
- B. pensylvanicus: A thatch nester, vulnerable to mowing; previously common in Champlain Valley, now appears extirpated in Vermont; has disappeared from northern part of range.
- B. perplexus: More prevalent in the historic record.
- B. rufocinctus: Possible recent declines compared to historic collections.
- B. terricola: Historically, appeared to be a common component of the Vermont bee fauna. Regional data suggest that it was probably found throughout the entire state. It represented about 13% of the 1915- 2011 records, the 2nd most common of 17 known species in Vermont. A severe population decline was noted in 2000 with few observations of the species until 2007 when perhaps a slight recovery began. In 2012 and 2013, this species represented less than 1% of specimens collected during the Vermont Bumble Bee Survey. It was encountered rarely in southern Vermont, in widely scattered locations in the Champlain Valley and central Vermont, and was more widespread in the Northeast Kingdom region.

Distribution

- B. pensylvanicus - Previously known from Champlain Valley; due to limited historical survey effort, may have been present in Northern and Southern VT Piedmont, Northeastern Highlands, Vermont Valley.
- B. affinis - previously widespread; no recent records.
- B. ashtoni - previously widespread; no recent records.
- B. citrinus - previously widespread; most recent records are from Addison Co.
- B. fernaldae - one historic record from Essex Co. Always rare throughout range.
- B. fervidus - previously widespread; recent records from Franklin, Chittenden, Addison, Bennington counties.
- B. perplexus - widespread, though less common than historically.
- B. rufocinctus - previously widespread; now collected mostly in the Champlain Valley.
- B. terricola - widespread, though now much more rare than historically. Nearly disappeared rangewide from 1999 until 2012. Vermont currently has greater numbers in recent collections than other similar latitudes and to the south.



Common Name: **Bumble Bee Group**
Scientific Name: **Bumble Bee Group**
Species Group: **Invert**

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Many species require open, unmanaged grasslands for nesting and foraging. Change in microhabitat features could impact overwintering queens. Climate change and local land-use may affect this critical stage. Bumble bees need diversity (taxonomic and phenologic) of nectar sources.

B. pennsylvanicus - thatch nester in large, unmanaged grasslands.

Habitat Types:

Outcrops and Alpine
Oak-Pine Northern Hardwood
Open Peatlands
Marshes and Sedge Meadows
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops
Wet Shores
Shrub Swamps
Building or Structure
Mine
Other Cultural
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Habitat Fragmentation



Common Name: **Bumble Bee Group**
 Scientific Name: **Bumble Bee Group**
 Species Group: **Invert**

Invasion by Exotic Species

Climate Change

Unknown Habitat Threats

Description of habitat threat(s): *Apis mellifera* is a foraging competitor and possibly disease vector. Changes in land management can alter microhabitat conditions. Climate change may affect spring ephemeral flowering phenology.

Non-Habitat Threats:

Genetics

Competition

Disease

Parasites

Pollution

Reproductive Traits

Loss of Prey Base

Loss of Relationship with Other Species

Description of non-habitat threat(s): Low numbers of individuals may cause decline in genetic health. Bumble bee trade and transport for agriculture is considered to have introduced parasites and diseases to native populations. This trade is poorly regulated. Pesticides, introduced diseases, and competition with non-native honey bee all may be impacting these species.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Define foraging and nesting requirements of each species within Vermont, utilizing current knowledge of researchers and field investigations.
Research	Basic Life History	High	Determine over wintering habitat needs and survival rate of queens.
Research	Other Research	High	Study the affects of land use changes on <i>Bombus</i> species. This may include mowing, development, conversion of open grass areas, succession to forest land.
Monitoring	Population Change	High	1) Monitor known SGCN bumble bee populations. 2) Population monitoring could be employed to track population trends at distinct locations. This would follow field surveys and assessments to identify populations judged to be large and viable. Focusing on such large populations would offer greater probability of detecting population shifts.
Monitoring	Habitat Change	Medium	Track the change in open grassland habitat in Vermont.
Monitoring	Monitor Threats	High	Assess and monitor the use of domesticated <i>Bombus</i> in agriculture.
Monitoring	Other Monitoring Needs	High	Monitor the use of insecticides in Vermont that may be problematic to <i>Bombus</i> . Agricultural use and home use probably represent different threats.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Bumble Bee Group**
 Scientific Name: **Bumble Bee Group**
 Species Group: **Invert**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Alliance Development	High	Participate in region-wide approach to bumble bee conservation.	Development of regional conservation plan or management guidelines.	FWD, Other Northeastern States, USFWS	SWG, RCN, LLC, USFWS
Species Restoration	High	Develop species-specific restoration and reintroduction plans.	Number of plans produced.	FWD, VCE, UVM	SWG
Technical Assistance, Training, Learning Networks	High	Reduce the use of neonicotinoids and other insecticides that bees are vulnerable to, in agricultural, residential, and other settings. Use of education to accomplish this may be best approach.	Number of people reached through outreach efforts.	FWD, AAFM, EPA, USFWS, VCE, USDA	SWG, EPA, AAFM, USFWS
Compatible Resource Use	High	Experiment with and encourage use of pollinator-friendly buffers in agriculture and other areas where Bombus foraging can be enhanced.	Gain enough information to develop a landowner management guide.	FWD, AAFM, NRCS, USFWS, UVM, Middlebury College, VCE	SWG, NRCS, LCC, RCN
Research	High	Study the level and pattern of use of Bombus in agriculture within Vermont; determine whether this is associated with the pattern of decline in Bombus species.	Develop a map of current use of Bombus in hot houses and other agriculture; to be used in analysis of population declines.	FWD, AAFM, VCE, USDA	SWG, AAFM
Research	High	Studying the affects of various grassland management practices on bumble bee diversity and abundance.	Gain enough information to develop a landowner management guide.	FWD, USFWS, UVM, VCE, AAFM, NRCS, Middlebury College	SWG, NRCS, LCC, RCN

Bibliography

Hatfield, R., S. Jepsen, E. Mader, S. H. Black, and M. Shepherd. 2012. Conserving Bumble Bees. Guidelines for Creating and Managing Habitat for America's Declining Pollinators. The Xerces Society for Invertebrate Conservation. Portland, OR.

Kerr, J.T., L.L. Richardson. Et al. 2015. Climate change impacts on bumblebees converge across continents. Science 10 July 2015: 349 (6244), 177-180. DOI:10.1126/science.aaa7031

McFarland, K., and S. Zahendra. 2010. Vermont Butterfly Survey 2002-2007; final report to the Natural Heritage Program of the Vermont Department of Fish and Wildlife. Vermont Center for Ecostudies.

Williams, P., R. Thorp, L. Richardson, and S. Colla. 2014. Bumblebees of North America, an identification guide. Princeton University Press, Princeton.



Common Name: **Beetles-Carabid Group**
Scientific Name: **Beetles-Carabid Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Unknown

Extirpated in VT?

Regional SGCN?

Assessment Narrative:

This group contains a great number of species that are ranked as rare, but for which more information is needed before conservation strategies can be developed. Compilation of existing information as well as gathering new data is required. Our understanding of distribution, abundance, and status of the many rare species in this group is limited. Existing information is currently being gathered and compiled. These beetles vary in their distribution and habitat requirements. This group includes 30 species:

- Agonum crenistriatum*: two VT locations. Uses vulnerable habitat.
- Agonum darlingtoni*: 4 locations. Habitat specialist; uses vulnerable habitat.
- Agonum decorum*: nineteen locations. Occurs along much of Lake Champlain shoreline. Remove
- Agonum moerens*: three locations. Low elevation species near Lake Champlain.
- Agonum picicornoides*: six locations. Habitat specialist.
- Agonum punctiforme*: one location. Uses human-influenced areas.
- Agonum superioris*: ten locations. Variety of locations in state. Northern species; may be vulnerable to climate change.
- Amara erratica*: one location. Uses vulnerable habitat.
- Amara laevipennis*: ten locations. Uses forest openings, including human influenced sites.
- Apristus latens*: two locations. Specialist of habitat along rivers (dry, unshaded sand).
- Atranus pubescens*: two locations. Found in beaver lodges.
- Bembidion affine*: three locations. Southern species that extends well to south of VT.
- Bembidion cordatum*: one location. May be recent "irruption" from western populations.
- Bembidion grapii*: seven locations. Uses vulnerable habitat.
- Bembidion mutatum*: six locations. High mountain, relict populations in VT.
- Bembidion quadratum*: two locations. Uses vulnerable habitat. Northern species on edge of range in VT.
- Bembidion robusticolle*: one location. On northeast edge of range in VT; uses common habitat.
- Bembidion rolandi*: ten locations. Sites include many on Lake Champlain.
- Bembidion rufotinctum*: six locations. Habitat specialist.
- Blethisa hudsonica*: six locations. Status uncertain.
- Blethisa julii*: one location. Found only at one somewhat unique site.
- Blethisa quadricollis*: two locations. Uses specialized habitat.
- Carabus goryi*: thirteen locations. Appears to be advancing northward.
- Carabus maeander*: six locations. Many locations on or near Lake Champlain.
- Dicaelus dilatatus dilatatus*: three locations. Habitat specialist.
- Dicaelus teter*: three locations. Uses localized habitat.
- Dicheirotrichus cognatus*: four locations. Uses rare habitat.
- Diplocheila impressicollis*: five locations. Uses common habitat over a large area of VT.
- Diplocheila striatopunctata*: six locations. Uses specialized habitat.
- Dyschirius brevispinus*: one location. On northeast edge of range; uses human-influenced habitat.
- Dyschirius erythrocerus*: six locations. On northeast edge of range; status uncertain.
- Dyschirius politus politus*: seven locations. Northern species on edge of range.
- Elaphropus dolosus*: three locations. Western species on edge of range in VT; possible recent arrival.
- Elaphropus levipes*: one location. Status uncertain.
- Elaphrus fuliginosus*: five locations. Rare even though VT is within the general range.
- Geopinus incrassatus*: four locations. Sites are localized along one river.



Common Name: Beetles-Carabid Group
Scientific Name: Beetles-Carabid Group
Species Group: Invert

- Harpalus fulvilabris: four locations. Limited occurrences, including high elevation sites; unclear whether a habitat specialist.
- Harpalus indigenus: six locations. Found over large area in VT in generalized habitat.
- Harpalus providens three locations. Appears to be somewhat of a habitat generalist over a large area of VT.
- Lophoglossus scrutator: four locations. Localized in VT.
- Nebria suturalis: one location. Highly localized in specialized habitat (Mt. Mansfield).
- Notiobia sayi: three locations. Though limited area of occurrence, it utilizes common habitat.
- Notiophilus aquaticus: two locations. Although one occurrence is high elevation, it may utilize more common habitats.
- Notiophilus borealis: one location. Apparently limited to one site on rare habitat (Mt. Mansfield).
- Notiophilus nemoralis: nine locations. High elevation specialist; habitat may be vulnerable to climate change.
- Notiophilus novemstriatus: one location. Though localized, it is a southern species on edge of range.
- Olisthopus micans: four locations. Specialist of habitat that is limited along Lake Champlain.
- Patrobus foveocollis: two locations. Specialist of high elevation sites. Northern species on edge of range; may be vulnerable to climate change.
- Pentagonica picticornis: four locations. Habitat specialist, but on fairly common habitat. Southern species on edge of range.
- Pericompsus ephippiatus: four locations. Southern species on edge of range.
- Philodes alternans: one VT location. Rare species, though not on edge of range.
- Philodes rectangularis: two VT locations. Northern species on edge of range. Uses common habitat by Lake Champlain.
- Platynus cincticollis: three locations. Utilizes variety of habitats. Southern species.
- Platypatrobus lacustris: three locations. Probably more widespread than occurrences indicate, due to difficulty of collection.
- Pseudamara arenaria: six locations. Most specimens from mid-, high elevation; but some habitat uncertainty.
- Pterostichus brevicornis brevicornis: seven locations. High elevation specialist; mountain crests. Northern species; on edge of range.
- Pterostichus castor: six locations. Common habitat type.
- Pterostichus pinguedineus: three locations. High elevation and habitat specialist. Northern species on edge of range.
- Pterostichus punctatissimus: fourteen locations. Several occurrences, but specialized on rare habitats in limited area.
- Scaphinotus bilobus: three locations. Status uncertain.
- Schizogenius ferrugineus: two locations. Few occurrences, but in common habitat. Southern species.
- Sericoda obsoleta: two locations. Status uncertain.
- Sericoda quadripunctata: three locations. Widespread species; status uncertain.
- Sphaeroderus nitidicollis: six locations. Northern species with relict populations in Adirondacks and New England mountains.
- Tachys oblitus: six locations. Occurs over wide area of VT; uses common habitat.
- Tachys rhodeanus: two locations. Only two occurrences despite use of common habitat and VT being in interior of range.
- Tetragonoderus fasciatus: four locations. Widespread species to south and west of VT. On edge of range.
- Tetraleucus picticornis: one location. Widespread species to south and west of VT. On edge of range.



Common Name: **Beetles-Carabid Group**
Scientific Name: **Beetles-Carabid Group**
Species Group: **Invert**

Distribution

Information is being gathered and compiled regarding known occurrences of these species. Biophysical regions of known collection sites are reflected below for each species. Additional collection effort is needed to gain a more complete picture of distributions.

- Agonum crenistriatum: Champlain Valley
- Agonum darlingtoni: Champlain Valley, Northern Green Mountains, Southern Green Mountains
- Agonum moerens: Champlain Valley
- Agonum picicornoides: Northern Green Mountains, Southern Green Mountains
- Agonum punctiforme: Vermont Valley
- Agonum superioris: Champlain Valley, Northern Highlands, Northern Green Mountains, Southern Green Mountains, Southern Vermont Piedmont
- Amara erratica: Northern Green Mountains
- Amara laevipennis: Northern Highlands, Northern Green Mountains, Taconic Mountains, Southern Vermont Piedmont
- Aristus latens: Northern Green Mountains, Southern Vermont Piedmont
- Atranus pubescens: Northern Green Mountains
- Bembidion affine: Southern Green Mountains, Southern Vermont Piedmont
- Bembidion cordatum: Champlain Valley
- Bembidion grapii: Northern Green Mountains, Southern Green Mountains
- Bembidion mutatum: Northern Green Mountains
- Bembidion quadratum: Northern Highlands, Northern Green Mountains
- Bembidion robusticollis: Champlain Valley
- Bembidion rolandi: Champlain Valley, Southern Green Mountains, Taconic Mountains
- Bembidion rufotinctum: Champlain Valley, Northern Vermont Piedmont, Southern Vermont Piedmont
- Blethisa hudsonica: Champlain Valley
- Blethisa julii: Northern Green Mountains
- Blethisa quadricollis: Northern Highlands, Northern Green Mountains
- Carabus goryi: all biophysical regions possible except Northern Highlands
- Carabus maeander: Champlain Valley, Champlain Hills
- Dicaelus dilatatus dilatatus: Champlain Valley, Northern Vermont Piedmont
- Dicaelus teter: Champlain Valley
- Dicheirotichus cognatus: Northern Highlands, Northern Green Mountains
- Diplocheila impressicollis: Champlain Valley, Northern Vermont Piedmont
- Diplocheila striatopunctata: Champlain Valley
- Dyschirius brevispinus: Champlain Valley
- Dyschirius erythrocerus: Champlain Valley, Vermont Valley
- Dyschirius politus politus: Champlain Valley, Northern Highlands
- Elaphropus dolosus: Champlain Valley, Southern Vermont Piedmont
- Elaphropus levipes: Champlain Valley
- Elaphrus fuliginosus: Champlain Valley, Northern Highlands, Northern Green Mountains, Northern Vermont Piedmont
- Geopinus incrassatus: Champlain Valley
- Harpalus fulvilabris: Northern Green Mountains, Southern Green Mountains
- Harpalus indigenus: Northern Highlands, Northern Green Mountains, Northern Vermont Piedmont, Southern Vermont Piedmont, Taconic Mountains, Champlain Valley
- Harpalus providens: Champlain Valley, Vermont Valley
- Lophoglossus scrutator: Champlain Valley
- Nebria suturalis: Northern Green Mountains
- Notiobia sayi: Champlain Valley

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Beetles-Carabid Group**
Scientific Name: **Beetles-Carabid Group**
Species Group: **Invert**

--Notiophilus aquaticus: Champlain Valley, Northern Green Mountains
--Notiophilus borealis: Northern Green Mountains
--Notiophilus nemoralis: Northern Green Mountains, Southern Green Mountains, Taconic Mountains
--Notiophilus novemstriatus: Champlain Valley
--Olisthopus micans: Champlain Valley
--Patrobus foveocollis: Northern Green Mountains
--Pentagonica picticornis: Champlain Valley, Northern Green Mountains
--Pericompso ephippiatus: Southern Vermont Piedmont
--Philodes alternans: Taconic Mountains
--Philodes rectangularis: Champlain Valley
--Platynus cincticollis: Champlain Valley, Northern Green Mountains
--Platypatrobus lacustris: Champlain Valley, Northern Green Mountains
--Pseudamara arenaria: Champlain Valley, Northern Green Mountains
--Pterostichus brevicornis brevicornis: Northern Green Mountains
--Pterostichus castor: Northern Green Mountains
--Pterostichus pinguedineus: Northern Green Mountains
--Pterostichus punctatissimus: Northern Green Mountains
--Scaphinotus bilobus: Northern Highlands, Northern Green Mountains, Southern Vermont Piedmont
--Schizogenius ferrugineus: Champlain Valley, Northern Highlands
--Sericoda obsoleta: Champlain Valley, Northern Green Mountains
--Sericoda quadripunctata: Champlain Valley, Northern Green Mountains, Southern Vermont Piedmont
--Sphaeroderus nitidicollis: Northern Highlands, Northern Green Mountains, Northern Vermont Piedmont
--Tachys oblitus: Champlain Valley, Northern Green Mountains, Southern Vermont Piedmont
--Tachys rhodeanus: Champlain Valley, Southern Vermont Piedmont
--Tetragonoderus fasciatus: Champlain Valley, Southern Vermont Piedmont
--Tetraleucus picticornis: Champlain Valley

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Existing information on habitat use and requirements of these beetles is being gathered and compiled. Much work is still needed to better defined habitat use and needs. Some are known to use specialized habitats and natural communities.

--Agonum crenistriatum: One from an area of sand dunes (since destroyed). The other were on limestone pavements (alvars). Elsewhere this species has been collected from other hot, dry habitats, such as gravel pits, sandy fields and croplands, sea and lake shores.



Common Name: Beetles-Carabid Group
Scientific Name: Beetles-Carabid Group
Species Group: Invert

- Agonum darlingtoni*: Sphagnum bogs.
- Agonum moerens*: Soft, wet mud by ponds, streams, and fens. In VT restricted to very low elevations, below 35 m near Lake Champlain.
- Agonum picicornoides*: In VT, usually by beaver ponds, bare mud in willow, alder thickets. Usually at 300 m or above (never at Sphagnum bogs).
- Agonum punctiforme*: Open areas such as croplands, pastures, also forests.
- Agonum superioris*: Among emergent vegetation in marshes, bogs, and swamps, usually above 300 m.
- Amara erratica*: Grasslands on high mountains.
- Amara laevipennis*: Clearings in forests, such as beaver ponds, clearcuts, roadsides.
- Apristus latens*: Dry, unshaded sand along rivers.
- Atranus pubescens*: In VT, taken from an abandoned beaver house. Elsewhere, it has been found in heaps of leaves, sticks, and mud along rivers and brooks (flood debris) as well as beaver houses. The larva has been found in beaver houses.
- Bembidion affine*: VT specimens were taken on bare, wet mud on the margins of marshy pools.
- Bembidion cordatum*: Mud or muddy sand beside lakes, ponds, impounded sections of streams.
- Bembidion grapii*: Restricted to high mountains where it is found on rocky summits which are bare or have only small or stunted trees. Most records are from 1200 m or higher and are associated with some tundra plants.
- Bembidion mutatum*: High mountain relict populations. In VT, collected under dry clumps of moss on barren, smooth bedrock. Records are mostly from alpine tundra. Further north, it has been recorded from barren spots on dry moraines. More northern records from QC are from roadsides, fields, and sand pits.
- Bembidion quadratum*: Sphagnum mats of bogs.
- Bembidion robusticollis*: Sand banks by rivers.
- Bembidion rolandi*: Gravelly areas along lakes and rivers, especially in the angular shale gravel below bluffs along Lake Champlain.
- Bembidion rufotinctum*: On rock ledges along big rivers by rapids or falls. Usually they are within a meter of the water's edge where spray moistens and cools the rocks. Often there are thin mats of hair-like green algae. They can be found on isolated rocks or islets within the rapids.
- Blethisa hudsonica*: Floating mats of vegetation in lakes and still portions of rivers.
- Blethisa julii*: In NH, occurs in moss and grass beside small high elevation lakes. VT location is a lower (220m) elevation pond, which is shaded most of the day.
- Blethisa quadricollis*: Sphagnum mats in acid bogs.
- Carabus goryi*: Deciduous forest at least to 300 m elevation.
- Carabus macander*: An amphibious species inhabiting swampy spots with shallow water usually with cattails (Typha) or sedges (Carex).
- Dicaelus dilatatus dilatatus*: Dry deciduous forests and sand areas. Adapted to dry conditions.
- Dicaelus teter*: Deciduous forests, especially oaks growing on limestone. Forages at night on fallen logs and climbs standing trees. Recorded as feeding on snails and caterpillars.
- Dicheirotichus cognatus*: Open areas just below the tree line, and in the alpine tundra. Has been taken several times in beaver houses.
- Diplocheila impressicollis*: In cattail (Typha) marshes and other wetlands.
- Diplocheila striatopunctata*: In VT, only in bottomland, swamp forests by Lake Champlain.
- Dyschirius brevispinus*: VT specimens were taken under small stones on bare cultivated soil. Elsewhere it has been collected in gravel pits.
- Dyschirius erythrocerus*: Recorded habitats include river banks, lake shore, and sea beaches. Most of our specimens however, were caught in light traps.
- Dyschirius politus politus*: According to general literature, it is found in sand by rivers and lakes but also in sand pits. It is rarely found in riverside or lakeside sand in the Burlington area where the similar *D. sphaericollis* is abundant. Relatively common in light traps despite its apparent rarity along rivers.
- Elaphropus dolosus*: Bare sand along rivers and lake shores.



Common Name: Beetles-Carabid Group
Scientific Name: Beetles-Carabid Group
Species Group: Invert

- Elaphropus levipes: River banks on muddy sand. Shelters by day in soil or under bark.
- Elaphrus fuliginosus: Habitat difficult to characterize and variously described in literature, but generally open places with sparse vegetation on wet, sandy or fine mud soils; sometimes with mosses such as sphagnum.
- Geopinus incrassatus: Sand or very sandy soil, especially in the higher, drier parts of sand banks along rivers. A specialized burrower, usually deeply buried by day.
- Harpalus fulvilabris: Open or partially shaded areas in the mountains, including exposed bedrock.
- Harpalus indigens: Open areas on poor soil, usually sand, but also found on a sterile, sloping field of clay and gravel.
- Harpalus providens: A forest species, with records from a forestry plantation on sand, and from an open oak forest on a limestone ridge.
- Lophoglossus scrutator: In VT, on very soft mud on natural levees within river delta; most found under large logs embedded in the soft, shaded mud. Reported elsewhere from open marsh habitats among dense vegetation.
- Nebria suturalis: In VT, confined to above 1200 m elevation on Mount Mansfield. Confined to series of deep joint crevices in warm or dry weather; in cool, wet weather may extend to nearby fell field and talus. Larva has been found in crevices where snow lingers until early summer.
- Notiobia sayi: Sand areas near Lake Champlain among sparse vegetation, including croplands.
- Notiophilus aquaticus: Relatively dry, open ground in alpine tundra on Mt. Mansfield, but also found on bare soil in a low elevation apple orchard near Lake Champlain.
- Notiophilus borealis: Alpine tundra. Reported from dry moss.
- Notiophilus nemoralis: Found amid litter and mosses in spruce-fir forests, 900 – 1200 m elevation.
- Notiophilus novemstriatus: General habitat is dry forest edges.: VT specimens found in tufts of grass on quartzite ledges above cliffs. Reported from IL in oak forest at edge of shale bluffs along river valleys.
- Olisthopus micans: By river mouths, in forests that are flooded by Lake Champlain in the springtime.
- Patrobus foveocollis: Dry openings in spruce-fir forest of high elevation (900-1200 m).
- Pentagonica picticornis: In VT, quartzite and schist rock ledges. Also reported as found under moss clumps on boulders.
- Pericompso ephippiatus: Found only on short stretch of Connecticut River in southeastern VT.: Barren or sparsely vegetated sand bars where sand is usually dry but close to the water.
- Philodes alternans: Found under a large flat stone at the margin of the Battenkill, below an eroding bank about 1.3 m height.
- Philodes rectangularis: records are from the wet mud banks by Lake Champlain at the mouths of two small rivers, elevation about 30 m.
- Platynus cincticollis: Floodplain forests and forests bordering ponds and slow streams. Occasionally found in beaver houses, tree cavities, or under plant debris on beaches.
- Platypatrobus lacustris: Active beaver huts (beaver present).
- Pseudamara arenaria: Usually above 400 m elevation, in mountain forests.: Some habitat uncertainty.
- Pterostichus brevicornis brevicornis: High, cold parts of the coniferous forest; 750-1130 m elevation.
- Pterostichus castor: Beaver houses, both active and abandoned.
- Pterostichus pinguedineus: Most specimens collected in deep rock crevices at or above tree line on mountain tops. Share habitat with Nebria suturalis. Lowest VT record is 750 m elevation; others all above 1200 m.
- Pterostichus punctatissimus: Boreal species. Two habitat types: (1) in higher mountains above 900 m, found under cover (especially mosses) in fir and spruce forests; (2) near some bogs where cold air accumulates
- Scaphinotus bilobus: Generally spruce forests.
- Schizogenius ferrugineus: Clean sand or sandy fields, often near water. Found under woody debris or in grass tufts.
- Sericoda obsoleta: Found in wood ashes. Have been taken under bark of standing dead trees, in fire places in campgrounds, and in houses which have wood stoves or furnaces. May arrive at sites while fires are still burning.
- Sericoda quadripunctata: Found in wood ashes. Have been taken under bark of standing dead trees, in fire places in campgrounds, and in houses which have wood stoves or furnaces. May arrive at sites while fires are



Common Name: Beetles-Carabid Group
Scientific Name: Beetles-Carabid Group
Species Group: Invert

still burning.

--Sphaeroderus nitidicollis: Coniferous and mixed forest, generally higher elevation in VT, 600-750 m.

--Tachys oblitus: Muddy borders of lakes and rivers, usually in vegetation.

--Tachys rhodeanus: On margins of slow rivers, lakes, and marshes; on very moist soil which is bare or with sparse vegetation.

--Tetragonoderus fasciatus: Dry sand areas near large rivers and lakes; found in sunny spots, but usually near shade. Less than 90 m elevation.

--Tetraleucus picticornis: South of VT, reported from cypress swamps and along swampy shore of rivers.: VT specimens were found beneath driftwood along a seasonally flooded ditch beside a dirt road within forest.

Habitat Types:

Upland Shores

Outcrops and Alpine

Cliffs and Talus

Spruce Fir Northern Hardwood

Northern Hardwood

Oak-Pine Northern Hardwood

Open Peatlands

Marshes and Sedge Meadows

Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Lawns, Gardens, and Row Crops

Wet Shores

Shrub Swamps

Building or Structure

Other Cultural

Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat

Energy Infrastructure and Development

Habitat Succession

Habitat Alteration

Sedimentation

Habitat Fragmentation

Impacts of Roads or Transportation Systems

Climate Change

Incompatible Recreation



Common Name: **Beetles-Carabid Group**
Scientific Name: **Beetles-Carabid Group**
Species Group: **Invert**

Description of habitat threat(s): Habitat problems are known for some species, and are related to habitat loss, change, and degradation. Being rare species, habitat fragmentation would lead to smaller, more vulnerable populations.

Non-Habitat Threats:

Genetics

Pollution

Trampling or Direct Impacts

Unknown Non-Habitat Threats

Description of non-habitat threat(s): The problems not related to habitat are poorly known for these beetles and need study. As rare species with often small populations, loss of metapopulation structure and function would be a problem. Some species are alpine, where heavy recreational use can result in trampling.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Gather information on habitats in which each SGCN carabid species reportedly occurs (literature research, consult researchers and hobbyists, etc.); this will be needed to refine distributional field surveys.
Research	Basic Life History	High	Life history information is needed for all species
Research	Distribution and Abundance	High	Conduct literature research and field surveys to update information on distribution of SGCN carabid species in Vermont.
Research	Threats and Their Significance	High	Research is needed on the vulnerability of species to various significant limiting factors to each habitat type.
Monitoring	Population Change	High	Revisit and survey sites with previous records of SGCN carabids to determine presence/absence; where present, determine
Monitoring	Range Shifts	High	Document changes in distribution, utilizing historic and recent records.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Beetles-Carabid Group**
 Scientific Name: **Beetles-Carabid Group**
 Species Group: **Invert**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	High	Visit known sites of occurrences which has undergone recent development or other disturbance.	Number of sites visited.	FWD, VMC	SWG
Technical Assistance, Training, Learning Networks	High	Sponsor training workshops for carabid identification, survey techniques, web database use	Number of workshops; number of participants	FWD, FPR, VCE, Eagle Hill	SWG
Research	High	Revisit historic sites to update records and monitor species.	Number of historic sites visited.	FWD	SWG
Alliance Development	High	Develop web-based database accessible to professionals and site record providers.	Publication on web site	FPR, FWD, VCE, VT Entomological Society, Carabid specialists	SWG, Lintillac Foundation
Awareness Raising and Communications	High	Publish "Carabidae of Vermont and New Hampshire", which is currently in final draft, as a hard copy and on-line resource.	Availability of Carabidae information	FPR, FWD, VCE, VMC, VT Entomological Society	SWG, Lintillac Foundation, UVM



Common Name: Beetles-Carabid Group
Scientific Name: Beetles-Carabid Group
Species Group: Invert

Bibliography

- Allegro, G., and R. Sciaky. 2003. Assessing the potential role of ground beetles (Coleoptera, Carabidae) as bioindicators in poplar stands, with a newly proposed ecological index (FAI). *Forest Ecology and Management* 175:275-284.
- Andersen, J. 1985. Humidity responses and water balance of riparian species of Bembidiini (Col., Carabidae). *Ecological Entomology* 10:363-375.
- Andersen, J. 1995. A comparison of pitfall trapping and quadrat sampling of Carabidae (Coleoptera) on river banks. *Entomologica Fennica* 6:65-77.
- Anonymous. 1991. Endangered and threatened wildlife and plants; animal candidate review for listing as endangered or threatened species, proposed rule. Federal register. Department of the Interior. Fish and Wildlife Service.
- Antvogel, H., and A. Bonn. 2001. Environmental parameters and microspatial distribution of insects: a case study of carabids in an alluvial forest. *Ecography* 24:470-482.
- Baars, M. A. 1979a. Catches in pitfall traps in relation to mean densities of carabid beetles. *Oecologia* 41:25-46.
- Ball, G.E. 1979. Carabidae. Pp. 364-366 in H.V. Danks (Ed.). *Canada and its insect fauna. Memoirs of the Entomological Society of Canada* No. 108. 573 pp.
- Belaoussoff, S., P. G. Kevan, S. Murphy, and C. Swanton. 2003. Assessing tillage disturbance on assemblages of ground beetles (Coleoptera: Carabidae) by using a range of ecological indices. *Biodiversity and Conservation* 12:851-882.
- Bousquet, Y. 1986. Observations on the life cycle of some species of Pterostichus (Coleoptera: Carabidae) occurring in northeastern North America. *Le Naturaliste canadien* 113 : 295-307.
- Bousquet, Y. 1987. The carabid fauna of Canada and Alaska: range extensions, additions and descriptions of two new species of Dyschirius (Coleoptera: Carabidae). *The Coleopterists Bulletin* 41 : 111-135.
- Bousquet, Y. (Ed.). 1991. Checklist of beetles of Canada and Alaska. Agriculture Canada Publication 1861/E. 430 pp.
- Burel, F. 1989. Landscape structure effects on carabid beetles spatial patterns in western France. *Landscape Ecology* 2:215-226.
- Butterfield, J. 1996. Carabid life-cycle strategies and climate change: a study on an altitude transect. *Ecological Entomology* 21:9-16.
- Campbell, J.M. 1980. Distribution patterns of Coleoptera in eastern Canada. *The Canadian Entomologist* 112 : 1161-1175.
- Darlington, P.J. 1938. The American Patrobini. *Entomologica Americana* 18 : 135-183.
- Davidson, R. 1975. Harpalus (Ophonus) puncticeps Stephens (Coleoptera: Carabidae) in New York and Vermont. *The Coleopterists Bulletin* 29 : 256.
- Davies, K. F., and C. R. Margules. 1998. Effects of habitat fragmentation on carabid beetles: experimental evidence. *Journal of Animal Ecology* 67:460-471.
- Dietrich, H. 1957. Harpalus puncticeps Steph. On Long Island, N.Y. *The Coleopterists Bulletin* 11 : 46.
- Duval, P. & J. Duval. 1977. Première mention de l'Harpalus puncticeps Stephens (Coleoptera: Carabidae) pour le Maine, Etats-Unis d'Amérique. *Cordulia* 3 : 165-166.
- Epstein, M. E., and H. M. Kulman. 1990. Habitat distribution and seasonal occurrence of carabid beetles in East-central Minnesota. *The American Midland Naturalist* 123:209-225.
- Goulet, H. 1965. The habitat of Platypatrobus Darlington (Coleoptera: Carabidae). *Psyche* 72 : 305-306.
- Goulet, H. 1977. Contributions of characters of larvae to systematics of Carabidae. Pp. 205-208 in T.L. Erwin, G.E. Ball, D.R. Whitehead and A.L. Halpern (eds), *Carabid beetles: their evolution, natural history, and classification*. Junk, The Hague.
- Greene, A. 1975. Biology of the five species of Cychrini (Coleoptera: Carabidae) in the steppe region of southeastern Washington. *Melanderia* 19:1-43.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Beetles-Carabid Group
Scientific Name: Beetles-Carabid Group
Species Group: Invert

Halme, E., and J. Niemälä. 1993. Carabid beetles in fragments of coniferous forest. *Annales Zoologici Fennici* 30:17-30.

Halsall, N. B., and S. D. Wratten. 1988. The efficiency of pitfall trapping for polyphagous predatory Carabidae. *Ecological Entomology* 13:293-299.

Heliola, J., M. Koivula, and J. Niemälä. 2001. Distribution of carabid beetles (Coleoptera, Carabidae) across a boreal forest-clearcut ecotone. *Conservation Biology* 15:370-377.

Honek, A. 1997. The effect of temperature on the activity of Carabidae (Coleoptera) in a fallow field. *European Journal of Entomology* 94:97-104.

Joyce, K. A., J. M. Holland, and C. P. Doncaster. 1999. Influences of hedgerow intersections and gaps on the movement of carabid beetles. *Bulletin of Entomological Research* 89:523-531.

Koivula, M. 2002. Alternative harvesting methods and boreal carabid beetles (Coleoptera, Carabidae). *Forest Ecology and Management* 167:103-121.

LaBonte, J. R. 2002. Habitat associations of riparian beetles (Coleoptera) at Big Beaver Creek Research Natural Area, North Cascades National Park, Washington. M.S. thesis. Oregon State University, Corvallis, Oregon.

Larochelle, A. 1975. Les Carabidae du Québec et du Labrador. *Bulletin du Département de Biologie du Collège Bourget, Rigaud* 1 : 1-255.

Larochelle, A., and M.-C. Larivière. 2003. A natural history of the ground-beetles (Coleoptera: Carabidae) of America north of Mexico. *Pensoft Series Faunistica* No. 27, Sofia, Bulgaria.

Lindroth, C.H. 1957. The faunal connections between Europe and North America. Stockholm. 344 pp.

Lindroth, C. 1961. The ground-beetles (Carabidae excl. Cicindelinae) of Canada and Alaska. Part 2. *Opuscula Entomologica Supplementum* 20 : 1-200.

Lindroth, C. 1963. The ground-beetles (Carabidae excl. Cicindelinae) of Canada and Alaska. Part 3. *Opuscula Entomologica Supplementum* 24 : 201-408.

Lindroth, C. 1966. The ground-beetles (Carabidae excl. Cicindelinae) of Canada and Alaska. Part 4. *Opuscula Entomologica Supplementum* 29 : 409-648.

Lindroth, C. 1968. The ground-beetles (Carabidae excl. Cicindelinae) of Canada and Alaska. Part 5. *Opuscula Entomologica Supplementum* 33 : 649-944.

Lindroth, C. 1969. The ground-beetles (Carabidae excl. Cicindelinae) of Canada and Alaska. Part 6. *Opuscula Entomologica Supplementum* 34 : 945-1192.

Magura, T., B. Tothmeresz, and T. Molnar. 2001. Forest edge and diversity: carabids along forest-grassland transects. *Biodiversity and Conservation* 10:287-300.

Neve, G. 1994. Influence of temperature and humidity on the activity of three Carabus species. Pages 189-192 in K. Desender, M. Dufrene, M. Loreau, M. L. Luff, and J.-P. Maelfait, editors. *Carabid beetles: ecology and evolution*. Kluwer Academic Publishers, Dordrecht, The Netherlands.

Niemälä, J., Y. Haila, E. Halme, T. Lahti, T. Pajunen, and P. Punttila. 1988. The distribution of carabid beetles in fragments of old coniferous taiga and adjacent managed forests. *Zoologici Fennici* 25:107-119.

Niemälä, J., D. Langor, and J. R. Spence. 1993a. Effects of clear-cut harvesting on boreal ground-beetle assemblages (Coleoptera: Carabidae) in western Canada. *Conservation Biology* 7:551-561.

Niemälä, J., J. R. Spence, D. Langor, Y. Haila, and H. Tukia. 1993b. Logging and boreal ground-beetle assemblages on two continents: implications for conservation. Pages 29-50 in K. J. Gaston, T. R. New, and M. J. Samways, editors. *Perspectives on insect conservation*. Intercept, Ltd., Andover, United Kingdom.

Rykken, J. J., D. E. Capen, and S. P. Mahabir. 1997. Ground beetles as indicators of land type diversity in the Green Mountains



Common Name: Beetles-Carabid Group
Scientific Name: Beetles-Carabid Group
Species Group: Invert

of Vermont. Conservation Biology 11:522-530.

Spence, J. R. 1979. Riparian carabid guilds--a spontaneous question generator. Pages 525-537 in T. L. Erwin, G. E. Ball, and D. W. Whitehead. Carabid beetles: their evolution, natural history, and classification. W. Junk Publishers, The Hague, The Netherlands.

Spence, J. R., D. W. Langor, J. Niemälä, H. A. Carcamo, and C. R. Currie. 1996. Northern forestry and carabids: the case for concern about old-growth species. *Annales Zoologici Fennici* 33:173-184.

Spence, J. R., and J. K. Niemälä. 1994. Sampling carabid assemblages with pitfall traps: the madness and the method. *The Canadian Entomologist* 126:881-894.

Spence, J.R. and D.H. Spence. 1988. Of ground-beetles and man: introduced species and the synanthropic fauna of western Canada. *Memoirs of the Entomological Society of Canada* 144 : 151-168.

Strong, A. M., C. A. Dickert, and R. T. Bell. 2002. Ski trail effects on a beetle (Coleoptera: Carabidae, Elateridae) community in Vermont. *Journal of Insect Conservation* 6: 149-159.

Sustek, Z. 1994a. Classification of the carabid assemblages in the floodplain forests in Moravia and Slovakia. Pages 371-376 in K. Desender, M. Dufrene, M. Loreau, M. L. Luff, and J.-P. Maelfait, editors. Carabid beetles: ecology and evolution. Kluwer Academic Publishers, Dordrecht, The Netherlands.

Thiele, H.-U. 1977. Carabid beetles in their environments. A study on habitat selection by adaptations in physiology and behaviour. Springer-Verlag, Berlin. Xvii + 369 pp.

Thompson, R.G. 1977. A synoptic list of the described ground beetle larvae of North America (Coleoptera: Carabidae). *Proceedings of the Biological Society of Washington* 90 : 99-107.

Trudel-Levesque, C. 1971. Etude des Coléoptères Carabidae de forêts décidues des Laurentides (Québec), particulièrement en fonction de l'activité locomotrice. M.Sc. Thesis, Université de Montréal. 86 pp.

Zulka, K. P. 1994. Carabids in a Central European floodplain: species distribution and survival during inundations. Pages 399-405 K. Desender, M. Dufrene, M. Loreau, M. L. Luff, and J.-P. Maelfait, editors. Carabid beetles: ecology and evolution. Kluwer Academic Publishers, Dordrecht, The Netherlands



Common Name: **Beetles-Tiger Beetle Group**
Scientific Name: **Beetles-Tiger Beetle Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

This group includes three state-threatened species, one of which is also federally threatened, and four rare species, three of which are known from only one or two collections and includes:

--Boulder-beach Tiger Beetle (*Cicindela ancocisconensis*): This rarely observed beetle is known from only single collections on two widely separated rivers. Globally rare (G3), RSGCN*. Species appears to be in decline globally.

--Hairy-necked Tiger Beetle (*Cicindela hirticollis*): This state-threatened beetle's habitat and range has been greatly reduced and fragmented in Vermont by lakeshore development. It was formerly known from several sites along the northern Lake Champlain shores, but is now reduced to a single site. This habitat is protected by the Winooski Valley Park District.

--Boreal Long-lipped Tiger Beetle (*Cicindela longilabris*): There are few records of this little known beetle in VT. It is a northern species found in VT at moderately high elevations. More survey work is needed.

--Cobblestone Tiger Beetle (*Cicindela marginipennis*): There are few records of this state-threatened species scattered around the state in uncommon habitat. It has been studied in VT to a greater degree than other *Cicindela*. At least one site appears to no longer support the species. Habitat losses along the Connecticut River and possibly other rivers have been significant due to impoundments. Globally rare (G2), RSGCN*.

--Northern Barrens Tiger Beetle (*Cicindela patruela*): This is a very rare species throughout the Northeast; known in VT from a single historic collection. Globally rare (G3), RSGCN*. It uses restricted habitat (sand plains), which has been extensively destroyed in VT and elsewhere; global occurrences are now highly fragmented. It is now a rare, relict species.

--Puritan tiger beetle (*Cicindela puritana*): This federally threatened species is known from a single historic VT collection, although other historic records were known along the New Hampshire side of the river. Impoundments along the Connecticut River likely caused the extirpation of this species. Other habitat losses may have also been a factor. Reintroduction could be considered if sufficient habitat improvements are made. Riverside recreational use has had a significant impact on populations at other New England sites. RSGCN*

--Eastern Red-bellied Tiger Beetle (*Cicindela rufiventris*): Known from a single VT location. Its status is unknown.

*Regional Species of Greatest Conservation Need (RSGCN) among the 13 Northeastern states

Distribution

--Boulder-beach Tiger Beetle (*Cicindela ancocisconensis*): Reported from Underhill historically. More recently from West River and Third Branch of the White River. Biophysical regions: Southern Green Mountains, Northern Green Mountains.

--Hairy-necked Tiger Beetle (*Cicindela hirticollis hirticollis*): Historically six locations on Lake Champlain, three of which probably represented dispersing individuals. Only a single extant breeding population now known in Colchester. Biophysical regions: Champlain Valley.

--Boreal Long-lipped Tiger Beetle (*Cicindela longilabris*): Three locations. Restricted to the mountains and northern plateau in VT. Biophysical regions: Northern Green Mountains, Northern Highlands.



Common Name: **Beetles-Tiger Beetle Group**
Scientific Name: **Beetles-Tiger Beetle Group**
Species Group: **Invert**

--Cobblestone Tiger Beetle (*Cicindela marginipennis*): Known from West, White, and Winooski rivers. It is reported that populations along the Connecticut River are believed to use islands (NH) for larval sites, though adults will forage on the west (VT) shore. Biophysical regions: Southern Vermont Piedmont, Northern Green Mountains, Champlain Hills.

--Northern Barrens Tiger Beetle (*Cicindela patruela*): Single historic record from Burlington. Sand plains in and around Burlington have largely been destroyed by urban growth. Biophysical regions: Champlain Valley.

--Puritan Tiger Beetle (*Cicindela puritana*): On historic location in Hartland. Biophysical regions: Southern Vermont Piedmont.

--Eastern Red-bellied Tiger Beetle (*Cicindela rufiventris*): One location in Sandgate. Biophysical regions: Taconic Mountains.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Not Probable	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Known Watersheds

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐



Common Name: **Beetles-Tiger Beetle Group**
Scientific Name: **Beetles-Tiger Beetle Group**
Species Group: **Invert**

Habitat Types:

Upland Shores
Outcrops and Alpine
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Open Peatlands
Wet Shores

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Inadequate Disturbance Regime
Climate Change
Sedimentation
Invasion by Exotic Species
Incompatible Recreation

Description of habitat threat(s): Regular scouring of river shores by high water events helps keep habitat open by reducing vegetation; damming of rivers impacts this process downstream of these structures and may degrade habitat. Replenishment of substrates is also reduced downstream of dams, which can alter substrate composition along shores. Excessive fine sediments that enter streams and rivers can alter the substrate composition along shores, thereby reducing the suitability of habitat. Development along shores of Lake Champlain and rivers has reduced the availability of habitat. Loss of sand plain habitat has probably caused extirpation of one species. Northern species on the southern edge of their range in Vermont may be impacted by climate change. Invasive plants that colonize river shores could eliminate tiger beetle populations.

Non-Habitat Threats:

Genetics
Trampling or Direct Impacts

Description of non-habitat threat(s): Small, isolated populations may be at risk genetically. 4-wheelers on rivershores and islands can crush larvae and make habitat unsuitable for sustaining burrows. Use of beaches and sand shores can also cause trampling of areas used by larvae.



Common Name: **Beetles-Tiger Beetle Group**
 Scientific Name: **Beetles-Tiger Beetle Group**
 Species Group: **Invert**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Define particular habitat requirements of each species within Vermont, utilizing current knowledge of researchers and field investigations.
Research	Basic Life History	Medium	Need to rear and describe the larvae of <i>C. marginipennis</i> .
Research	Distribution and Abundance	High	1) Conduct inventories to detect and gather information on new SGCN tiger beetles populations. 2) Obtain baseline distributional and abundance data for all species in group by conducting surveys throughout the state. Efforts should be focused on particular habitats required by each. Abundance information should be collected at sites of known occurrence. Determine if <i>C. marginipennis</i> colonies are disappearing and/or new colonies appearing.
Research	Threats and Their Significance	High	1) Assess potential and existing impacts of limiting factors to habitat. Such limiting factors as habitat loss and degradation, exotic invasive plants, incompatible recreation, and dams should be examined. 2) Investigate how rivershore tiger beetle populations are being affected by dams, and actions that can be taken to restore or mimic natural processes that maintain habitat.
Monitoring	Population Change	High	Monitor known SGCN tiger beetle populations. Track population trends at distinct locations. This would follow field surveys and assessments to identify populations judged to be large and viable. Focusing on such large populations would offer greater probability of detecting population shifts.
Monitoring	Habitat Change	High	Monitor change in available habitat for each species' specific requirements. Loss, restoration, and other changes to local habitat sites recognized as important to these species should be tracked.
Monitoring	Monitor Threats	High	Recreational use of shoreline habitat needs to be monitored, as it can affect several species.



Common Name: **Beetles-Tiger Beetle Group**
Scientific Name: **Beetles-Tiger Beetle Group**
Species Group: **Invert**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Easements	High	Acquisition/easement of high priority SGCN tiger beetle sites	Number of SGCN tiger beetle sites protected	FWD, FPR, USFS, NRCS, VLT, other land trusts	VHCB, SWG, GMNF, EQUIP, USFWS
Compatible Resource Use	High	Work with land owners to direct recreational use away from necessary rivershore tiger beetle habitat	Number of monitored sites where trampling of habitat is eliminated	FWD, watershed groups, local landowners	
Compatible Resource Use	High	Work to restrict recreational vehicles from accessing riverbank and lakeshore SGCN tiger beetle habitat	Number of sites that have eliminated motorized access to SGCN tiger beetle habitat	FWD, NRCS, watershed groups, local landowners	

Bibliography

Bell, R. T. Carabidae of Vermont and New Hampshire. In draft.

Leonard, J. G., and R. T. Bell. 1999. Northeastern Tiger Beetles, A Field Guide to Tiger Beetles of New England and Eastern Canada. CRC Press. 176 pp.



Common Name: **Butterflies-Grassland Group**
 Scientific Name: **Butterflies-Grassland Group**
 Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Declining

Extirpated in VT?

Regional SGCN?

Assessment Narrative:

This group consists of four species. The first three are very rare in Vermont and their habitat requirements vary within the general grasslands category. The fourth, Regal fritillary, is extirpated in Vermont and almost all of the Northeast.

--Cobweb Skipper (*Hesperia metea*): Two recent sight records only; needs further documentation.

--Dusted Skipper (*Atrytonopsis hianna*): First records from Vermont in 2004. Only in Southern Vermont; highest density populations along I-91 where bluestem grasses planted. May benefit and expand northward from additional plantings in appropriate areas.

--Monarch (*Danaus plexippus*): On-going and sharp decline of eastern North American population has led to recent proposal to list the Monarch as threatened under the U.S. Endangered Species Law. A large factor in the species decline may be habitat loss, particularly of milkweed (*Asclepias*), which is the host plant.

--Regal fritillary (*Speyeria idalia*): Extirpated from Vermont; ability to re-establish uncertain.

Distribution

2002-2007 butterfly survey (VBS) records by biophysical region:

--Cobweb skipper: 2 sight records in Taconic Mountains and Vermont Valley

--Regal fritillary: extirpated. Historically from collections in Southern Vermont Piedmont and Vermont Valley

--Dusted skipper: Taconic Mountains, Southern Vermont Piedmont

--Monarch: Northern Highlands, Northern Vermont Piedmont, Northern Green Mountains, Champlain Hills, Champlain Valley, Taconic Mountains, Vermont Valley, Southern Green Mountains, Southern Vermont Piedmont

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:



Common Name: **Butterflies-Grassland Group**
Scientific Name: **Butterflies-Grassland Group**
Species Group: **Invert**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

This group consists of three species, the habitat requirements of which vary within the general grasslands category. Regal fritillary is extirpated in Vermont and all of North America east of the Mississippi River except for two populations in PA and WV. The other species are very rare in Vermont or in the case of the Monarch, may still be somewhat common in optimal years..

--Cobweb Skipper (*Hesperia metea*) Grasslands, old dry fields, and open barrens. Host plants are Little Bluestem (*Schizachyrium scoparius*) and Big Bluestem (*Andropogon gerardi*). Adults prefer nectaring on low-growing plants such as Labrador Tea (*Ledum groenlandicum*), Wild Strawberry (*Fragaria virginiana*), Blackberry (*Rubus allegheniensis*), Winter Cress (*Barbarea vulgaris*), and Red Clover (*Trifolium pratense*).

--Regal fritillary (*Speyeria idalia*) In Massachusetts, seems to have preferred extensive open areas with a combination of wetlands and upland fields containing an abundance of nectaring plants. Host plants are violets (*Viola* sp.).

--Dusted Skipper (*Atrytonopsis hianna*) Open, dry habitats in far southern Vermont valleys with bluestem grasses. Often found in the same habitat as the Cobweb Skipper. Host plants are Little Bluestem (*Schizachyrium scoparius*) and Big Bluestem (*Andropogon gerardi*). Adults nectar from flowers including Japanese Honeysuckle (*Lonicera japonica*), Wild Strawberry (*Fragaria virginiana*), Blackberry (*Rubus allegheniensis*), Phlox (*Phlox*), Vervain (*Verbena*) and Red Clover (*Trifolium pratense*).

--Monarch (*Danaus plexippus*) Prefers open meadows, weedy areas, marshes, roadsides and disturbed habitats with milkweed. Caterpillars feed on Common Milkweed (*Asclepias syriaca*), Swamp Milkweed (*Asclepias incarnata*), and Showy Milkweed (*Asclepias speciosa*). Blooming later summer/early fall clover fields are important stopover habitat in the Champlain and Connecticut valleys. Monarchs occur in Vermont from as early as mid-May to early November; adults migrate south in the fall to reach overwintering habitat in Mexico. Multiple generations are necessary to reach Vermont in the spring/summer.

Habitat Types:

Oak-Pine Northern Hardwood

Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Other Cultural

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Succession

Habitat Alteration

Impacts of Roads or Transportation Systems

Inadequate Disturbance Regime



Common Name: **Butterflies-Grassland Group**
Scientific Name: **Butterflies-Grassland Group**
Species Group: **Invert**

Habitat Fragmentation

Description of habitat threat(s): Loss of grasslands (anthropogenic and natural) and host plants is a threat to members of this group

Non-Habitat Threats:

Pollution

Loss of Prey Base

Trampling or Direct Impacts

Loss of Relationship with Other Species

Description of non-habitat threat(s): --Loss of and impacts to host plants have a negative impact on these grassland butterflies. Increased use of herbicides in agricultural fields may be reducing host plant (milkweeds) density for Monarch in Midwest; impacts in Vermont are unknown.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Define particular habitat requirements of species for which we still have inadequate information, utilizing current knowledge of researchers and field investigations. This has been completed for most SGCN grassland butterflies.
Research	Threats and Their Significance	High	Assess potential and existing impacts of threats to habitat, host plants, and individual butterflies. Such threats as habitat loss and degradation, exotic invasive plants, disease, and host plant loss should be examined.
Research	Population Genetics	Low	Reintroduction of regal fritillary would need to identify the source populations that are likely most similar to those originally occurring in Vermont.
Monitoring	Population Change	Medium	Monitor known SGCN butterfly populations. Track population trends at distinct locations. This would follow field surveys and assessments to identify populations judged to be large and viable. Focusing on such large populations would offer greater probability of detecting population shifts.
Monitoring	Habitat Change	High	Grasslands are some of the habitats most vulnerable to loss due to development and intensive agriculture. Landscape level changes in this general habitat type should be monitored. Loss, restoration, and other changes to local habitat sites recognized as important to these species should be tracked.



Common Name: **Butterflies-Grassland Group**
Scientific Name: **Butterflies-Grassland Group**
Species Group: **Invert**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Workshops	High	Incorporate butterfly SGCN occurrence information into environmental review and technical assistance	Number of sites with butterfly SGCN that received conservation benefits to this group	FWD, USFWS, DEC, ANR, VTrans, NRCS	SWG
Easements	High	Acquisition/easement of high priority SGCN butterfly grassland sites	Number of SGCN butterfly sites protected	FWD, VLT, other land trusts	VHCB, SWG, EQIP, USFWS

Bibliography

- Glassberg, J. 1999. Butterflies through binoculars. A field guide to the butterflies of Eastern North America. Oxford University Press, New York, NY. 242 pp.
- Layberry, R.A., P. W. Hall, and D. J. Lafontaine. 1998. The Butterflies of Canada. University of Toronto Press, Toronto, ON. 280 pp.
- McFarland, K., and S. Zahendra. 2010. Vermont Butterfly Survey 2002-2007; final report to the Natural Heritage Information Project of the Vermont Department of Fish and Wildlife. Vermont Center for Ecostudies.
- NatureServe. 2004. NatureServe Explorer: an online encyclopedia of life [web application]. Version 4.0. NatureServe, Arlington, VA. Available <http://www.natureserve.org/explorer>.
- Opler, P. A., H. Pavulaan, and R. E. Stanford (coordinators). 1995. Butterflies of North America. Jamestown, ND: Northern Prairie Wildlife Research Center Home Page. <http://www.npwrc.usgs.gov/resource/distr/lepid/bflyusa/bflyusa.htm> (Version 12 DEC2003).
- Scott, J. A. 1986. The butterflies of North America. Stanford University Press, Stanford, CA. 583 pp.
- Shapiro, A.M. 1974. Butterflies and Skippers of New York State. Cornell Univ. Agricultural Experimental Station, Ithaca, N.Y. Search 4:1-60.
- Unpublished data by the Vermont Butterfly Survey database as of March 23, 2005. Vermont Institute of Natural Science.
- Wagner, D.L., Nelson, M.W., Schweitzer, D.F. 2003. Shrubland lepidoptera of Southern New England and southeastern New York: ecology, conservation, and management. For. Ecol. Manage. 185: 95-112.
- Williams, B. L. 2002. Conservation genetics, extinction and taxonomic status: a case history for the regal fritillary. Conservation Biology 16 (1): 148-157.



Common Name: **Butterflies-Hardwood Forest Group**
 Scientific Name: **Butterflies-Hardwood Forest Group**
 Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

This group consists of four species, the habitat requirements of which vary within the general hardwood forest category.

--West Virginia white (*Pieris virginiensis*): Threatened by exotic garlic mustard, disease, and poor weather conditions; colonies easily extirpated; does not recolonize isolated sites well.

--Early hairstreak (*Erora laeta*): Widely scattered and localized populations. Beech bark disease is killing off large stands of beech in Vermont, often leaving only 1% of trees remaining. The future for the beech and the early hairstreak in Vermont and much of northeastern North America is uncertain. The beaked hazelnut has also been reported as a host plant further west; the extent to which early hairstreak would adapt to this plant in Vermont is unknown.

--Hackberry emperor (*Asterocampa celtis*): First VT record in 2002. Likely to increase with climate change.

--Tawny emperor (*Asterocampa clyton*): First VT record in 2002. Likely to increase with climate change.

Distribution

2002-2007 butterfly survey (VBS) records by biophysical region:

--West Virginia white: Champlain Valley, Taconic Mountains, Vermont Valley, Southern Green Mountains, Southern Vermont Piedmont

--Early hairstreak: Champlain Valley, Northern Green Mountains, Northern Vermont Piedmont, Southern Green Mountains

--Hackberry emperor: Champlain Valley, Taconic Mountains, Northern Green Mountains; possibly Southern Vermont Piedmont

--Tawny emperor: Champlain Valley, Vermont Valley, Northern Green Mountains, Southern Vermont Piedmont

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		



Common Name: **Butterflies-Hardwood Forest Group**
Scientific Name: **Butterflies-Hardwood Forest Group**
Species Group: **Invert**

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

This group consists of several species, the habitat requirements of which vary within the general hardwood forest category.

--West Virginia White (*Pieris virginiensis*) Requires mature, relatively undisturbed rich hardwood forests with large populations of the host plants, Two-leaved Toothwort (*Dentaria diphylla*) and Cut-leaved Toothwort (*Cardamine concatenata*). Adults nectar from Toothworts, Spring Beauty (*Claytonia virginica*), Violets (*Viola*), and other spring wildflowers.

--Early Hairstreak (*Erora laeta*) requires stands of American Beech (*Fagus grandifolia*), the host plant; failure of the beechnut crop, even for a single year, may seriously impact populations. Adults nectar on fleabane (*Erigeron* sp.) and Ox-eyed Daisy (*Chrysanthemum leucanthemum*). Often found on bare ground puddling.

--Hackberry Emperor (*Astrocampa celtis*) Found in floodplain forests with Northern Hackberry (*Celtis occidentalis*), the host plant. Also reported from suburban Hackberry plantings in Burlington. Adults feed on sap, mud, rotting fruit, and excrement, which can sometimes take them outside of their regular habitat.

--Tawny Emperor (*Astrocampa clyton*) Found in floodplain forests with Northern Hackberry (*Celtis occidentalis*), the host plant. Also reported from suburban Hackberry plantings in Burlington. Adults feed on tree sap, rotting fruit, dung, and carrion.

Habitat Types:

Softwood Swamps
Spruce Fir Northern Hardwood
Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Habitat Fragmentation
Invasion by Exotic Species
Climate Change

Description of habitat threat(s): Early hairstreak is limited by the loss of American beech stands due to beech bark disease. Caterpillars feed on the fruits of this tree, which are only produced by individuals 40 years old or more. Invasion of garlic mustard may threaten the West Virginia White, as the adults will lay



Common Name: Butterflies-Hardwood Forest Group
Scientific Name: Butterflies-Hardwood Forest Group
Species Group: Invert

eggs on it. The plant is toxic to eggs and larvae. West Virginia White will not recolonize an isolated site once extirpated because it doesn't fly across open areas. Logging activities may impact West Virginia White and Early Hairstreak.

Non-Habitat Threats:

Disease

Pollution

Loss of Prey Base

Loss of Relationship with Other Species

Description of non-habitat threat(s): Disease and invasives threaten host plants that are required for egg laying and larval development. Build up of granulosis virus in soil causes premature death of larval West Virginia White. Forest pest control spraying may impact West Virginia White and Early Hairstreak. Beech bark disease poses a significant threat to Early Hairstreak.

--West Virginia White is a poor recolonizer of isolated sites that become extirpated; colonies may be easily lost due to disturbance of habitat.

--Early Hairstreak populations are scattered and highly localized in VT; genetic exchange and recolonization would be unlikely among known sites.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Define particular habitat requirements of species for which we still have inadequate information, utilizing current knowledge of researchers and field investigations. This has been completed for most SGCN hardwood forest butterflies.
Research	Basic Life History	Medium	It is possible that early hairstreak uses beaked hazelnut in Vermont, as well as American beech; but this is unknown.
Research	Threats and Their Significance	High	Assess potential and existing impacts of limiting factors to habitat, host plants, and individual butterflies. Such limiting factors as habitat loss and degradation, exotic invasive plants, disease, and host plant loss should be examined.
Monitoring	Population Change	High	Monitor known SGCN butterfly populations. Track population trends at distinct locations. This would follow field surveys and assessments to identify populations judged to be large and viable. Focusing on such large populations would offer greater probability of detecting population shifts.
Monitoring	Habitat Change	High	Monitor change in available habitat for each species' specific requirements. Loss, restoration, and other changes to local habitat sites recognized as important to these species should be tracked.

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Butterflies-Hardwood Forest Group**
Scientific Name: **Butterflies-Hardwood Forest Group**
Species Group: **Invert**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Workshops	High	Incorporate butterfly SGCN occurrence information into environmental review and technical assistance	Number of sites with butterfly SGCN that received conservation benefits to this group	FWD, USFWS, DEC, ANR, VTrans, NRCS	SWG
Easements	High	Acquisition/easement of high priority SGCN butterfly hardwood forest sites	Number of SGCN butterfly sites protected	FWD, FPR, TNC, VLT, other land trusts	VHCB, SWG, EQUIP, USFWS
Standards	High	Work with foresters to avoid significant impacts to SGCN butterfly populations and habitats during forest management activities	Number of SGCN butterfly locations indicated and protected in forest management plans (including mature beech stands)	FWD, FPR, USFS, private landowners	

Bibliography

Glassberg, J. 1999. Butterflies through binoculars. A field guide to the butterflies of Eastern North America. Oxford University Press, New York, NY. 242 pp.

McFarland, K., and S. Zahendra. 2010. Vermont Butterfly Survey 2002-2007; final report to the Natural Heritage Information Project of the Vermont Department of Fish and Wildlife. Vermont Center for Ecostudies.



Common Name: **Butterflies-Wetland Group**
Scientific Name: **Butterflies-Wetland Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend:

Extirpated in VT?

Regional SGCN? Yes

Assessment Narrative:

This group consists of six species, the habitat requirements of which vary within the general wetland category. Their caterpillars require specific food plants.

--Bog copper (*Lycaena epixanthe*): Found only in acidic bogs with cranberries, which are the host plants; few colonies; a weak flier (limited dispersal capability); six VBS survey blocks

--Jutta arctic (*Oeneis jutta*) Only 3 colonies known, all in Northeast Highlands. Restricted to acidic bogs in Northeast Highlands.

--Dion skipper (*Euphyes dion*): Strong flier; a good short-distance colonizer. Typically low densities in colonies. 12 VBS blocks.

--Black dash (*Euphyes conspicua*): First VT record in 2002; found in southern-most VT sedge wetlands, except one possible sighting in the Champlain Valley. Nine VBS blocks.

--Two-spotted skipper (*Euphyes bimacula*) Low numbers observed in widely scattered colonies; may disappear from a location for several years, then reappear. Only four colonies known; widely separated.

--Mulberry wing (*Poanes massasoit*): Known from limited area Taconic Mountains and Vermont Valley, with two possible observations in Champlain Valley and Southern Green Mountains. Eight VBS blocks.

Distribution

Biophysical regions recorded during 2002-2007 butterfly survey (VBS):

--Bog copper: Northern Highlands, Champlain Hills, Northern Vermont Piedmont, Vermont Valley

--Jutta arctic: Northern Highlands

--Dion skipper: Champlain Valley, Northern Green Mountains, Taconic Mountains, Vermont Valley, Southern Vermont Piedmont

--Two-spotted skipper: Northern Highlands, Champlain Valley, Northern Vermont Piedmont, Southern Green Mountains

--Black dash: Taconic Mountains, Southern Vermont Piedmont; possibly Champlain Valley

--Mulberry wing: Taconic Mountains, Vermont Valley; possibly Champlain Valley, Southern Green Mountains

--Broad-winged skipper: 11 survey blocks in Bennington, Grand Isle, Addison, and Rutland counties



Common Name: **Butterflies-Wetland Group**
Scientific Name: **Butterflies-Wetland Group**
Species Group: **Invert**

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☐ General Literature ☒

This group consists of several species, the habitat requirements of which vary within the general hardwood forest category. Caterpillars require specific food plants, which are referenced below.

--Bog Copper (*Lycaena epixanthe*): Found only in acidic bogs with cranberries (*Vaccinium* sp.), which are the host plants.

--Jutta Arctic (*Oeneis jutta*): Restricted to Black Spruce bogs in the Northeast Highlands; host plants are Dense Cottongrass (*Eriophorum spissum*), *Carex geyeri*, and *C. confine*. Adults nectar at bog flowers such as Labrador Tea (*Ledum groenlandicum*).

--Dion Skipper (*Euphys dion*): Calcareous sedge wetlands; host plants are narrow-leaved sedges such as Tussock Sedge (*Carex stricta*); adults nectar on Buttonbush (*Cephalanthus occidentalis*), jewelweed (*Impatiens* sp.), And Swamp Thistle (*Cirsium muticum*).

--Black Dash (*Euphys conspicua*): Sedge wetlands in southern-most VT, except also one possible sighting in the Champlain Valley. Host plants are narrow-leaved sedges, predominantly Tussock Sedge (*Carex stricta*), though others are possible. Adults rely on nectar from Buttonbush (*Cephalanthus occidentalis*), jewelweed (*Impatiens* sp.), and Swamp Thistle (*Cirsium pumilum*). Associated with the Mulberry Wing (*Poanes massasoit*).

--Two-spotted Skipper (*Euphys bimacula*): Prefers spruce bogs and sedge wetlands. Larval host plants are sedges, especially Hairy-fruited Sedge (*Carex trichocarpa*) and Tussock Sedge (*C. stricta*); adults nectar on Pickerelweed (*Pontederia cordata*), blue flag iris (*Iris* sp.), Common Milkweed (*Asclepias syriaca*), and spireas (*Spirea* sp.).

--Mulberry Wing (*Poanes massasoit*): Sedge wetlands in southwestern VT, often with Black Dash and Dion Skipper; sometimes found in bogs, fens, and wet meadows. Known host plant is Tussock Sedge (*Carex stricta*), but there are likely others. Adults nectar on Swamp Milkweed (*Asclepias incarnata*), Common Milkweed (*A. syriaca*), and possibly other wetland flowers..



Common Name: **Butterflies-Wetland Group**
Scientific Name: **Butterflies-Wetland Group**
Species Group: **Invert**

Habitat Types:

Open Peatlands
Marshes and Sedge Meadows
Shrub Swamps

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Climate Change
Habitat Fragmentation
Invasion by Exotic Species

Description of habitat threat(s): Wetlands are particularly vulnerable to invasive exotic plants. Invasives threaten to replace native flora, including larval host plants of wetland SGCN butterflies. Wetland impacts due to development can also impact these butterflies.

Non-Habitat Threats:

Trampling or Direct Impacts
Loss of Relationship with Other Species

Description of non-habitat threat(s): Loss or reduction of native wetland flora can include SGCN host plants.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Define particular habitat requirements of species for which we still have inadequate information, utilizing current knowledge of researchers and field investigations. This has been completed for most SGCN wetland butterflies.
Research	Threats and Their Significance	High	Assess potential and existing impacts of limiting factors to habitat, host plants, and individual butterflies. Such limiting factors as habitat loss and degradation, exotic invasive plants, disease, pesticides, and host plant loss should be examined.
Monitoring	Population Change	High	Monitor known SGCN butterfly populations. Track population trends at distinct locations. This would follow field surveys and assessments to identify populations judged to be large and viable. Focusing on such large populations would offer greater probability of detecting population shifts.
Monitoring	Habitat Change	High	Monitor change in available habitat for each species' specific requirements. Loss, restoration, and other changes to local habitat sites recognized as important to these species should be tracked.
Monitoring	Monitor Threats	Medium	Monitor the spread (and control) of wetland invasive species that can impact the habitat and host plants of these butterfly species.

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Butterflies-Wetland Group**
Scientific Name: **Butterflies-Wetland Group**
Species Group: **Invert**

Species Strategies

<i>Strategy Type</i>	<i>Strategy Priority</i>	<i>Strategy Description</i>	<i>Performance Measure</i>	<i>Potential Partners</i>	<i>Potential Funding Sources</i>
Technical Assistance, Training, Workshops	High	Incorporate butterfly SGCN occurrence information into environmental review and technical assistance	Number of sites with butterfly SGCN that received conservation benefits to this group	FWD, USFWS, DEC, ANR, VTrans, NRCS	SWG
Easements	High	Acquisition/easement of high priority SGCN butterfly wetland sites	Number of SGCN butterfly sites protected	FWD, TNC, VLT, other land trusts	VHCB, SWG, EQUIP, USFWS

Bibliography

McFarland, K., and S. Zahendra. 2010. Vermont Butterfly Survey 2002-2007; final report to the Natural Heritage Information Project of the Vermont Department of Fish and Wildlife. Vermont Center for Ecostudies.

Glassberg, J. 1999. Butterflies through binoculars: the East. New York: Oxford University Press. 242 pp.



Common Name: **Moths Group**
Scientific Name: **Moths Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend:

Extirpated in VT?

Regional SGCN? Yes

Assessment Narrative:

Numerous species of moths are considered rare, declining, or extirpated. Very little information is available on the distribution of these species, and there is even less known about their trends throughout the region. This group includes the following 17 species:

- Eacles imperialis pini, Imperial Moth: Thought to be extirpated until collected in 2001. Declined drastically or disappeared from VT in 1950's. Probable cause of decline was pesticide spraying and release of Compsilura concinnata for gypsy moth control.
- Eana georgiella, A tortricid moth: One recent collection from high elevation site. Status unknown.
- Hemileuca lucina, New England Buckmoth: Restricted to New England; found in SE VT. Have increased in abundance globally.
- Lasionycta taigata, A noctuid moth: One collection in 1975. A bog associate. Status uncertain.
- Lemmeria digitalis, A noctuid moth: Several collected in 1991-93. Status uncertain.
- Lithophane franclemonti, Franclemont's Lithophane: One VT collection. Globally rare, with evidence of decline.
- Pachypolia atricornis, An autumnal noctuid moth: Reported from Chittenden Co. (Proctor Maple Research Forest). Very rarely collected throughout range.
- Papaipema sp. 2 nr. pterisii, Ostrich Fern Borer moth: Rare outside of VT. Responsibility species. Metapopulation structure is needed for long-term viability. Globally rare (G3G4).
- Properigea costa, A noctuid moth: Collected in Chittenden Co. Associated with shale, granite, or limestone barrens. More common globally than previously believed.
- Speranza ribearia, Currant Spanworm: Collected Bakersfield, 1991 (Franklin Co). Formerly widespread, now rare through much of range due to eradication of currant in 1920's-'60's (alternate host of white pine blister rust).
- Sphinx drupiferarum, Plum Sphinx or Wild Cherry Sphinx: Severe populations declining in Eastern NA; now uncommon to rare throughout range. Reasons for decline unknown, but may include introduction of parasitoid Compsilura concinnata and historic aerial (DDT) spraying.
- Sphinx luscitiosa, Clemens' Sphinx: Populations declining rangewide. Uncommon or rare throughout Northeast.
- Sthenopis thule, Willow Ghost moth: Single specimen, South Hero in 1992. Only reported from VT and ON.
- Xestia fabulosa, A noctuid moth: Range not well-known. Recorded from VT, NH, and ON. Status uncertain.
- Xestia homogena, A noctuid moth: Found in alpine habitat (high elevation; mountain peaks). Habitat specialist.
- Zale submediana, Gray Spring Zale: Associated with rare, highly impacted habitat. Very rare in VT; single report; probably was previously common in sand plains.
- Zanclognatha martha, Pine Barrens Zanclognatha: Associated with rare habitat. Somewhat rare outside of NJ; single VT report.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Moths Group**
Scientific Name: **Moths Group**
Species Group: **Invert**

Distribution

- Eacles imperialis pini, Imperial Moth: One collected Grand Isle, 2001; Chittenden Co. historically. Also known from northern NY and southern Canada.
- Eana georgiella A tortricid moth: One recent collection from Jay Peak.
- Hemileuca lucina New England Buckmoth: Restricted to New England; found in SE VT.
- Lasionycta taigata A noctuid moth: Reported from Essex Co. '75 (Moose Pond).
- Lemmeria digitalis A noctuid moth: Several collected in Chittenden Co., 1991-93.
- Lithophane franclemonti Franclemont's Lithophane: Known from one Grand Isle specimen.
- Pachypolia atricornis An autumnal noctuid moth: Reported from Chittenden Co. (Proctor Maple Research Forest).
- Papaipema sp. 2 nr. pterisii Ostrich Fern Borer Moth: Several collections along rivers in VT, including Huntington River. Rare outside of VT.
- Properigea costa A noctuid moth: Collected in Chittenden Co.
- Speranza ribearia Currant Spanworm: Collected Bakersfield, 1991 (Franklin Co).
- Sphinx drupiferarum Plum Sphinx or Wild Cherry Sphinx: Reported from Franklin, Chittenden, Bennington, Orleans, and Lamoille counties.
- Sphinx luscitiosa Clemens' Sphinx: VT distribution unknown.
- Sthenopsis thule A ghost moth: Single specimen, South Hero in 1992.
- Xestia fabulosa A noctuid moth: Range not well-known. Recorded from VT, NH, and ON.
- Xestia homogena A noctuid moth: Found in alpine habitat (high elevation; mountain peaks).
- Zale submediana Gray Spring Zale: Reported from Jericho Research Forest (Chittenden Co.) in '90. Probably was previously common in sand plains.
- Zanclognatha martha Pine Barrens Zanclognatha: Reported from Jericho Research Forest (Chittenden Co.) in '90.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Probable
Champlain Hills	Confident	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Probable
Northern VT Piedmont	Probable	Taconic Mtns	Probable
Northeastern Highlands	Confident		



Common Name: **Moths Group**
Scientific Name: **Moths Group**
Species Group: **Invert**

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

- Eacles imperialis pini, Imperial Moth: White pine specialist; possibly also pitch pine. Found in coniferous forest.
- Eana georgiella, A tortricid moth: High elevation species. VT specimen collected from summit in a "disturbed area."
- Hemileuca lucina, New England Buckmoth: Found in wet meadows and open fields. May use powerline corridors. Host plants: oak, Prunus serotina, willows, gray birch, Vaccinium; early instars on meadowsweet.
- Lasionycta taigata, A noctuid moth: Bog associate.
- Lemmeria digitalis, A noctuid moth: Reported from wetland habitats in PA. Host plants unknown.
- Lithophane franclemonti, Franclemont's Lithophane: Host plants and habitat unknown.
- Pachypolia atricornis, An autumnal noctuid moth: Found in mesic northern hardwood or mixed forests with non-acidic soils.
- Papaipema sp. 2 nr. pterisii, Ostrich Fern Borer Moth: May occur where there are large areas of ostrich fern, the host plant; particularly along floodplain forests.
- Properigea costa, A noctuid moth: Associated with shale, granite, or limestone barrens.
- Speranza ribearia, Currant Spanworm: Host plants are currant, gooseberry.
- Sphinx drupiferarum, Plum Sphinx or Wild Cherry Sphinx: Found in wooded habitats and suburbs. Host plants: cherry, plum, apple; also lilac and hackberry.
- Sphinx luscitiosa, Clemens' Sphinx: Found in clearings, edges, and meadows in wooded areas. Host plants: willow, poplar, birch, apple, ash. Has been reportedly seen obtaining nourishment from dead fish.
- Sthenopsis thule, Willow Ghost Moth: Host plant: reported on willows, but not well known.
- Xestia fabulosa, A noctuid moth: Reported elsewhere from boreal and montane spruce-fir forests, and high-elevation subalpine forests. Host plants: Vaccinium sp.
- Xestia homogena, A noctuid moth: Found in alpine habitat (high elevation; mountain peaks).
- Zale submediana, Gray Spring Zale: probably was previously common in sand plains. Host plants: jack, pitch, red, and probably other hard pines.
- Zanclognatha martha, Pine Barrens Zanclognatha: Found in pitch pine/ scrub oak barrens in PA northward; Jericho population may have colonized pines from a remnant pine barren.

Habitat Types:

Outcrops and Alpine
Spruce Fir Northern Hardwood
Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops



Common Name: **Moths Group**
Scientific Name: **Moths Group**
Species Group: **Invert**

Other Cultural

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Climate Change
Inadequate Disturbance Regime
Habitat Fragmentation
Invasion by Exotic Species

Description of habitat threat(s): Hemileuca lucina, New England Buckmoth Habitat loss, habitat succession.

- Papaipema sp. 2 nr. pterisii Ostrich Fern Borer Moth Metapopulation structure is needed for long-term viability (habitat loss, habitat fragmentation, invasion by exotic species).
- Zale submediana Gray Spring Zale Habitat loss and conversion.
- Zanclognatha martha Pine Barrens Zanclognatha Habitat loss and conversion.
- Habitat threats for other species are not known.

Non-Habitat Threats:

Parasites
Pollution
Reproductive Traits
Loss of Prey Base
Trampling or Direct Impacts

Description of non-habitat threat(s): Eacles imperialis pini, Imperial Moth Declined drastically or disappeared from VT in 1950's. Probable cause of decline was pesticide spraying and release of Compsilura concinnata for gypsy moth control. Use of BTK is also potential problem if used wrong time of year (late June or later).

- Speranza ribearia Currant Spanworm Formerly widespread, now rare through much of range due to eradication of currant in 1920's-'60's (alternate host of white pine blister rust).
- Sphinx drupiferarum Plum Sphinx or Wild Cherry Sphinx Introduced parasitoid Compsilura concinnata was probably involved in the decline of this species, but was not the only factor.
- Xestia homogena A noctuid moth Trampling or disturbance of host vegetation in alpine areas is a potential threat.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Moths Group**
 Scientific Name: **Moths Group**
 Species Group: **Invert**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Define Vermont-specific habitat requirements of species for which this information is lacking; utilize field investigations and current knowledge of researchers.
Research	Basic Life History	Medium	Larval host plant requirements need to be studied and described or refined.
Research	Distribution and Abundance	High	1) Obtain baseline distributional and abundance data for all species in group by conducting surveys throughout the state. Efforts should be focused on particular habitats required by each and, where appropriate, on regions of expected occurrence within the state (e.g., a northern peripheral species might be expected in the northern tier counties). Abundance information should be collected at sites of known occurrence. 2) Conduct inventories to detect and gather information on new SGCN moth populations.
Research	Threats and Their Significance	High	Assess potential and existing impacts of limiting factors to habitat, host plants, and individual moths. Such limiting factors as habitat loss and degradation, exotic invasive plants, diseases, parasitoids, pest control, and host plant loss should be examined.
Research	Taxonomy	Medium	Taxonomic uncertainty of some species, such as <i>Eacles imperialis pini</i> , needs to be resolved.
Monitoring	Population Change	High	Monitor known SGCN moth populations. Many of these species are declining regionally; trends need to be monitored.
Monitoring	Habitat Change	Medium	Monitor change in available habitat for each species' specific requirements. Loss, restoration, and other changes to local habitat recognized as important to these species should be tracked.
Monitoring	Monitor Threats	Medium	Several species have been negatively impacted by gypsy moth control methods, including release of exotic parasitoids. Gypsy moth and other species-targeted control needs to be tracked and considered in managing for SGCN moths. Exotic parasitoid populations and distributions need to be assessed also.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Easements	High	Acquisition/easement/conservation/restoration of high priority SGCN moth sites	Number of SGCN moth sites protected/restored	FWD, FPR, USFWS, USFS, NRCS, VLT, other land trusts	VHCB, SWG, EQIP, USFWS
Standards	High	Work with landowners and those who manage forest pests to reduce/eliminate the use of pesticides and exotic species where they may negatively impact SGCN moth species	Area of land where methods detrimental to SGCN moths have been eliminated through management planning	FWD, FPR, USFWS, VT Entomological Society, towns, private landowners	SWG

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Moths Group**
Scientific Name: **Moths Group**
Species Group: **Invert**

Bibliography

- Boettner, G. H., J. S. Elington, and C. J. Boettner. 2000. Effects of a biological control introduction on three nontarget native species of saturniid moths. *Conservation Biology* 14: 1798-1806.
- Boone, J. H., J. R. Grehan, B. L. Parker. 2000. A Checklist of Diptera, Hymenoptera and Coleoptera from Mount Mansfield, Vermont. Agricultural Experiment Station, University of Vermont Miscellaneous Publication No. 118.
- Covell, C. V., Jr. 1984. Moths of Eastern North America. A Roger Tory Peterson Field Guide. Easton Press, Norwalk, CT. 406 pp.
- Danks, H. V. (ed.). 1979. Canada and its insect fauna. *Memoirs of the Entomological Society of Canada*, No. 108. 573 pp.
- Ferguson, D. C., C. E. Harp, P. A. Opler, R. S. Peigler, M. Pogue, J. A. Powell, and M. J. Smith. 1999. Moths of North America. Jamestown, ND: Northern Prairie Wildlife Research Center Home Page.
<http://www.npwrc.usgs.gov/resource/distr/lepid/moths/mothsusa.htm> (version 12DEC2003). (Accessed November 11, 2004)
- Forbes, W. T. M. 1923. Lepidoptera of New York and Neighboring States. Part I (Primitive Forms, Microlepidoptera, Pyralids, Bombyces). Cornell University Agricultural Experiment Station Memoir 68. Ithaca, NY. 729 pp.
- Forbes, W. T. M. 1948. Lepidoptera of New York and Neighboring States. Part II (Geometrida, Lymantriidae, Notodontidae, Sphingidae). Cornell University Agricultural Experiment Station Memoir 274. Ithaca, NY. 263 pp.
- Forbes, W. T. M. 1954. Lepidoptera of New York and Neighboring States. Part III (Noctuidae). Cornell University Agricultural Experiment Station Memoir 329. Ithaca, NY. 433 pp.
- Grehan, J. R., B. L. Parker, G. R. Nielsen, D. H. Miller, J. D. Hedbor, M. Sabourin, and M. S. Griggs. 1995. Moths and butterflies of Vermont (Lepidoptera); a faunal checklist. Agricultural Experiment Station, University of Vermont; Department of Forests, Parks and Recreation, State of Vermont. Miscellaneous Publication 116. Vermont Monitoring Cooperative Bulletin No. 1.
- Grehan, J. R., M. Sabourin. 1995. Informal State Ranks for Vermont Moths and Butterflies (Lepidoptera). For the Nongame and Natural Heritage Program. Vermont Fish and Wildlife Department.
- Griggs, M. S., J. R. Grehan. 2000. Biodiversity of the Moth Fauna from a Light Trap Survey at Ethan Allen Firing Range. For the Vermont Army National Guard.
- Griggs, M. S., J. R. Grehan. 2001. Light Trap Survey of Moth Biodiversity of Mount Mansfield for 2000. For the Vermont Monitoring Cooperative.
- Handfield, L. 1997. Liste des Lepidopteres du Quebec et du Labrador. *Fabriques*, Supplement 7: 1-155.
- Holland, W. J. 1968. The Moth Book. Dover Publications, Inc. New York, NY. 479 pp.
- Lafontaine, J.D. 1998. Noctuoidea, Noctuidae (part), Noctuinae (part-Noctuini). The moths of America north of Mexico, including Greenland. Fascicle 27.3. Washington, DC: The Wedge Entomological Research Foundation; 348 pp.
- Maier, C. T., C. R. Lemmon, J. M. Fengler, D. F. Schweitzer, and R. C. Reardon. 2004. Caterpillars on the foliage of conifers in the northeastern United States. Forest Health Technology Enterprise Team, Morgantown, WV. USDA Forest Service and Connecticut Agricultural Experimental Station. FHTET-2004-01.
- Munroe, E. G. 1972. Pyraloidea: Pyralidae (in part). Pp. 1-304, Fasc. 13.1, A-C in Dominick, R. B., ed. The moths of America north of Mexico. E. W. Classey Ltd, London, UK.
- NatureServe. 2004. NatureServe Explorer: an online encyclopedia of life [web application]. Version 4.0. NatureServe, Arlington, VA. Available <http://www.natureserve.org/explorer>.
- Neunzig, H. H. 1986. Phycitinae; Acrobasis and allies. Fasc. 15.2, in Dominick, R. B., et al. Moths of American North of Mexico. Wedge Entomological Research Foundation, Washington, D. C. 112 pp.
- Prentice, R. M. 1962. Forest Lepidoptera of Canada recorded by the Forest Insect Survey. Vol. 2. Nycteolidae, Notodontidae, Noctuidae, Liparidae. Forest Entomology and Pathology Branch, Canada Department of Forestry. Bulletin 128. pp. 77-281.
- Prentice, R. M. 1963. Forest Lepidoptera of Canada recorded by the Forest Insect Survey. Vol. 3. Lasiocampidae, Drepanidae, Thyrididae, Geometridae. Forest Entomology and Pathology Branch, Canada Department of Forestry. Publ. 1013. pp. 282-542.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Moths Group**
Scientific Name: **Moths Group**
Species Group: **Invert**

Rockburne, E. W., and J. D. Lafontaine. 1976. The Cutworm Moths of Ontario and Quebec. Canada Department of Agriculture, Research Branch Publication 1593, 164 pp.

Schweitzer, Dale W. 2004 personal communications. NatureServe.

Thomas, A. W. 1996. Light-trap catches of moths within and above the canopy of a northeastern Forest. Journal of the Lepidopterists' Society 50: 21-45.

Thomas, A. W. 2001. Moth diversity in a northeastern North America red spruce forest. Natural Resources Canada, Information Report M-X-210E. 42 pp.

Wagner, D. L., D. C. Ferguson, T. L. McCabe, and R. C. Reardon. 2001. Geometroid caterpillars of northeastern and Appalachian forests. Forest Health Technology Enterprise Team, Morgantown, WV. USDA Forest Service. FHTET-2001-10.

Wagner, D. L., V. Giles, R. C. Reardon, and M. L. McManus. 1997. Caterpillars of eastern forests. Forest Health Technology Enterprise Team, Morgantown, WV. USDA Forest Service. FHTET-96-34.

Wagner, D. L., Nelson, M. W., Schweitzer, D.F. 2003. Shrubland lepidoptera of Southern New England and southeastern New York: ecology, conservation, and management. For. Ecol. Manage. 185: 95-112.

Webster, R. P., A. W. Thomas. 1999. A new species of Lithophane (Lepidoptera: Noctuidae: Cuculliinae) from Northeastern North America. Journal of the Lepidopterists' Society 53(2): 55-59.

Additional information on Vermont-specific information and collection records gathered through personal communications with Dale Schweitzer (NatureServe) and Jim Hedbor.



Common Name: **Mayflies/Stoneflies/Caddisflies Group**
Scientific Name: **Mayflies/Stoneflies/Caddisflies Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

This group consists eleven species with all but the *Rhyacophila brunnea* considered globally rare (G1-G3). Very little information is available on Vermont populations, making status determination difficult.

--A Mayfly (*Ameletus browni*): This is a globally rare species (G3/G4). Insufficient information is available to determine VT status.

--Tomah Mayfly (*Siphonisca aerodromia*): This is a globally rare species (G2G3). Although not yet recorded from VT, insufficient information is available to determine VT status.

--A Mayfly (*Siphonurus demaryi*): This is a globally rare species (G2G3). Insufficient information is available to determine VT status.

--Roaring Brook Mayfly (*Epeorus frisoni*): This is a globally rare species (G1). Known only recently from a single location in VT. To date, only four small stream populations are known worldwide.

--A Mayfly (*Eurylophella bicoloroides*): This is a globally rare species (G3). Insufficient information is available to determine VT status.

--A Mayfly (*Baetisca rubescens*): This is a globally rare species (G3/G4). Insufficient information is available to determine VT status.

--A Stonefly (*Alloperla voinae*): This is a globally rare species (G3) which is rarely collected. Insufficient information is available to determine VT status.

--Appalachian Stone (*Hansonoperla appalachia*): This is a globally rare species (G3). Insufficient information is available to determine VT status.

--Spiny Salmonfly (*Pteronarcys comstocki*): This is a globally rare species (G3). Insufficient information is available to determine VT status.

--A Caddisfly (*Rhyacophila brunnea*): This species is known from fewer than 10 sites in Vermont.

--A Caddisfly (*Rhyacophila amicus*): This is a globally rare species (G2). Insufficient information is available to determine VT status.

Distribution

--Roaring Brook mayfly (*Epeorus frisoni*): Known in Vermont only from a Battenkill tributary near Dorset. Globally, known from only from NH, VT, and ME.

--*Ameletus browni*: This species is not well documented, but has been reported from Bennington County.

--*Eurylophella bicoloroides*: This species is not well documented, but has been reported from Bennington County.

--Tomah mayfly (*Siphonisca aerodromia*): This mayfly has not yet been documented in VT. It is known only from a few sites in NY, ME, QE, and Labrador. Noted as one of the rarest mayflies in the world.

--*Baetisca rubescens*: Distribution of this species in VT is not known. Only a few U.S and Canadian records.

--*Alloperla voinae*: Distribution of this species in VT is not known. Otherwise known from NY to NS, south to MA.

--*Rhyacophila brunnea*: Known from Killington, Shrewsbury, Mendon, Orange, and Stowe.

--*Siphonurus demaryi*: This species is not well documented, but has been reported from Bennington County. A regional endemic to northeastern U.S and southeastern Canada.

--*Hansonoperla appalachia*: This species is not well documented in Vermont.

--*Pteronarcys comstocki*: This species is not well documented in Vermont.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Mayflies/Stoneflies/Caddisflies Group**
Scientific Name: **Mayflies/Stoneflies/Caddisflies Group**
Species Group: **Invert**

--Rhyacophila amicus: This species is not well documented in Vermont.

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Probable
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Known Watersheds

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

--Tomah mayfly (*Siphonisca aerodromia*) This mayfly inhabits rivers with broad, seasonally flooded sedge-dominated floodplains.

--*Siphonurus demaryi* Larvae have been collected in lake littoral zones near stream inlets and along margins of larger streams among vegetation.

--Roaring Brook mayfly (*Epeorus frisoni*) Found in small, first and second order tributaries of high elevation streams. May be associated with conditions of pristine water quality and minimally or undisturbed riparian habitat.

--*Ameletus browni* Restricted to cold, high elevation, first order streams in undisturbed habitat.

--*Eurlophella bicoloroides* Found in small (2nd order) streams up to medium sized rivers (6th or 7th order). In large streams, has been reported in reaches below reservoirs with hypolimnetic release.

--*Baetisca rubescens* A northern species, restricted to cold, high elevation streams.

--*Alloperla voinae* Habitat is unknown for this species.

--*Rhyacophila brunnea* This species inhabits small, high-elevation streams; these are acid-sensitive streams.

--*Rhyacophila amicus* A coldwater montane stream caddisfly.

--*Hansonoperla appalachia* Occurs in pristine medium-sized streams of the elevated Appalachians. Nymphs were collected in WV from undercut banks of riffle areas where roots of riparian vegetation trapped coarse detritus and caused deposits of sand to accumulate.

--*Pteronarcys comstocki* Prefers medium-sized to large cold streams. An Appalachian species.

Habitat Types:

Marshes and Sedge Meadows

Aquatic: Fluvial

Current Threats

Habitat Threats:

Conversion of Habitat

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Mayflies/Stoneflies/Caddisflies Group**
 Scientific Name: **Mayflies/Stoneflies/Caddisflies Group**
 Species Group: **Invert**

Habitat Alteration

Sedimentation

Habitat Fragmentation

Description of habitat threat(s):

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Several of these species are known globally from very few sites, which are often widely separated. Recolonization may not be possible if individual populations are lost.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Need to determine details of habitat requirements in order to refine distributional searches.
Research	Distribution and Abundance	High	1) Conduct inventories to detect and gather information on SGCN stoneflies, mayflies, and caddisflies. 2) Need statewide surveys to provide basic understanding of distribution for all species.
Research	Threats and Their Significance	Medium	Assess threat of high elevation and headwater stream acidification to mayflies, caddisflies, and stoneflies that occupy these waters.
Monitoring	Population Change	Medium	1) Monitor known SGCN stonefly/mayfly/caddisfly populations. 2) Populations should be monitored for presence/absence now; monitor for population changes after baseline abundance data is available.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Protect and restore habitats on which SGCN stoneflies, mayflies, and caddisflies are dependent through pollution abatement, riparian buffers, flow regulation, easements, etc.	Number of acres of riparian habitat protected and/or restored	FWD, USFS, Trout Unlimited, Watershed groups, Landowners, NRCS	SWG, EQIP, LCLT, VLT

Bibliography

Vermont Department of Environmental Conservation, Biomonitoring Database.
 Maine 2015 Wildlife Action Plan Revision.

Natureserve. 2015. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.0. NatureServe, Arlington, VA. U.S.A. Available <http://explorer.natureserve.org>. (Accessed: 01/21/2015)

Funk, D. H. and B. W. Sweeney. 1994. The Larvae of Eastern North American Eurylophella Tiensuu (Ephemeroptera: Ephemerellidae). Transactions of the American Entomological Society 120(3): 209-286.

Kirchner, R. F., and B. C. Kondratieff. 1985. The Nymph Of Hansonoperla appalachia Nelson (Plecoptera, Perlidae). Proceedings of the Entomological Society of Washington 87: 593-596.



Common Name: Odonates-Bog/Fen/Swamp/Marshy Pond Group
Scientific Name: Odonates-Bog/Fen/Swamp/Marshy Pond Group
Species Group: Invert

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

Individual species in this group are rare in Vermont; several are regionally or globally rare.

This group consists of 15 species, the habitat requirements of which vary within the general wetland category. Habitats given for each species refer primarily to reproduction and nymphal requirements; adults of many species require various nearby terrestrial habitats as well.

- Subarctic Bluet (*Coenagrion interrogatum*): Bogs, boggy-edged ponds
- Comet Darner (*Anax longipes*): On northern edge of range in Vermont; one larval record. Vernal pools and semi-permanent ponds (no fish)
- Mottled Darner (*Aeshna clepsydra*): Boggy/marshy edges of lakes
- Zigzag Darner (*Aeshna sitchensis*): Boggy ponds, small bog pools, fen puddles
- Subarctic Darner (*Aeshna subarctica*): Bogs with saturated sphagnum
- Spatterdock Darner (*Rhionaeschna mutata*): Southern species, could advance north with climate change; vegetated ponds
- Swamp Darner (*Epiaeschna heros*): Hardwood swamps
- Cyrano Darner (*Nasiaeschna pentacantha*): Vegetated ponds
- Petite Emerald (*Dorocordulia lepida*): Boggy ponds and lakes, marshes, cedar swamp streams
- Ski-tailed Emerald (*Somatochlora elongata*): Marshy ponds, peatlands
- Forcipate Emerald (*Somatochlora forcipata*): Boggy rills, bogs, and small forested streams
- Delicate Emerald (*Somatochlora franklini*): Bogs, boggy-edged ponds
- Kennedy's Emerald (*Somatochlora kennedyi*): Boggy streams; bogs, fens, and swamps often with flowing water.
- Ebony Boghaunter (*Williamsonia fletcheri*): Bogs, fens
- Black Meadowhawk (*Sympetrum danae*): Marshy ponds, bogs, fens

Distribution

Distributions by biophysical region for bog/fen/swamp/marshy pond odonates are as follows:

- Subarctic Bluet (*Coenagrion interrogatum*): Northern Highlands, Southern Green Mountains
- Comet Darner (*Anax longipes*): Southern Vermont Piedmont
- Mottled Darner (*Aeshna clepsydra*): Champlain Valley, Taconic Mountains, Southern Green Mountains, Southern Vermont Piedmont
- Zigzag Darner (*Aeshna sitchensis*): Northern Highlands
- Subarctic Darner (*Aeshna subarctica*): Northern Highlands, Northern Green Mountains
- Spatterdock Darner (*Rhionaeschna mutata*): Champlain Valley, Northern Green Mountains, Southern Vermont Piedmont
- Swamp Darner (*Epiaeschna heros*): Champlain Valley, Northern Green Mountains, Taconic Mountains, Southern Green Mountains
- Cyrano Darner (*Nasiaeschna pentacantha*): Southern Vermont Piedmont
- Petite Emerald (*Dorocordulia lepida*): Northern Highlands, Northern Vermont Piedmont, Southern Vermont Piedmont, Vermont Valley
- Ski-tailed Emerald (*Somatochlora elongata*): Northern Highlands, Northern Vermont Piedmont, Northern



Common Name: Odonates-Bog/Fen/Swamp/Marshy Pond Group
Scientific Name: Odonates-Bog/Fen/Swamp/Marshy Pond Group
Species Group: Invert

Green Mountains, Taconic Mountains, Southern Green Mountains, Southern Vermont Piedmont
 --Forcipate Emerald (*Somatochlora forcipata*): Northern Highlands, Northern Vermont Piedmont, Vermont Valley, Southern Green Mountains
 --Delicate Emerald (*Somatochlora franklini*): Northern Highlands, Northern Vermont Piedmont
 --Kennedy's Emerald (*Somatochlora kennedyi*): Northern Highlands
 --Ebony Boghaunter (*Williamsonia fletcheri*): Northern Highlands, Northern Vermont Piedmont, Vermont Valley
 --Black Meadowhawk (*Sympetrum danae*): Northern Highlands, Northern Vermont Piedmont, Champlain Valley

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

This group consists of 15 species, the habitat requirements of which vary within the general wetland category. Habitats given for each species refer primarily to reproduction and nymphal requirements; adults of many species require various nearby terrestrial habitats as well. Some dragonflies are known to exhibit "hilltopping" behavior, in which they congregate on tops of hills, which may be fairly distant from egg-laying sites. This would increase the home range requirements for such species.

- Subarctic Bluet (*Coenagrion interrogatum*): Bogs, boggy-edged ponds
- Comet Darner (*Anax longipes*): Vernal pools and semi-permanent ponds (no fish)
- Mottled Darner (*Aeshna clepsydra*): Boggy/marshy edges of lakes
- Zigzag Darner (*Aeshna sitchensis*): Boggy ponds, small bog pools, fen puddles
- Subarctic Darner (*Aeshna subarctica*): Bogs with saturated sphagnum
- Spatterdock Darner (*Rhionaeschna mutata*): Vegetated ponds
- Swamp Darner (*Epiaeschna heros*): Hardwood swamps
- Cyrano Darner (*Nasiaeschna pentacantha*): Vegetated ponds
- Petite Emerald (*Dorocordulia lepida*): Boggy ponds and lakes, marshes, cedar swamp streams
- Ski-tailed Emerald (*Somatochlora elongata*): Marshy ponds, peatlands
- Forcipate Emerald (*Somatochlora forcipata*): Boggy rills, bogs, and small forested streams
- Delicate Emerald (*Somatochlora franklini*): Bogs, boggy-edged ponds
- Kennedy's Emerald (*Somatochlora kennedyi*): Boggy streams; bogs, fens, and swamps often with flowing water.
- Ebony Boghaunter (*Williamsonia fletcheri*): Bogs, fens
- Black Meadowhawk (*Sympetrum danae*): Marshy ponds, bogs, fens



Common Name: **Odonates-Bog/Fen/Swamp/Marshy Pond Group**
Scientific Name: **Odonates-Bog/Fen/Swamp/Marshy Pond Group**
Species Group: **Invert**

Habitat Types:

Hardwood Swamps
Softwood Swamps
Seeps and Pools
Open Peatlands
Marshes and Sedge Meadows
Shrub Swamps
Aquatic: Fluvial
Aquatic: Lacustrine

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Sedimentation
Habitat Fragmentation
Invasion by Exotic Species
Climate Change

Description of habitat threat(s): Many of these wetlands are small, fragile, and could be easily degraded by disturbances within their watershed or groundwater source (e.g., development, clearcutting, hydrologic alterations). There is little specific information available citing negative impacts on these odonates. Several or all of these species utilize a combination of wetland and upland habitat to complete their life cycle; fragmentation of this complex could potentially have a negative effect on these odonates. Climate change may affect bog communities and the odonates that reside in them. Exotic plants such as Phragmites and purple loosestrife can dominate the floral community and make odonate habitat less suitable.

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Non-point source nutrients can alter water chemistry, nutrient availability, and benthic habitat.



Common Name: **Odonates-Bog/Fen/Swamp/Marshy Pond Group**
Scientific Name: **Odonates-Bog/Fen/Swamp/Marshy Pond Group**
Species Group: **Invert**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	Define particular habitat requirements of each species within Vermont, utilizing current knowledge of researchers and field investigations.
Research	Distribution and Abundance	High	1) Obtain baseline distributional and abundance data for all species in group by conducting surveys throughout the state. Efforts should be focused on particular habitats required by each and, where appropriate, on regions of expected occurrence within the state (e.g., a northern peripheral species might be expected in the northern tier counties). Abundance information should be collected at sites of known occurrence. 2) Conduct inventories to detect and gather information on new SGCN odonate populations.
Research	Threats and Their Significance	Medium	Assess the vulnerability of nymphs of each species to wetland perturbations, such as siltation, temperature and water quality shifts, chemical pollution, and changes in vegetation. Investigate the upland habitat needs of the adults and the effects of such impacts as fragmentation and reduction.
Monitoring	Population Change	High	Population monitoring could be employed to track population trends at distinct locations. This would follow field surveys and assessments to identify populations judged to be large and viable. Focusing on such large populations would offer greater probability of detecting population shifts.

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Odonates-Bog/Fen/Swamp/Marshy Pond Group**
Scientific Name: **Odonates-Bog/Fen/Swamp/Marshy Pond Group**
Species Group: **Invert**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications	Medium	Educate public and improve appreciation of vulnerable odonate habitats	Number of participants at events/workshops	FWD, VCE, town Cons Comms, media outlets	SWG
Technical Assistance, Training, Learning Networks	High	Incorporate odonate SGCN occurrence information into environmental review and technical assistance	Number of sites with odonate SGCN that received conservation benefits to this group.	FWD, USFWS, DEC, ANR, VTrans, NRCS	SWG
Compatible Resource Use	Medium	Manage protected areas (wetlands) for odonate conservation	Number of protected sites with odonate protective strategies in place.	FWD, FPR, USFS, USFWS, DEC, NRCS, VLT, other land trusts, town conservation and planning commissions.	SWG, FPR, USFS
Easements	High	Acquisition/easement of high priority SGCN odonate wetland sites	Number of SGCN odonate sites protected	FWD, FPR, TNC, VLT, other land trusts	VHCB, SWG, EQIP, USFWS



Common Name: **Odonates-Bog/Fen/Swamp/Marshy Pond Group**
Scientific Name: **Odonates-Bog/Fen/Swamp/Marshy Pond Group**
Species Group: **Invert**

Bibliography

- Brunelle, P.M. 1999. Distribution of damselflies and dragonflies (Odonata) of Maine, United States. *Northeastern Naturalist* 6(1):35-38.
- Carle, F. L. 1994. Dragonflies and damselflies (Odonata) known to or likely to occur in Vermont. A report to the Nongame and Natural Heritage Program, Vermont Fish & Wildlife Department. 22 pp.
- Carle, F. L. 1997. Odonates of the Green Mountain National Forest A report to the National Forest Service, Manchester Ranger District.
- Carpenter, V.A. 1997. Dragonflies and damselflies of Cape Cod. Edition 2. The Cape Cod Museum of Natural History, Brewster, Massachusetts. 80 pp.
- Dunkle, S.W. 2000. Dragonflies through binoculars: a field guide to dragonflies of North America. Oxford University Press, New York, NY. 266pp.
- Dunkle, S.W. 1990. Damselflies of Florida, Bermuda and the Bahamas. Scientific Publishers Nature Guide # 3.
- Dunkle, S.W. 1989. Dragonflies of the Florida Peninsula, Bermuda and the Bahamas. Scientific Publishers Nature Guide # 1.
- Lam, E. 2004. Damselflies of the Northeast. Biodiversity Books, Forest Hills, N.Y. 96 pp.
- NatureServe. 2004. NatureServe Explorer: an online encyclopedia of life [web application]. Version 4.0. NatureServe, Arlington, VA. Available <http://www.natureserve.org/explorer>.
- Needham, J. G., M. J., Jr., Westfall, and M. L. May. 2000. Dragonflies of North America. Scientific Publishers, Gainesville FL. 940 pp.
- Nikula, B., J.L. Loose, and M.R. Burne. 2003. A field guide to dragonflies and damselflies of Massachusetts. Natural Heritage and Endangered Species Program. Massachusetts Division of Fisheries and Wildlife. Boston, MA.
- Nothnagle, P. 1988. A provisional list of the odonates (dragonflies and damselflies) of Vermont. Technical Report 1. Nongame and Natural Heritage Program, Vermont Fish and Wildlife Department.
- Pfeiffer, B. M. 2002. *Williamsonia fletcheri* Williamson (Anisoptera: Corduliidae) at Chickering Bog Natural Area, Calais, Vermont: an encounter on June 4, 2002. A report to the Vermont Chapter of The Nature Conservancy. 4 pp.
- Westfall, M.J. and M.L. May. 1996. Damselflies of North America. Scientific Publishers, Gainesville, Florida.



Common Name: **Odonates-Lakes/Ponds Group**
Scientific Name: **Odonates-Lakes/Ponds Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

This group consists of seven species, the habitat requirements of which vary within the general lakes/ponds category. Habitats given for each species refer primarily to reproduction and nymphal requirements; adults of many species depend on various nearby terrestrial habitats as well. Some dragonflies are known to exhibit "hilltopping" behavior, in which they congregate on tops of hills, which may be fairly distant from egg-laying sites. This would increase the home range requirements for such species.

--New England Bluet (*Enallagma laterale*): Vegetated ponds.

--Slender Bluet (*Enallagma traviatum*): Lakes and ponds with vegetation

--Lilypad Forktail (*Ischnura kellicotti*): Ponds with lilypads

--Ringed Emerald (*Somatochlora albicincta*): Cold ponds

--Lake Emerald (*Somatochlora cingulata*): Shallow cold lakes, sluggish rivers and streams

--Banded Pennant (*Celithemis fasciata*): Newly discovered in Vermont. A southern species that may continue to move northward. Vegetated ponds and lakes

--Carolina Saddlebags (*Tamea carolina*): Newly discovered in Vermont. A southern species that may continue to move northward. Vegetated ponds and lakes.

Distribution

Distribution records by biophysical region for lakes/ponds odonates:

--New England Bluet (*Enallagma laterale*): Southern Vermont Piedmont

--Slender Bluet (*Enallagma traviatum*): Champlain Valley, Taconic Mountains, Southern Vermont Piedmont

--Lilypad Fork tail (*Ischnura kellicotti*): Taconic Mountains, Southern Vermont Piedmont

--Ringed Emerald (*Somatochlora albicincta*): Northern Highlands, Northern Green Mountains

--Lake Emerald (*Somatochlora cingulata*): Northern Highlands, Northern Vermont Piedmont, Southern Green Mountains

--Banded Pennant (*Celithemis fasciata*): Southern Green Mountains

--Carolina Saddlebags (*Tamea carolina*): Vermont Valley, Southern Vermont Piedmont



Common Name: **Odonates-Lakes/Ponds Group**
Scientific Name: **Odonates-Lakes/Ponds Group**
Species Group: **Invert**

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

This group consists of seven species, the habitat requirements of which vary within the general lakes/ponds category. Habitats given for each species refer primarily to reproduction and nymphal requirements; adults of many species require various nearby terrestrial habitats as well. Some dragonflies are known to exhibit "hilltopping" behavior, in which they congregate on tops of hills, which may be fairly distant from egg-laying sites. This would increase the home range requirements for such species.

- New England Bluet (*Enallagma laterale*) Vegetated ponds
- Slender Bluet (*Enallagma traviatum*) Lakes and ponds with vegetation
- Lilypad Forktail (*Ischnura kellicotti*) Ponds with lily pads
- Ringed Emerald (*Somatochlora albicincta*) Cold ponds and lakes
- Lake Emerald (*Somatochlora cingulata*) Shallow cold lakes, sluggish rivers and streams
- Banded Pennant (*Celithemis fasciata*) Vegetated lakes and ponds
- Carolina Saddlebags (*Tamea carolina*) Vegetated lakes and ponds

Habitat Types:

Open Peatlands
Marshes and Sedge Meadows
Wet Shores
Aquatic: Fluvial
Aquatic: Lacustrine
Aquatic: Man-Made Water Bodies



Common Name: **Odonates-Lakes/Ponds Group**
Scientific Name: **Odonates-Lakes/Ponds Group**
Species Group: **Invert**

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Sedimentation
Incompatible Recreation

Description of habitat threat(s): Shoreline development and fill can degrade and eliminate suitable aquatic habitat. Exotic invasive aquatic plants such as Eurasian milfoil may change the habitat available to lake and pond odonates.

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Wakes caused by boating can wash over and kill emerging adults.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	Define particular habitat requirements of each species within Vermont, utilizing current knowledge of researchers and field investigations.
Research	Distribution and Abundance	Medium	1) Conduct inventories to detect and gather information on new SGCN odonate populations. 2) Obtain baseline distributional and abundance data for all species in group by conducting surveys throughout the state. Efforts should be focused on particular habitats required by each and, where appropriate, on regions of expected occurrence within the state (e.g., a northern peripheral species might be expected in the northern tier counties). Abundance information should be collected at sites of known occurrence.
Research	Threats and Their Significance	Medium	Assess the vulnerability of nymphs of each species to habitat perturbations, such as substrate alteration, temperature and water quality shifts, chemical pollution, and changes in vegetation. Investigate the upland habitat needs of the adults and the effects of such impacts as fragmentation and reduction.
Monitoring	Population Change	High	Monitor known SGCN odonate populations. Track population trends at distinct locations. This would follow field surveys and assessments to identify populations judged to be large and viable. Focusing on such large populations would offer greater probability of detecting population shifts. Particularly important monitoring sites would include Lily Pond (Windham Co.) and those sites with isolated populations or scattered distributions.
Monitoring	Range Shifts	Medium	Monitor particular northern peripheral species (northern Vermont) to detect shifts in range over time.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Odonates-Lakes/Ponds Group**
 Scientific Name: **Odonates-Lakes/Ponds Group**
 Species Group: **Invert**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Workshops	High	Incorporate odonate SGCN occurrence information into environmental review and technical assistance	Number of sites with odonate SGCN that received conservation benefits to this group	FWD, USFWS, DEC, ANR, VTrans, NRCS	SWG
Compatible Resource Use	High	Develop stewardship program for sites with high priority odonate SGCN	Number of stewardship sites established	FWD, Federation of VT Lakes & Ponds, VCE, DEC, CLF, town Cons Comms	SWG, conservation license plate funds, corporate funding
Awareness Raising and Communications	High	Educate public and improve appreciation of vulnerable odonate habitats	Number of participants at events/workshops	FWD, VCE, Federation of VT Lakes & Ponds, town Cons Comms, media outlets	SWG
Awareness Raising and Communications	High	Raise awareness of Vermont Shoreland Protection Act with landowners	Number of lakeshore landowners contacted	DEC, FWD, ANR, Federation of VT Lakes and Ponds, media outlets	DEC, ANR
Compliance & Enforcement	High	Enforcement of the new (2014) Vermont Shoreland Protection Act	Number of permit requests annually	DEC, ANR	DEC, ANR
Easements	High	Acquisition/easement of high priority SGCN odonate lake and pond sites	Number of SGCN odonate sites protected	FWD, VLT, other land trusts	VHCB, SWG, EQUIP, USFWS



Common Name: **Odonates-Lakes/Ponds Group**
Scientific Name: **Odonates-Lakes/Ponds Group**
Species Group: **Invert**

Bibliography

- Brunelle, P.M. 1999. Distribution of damselflies and dragonflies (Odonata) of Maine, United States. *Northeastern Naturalist* 6(1):35-38.
- Carle, F. L. 1994. Dragonflies and damselflies (Odonata) known to or likely to occur in Vermont. A report to the Nongame and Natural Heritage Program, Vermont Fish & Wildlife Department. 22 pp.
- Carle, F. L. 1997. Odonates of the Green Mountain National Forest A report to the National Forest Service, Manchester Ranger District.
- Carpenter, V.A. 1997. Dragonflies and damselflies of Cape Cod. Edition 2. The Cape Cod Museum of Natural History, Brewster, Massachusetts. 80 pp.
- Dunkle, S.W. 2000. Dragonflies through binoculars: a field guide to dragonflies of North America. Oxford University Press, New York, NY. 266pp.
- Dunkle, S.W. 1990. Damselflies of Florida, Bermuda and the Bahamas. Scientific Publishers Nature Guide # 3.
- Dunkle, S.W. 1989. Dragonflies of the Florida Peninsula, Bermuda and the Bahamas. Scientific Publishers Nature Guide # 1.
- Lam, E. 2004. Damselflies of the Northeast. Biodiversity Books, Forest Hills, N.Y. 96 pp.
- NatureServe. 2004. NatureServe Explorer: an online encyclopedia of life [web application]. Version 4.0. NatureServe, Arlington, VA. Available <http://www.natureserve.org/explorer>.
- Needham, J. G., M. J., Jr., Westfall, and M. L. May. 2000. Dragonflies of North America. Scientific Publishers, Gainesville FL. 940 pp.
- Nikula, B., J.L. Loose, and M.R. Burne. 2003. A field guide to dragonflies and damselflies of Massachusetts. Natural Heritage and Endangered Species Program. Massachusetts Division of Fisheries and Wildlife. Boston, MA.
- Nothnagle, P. 1988. A provisional list of the odonates (dragonflies and damselflies) of Vermont. Technical Report 1. Nongame and Natural Heritage Program, Vermont Fish and Wildlife Department.
- Westfall, M.J. and M.L. May. 1996. Damselflies of North America. Scientific Publishers, Gainesville, Florida.



Common Name: **Odonates-River/Stream Group**
Scientific Name: **Odonates-River/Stream Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

This group consists of 17 species, the habitat requirements of which vary within the general rivers/streams category. Habitats given for each species refer primarily to reproduction and nymphal requirements; adults of many species depend on various nearby terrestrial habitats as well. Riparian vegetation is important to adults as habitat.

- American Rubyspot (*Hetaerina americana*): Streams and rivers with emergent vegetation
- Blue-fronted Dancer (*Argia apicalis*): Small to large rivers with sand or mud; occasionally ponds and lakes
- River Bluet (*Enallagma anna*): First Vermont record in 2014; rivers
- Rainbow Bluet (*Enallagma antennatum*): Slow streams and big rivers
- Big Bluet (*Enallagma durum*): Discovered in 2005 in southern Vermont; two known sites; slow rivers
- Spine-crowned Clubtail (*Gomphus abbreviatus*): Rivers
- Midland Clubtail (*Gomphus fraternus*): Large rivers
- Rapids Clubtail (*Gomphus quadricolor*): Large streams and rivers
- Skillet Clubtail (*Gomphus ventricosus*): Large rivers; mud or sand bottom
- Cobra Clubtail (*Gomphus vastus*): Large rivers with mud bottom; sometimes large streams and lakes
- Brook Snaketail (*Ophiogomphus aspersus*): Sandy streams
- Riffle Snaketail (*Ophiogomphus carolus*): Rapid, rocky or sandy streams and rivers
- Maine Snaketail (*Ophiogomphus mainensis*): Clear, rocky forested streams
- Rusty Snaketail (*Ophiogomphus rupinsulensis*): Large streams and rivers
- Riverine Clubtail (*Stylurus amnicola*): Big rivers
- Zebra Clubtail (*Stylurus scudderi*): Rivers
- Stygian Shadowdragon (*Neurocordulia yamaskanensis*): Large rivers and lakes; often rocky

Distribution

Biophysical region records for river/stream odonates:

- American Rubyspot (*Hetaerina americana*): Champlain Valley, Southern Vermont Piedmont
- Blue-fronted Dancer (*Argia apicalis*): Champlain Valley, Taconic Mountains, Southern Vermont Piedmont
- River Bluet (*Enallagma anna*): Southern Vermont Piedmont
- Rainbow Bluet (*Enallagma antennatum*): Champlain Valley, Northern Green Mountains, Taconic Mountains, Southern Vermont Piedmont.
- Big Bluet (*Enallagma durum*): Champlain Valley, Taconic Mountains
- Spine-crowned Clubtail (*Gomphus abbreviatus*): Champlain Valley, Taconic Mountains, Southern Vermont Piedmont
- Midland Clubtail (*Gomphus fraternus*): Champlain Valley
- Rapids Clubtail (*Gomphus quadricolor*): Champlain Valley, Champlain Hills, Taconic Mountains, Southern Green Mountains, Southern Vermont Piedmont.
- Skillet Clubtail (*Gomphus ventricosus*): Southern Vermont Piedmont
- Cobra Clubtail (*Gomphus vastus*): Southern Vermont Piedmont
- Brook Snaketail (*Ophiogomphus aspersus*): Northern Highlands, Northern Vermont Piedmont, Northern



Common Name: Odonates-River/Stream Group
Scientific Name: Odonates-River/Stream Group
Species Group: Invert

Green Mountains, Champlain Valley, Taconic Mountains, Southern Vermont Piedmont.
 --Riffle Snaketail (*Ophiogomphus carolus*): Northern Highlands, Northern Green Mountains, Champlain Valley, Taconic Mountains, Southern Green Mountains, Southern Vermont Piedmont.
 --Maine Snaketail (*Ophiogomphus mainensis*): Northern Highlands, Northern Vermont Piedmont, Northern Green Mountains, Southern Green Mountains, Southern Vermont Piedmont.
 --Rusty Snaketail (*Ophiogomphus rupinsulensis*): Northern Green Mountains, Taconic Mountains, Southern Vermont Piedmont
 --Riverine Clubtail (*Stylurus amnicola*): Southern Vermont Piedmont
 --Zebra Clubtail (*Stylurus scuderi*): Northern Highlands, Northern Vermont Piedmont, Northern Green Mountains, Champlain Valley, Taconic Mountains, Vermont Valley, Southern Vermont Piedmont.
 --Stygian Shadowdragon (*Neurocordulia yamaskanensis*): Northern Vermont Piedmont, Champlain Valley, Taconic Mountains, Southern Green Mountains, Southern Vermont Piedmont.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

This group consists of several species, the habitat requirements of which vary within the general rivers/streams category. Habitats given for each species refer primarily to reproduction and nymphal requirements; adults of many species depend on various nearby terrestrial habitats as well. Riparian vegetation is important to adults as habitat.

- American Rubyspot (*Hetaerina americana*): Streams and rivers with emergent vegetation
- Blue-fronted Dancer (*Argia apicalis*): Small to large rivers with sand or mud; occasionally ponds and lakes
- River Bluet (*Enallagma anna*): Rivers
- Rainbow Bluet (*Enallagma antennatum*): Slow streams and big rivers
- Big Bluet (*Enallagma durum*): Slow rivers
- Spine-crowned Clubtail (*Gomphus abbreviatus*): Rivers
- Midland Clubtail (*Gomphus fraternus*): Big rivers
- Rapids Clubtail (*Gomphus quadricolor*): Large streams and rivers



Common Name: Odonates-River/Stream Group
Scientific Name: Odonates-River/Stream Group
Species Group: Invert

- Skillet Clubtail (*Gomphus ventricosus*): Large rivers; mud or sand bottom
- Cobra Clubtail (*Gomphus vastus*): Large rivers with mud bottom; sometimes large streams and lakes
- Brook Snaketail (*Ophiogomphus aspersus*): Sandy streams
- Riffle Snaketail (*Ophiogomphus carolus*): Rapid, rocky or sandy streams and rivers
- Maine Snaketail (*Ophiogomphus mainensis*): Clear, rocky forested streams
- Rusty Snaketail (*Ophiogomphus rupinsulensis*): Large streams and rivers
- Riverine Clubtail (*Stylurus amnicola*): Big rivers
- Zebra Clubtail (*Stylurus scudleri*): Rivers
- Stygian Shadowdragon (*Neurocordulia yamaskanensis*): Large rivers and lakes; often rocky.

Habitat Types:

Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lower CT River
Aquatic: Lacustrine

Current Threats

Habitat Threats:

Energy Infrastructure and Development
Habitat Alteration
Sedimentation
Incompatible Recreation

Description of habitat threat(s): Bank armoring (riprap) and other disturbance that alters the river bed can kill individuals and permanently reduce habitat suitability. New dam construction alters habitat from riverine to impoundment. Dam operation alters water velocity, water level, and other hydrologic factors.

Non-Habitat Threats:

Pollution

Description of non-habitat threat(s): Wakes caused by boating can wash over and kill emerging adults.



Common Name: **Odonates-River/Stream Group**
 Scientific Name: **Odonates-River/Stream Group**
 Species Group: **Invert**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Define particular habitat requirements of each species within Vermont, utilizing current knowledge of researchers and field investigations.
Research	Distribution and Abundance	High	1) Obtain baseline distributional and abundance data for all species in group by conducting surveys throughout the state. Efforts should be focused on particular habitats required by each and, where appropriate, on regions of expected occurrence within the state (e.g., a northern peripheral species might be expected in the northern tier counties). Abundance information should be collected at sites of known occurrence. 2) Conduct inventories to detect and gather information on new SGCN odonate populations.
Research	Threats and Their Significance	Medium	Assess the vulnerability of nymphs of each species to habitat perturbations, such as siltation, temperature and water quality shifts, chemical pollution, hydropower development and operation, and changes in vegetation. Investigate the upland habitat needs of the adults and the effects of such impacts as fragmentation and reduction.
Research	Other Research	Medium	Conduct cage study of Hetaerina americana during lampricide treatment of Lewis Creek to provide information on vulnerability of the species to these chemicals.
Monitoring	Population Change	High	Monitor known SGCN odonate populations. Track population trends at distinct locations. This would follow field surveys and assessments to identify populations judged to be large and viable. Focusing on such large populations would offer greater probability of detecting population shifts.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Workshops	High	Incorporate odonate SGCN occurrence information into environmental review and technical assistance	Number of sites with odonate SGCN that received conservation benefits to this group	FWD, USFWS, DEC, ANR, VTrans, NRCS	SWG
Awareness Raising and Communications	High	Educate public and improve appreciation of vulnerable odonate habitats	Number of participants at events/workshops	FWD, VCE, watershed groups, VT River Conservancy, TU, town Cons Comms, media outlets	SWG
Easements	High	Acquisition/easement of high priority SGCN odonate riverine sites	Number of SGCN odonate sites protected	FWD, TNC, VLT, other land trusts, watershed groups	SWG, EQIP, USFWS



Common Name: **Odonates-River/Stream Group**
Scientific Name: **Odonates-River/Stream Group**
Species Group: **Invert**

Bibliography

- Brunelle, P.M. 1999. Distribution of damselflies and dragonflies (Odonata) of Maine, United States. *Northeastern Naturalist* 6(1):35-38.
- Carle, F. L. 1994. Dragonflies and damselflies (Odonata) known to or likely to occur in Vermont. A report to the Nongame and Natural Heritage Program, Vermont Fish & Wildlife Department. 22 pp.
- Carle, F. L. 1997. Odonates of the Green Mountain National Forest A report to the National Forest Service, Manchester Ranger District.
- Carpenter, V.A. 1997. Dragonflies and damselflies of Cape Cod. Edition 2. The Cape Cod Museum of Natural History, Brewster, Massachusetts. 80 pp.
- Dunkle, S.W. 2000. Dragonflies through binoculars: a field guide to dragonflies of North America. Oxford University Press, New York, NY. 266pp.
- Dunkle, S.W. 1990. Damselflies of Florida, Bermuda and the Bahamas. Scientific Publishers Nature Guide # 3.
- Dunkle, S.W. 1989. Dragonflies of the Florida Peninsula, Bermuda and the Bahamas. Scientific Publishers Nature Guide # 1.
- Lam, E. 2004. Damselflies of the Northeast. Biodiversity Books, Forest Hills, N.Y. 96 pp.
- NatureServe. 2004. NatureServe Explorer: an online encyclopedia of life [web application]. Version 4.0. NatureServe, Arlington, VA. Available <http://www.natureserve.org/explorer>.
- Needham, J. G., M. J., Jr., Westfall, and M. L. May. 2000. Dragonflies of North America. Scientific Publishers, Gainesville FL. 940 pp.
- Nikula, B., J.L. Loose, and M.R. Burne. 2003. A field guide to dragonflies and damselflies of Massachusetts. Natural Heritage and Endangered Species Program. Massachusetts Division of Fisheries and Wildlife. Boston, MA.
- Nothnagle, P. 1988. A provisional list of the odonates (dragonflies and damselflies) of Vermont. Technical Report 1. Nongame and Natural Heritage Program, Vermont Fish and Wildlife Department.
- Westfall, M.J. and M.L. May. 1996. Damselflies of North America. Scientific Publishers, Gainesville, Florida.



Common Name: **Crustaceans Group**
 Scientific Name: **Crustaceans Group**
 Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Unknown

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

This group includes the following three species:

--An amphipod (*Diporeia hoyi*): This deep-water species is rarely collected. It has been reported from Lake Champlain and Lake Memphremagog. It may occur in other deep lakes also. This species is cited as being very sensitive to pollution. It is likely to be one of the first species to disappear when a lake is polluted.

--Taconic Cave amphipod (*Stygobromus borealis*): This state-endangered species is known from only a single cave in Vermont, which is a popular spelunking cave. The status of this population is not known.

--Appalachian Mountain crayfish (*Cambarus bartonii*): This rare crayfish is only found in small cold water streams and is threatened by development (stormwater sedimentation), acid rain, climate change, and introduction of the rusty crayfish (*Orconectes rusticus*).

Distribution

--An amphipod (*Diporeia hoyi*): This species is known in Vermont only from (1) deep water in Lake Champlain from Crown Pt. to Rouses Pt., including Inner Mallets Bay and Missisquoi Bay; and (2) possibly Lake Memphremagog. It may also occur in other large lakes with similar deep water habitat. It was very rare in Lake Champlain 1992-1996 samples.

--Taconic Cave amphipod (*Stygobromus borealis*): This amphipod is known in Vermont only from Morris Cave in Danby; it may also occur in Dorset (Aeolus) Cave in Dorset, but that record is unclear. It has also been reported from MA and NY. It appears to be limited to subterranean drainage systems of karst terrain in the Taconic Mountains. Three single locations comprise the entire global distribution known for this species.

--Appalachian Mountain crayfish (*Cambarus bartonii*): This crayfish is known from the West River, Hudson drainage (Battenkill), and the Champlain Basin. It occurs both in the mountains and in small, valley streams.

Distribution by Biophysical Region:

Champlain Valley	Southern VT Piedmont
Champlain Hills	Vermont Valley
Northern Green Mtns	Southern Green Mtns
Northern VT Piedmont	Taconic Mtns
Northeastern Highlands	

Distribution by Watershed:

Known Watersheds

West

Probable Watersheds

St. Francois River



Common Name: **Crustaceans Group**
Scientific Name: **Crustaceans Group**
Species Group: **Invert**

Hudson-Hoosic
Lake Champlain
Lamoille River
Missiquoi River
Otter Creek
Winooski River
Metawee River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

--An amphipod (*Diporeia hoyi*) This is a deep-water species of large lakes, where it undergoes vertical migrations. It has been collected as deep as 900 feet in Lake Superior. It requires cold, deep water with a good oxygen supply. It is likely to be one of the first species to disappear when a lake is polluted.

--Taconic Cave amphipod (*Stygobromus borealis*) In Vermont, this amphipod is found in a cave of marble bedrock with a deep (probably > 9 meters) pool of water with a silt and/or sand bottom. In MA, it was reported from a springhouse.

--Appalachian Mountain crayfish (*Cambarus bartonii*) This rare crayfish is only found in small cold water streams. It occurs both in the mountains and in small, valley streams.

Habitat Types:

Subterranean
Aquatic: Lacustrine
Aquatic: Fluvial
Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Habitat Alteration
Sedimentation
Invasion by Exotic Species
Climate Change

Description of habitat threat(s): *D. hoyi* is limited by shifts in food web dynamics caused by zebra mussels. The invasive quagga mussel is displacing *D. hoyi* in the Great Lakes and could cause similar effects if it reaches Lake Champlain. Spiny waterflea now poses a new threat in Lake Champlain. Long-term deposition of silt is likely altering the benthic habitat this amphipod uses.

--*S. borealis*, due to the nature of its habitat, is vulnerable to hydrologic alterations that may originate away



Common Name: **Crustaceans Group**
 Scientific Name: **Crustaceans Group**
 Species Group: **Invert**

from the occupied site.

--Cambarus bartonii is limited by development (stormwater sedimentation), acid rain, climate change, and introduction of the rusty crayfish (Orconectes rusticus).

Non-Habitat Threats:

Pollution

Reproductive Traits

Trampling or Direct Impacts

Description of non-habitat threat(s): D. hoyi is pollution sensitive, needing clean, cold, well-oxygenated water.

--S. borealis, due to the nature of its habitat, is vulnerable to such problems as pollution and nutrient shift. It is also vulnerable to direct loss of individuals, due to the small size and isolated nature of the population. The reproductive capability of this species is likely much more limited than that of surface-water inhabiting amphipods due to limited food supply. This makes rebounding from population losses or poor reproductive years difficult. The threat posed by cave recreation is uncertain.

--C. bartonii lives in streams which can be greatly affected by acid rain.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	The habitat requirements for S. borealis needs to be studied and refined.
Research	Basic Life History	Medium	The life history of S. borealis is virtually unknown. Study is needed.
Research	Distribution and Abundance	High	The distribution and abundance of all three species has not been sufficiently investigated. Further survey work is needed to accurately determine the extent and status of these species in Vermont.
Research	Threats and Their Significance	High	The potential impacts of recreational and other human uses of caves supporting S. borealis need to be assessed. The potential for negative impacts due to manipulation or contamination of groundwater feeding these cave streams needs to be determined. The affects of fine sediments on the benthic habitat of D. hoyi should be studied. The effects on food web dynamics caused by zebra mussels and spiny waterflea need to be studied with regard to D. hoyi.
Monitoring	Population Change	High	1) Known S. borealis population should be monitored; methods that do not impact individual amphipods will be required. 2) Monitor known SGCN crustacean populations.
Monitoring	Habitat Change	Medium	Water quality and volume in known S. borealis cave stream should be monitored.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Crustaceans Group**
 Scientific Name: **Crustaceans Group**
 Species Group: **Invert**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration	High	Protect and restore habitats on which SGCN crustaceans are dependent through pollution abatement, riparian buffers, groundwater protection, etc.	Number of acres of land protected for river buffers, groundwater recharge, etc.	Watershed groups, USFWS, DEC, FWD, LCLT, VLT, TNC	SWG, EQUIP, LCLT, VLT, NRCS, USFWS
Easements	High	Acquisition/easement of any newly discovered <i>Stygobromus borealis</i> sites	Number of newly discovered sites protected	FWD, USFS, TNC, LCLT, VLT, USFWS	VHCB, SWG, GMNF, EQUIP, USFWS

Bibliography

Bell, R.T. 1971. Handbook of the Malacostraca of Vermont and neighboring regions (crayfish, sowbugs and their relatives). Published by author. Burlington, VT. 65 pp.

Biomonitoring Database 2005. Vermont Department of Environmental Conservation. Biomonitoring Section.

Bousfield, E.L. 1973. Shallow-water gammaridean Amphipoda of New England. Cornell University Press, Ithaca, NY. 312pp.

Dermott, R. 2001. Sudden disappearance of the amphipod *Diporeia* from eastern Lake Ontario, 1993-1995. J. Great Lakes Res. 27: 423-433.

NatureServe. 2004. NatureServe Explorer: an online encyclopedia of life [web application]. Version 4.0. NatureServe, Arlington, VA. Available <http://www.natureserve.org/explorer>.

Smith, D.G. 1995. Keys to the freshwater macroinvertebrates of Massachusetts, Second Ed. Privately published. Douglas G. Smith, 30 Montague Road, Sunderland. MA. 243pp.

Nalepa, T. F.; D.L. Fanslow, and G. Messick. 2005. Characteristics and potential causes of declining *Diporeia* spp. populations in southern Lake Michigan and Saginaw Bay, Lake Huron. Page 157-188. Publications, Agencies and Staff of the U.S. Department of Commerce. Paper 414.

Smith, D.G. 2001. Pennak's freshwater invertebrates of the United States: Porifera to Crustacea. 4th Ed. John Wiley & Sons. 638pp.

The Amphipod Homepage. [Http://www.imv.uit.no/amphipod/index.html](http://www.imv.uit.no/amphipod/index.html).



Common Name: **Freshwater Mussels Group**
Scientific Name: **Freshwater Mussels Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

This group includes 13 species. Seven are state-endangered species and three state-threatened species. One is also federally endangered. Two of the three remaining species are each known from only a single short stretch of river. Freshwater mussels are recognized as one of the most endangered groups in the state.

- Eastern pearlshell (*Margaritifera margaritifera*): State threatened. Regional SGCN*
- Dwarf wedgemussel (*Alasmidonta heterodon*): State and federal endangered. Regional SGCN*
- Brook floater (*Alasmidonta varicosa*): State threatened. Only one population, occupying only one river stretch. Regional SGCN*
- Elktoe (*Alasmidonta marginata*): Very rare. Occupies a single short river stretch. Regional SGCN*
- Alewife floater (*Anodonta implicata*): Rare. Occupies a single river stretch. Regional SGCN*
- Cylindrical papershell (*Anodontoides ferussacianus*): State endangered. Regional SGCN*
- Pocketbook (*Lampsilis ovata*): State endangered. Regional SGCN*
- Fluted-shell (*Lasmigona costata*): State endangered.
- Creek heelsplitter (*Lasmigona compressa*): Rare. Regional SGCN*
- Fragile papershell (*Leptodea fragilis*): State endangered. Regional SGCN*
- Black sandshell (*Ligumia recta*): State endangered. Probably most endangered Vermont mussel. Regional SGCN*
- Pink heelsplitter (*Potamilus alatus*): State endangered
- Giant floater (*Pyganodon grandis*): State threatened

*Regional Species of Greatest Conservation Need (RSGCN) among the 13 Northeastern states

Distribution

Eastern pearlshell (*Margaritifera margaritifera*): Reported from the upper Winooski River and Lewis Creek systems of the Champlain basin, and the Passumpsic River, West River, and Nulhegan River systems of the Connecticut River basin.

--Dwarf wedgemussel (*Alasmidonta heterodon*): Connecticut River mainstem, and slightly upstream into some large tributaries. Historically found from Bloomfield to Brattleboro. More recently known from Hartland to Springfield, and from Guildhall to Lunenburg.

--Brook floater (*Alasmidonta varicosa*): Known only from West River (Connecticut River tributary). One historic report from the Connecticut River mainstem.

--Elktoe (*Alasmidonta marginata*): Known only from a short stretch (~ 5 miles) of the Lamoille River.

--Alewife floater (*Anodonta implicata*): Occurs in Connecticut River downstream of Bellows Falls.

--Cylindrical papershell (*Anodontoides ferussacianus*): Known from Missisquoi, Lamoille, and Poultney river systems, and also Stone Bridge Brook (Milton). May occur in other Lake Champlain rivers and the main lake itself. One historic report from the Clyde River.



Common Name: Freshwater Mussels Group
Scientific Name: Freshwater Mussels Group
Species Group: Invert

--Pocketbook (*Lampsilis ovata*): Champlain basin species; reported from Lake Champlain, Missisquoi River, Lamoille River, Winooski River, LaPlatte River, Lewis Creek, Otter Creek, and Poultney River. Only found below principal fall line.

--Fluted-shell (*Lasmigona costata*): Champlain basin species; reported from Lamoille River, Winooski River, Otter Creek, Lewis Creek, and Poultney River. Historically in the Missisquoi River (shells collected), but no live specimens have been observed. Observed above the principal fall line only in Otter and Lewis creeks.

--Creek heelsplitter (*Lasmigona compressa*): Reported from several small to large tributaries to Lake Champlain. Primarily a headwater to medium-sized creek species, but does occur in larger habitats (e.g., below fall line in Winooski River). One record is from outside the Champlain basin, the Coaticook River.

--Fragile papershell (*Leptodea fragilis*): Lake Champlain basin only; reported from Lake Champlain, Missisquoi River, Lamoille River, Winooski River, Poultney River, and Otter Creek system. Only found below principal fall line.

--Black sandshell (*Ligumia recta*): Champlain basin species; reported from Missisquoi River, Otter Creek, Poultney River, and Hospital Creek, and shallow areas in Lake Champlain near the mouths of these rivers. Appears extirpated from Hospital Creek; most recent survey did not find it in Otter Creek. Only found below principal fall line.

--Pink heelsplitter (*Potamilus alatus*): Champlain basin species; reported from Lake Champlain, Missisquoi River, Lamoille River, Winooski River, Otter Creek, Lewis Creek, Hospital Creek, and Poultney River. Only found below principal fall line.

--Giant floater (*Pyganodon grandis*): Champlain basin species; reported in Lake Champlain, Missisquoi River, Lamoille River, Winooski River, East Creek, Poultney River, and Otter Creek system. Reported from above the principal fall line only in Otter Creek and Lamoille River. A high elevation pond population in Chittenden Co. is believed to be the result of an unintentional stocking.

Distribution by Biophysical Region:

Champlain Valley	Southern VT Piedmont
Champlain Hills	Vermont Valley
Northern Green Mtns	Southern Green Mtns
Northern VT Piedmont	Taconic Mtns
Northeastern Highlands	

Distribution by Watershed:

Known Watersheds	Probable Watersheds
Metawee River	St. Francois River
Waits	Middle Connecticut
Upper Connecticut-Mascoma	West
Black-Ottawquechee	
Lake Champlain	



Common Name: **Freshwater Mussels Group**
Scientific Name: **Freshwater Mussels Group**
Species Group: **Invert**

Lamoille River
Missiquoi River
Otter Creek
Passumpsic
Upper Connecticut
Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒

--Eastern pearlshell (*Margaritifera margaritifera*) This is a coldwater species, found in streams that support trout populations. It inhabits firm sand substrates, often amidst gravel and cobbles, and occasionally tightly packed cobbles and gravel. Salmonids are the fish hosts.

--Dwarf wedgemussel (*Alasmidonta heterodon*) A large river species, this mussel is found in stable mud, silty sand, sand, or gravel where the current is sufficient to keep the substrate free of surficial silt. Fish hosts may include the tessellated darter and slimy sculpin.

--Elktoe (*Alasmidonta marginata*) This is a mussel of medium-sized rivers, where it occupies riffles or rapids on stable gravel or rocky bottoms. It burrows securely into the sand-filled spaces between stones.

--Brook floater (*Alasmidonta varicosa*) This mussel occupies small rivers, where it is found in broad and shallow pools, runs, and glides, in sand and gravel substrates, often among or near cobble and boulder. Mussel beds usually support some submerged aquatic plants.

--Pocketbook (*Lampsilis ovata*) This relatively mobile mussel is found primarily in stretches of larger rivers from Lake Champlain to the first major waterfall. It also can be found in shallow areas of the main lake near deltas of these rivers. Mussels occupy firmly packed sand, sand and gravel, or silty sand.

--Fluted-shell (*Lasmigona costata*) This mussel is found primarily in medium-sized creeks to larger rivers from Lake Champlain to the first major waterfall, but also occurs above this fall line in some streams. It inhabits a variety of substrates, including mud, sand, gravel, and aggregates of cobble, gravel, and sand.

--Creek heelsplitter (*Lasmigona compressa*) This is a mussel of small creeks to small rivers, but occurs in small numbers in large river sections above Lake Champlain as well. It is found in gravel, sand, or mud.

--Pink heelsplitter (*Potamilus alatus*) This mussel occurs in large rivers only between Lake Champlain and the first major waterfall. It also occupies areas of Lake Champlain near the deltas of these rivers. It is found in clay, clayey silt, sand, gravel and sand, or mixtures of cobble, sand, and silt. Pink heelsplitters usually bury themselves nearly completely into the substrate, their shape anchoring them securely in place.

--Fragile papershell (*Leptodea fragilis*) This mussel occurs in large rivers only between Lake Champlain and the first major waterfall. It also occupies areas of Lake Champlain near the deltas of these rivers. It is found in sand, clayey silt, silty sand, or gravel and sand. Fragile papershells usually bury themselves nearly completely into the substrate, their shape anchoring them securely in place.



Common Name: Freshwater Mussels Group
Scientific Name: Freshwater Mussels Group
Species Group: Invert

--Black sandshell (*Ligumia recta*) This mussel occurs only in large rivers between Lake Champlain and the first major waterfall. Substrates include sand, sand and gravel, and mud.

--Giant floater (*Pyganodon grandis*) A relatively mobile species of large rivers and lakes, it is found in sand, sand and gravel, silty sand, and clay.

--Cylindrical papershell (*Anodontoidea ferussacianus*) This mussel is found in creeks and rivers, but likely occupies shallow lake habitat as well (Lake Champlain). Substrates include silt and silty sand in slow currents. It is known from both above and below the first waterfall upstream of Lake Champlain.

--Alewife floater (*Anodonta imbecilis*) This is a riverine mussel in Vermont found in slow to fast waters. It occurs in a variety of substrates, including sand, sand and gravel, and silt.

POTENTIAL FISH HOSTS REPORTED*:

--Eastern pearlshell: Rainbow trout, Atlantic salmon, brook trout, brown trout

--Dwarf wedgemussel: tessellated darter, slimy sculpin, banded killifish, Atlantic salmon

--Elktoe: white sucker, shorthead redhorse, rock bass

--Brook floater: slimy sculpin, longnose dace, blacknose dace, golden shiner, pumpkinseed, yellow perch, tessellated darter

--Pocketbook: smallmouth bass, white crappie, largemouth bass, bluegill, sauger, yellow perch

--Fluted-shell: common carp, spotfin shiner, longnose dace, creek chub, slimy sculpin, black crappie, yellow perch, bowfin, northern pike, bluegill, pumpkinseed, largemouth bass, walleye

--Creek heelsplitter: slimy sculpin, black crappie, spotfin shiner, yellow perch, yellow bullhead, brook stickleback, brassy minnow, bluegill, smallmouth bass, emerald shiner, mimic shiner, longnose dace, creek chub, bluntnose minnow

--Pink heelsplitter: freshwater drum

--Fragile papershell: freshwater drum

--Black sandshell: sauger, banded killifish, rosyside shiner, bluegill, redbreast sunfish, pumpkinseed, rock bass, American eel, common carp, white perch, white crappie, black crappie, largemouth bass, yellow perch, walleye

--Giant floater: banded killifish, blackchin shiner, blacknose shiner, black crappie, blacknose dace, bluegill, bluntnose minnow, brook silverside, common carp, brook stickleback, common shiner, creek chub, freshwater drum, gizzard shad, golden shiner, largemouth bass, longnose gar, pearl dace, pumpkinseed, rock bass, white crappie, white sucker, yellow bullhead, yellow perch

--Cylindrical papershell: spotfin shiner, black crappie; possibly sea lamprey, mottled sculpin, brook stickleback, white sucker, common shiner, blacknose shiner, bluntnose minnow, fathead minnow, bluegill,



Common Name: **Freshwater Mussels Group**
Scientific Name: **Freshwater Mussels Group**
Species Group: **Invert**

largemouth bass

--Alewife floater: American shad, alewife, blueback herring, white sucker, pumpkinseed, white perch

*Primarily from lab studies; this list does not indicate that fish hosts have been demonstrated in natural environment.

Habitat Types:

Aquatic: Fluvial

Aquatic: Large Lake Champlain Tribs Below Falls

Aquatic: Lower CT River

Aquatic: Man-Made Water Bodies

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Alteration

Sedimentation

Habitat Fragmentation

Invasion by Exotic Species

Description of habitat threat(s): --Zebra mussels are decimating Lake Champlain mussel populations, and are found in the lower portions of some of the lake's tributaries.

--Asiatic clams now occur in Lake George, which is upstream of Lake Champlain; these pose a future threat to Vermont's mussels.

--Bridge construction and road-related river bank stabilization have been common direct impacts. Streambank erosion and stormwater runoff can cover mussel habitat in intolerable levels of sediment.

--Dams have been responsible for large losses of habitat, particularly with the Connecticut River. Dams have converted river habitat to reservoirs, altered and degraded downstream habitats, and created barriers to movement. Hydropower dams create unnatural frequency of water level and velocity changes.

Non-Habitat Threats:

Genetics

Pollution

Harvest or Collection

Reproductive Traits

Predation or Herbivory

Trampling or Direct Impacts

Description of non-habitat threat(s): --Loss of specific fish hosts can result in reduced/eliminated reproductive success.

--Low local mussel abundance can reduce success rate of external fertilization.

--Muskrats are mussel predators and can decimate local populations when their numbers are too high or



Common Name: **Freshwater Mussels Group**
Scientific Name: **Freshwater Mussels Group**
Species Group: **Invert**

when a mussel species is particularly vulnerable.

--Low mussel densities can have genetic consequences.

--Mussels have been shown to be sensitive to a variety of pesticides and other anthropogenic chemicals.

--Wastewater effluents, stormwater runoff, and agricultural runoff can carry these pollutants into rivers where mussels reside.

--Use of mussels as bait by anglers has been problematic in some other states. This has not been investigated in Vermont.

--Damage to/death of mussels due to trampling by stream users may occur regularly, but has not been investigated. Thin-shelled species are often found dead due to breakage.

--Fisheries sampling also has the potential of affecting young and adult mussels, where equipment is in contact with the substrate.



Common Name: **Freshwater Mussels Group**
 Scientific Name: **Freshwater Mussels Group**
 Species Group: **Invert**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Define particular habitat requirements of each species within Vermont, utilizing field investigations and knowledge of researchers.
Research	Basic Life History	High	Determine host fish requirements for each mussel species. This needs to be specific to Vermont populations.
Research	Distribution and Abundance	High	1) Obtain baseline information on distribution and abundance of each mussel species. This should include all existing information sources. 2) Conduct inventories of rivers and appropriate lake habitat to detect and gather information on SGCN mussel populations.
Research	Threats and Their Significance	High	1) Assess limiting factors and their potential impacts to the aquatic habitats of each mussel species. Assessment should be on a watershed scale, including upland sources of potential limiting factors. 2) Research is needed on how to protect native mussel populations that are being impacted by zebra mussels, and also those populations that are vulnerable to further zebra mussel colonization. 3) Determine how dams affect mussel populations, their habitats, and life history requirements. 4) Investigate impacts of Missouri trawl (fisheries sampling) operation on young-of-year and older juvenile SGCN mussels.
Research	Population Genetics	High	1) Certain species (e.g., black sandshell, elktote, brook floater) are highly isolated from nearest populations. Genetic comparisons with other populations will be needed before reintroduction options can be evaluated. 2) Determine genetic constraints that may be hampering the recovery of isolated SGCN mussel populations (particularly black sandshell and elktote).
Research	Taxonomy	Medium	There is uncertainty about the species assignment of <i>Lampsilis ovata</i> . Vermont populations may be <i>L. cardium</i> , or more than one species could be here. This needs to be determined.
Research	Other Research	High	Investigate the potential benefits and risks of dam removal to SGCN mussel populations.
Monitoring	Population Change	High	Monitor known SGCN mussel populations. Track population trends at distinct locations. This would follow field surveys and assessments to identify appropriate populations for monitoring. Focusing on large populations would offer greater probability of detecting population shifts. Areas of habitat where species have disappeared need to be tracked.
Monitoring	Habitat Change	High	Changes to habitat due to specific threats (e.g., hydrodams) need to be monitored.
Monitoring	Monitor Threats	High	Zebra mussel detection and monitoring is needed in watersheds that support SGCN mussels. Occupied rivers and boating lakes that occur upstream need to be monitored. The effects of dams on downstream habitat needs to be monitored, including de-watering, temperature regime, and silt releases.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Freshwater Mussels Group**
 Scientific Name: **Freshwater Mussels Group**
 Species Group: **Invert**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Species Restoration	High	Investigate possibility of developing a freshwater mussel propagation facility for population reintroduction and enhancement. Such a facility could provide research opportunities as well.	Completion of design and plans for a mussel propagation facility	FWD, USFWS, UVM, other Northeast states	FWD, SWG, Endangered Species Section 6, PR
Habitat Restoration	High	Reduce the use of riprap in aquatic habitat as a method of bank stabilization.	Track change in annual number of stream feet impacted by new riprap projects.	FWD, DEC, ANR, NRCS, FEMA, VTrans, FHWA	PR, DJ, NRCS
Natural Processes Restoration	High	Work with regulators and dam operators/owners to reduce the impacts of dam operations on SGCN mussel populations. FERC re-licensing should require run-of-river flows.	Number of operating dams on SGCN mussel rivers that modify operations to run-of-river flows through FERC re-licensing or other negotiations.	FWD, DEC, ANR, FERC, dam operators/owners, USACOE, towns, VNRC	PR, DJ, DEC, USFWS, Hydro funds
Invasive Species Control & Prevention	High	Prevent the introduction and spread of zebra mussels.	Monitor sites of potential occurrence	LCBP, DEC, Towns, FWD, USFWS	VT Watershed Grants, LCBP, DEC
Species Restoration	High	Remove muskrats and potentially other predators where local mussel populations are vulnerable and are likely to be impacted by predation.	Continued surveillance to determine whether muskrats are impacting critical areas.	FWD, Wildlife Services (USDA), TNC, local trappers, USFWS	FWD, SWG, Endangered Species Section 6, PR
Easements	Medium	Acquire conservation easements for the protection of necessary SGCN mussel habitats and maintenance or restoration of ecological functions.	Number of riparian habitat acres acquired/enrolled	LCLT, VLT, FWD, ANR, TNC, NRCS, USFWS	LCLT, VLT, TNC, SWG, NRCS, PR, DJ
Alliance Development	High	Enhance coordination between government agencies and partners to ensure consistency in respective program implementation and increased sensitivity to SGCN mussel requirements and problems.	Number of mussel-pertinent permit and project notifications that bridge between agencies/organizations annually.	ANR, USFWS, COE, FEMA, FHWA, NRCS, Wildlife Services, VTrans, others	PR, SWG

***Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report***



Common Name: **Freshwater Mussels Group**
Scientific Name: **Freshwater Mussels Group**
Species Group: **Invert**

Habitat Restoration	High	Protect and restore habitats on which SGCN mussels are dependent through pollution abatement, riparian buffers, flow regulation, etc.	Number of acres of riparian habitat protected and/or restored	LCLT, VLT, Watershed groups, USFWS, DEC, FWD	SWG, LCLT, VLT, NRCS



Common Name: Freshwater Mussels Group
Scientific Name: Freshwater Mussels Group
Species Group: Invert

Bibliography

- Bauer, G. 1983. Age structure, age specific mortality rates and population trend of the freshwater pearl mussel (*Margaritifera margaritifera*) in North Bavaria. - Archiv fuer Hydrobiologie 98:523-532.
- Bauer, G. 1987. Reproductive strategy of the freshwater pearl mussel *Margaritifera margaritifera*. Journal of Animal Ecology 56: 691-704.
- Bauer, G. 1988. Threats to the freshwater pearl mussel *Margaritifera margaritifera* L. in central Europe. Biological Conservation 45: 239-253.
- Baxter, R. M. 1977. Environmental effects of dams and impoundments. Annual Review of Ecology and Systematics 8: 255-283.
- Bogan, A.E. 1993. Freshwater bivalve extinctions (Mollusca: Unionoida): a search for causes. American Zoologist 33: 599-609.
- Bogan, A.E. 1996. Decline and decimation: the extirpation of the unionid freshwater bivalves of North America. Journal of Shellfish Research 15(2): 484.
- Fichtel, C., and D.G. Smith. 1995. The Freshwater Mussels of Vermont. Leahy Press, Montpelier, VT.
- Neddeau, E.J., M.A. McCollough, and B.I. Swartz. 2000. The Freshwater Mussels of Maine. Maine Department of Inland Fisheries and Wildlife, Augusta, ME.
- Neddeau E.J. (2008) Freshwater Mussels and the Connecticut River Watershed. Connecticut River Watershed Council, Greenfield, MA.
- Clarke, A. H. 1981. The Freshwater Molluscs of Canada. National Museum of Natural Science. Ottawa, ON.
- Strayer, D. L., and Jirka, K. J. 1997. The Pearly Mussels of New York State. University of the State of New York, State Education Department. 113p.
- Brim Box, J. and J. Mossa. 1999. Sediment, land use and freshwater mussels: prospects and problems. Journal of the North American Benthological Society 18:99-117.
- Clarke, A. H. 1981. The freshwater molluscs of Canada. National Museum of Natural Sciences/National Museums of Canada, Ottawa, Canada. 446 pp.
- Cope, W. G., and D. L. Waller. 1995. Evaluation of freshwater mussel relocation as a conservation and management strategy. Regulated Rivers: Research & Management. 11(2): 147-155.
- Cummings, K. S., and C. A. Mayer. 1992. Field guide to freshwater mussels of the Midwest. Illinois Natural History Survey, Manual 5. Champaigne, IL. 194 pp.
- Di Maio, J. and L. D. Corkum. 1995. Relationship between the spatial distribution of freshwater mussels (Bivalvia: Unionidae) and the hydrological variability of rivers. Canadian Journal of Zoology 73: 663-671.
- Dunn, H. L. and B. E. Sietman. 1997. Guidelines used in four geographically diverse unionid relocations. Pages 176-183 in Cummings, K. S., A. C. Buchanan, C. A. Mayer, and T. J. Naimo (eds.). Conservation and management of freshwater mussels II: initiatives for the future. Proceedings of a Upper Mississippi River Conservation Committee symposium, 16-18 October 1995, St. Louis, Missouri. Upper Mississippi River Conservation Committee, Rock Island, Illinois.
- Fichtel, C., and D. G. Smith. 1995. The freshwater mussels of Vermont. Nongame & Natural Heritage Program, Vermont Fish and Wildlife Department. 54 pp.
- Goudreau, S.E., R.J. Neves and R.J. Sheehan. 1993. Effects of wastewater treatment plant effluents on freshwater mollusks in the upper Clinch River, Virginia, USA. Hydrobiologia 252: 211-230.
- Haag, W. R. and M. L. Warren. 1998. Role of ecological factors and reproductive strategies in structuring freshwater mussel communities. Canadian Journal of Fisheries and Aquatic Sciences 55: 297-306.
- Hallac, D. E. 1999. Conservation of unionid mussels threatened by zebra mussels (*Dreissena polymorpha*) in Lake Champlain, VT: and examination of species vulnerability, conservation strategies, and causal mechanisms of stress. M.S. Thesis, UCM. 110 pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Freshwater Mussels Group
Scientific Name: Freshwater Mussels Group
Species Group: Invert

Johnson, R.I. 1980. Zoogeography of North American Unionacea (Mollusca: Bivalvia) north of the maximum Pleistocene glaciation. *Bulletin of the Museum of Comparative Zoology* 149(2): 77-189.

Jokela, J. and Mutikainen, P. Effect of size-dependent muskrat (*Ondatra zibethica*) predation on spatial distribution of a freshwater clam, *Anodonta piscinalis* Nilss. (Unionidae, Bivalvia). *Canadian Journal of Zoology* 73: 1085-1094.

Keller, A.E., and M. Lydy. 1997. Biomonitoring and the hazards of contaminants to freshwater mollusks. Unpublished report in: Freshwater mollusks as indicators of water quality: a workshop. U.S. Geological Survey Biological Resources Division and National Water Quality Assessment Program. 55 pp.

Lewis, J.B., and P.N. Riebel. 1984. The effect of substrate on burrowing in freshwater mussels (Unionidae). *Canadian Journal of Zoology* 62: 2023-2025.

Ligon, F. K., W. E. Dietrich, and W. J. Trush. 1995. Downstream ecological effects of dams: a geomorphic perspective. *Bioscience* 45 (3): 183-192.

Michaelson, D. and R. J. Neves. 1995. Life history and habitat of the endangered dwarf wedgemussel *Alasmidonta heterodon* (Bivalvia: Unionidae). *Journal of the North American Benthological Society* 14:324-340.

Mills, E. L., G. Rosenberg, A. P. Spidle, M. Ludyanskiy, Y. Pligin, and B. May. 1996. A review of the biology and ecology of the quagga mussel (*Dreissena bugensis*), a second species of freshwater dreissenid introduced to North America. *American Zoologist* 36: 271-286.

National Native Mussel Conservation Committee. 1998. National strategy for the conservation of native freshwater mussels. *Journal of Shellfish Research* 17: 1419-1428.

NatureServe. 2004. NatureServe Explorer: an online encyclopedia of life [web application]. Version 4.0. NatureServe, Arlington, VA. Available <http://www.natureserve.org/explorer>.

Neves, R. J., and M. C. Odum. 1989. Muskrat predation on endangered freshwater mussels in Virginia. *Journal of Wildlife Management* 53(4): 939-940.

Neddeau, E. J., M. A. McCollough, and B. I. Swartz. 2000. The freshwater mussels of Maine. Maine Department of Inland Fisheries and Wildlife. 118 pp.

Neddeau, E. J., and J. Victoria. 2003. A field guide to the freshwater mussels of Connecticut. Connecticut Department of Environmental Protection. 32 pp.

Richter, B.D., D.P. Braun, M. A. Mendelson, and L.L. Master. 1997. Threats to imperiled freshwater fauna. *Conservation Biology* 11:1081-1093.

Schloesser, D.W., T.F. Nalepa, and G.L. Mackie. 1996. Zebra mussel infestation of unionid bivalves (Unionidae) in North America. *American Zoologist* 36: 300-310.

Smith, D. G. 1985. Recent range expansion of the freshwater mussel *Anodonta imbecilis* and its relationship to clupeid fish restoration in the Connecticut River system. *Freshwater Invertebrate Biology* 4: 105-108.

Strand, M., and R. W. Merritt. 1999. Impacts of livestock grazing activities on stream insect communities and the riverine environment. *American Entomologist* 45(1): 13-29.

Strayer, D. L. 1999. Use of flow refuges by unionid mussels in rivers. *Journal of the North American Benthological Society* 18:468-476.

Strayer, D., and K. Jirka. 1997. The pearly mussels of New York State. New York State Museum, Memoir 26. 113 pp.

Watters, G. T. 1994. An annotated bibliography of the reproduction and propagation of the Unionoidea. (primarily of North America). Ohio Biological Survey Miscellaneous Contribution No. 1. vi+ 158pp.

Watters, G. T. 1996. Small dams as barriers to freshwater mussels (Bivalvia, Unionoidea) and their hosts. *Biological Conservation* 78: 79-85.

Williams, J. D., M. L. Warren, K. S. Cummings, J. L. Harris, and R. J. Neves. 1993. Conservation status of freshwater mussels of

***Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report***



Common Name: **Freshwater Mussels Group**
Scientific Name: **Freshwater Mussels Group**
Species Group: **Invert**

the United States and Canada. Fisheries 18: 6-22.

Yeager, B.L. 1994. Impacts of reservoirs on the aquatic environment of regulated rivers. Tennessee Valley Authority, Water Resources, Aquatic Biology Department, TVA/WR/AB-93/1, Norris, Tennessee.



Common Name: **Freshwater Snails Group**
Scientific Name: **Freshwater Snails Group**
Species Group: **Invert**

Conservation Assessment

Final Assessment: High Priority

Global Rank:

Global Trend:

State Rank:

State Trend: Declining

Extirpated in VT?

Regional SGCN?

Assessment Narrative:

A diverse group of 15 snail species whose general status ranges from extirpated to declining to rare. Much work is needed to refine status assessment. This group includes

- Fringed valvata (*Valvata lewisi*): Recorded from four locations in VT: St Catherine, and associated Little Pond, Lake Salem and Berlin Pond. Possibly occurs elsewhere. Only found in low numbers throughout range.
- Mossy valvata (*Valvata sincera*): Only found at four locations in VT: three in Lake Champlain, and Flag pond.
- Squat duskysnail (*Lyogyrus granum*): Only three locations: Lake St Catherine, Connecticut River, Lake Fairlee. An East Coast species.
- Pupa duskysnail (*Lyogyrus pupoideus*): Only six collections, all in Lake Champlain, may be susceptible to zebra mussel colonization.
- Canadian duskysnail (*Lyogyrus walkeri*): Only known from Lake St. Catherine.
- Buffalo pebblesnail (*Gillia altilis*): 12 locations in: Lake Champlain, Missisquoi River, Connecticut River, Crystal Lake, Indian Brook Reservoir, Hinkum Pond. Atlantic drainage species.
- Boreal marstonia (*Marstonia lustrica*): Found in Lake Champlain, Laplatte River and Joes Pond. Uncommon in northern part of range; more common southward. Reported as abundant at some Massachusetts sites.
- Liver elimia (*Goniobasis livescens*): Eleven sites, all in Lake Champlain. May be vulnerable to invasives such as *Bithynia tentaculata*, zebra mussel.
- Sharp hornsail (*Pleurocera acuta*): No recent occurrences from Vermont; may be extirpated. Reported historically from VT, QE, and NY in literature.
- Spindle lymnaea (*Acella haldemani*): Only known from one lake in VT: Beebee Pond, a lake periodically chemically treated and managed for invasive Eurasian Milfoil. Limited and localized in distribution. Appears to be greatly reduced from historical range. Often only one location is reported for a lake, but it is easily overlooked. Reproduction may be a limiting factor.
- Dusky ancylid (*Laevapex fuscus*): Reproductive traits may be a limiting factor (low colonization rate).
- Mammoth lymnaea (*Bulinnaea megasoma*): Reported from Lake Champlain and its tributaries in 19th century literature; no recent records. May be extirpated; only VT historic record, a small stream in Burlington area, has since been filled in.
- Country fossaria (*Fossaria rustica*): Only found in two streams in Champlain Valley: Little Otter and Lewis creeks. Locations have been resampled since 1997 and not found.
- Star gyro (*Gyraulus crista*): Limited distribution in VT. Found in seven rivers: White River, Calendar Brook tributary, East Branch Passumpsic River, Ranch Brook, Sunny Brook, East Creek-South Fork, Morehouse Brook. Holarctic (northern); distribution. Rare in NY.
- Thicklip rams-horn (*Planorbula armigera*): Found at four locations; three of these are in urban or agricultural watersheds with degraded water quality.



Common Name: **Freshwater Snails Group**
Scientific Name: **Freshwater Snails Group**
Species Group: **Invert**

Distribution

- Fringed valvata (*Valvata lewisi*): Recorded from four locations: St Catherine, and associated Little Pond, Lake Salem and Berlin Pond. Possibly occurs elsewhere.
- Mossy valvata (*Valvata sincera*): Recorded at four locations: three in Lake Champlain, and Flagg pond.
- Squat dusksnail (*Lyogyrus granum*): Three locations: Lake St Catherine, Connecticut River, Lake Fairlee.
- Pupa dusksnail (*Lyogyrus pupoideus*): Only six collections, all in Lake Champlain.
- Canadian dusksnail (*Lyogyrus walkeri*): Only known from Lake St. Catherine.
- Buffalo pebblesnail (*Gillia altilis*): 12 locations in: Lake Champlain, Missisquoi River, Connecticut River, Crystal Lake, Indian Brook Reservoir, Hinkum Pond.
- Boreal marstonia (*Marstonia lustrica*): Found in Lake Champlain, Laplatte River and Joes Pond.
- Liver elimia (*Goniobasis livescens*): Eleven sites, all in Lake Champlain.
- Sharp hornsnail (*Pleurocera acuta*): No recent occurrences from Vermont; may be extirpated. Reported historically from VT, QE, and NY in literature.
- Spindle lymnaea (*Acella haldemani*): Only known from one lake in VT: Beebee Pond.
- Dusky ancyliid (*Laevapex fuscus*): Known from two locations: Colchester Pond and Jones Mill Pond.
- Mammoth lymnaea (*Bulimnaea megasoma*): Reported from Lake Champlain and its tributaries in 19th century literature; no recent records. May be extirpated; only VT historic record, a small stream in Burlington area, has since been filled in.
- Country fossaria (*Fossaria rustica*): Only found in two streams in Champlain Valley: Little Otter and Lewis creeks. Locations have been resampled since 1997 and not found.
- Star gyro (*Gyraulus crista*): Limited distribution in VT. Found in seven rivers: White River, Calendar Brook tributary, East Branch Passumpsic River, Ranch Brook, Sunny Brook, East Creek-South Fork, Morehouse Brook.
- Thicklip rams-horn (*Planorbula armigera*): Found at four locations: Jewett Brook, Indian Brook, Pleasant Brook, and Leicester Sedge Marsh.

Distribution by Biophysical Region:

Champlain Valley	Southern VT Piedmont
Champlain Hills	Vermont Valley
Northern Green Mtns	Southern Green Mtns
Northern VT Piedmont	Taconic Mtns
Northeastern Highlands	

Distribution by Watershed:



Common Name: **Freshwater Snails Group**
Scientific Name: **Freshwater Snails Group**
Species Group: **Invert**

Known Watersheds

Middle Connecticut
Waits
West
Black - Ottauquechee
Upper Connecticut - Mascoma
Lake Champlain
Lamoille River
Missiquoi River
Otter Creek
Passumpsic
St. Francois River
Upper Connecticut
White
Winooski River
Metawee River

Probable Watersheds

Deerfield
Hudson-Hoosic

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

- Boreal marstonia (*Marstonia lustrica*) Found in well-buffered lakes and streams.
- Liver elimia (*Goniobasis livescens*) Grazes in shallow water on stones and gravel in lakes and clear rapid streams. In lakes, it burrows into the sand and feeds on algae and bacteria; this is not as good a food source as it is for other pulmonate snails.
- Sharp hornsnail (*Pleurocera acuta*) Found in a variety of habitats. In lakes, found on boulders on exposed shores or in mud and sand. In rivers, found on stones in rapid current. Snails remain in shallow water up to 1 m deep, where they burrow under the sand and layers of decaying leaves and other organic matter.
- Spindle lymnaea (*Acella haldemani*) Found on submerged logs, silt, sand, and mud; up to 2 m deep. Often is attached to leaves and stems of aquatic pondweed and other submerged vegetation. Reported to favor eutrophic lakes and ponds. Young don't travel far from where they hatched, leading to a clumped distribution.
- Mammoth lymnaea (*Bulinnaea megasoma*) Found in ponds, large and small lakes, and quiet embayment of rivers.
- Country fossaria (*Fossaria rustica*) Occupies rivers and streams, lakes, ponds, and occasionally ditches and canals. Can be in damp mud flats and bodies of water with fluctuations in water level.
- Star gyro (*Gyraulus crista*) Found in dense aquatic vegetation, water-logged wood, and rotting terrestrial leaves (in water).
- Fringed valvata (*Valvata lewisi*) A pool/pond/lake species. Found particularly on sand, but also mud and aquatic vegetation down to 7 m.



Common Name: Freshwater Snails Group
Scientific Name: Freshwater Snails Group
Species Group: Invert

--Mossy valvata (*Valvata sincera*) Generally limited to oligotrophic and mesotrophic situations, but occasionally in eutrophic waters. Associated with submerged aquatic vegetation. In MA, however, it is reported as requiring deep lakes with a pH of 7.6 or greater, where it is often associated with rooted vegetation.

--Squat dusksnail (*Lyogyrus granum*) Found on organic debris and vegetation in standing waters of larger lakes and ponds, oxbows, and major rivers. Highly tolerant of acidic conditions, but limited tolerance to sodium (e.g., road salt).

--Canadian dusksnail (*Lyogyrus walkeri*) Found in sluggish streams and quiet ponds where dead plants accumulate. Occupies lentic waters, ranging from oligotrophic to marl ponds.

--Buffalo pebblesnail (*Gillia altilis*) Large lakes and rivers. In Hudson River, it is found on mud and aquatic plants in the shallows.

--Pupa dusksnail (*Lyogyrus pupoideus*) Occurs in small to large ponds and large rivers. Found on organic debris and aquatic plants. Hybridization with *A. grana* reported.

--Dusky ancylid (*Laevapex fuscus*) Most commonly found in still waters, such as impoundments, back waters, and ponds; occasionally in temporary waters. Generally absent from mountainous regions. Absent from low diversity habitats and tends to have low colonization rates, but will compete well.

--Thicklip rams-horn (*Planorbula armigera*) Most likely to be found in slow streams, wetlands, temporary ponds, and ditches.

Habitat Types:

Seeps and Pools

Wet Shores

Aquatic: Fluvial

Aquatic: Large Lake Champlain Tribs Below Falls

Aquatic: Lacustrine

Aquatic: Lower CT River

Aquatic: Man-Made Water Bodies

Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Conversion of Habitat

Energy Infrastructure and Development

Habitat Alteration

Sedimentation

Impacts of Roads or Transportation Systems

Habitat Fragmentation

Invasion by Exotic Species

Climate Change

Description of habitat threat(s): --Sedimentation and chloride from road and impervious surface runoff.
--Lake/reservoir water level manipulation may be a problem to snails.



Common Name: **Freshwater Snails Group**
Scientific Name: **Freshwater Snails Group**
Species Group: **Invert**

- Acidification (acid rain) may be a problem for species sensitive to low pH, low calcium.
- Copper sulfate used for treating waters for algae and swimmers itch is a risk to snails (it is a molluscicide).
- Baylicide, used to reduce sea lamprey populations, is a molluscicide.
- Nuisance aquatic plant management can impact snails by removing snails and habitat (plants) and covering lake bottoms.
- Goniobasis livescens is at risk to invasive Bithynia tentaculata.
- Lake species vulnerable to zebra mussels.

Non-Habitat Threats:

Competition

Pollution

Reproductive Traits

- Description of non-habitat threat(s):** --Valvata lewisi - only found in very low densities throughout range; this may limit long-term viability of local populations.
- Valvata sincera - low reproductive rate. Only a few eggs (4-12) produced per individual.
 - Lyogyrus granum - low tolerance to sodium (e.g., road salt). This may impact populations anywhere they are found (particularly in rivers).
 - Lyogyrus pupoideus - reported to hybridize with Amnicola grana. This could jeopardize both species where they co-occur.
 - Goniobasis livescens - Likely being impacted by the exotic Bithynia tentaculata (snail) in Lake Champlain. Region mollusc expert Dr. Doug Smith (Umass) believes it will become extirpated from the lake for this reason.
 - Acella haldemani - Apparently greatly reduced in distribution and abundance from historical range for unknown reasons. The young don't travel far from where the hatch, which creates limited ability to disperse and colonize/recolonize other habitat patches.
 - Bulimnaea megasoma - may have been extirpated due to unknown causes.
 - Laevapex fuscus - reproductive traits may be a limiting factor (low colonization rate).
-
- Copper sulfate used for treating waters for algae and swimmers itch is a risk to snails (it is a molluscicide).
 - Baylicide, used to reduce sea lamprey populations, is a molluscicide.
 - Nuisance aquatic plant management can impact snails by removing snails and habitat (plants) and covering lake bottoms.
 - Fisheries reclamation in lakes (rotenone) is known to impact snail communities.



Common Name: **Freshwater Snails Group**
Scientific Name: **Freshwater Snails Group**
Species Group: **Invert**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Define particular habitat requirements of each species within Vermont, utilizing current knowledge of researchers and field investigations.
Research	Distribution and Abundance	High	1) Obtain baseline distributional and abundance data for all species in group by conducting surveys throughout the state. Abundance information should be collected at sites of known occurrence. 2) Conduct inventories to detect and gather information on new SGCN snail populations.
Research	Threats and Their Significance	High	Assess potential and existing impacts of limiting factors to habitat and individual species. Such limiting factors as habitat loss and degradation, exotic invasive snails, and use of pesticides should be examined.
Research	Population Genetics	High	Investigate genetic relationships of isolated populations to their larger population strongholds; potentially addressing taxonomic uncertainties.
Research	Taxonomy	Medium	Taxonomic questions regarding <i>Fossaria rustica</i> need to be resolved.
Monitoring	Population Change	Medium	1) Monitor known SGCN snail populations. 2) Population monitoring could be employed to track population trends at distinct locations. This would follow field surveys and assessments to identify populations judged to be large and viable. Focusing on such large populations would offer greater probability of detecting population shifts. Monitoring populations subjected to specific environmental perturbations should also be considered.
Monitoring	Habitat Change	High	Changes to habitat due to specific threats (e.g., dam removal) need to be monitored.
Monitoring	Monitor Threats	High	Number of waterbodies and areas chemically treated to control snails and algae needs to be tracked and used to assess the significance of this limiting factor to SGCN snails.
Monitoring	Other Monitoring Needs	High	Changes to populations due to specific threats (e.g., dam removal) need to be monitored.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Freshwater Snails Group**
 Scientific Name: **Freshwater Snails Group**
 Species Group: **Invert**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks	High	Work to reduce the problem and impacts of acidification on aquatic habitat		DEC, other state regulators, legislators, governor	
Alliance Development	Medium	Enhance coordination between government agencies and partners to ensure consistency in respective program implementation and increased sensitivity to SGCN snail requirements and threats.		ANR, USFWS	
Invasive Species Control & Prevention	High	Prevent the introduction and spread of invasive exotic species (particularly snails)		FWD, DEC, LCBP, USFWS	
Habitat Restoration	High	Protect and restore habitats on which SGCN snails are dependent through pollution abatement, riparian buffers, etc.	Number of acres of riparian and lakeshore natural vegetation protected and/or restored. Number of acres of lake habitat restored/protected	LCLT, VLT, Watershed groups, USFWS, DEC, FWD	SWG, LCLT, VLT, NRCS
Compatible Resource Use	High	Reduce the use of algicides, molluscicides, and other pesticides in waters where it may impact SGCN snails	Sustained reduction in the number of annual requests for use of pesticides in SGCN waters	FWD, DEC, ANR, lake associations, private landowners	

Bibliography

Vermont Department of Environmental Conservation, Biomonitoring Database 2014

Burch, J.B. 1982. Freshwater snails (Mollusca: Gastropoda) of North America. United States Environmental Protection Agency. Environmental Monitoring and Support Laboratory. EPA-600/3-82-026. 294pp.

Clarke, A. H. 1981. The freshwater molluscs of Canada. National Museum of Natural Sciences, National Museums of Canada, Ottawa, Canada. 446 pp.

Jokinen, E.H. 1983. The freshwater snails of Connecticut. State Geological and Natural History Survey of Connecticut. Survey Bull. 109. 83pp.

Jokinen, E. 1992. The freshwater snails (Mollusca: Gastropoda) of New York State. New York State Museum Bulletin 482, 1-112.

NatureServe. 2004. NatureServe Explorer: an online encyclopedia of life [web application]. Version 4.0. NatureServe, Arlington, VA. Available <http://www.natureserve.org/explorer>.

Vermont's Wildlife Action Plan

Mammal SGCN Conservation Reports

Appendix A5

Draft 10/1/2015

Species Page

Cinereus or Masked Shrew	1
Water Shrew	5
Smoky Shrew.....	8
Long-tailed or Rock Shrew	12
Pygmy Shrew.....	16
Hairy-tailed Mole	19
Little Brown Bat/Myotis.....	22
Indiana Bat.....	26
Small-footed Bat	31
Northern Long-eared Bat	36
Silver-haired Bat	42
Tri-colored bat.....	47
Big Brown Bat	51
Eastern Red Bat.....	56
Hoary Bat.....	61
New England Cottontail.....	66
Snowshoe Hare	69

Species Page

Southern Flying Squirrel.....	72
Northern Flying Squirrel	75
Rock Vole.....	79
Woodland Vole.....	84
Muskrat	87
Southern Bog Lemming.....	91
Northern bog lemming.....	96
Wolf.....	99
Gray Fox	105
Black Bear.....	111
American Marten.....	115
Long-tailed Weasel	120
Northern River Otter	123
Canada Lynx.....	126
Bobcat.....	131
Eastern Mountain Lion	135
Moose	140

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Cinereus or Masked Shrew**
Scientific Name: **Sorex cinereus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5

State Trend: Unknown

Extirpated in VT? NA

Regional SGCN? no

Assessment Narrative:

The trend of the cinereus (masked) shrew in Vermont is unknown. Historic records indicate a widespread distribution of the species in the state. Since 2008, during an effort to develop a small mammal atlas, the masked shrew was verified at 28 sites in Vermont further supporting the belief that the species continues to be widely distributed throughout the state. Although it may be the most common of the small shrews, it is still believed to be relatively rare at most locations and there is insufficient data on this species in Vermont to fully assess its status. The masked shrew may be more common at higher elevations but the overall role of elevation on the species distribution remains unclear. It is believed that masked shrews are more common in old growth or late successional forests.

Distribution

Considered rare though broadly distributed.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☐

Masked shrews are known to favor cool mesic deciduous and coniferous forests often at higher elevations. They are sometimes found in mixed habitat types such as edges of bogs and other cool and wet sites (seeps). The masked shrew uses grasses, rocks, and logs or stumps for cover. They are primarily carnivorous and insectivorous consuming worms, spiders, snails, slugs, and small amounts of vegetable matter. Dampness of site and depth of leaf litter, seems to be critical factors in determining habitat use. The species is known to utilize down woody debris. Its home range is understood to be less than 0.5ha.



Common Name: **Cinereus or Masked Shrew**
Scientific Name: **Sorex cinereus**
Species Group: **Mammal**

Habitat Types:

Upland Shores
Outcrops and Alpine
Cliffs and Talus
Northern Hardwood
Hardwood Swamps
Softwood Swamps
Seeps and Pools
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests
Open Peatlands
Early Succession Boreal Conifers
Early Succession Pine and Hemlock
Early Succession Upland Oak
Early Succession Other Types
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Wet Shores
Shrub Swamps
Early Succession Boreal Hardwoods
Early Succession Spruce-Fir
Early Succession Northern Hardwoods
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lower CT River

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development
Habitat Alteration
Climate Change

Description of habitat threat(s): Climate changes, habitat alterations/degradations, and/or habitat conversions resulting in overall drier conditions may significantly limit the species in Vermont. Although it is speculated that the direct impacts of energy infrastructure development (e.g. industrial wind power projects) could result in the reduction of upper elevation spruce fir habitats used by this species, more

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Cinereus or Masked Shrew**
 Scientific Name: **Sorex cinereus**
 Species Group: **Mammal**

research is needed to determine just how severe this impact would be on Vermont's population of masked shrews.

Non-Habitat Threats:

Competition

Pollution

Loss of Prey Base

Description of non-habitat threat(s): Previous studies indicate that competition from other shrews may pose a significant risk to masked shrews. Furthermore, there is concern that acid rain could alter the ecology of soil invertebrates resulting in adverse impacts to the prey base upon which this species depends.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Attempt to narrow down habitat requirements and determine critical habitat needs of the masked shrew.
Research	Basic Life History	Low	Determine basic life history and population demographics
Research	Distribution and Abundance	Medium	Determine distribution and abundance of the masked shrew and maintain a database of known locations.
Monitoring	Population Change	Medium	Analyze habitat conditions and local populations prior to construction of upper elevation wind generating facilities
Monitoring	Monitor Threats	Medium	Monitor known populations, e.g. Camel's Hump spruce-fir zone, to detect any significant population changes related to climate change

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	High	Conduct research to determine impact of warming, especially as it relates to drying of habitats	Amount of forest habitat protected	UVM, Middlebury, Johnson State College	SWG
Compatible Resource Use	High	Minimize habitat fragmentation	Amount of forest habitat protected from development	VFPR, GMP, UVM, TNC	EQIP, SWG
Standards	Medium	Minimize fragmentation (the permanent conversion of habitat as a result of development) between populations in core habitats. Maintain habitat mosaic	Number of travel corridors identified and protected	UVM, Middlebury, Johnson State College	SWG
Standards	Medium	Determine appropriate management strategies to improve and conserve habitat	Number of habitats identified and protected.	VFPR, USFS, Coverts	SWG



Common Name: Cinereus or Masked Shrew
Scientific Name: Sorex cinereus
Species Group: Mammal

Bibliography

- Buckner, C.h. 1966. Populations and ecological relationships of shrews in tamarack bogs of southeastern Manitoba. *Journal of Mammalogy* 47: 181-194.
- Brannon, M.P. 2000. Niche relationships of two syntopic species of shrews, *Sorex fumeus* and *S. cinereus* in the southern Appalachian Mountains. *Journal of Mammalogy*, 81:1053-1061.
- DeGraaf, Richard M. and Yamasaki, Mariko. *New England Wildlife: Habitat, Natural History, and Distribution*. University Press of New England, Hanover, NH, 2001.
- Doucet, J.G., and Riji Bider. 1974. The effects of weather on the activity of the masked shrew. *Journal of Mammalogy*, 55:348-363.
- Godin, A.J. 1977. *Wild Mammals of New England*. Johns Hopkins University Press. Baltimore. MD. 303pp.
- Hamilton, W.J., Jr. 1930. The food of soricidae. *Journal of Mammalogy*, 11:26-39.
- Innes, D.G.L., J.F. Bendall, and B.J. Naylor. 1990. High densities of the masked shrew, *Sorex cinereus*, in Jack pine plantations in northern Ontario.
- Kilpatrick, C. W., and J. Benoit. 2011. Small mammal project. University of Vermont/NorthWoods Stewardship Center, final report submitted to Vermont Fish and Wildlife Department
- Kirk, George L. 1916. The mammals of Vermont, *Joint Bulletin No. 2 Vermont botanical and Bird Club* 2: 28-34.
- Kirkland, G.L. Jr. and D.F. Schmidt. 1982. Abundance, habitat, reproduction and morphology of forest dwelling small mammals of Nova Scotia and southeastern New Brunswick, *Canadian field Naturalist* 96:156-162.
- Osgood, Frederick L. Jr., 1938. The mammals of Vermont. *J.Mammalogy* 19(4): 435-441.
- Saunders, D.Andrew. 1980. *Adirondack Mammals*. Adirondack Wildlife Program, State University of New York (College of Environmental Science and forestry -Syracuse) 216 pp.
- Whitaker, J.O. Jr. and W.J. Hamilton Jr. 1998. *Mammals of the Eastern United States*. 3rd edition. Cornell University Press. Ithaca. NY. 583 pp.
- Whitaker, J.O., Jr. 2004. *Sorex cinereus*. *Mammalian Species*, 743:1-9.



Common Name: **Water Shrew**
Scientific Name: **Sorex palustris**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S3

State Trend: unknown

Extirpated in VT? no

Regional SGCN? yes

Assessment Narrative:

There is very limited information regarding the water shrew in Vermont. Fifty seven specimens have been taken since 1915 from 21 different localities throughout the state including two specimens collected since 2008 as part of the effort to develop a statewide small mammal atlas. Historic records of this species' occurrence in the state indicate that there may be limited at risk populations.

The species is listed as a high priority because not a lot is known about it in the state and because of its very specific habitat requirements. Although there is no evidence of a decline in the state at this time, a number of potential limiting factors have been identified including changes in natural water flow regimes resulting from climate change, dams/flow regulation, inadequate riparian buffers, atmospheric deposition and acid rain, loss of habitat, and potential loss of prey base. Furthermore, there are taxonomic uncertainties and speculation exist that it may actually be more than one species.

Distribution

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☐

The water shrew is found in undercut banks of streams and beaver dams. It is assumed that wooded buffers on streams are desirable and there is a recognition that wooded wetlands and streams are utilized more often than cattail dominated wetlands. Mesic forests are believed to be important. Although the water shrew has been trapped on dry creekbeds, it may prefer streams that flow year-round. DeGraaf (2001) suggests that coniferous forests are preferred over deciduous forests. Whitaker & Hamilton (1998) found this species on mud flats of sluggish backwaters. It is believed that habitats adjacent to water, particularly fast cold streams, may hold the largest populations. Critical habitat appears to be undercut banks of streams and possibly beaver dams.



Common Name: **Water Shrew**
Scientific Name: **Sorex palustris**
Species Group: **Mammal**

Habitat Types:

Spruce Fir Northern Hardwood
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Wet Shores
Shrub Swamps
Aquatic: Fluvial

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Inadequate Disturbance Regime
Impacts of Roads or Transportation Systems

Description of habitat threat(s): Alteration or conversion of riparian buffers as a result of certain forest management and development practices could degrade key habitat requirements of the water shrew and impact its survival and productivity. Similarly, alterations of natural water flow regimes resulting from climate change could pose significant impacts on the species. The lack of baseline data on the distribution, abundance and basic life-history of water shrews in Vermont prevents a comprehensive assessment of the threats facing the species.

Non-Habitat Threats:

Pollution
Loss of Prey Base

Description of non-habitat threat(s): It is believed that the species' prey base could be impacted by the effects of acid rain.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Assess habitat at historical sites and sample for species.
Research	Basic Life History	High	Sample stomach contents to determine prey preferences.
Research	Distribution and Abundance	High	Develop baseline data on distribution and abundance
Research	Threats and Their Significance	High	Identify key limiting factors to this species.
Research	Population Genetics	High	Determine the extent of gene flow in the state.
Research	Taxonomy	High	Determine whether or not this is a single species.



Common Name: **Water Shrew**
Scientific Name: **Sorex palustris**
Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Compatible Resource Use	High	Minimize trail or road intrusion into wetlands or riparian buffers.	Miles of riparian buffers and acres of wetlands intact and protected	NRCS, VLT, Coverts	EQIP, SWG
Habitat Restoration	High	Enhance or restore degraded wetlands and repair buffers	Number of habitats maintained or improved	NRCS, DEC Wetlands, VT Rivers Conservancy	SWG, EQIP
Habitat Restoration	High	Identify and maintain rich wetland habitats and stream buffers.	Number of habitats identified and maintained	FVPR, NRCS, VLT, Coverts	EQIP, SWG

Bibliography

- Degraaf, R.M. and M. Yamasaki. 2001. New England Wildlife. Univ. Press of New England. Hanover. N.H.
- Kilpatrick, C. W., and J. Benoit. 2011. Small mammal project. University of Vermont/NorthWoods Stewardship Center, final report submitted to Vermont Fish and Wildlife Department
- Kirk, George L. 1916. The mammals of Vermont, Joint Bulletin No. 2 Vermont botanical and Bird Club 2: 28-34.
- Osgood, F.L. 1938. The mammals of Vermont J. of Mammalogy. 19(4): 435-441.
- Whitaker, J.O., Jr. and William J. Hamilton.1998. Mammals of the Eastern United States. Comstock Publishing, Ithaca.
- Van Zyll de Jong, C.G. 1985. Handbook of Canadian Mammals. Volume 2. Bats. National Museums of Canada, Ottawa, Ontario, Canada. 212 pp.



Common Name: **Smoky Shrew**
Scientific Name: **Sorex fumeus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S4

State Trend: unknown

Extirpated in VT? no

Regional SGCN? Yes

Assessment Narrative:

The smoky shrew is listed as a Regional Species of Greatest Conservation Need (RSGCN) among the 13 Northeastern states. Although the status of the smoky shrew in Vermont is poorly understood, the few records of its occurrence in the state indicate that the species is more limited in numbers than masked shrews. Compared with other relatively abundant shrews, the smoky shrew has more specific habitat requirements.

Distribution

Confident everywhere but Taconic Mountains and the Vermont Valley where it is unknown

Distribution by Biophysical Region:

Champlain Valley	Unknown	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

The smoky shrew often occupies damp, boulder-strewn, upland woods (DeGraff and Yamasaki 2001). It is found in cool mesic forests, often conifer, that are shady with deep, loose, leaf litter and is often associated with higher elevation sites with damp, moss covered rocks. Smoky shrews are also typically found along streams with moss covered banks (DeGraff and Yamasaki 2001). The dietary niche of the smoky shrew is broader than that of other shrews. Although it is 80 % insectivorous, it will also eat earthworms, spiders, snails, salamanders, small mammals, and birds. (Brannon 2000). The smoky shrew uses tunnels made by other animals for nesting as well as beneath stumps and rotten logs. It is also known to use downed woody debris for cover and foraging. Loose damp leaf litter may be critical to habitat use.



Common Name: **Smoky Shrew**
Scientific Name: **Sorex fumeus**
Species Group: **Mammal**

Habitat Types:

Outcrops and Alpine
Cliffs and Talus
Northern Hardwood
Softwood Swamps
Seeps and Pools
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Open Peatlands
Early Succession Boreal Conifers
Early Succession Spruce-Fir
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development

Description of habitat threat(s): The conversion of forest to non-forest habitat is thought to be a potential impact on smoky shrews. Similarly, the construction of upper elevation wind energy facilities causes major habitat conversion from areas of moist, boulder strewn, and loose humus conditions favored by these shrews to dry warm sites, including roadways, avoided by them.

Non-Habitat Threats:

Competition
Pollution
Loss of Prey Base

Description of non-habitat threat(s): Competition and predation from other shrews (*Blarina brevicauda*) may be a problem.
Acid rain may reduce invertebrate prey base.

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Smoky Shrew**
Scientific Name: **Sorex fumeus**
Species Group: **Mammal**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Determine critical habitat needs in Vermont. Narrow the habitat requirements. Are there areas within available habitats where species are concentrated.
Research	Basic Life History	Low	Determine the basic life history requirements.
Research	Distribution and Abundance	Medium	Determine distribution and abundance in Vermont.
Research	Threats and Their Significance	Medium	Determine the significance of potential impacts resulting from wind industry developments and other disturbances of preferred habitats, competition with other shrews, and pollution.
Monitoring	Population Change	Low	Monitor population changes in high elevation sites planned for wind energy development before and after construction.
Monitoring	Habitat Change	Low	Monitor changes to and availability of identified critical habitats in Vermont.
Monitoring	Range Shifts	Medium	Monitor distribution and abundance in Vermont to assess range shifts.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Standards	Medium	Protect stream buffers sufficient to maintain a mesic environment	Miles of riparian buffers intact and protected	NRCS, VLT, Coverts	EQIP, SWG
Habitat Restoration	High	Maintain prey base	Identification of prey use, abundance and distribution	UVM, Middlebury, Johnson State College	SWG
Compatible Resource Use	Medium	Identify and maintain rich mesic habitats	Number of habitats identified and maintained	Vermont Forest and Parks Dept., USFS, Coverts	SWG



Common Name: **Smoky Shrew**
Scientific Name: **Sorex fumeus**
Species Group: **Mammal**

Bibliography

- Brannon, M.P. 2000. Niche relationships of two syntopic species of shrews, *Sorex fumeus* and *S. cinereus* in the southern Appalachian Mountains. *Journal of Mammalogy*, 81:1053-1060.
- DeGraaf, Richard M. and Yamasaki, Mariko. *New England Wildlife: Habitat, Natural History, and Distribution*. University Press of New England, Hanover, NH, 2001.
- Godin, A.J. 1977. *Wild Mammals of New England*. Johns Hopkins University Press. Baltimore. MD. 303pp.
- Kilpatrick, C. W., and J. Benoit. 2011. Small mammal project. University of Vermont/NorthWoods Stewardship Center, final report submitted to Vermont Fish and Wildlife Department
- Kirk, George L. 1916. The mammals of Vermont, Joint Bulletin No. 2 Vermont botanical and Bird Club 2: 28-34.
- Owen, J.G. 1984. *Sorex fumeas*. *Mammalian species* 215:1-8
- Hamilton, W.J. Jr. 1940. The biology of the smoky shrew (*Sorex fumeas fumeas*) (Miller), *Zoologica*. 25: 473-492.
- Kirkland, G.L. Jr. and D.F. Schmidt. 1982. Abundance, habitat, reproduction and morphology of forest dwelling small mammals of Nova Scotia and southeastern New Brunswick, *Canadian field Naturalist* 96:156-162.
- Osgood, Frederick L. Jr., 1938. The mammals of Vermont . *J.Mammalogy* 19(4): 435-441.
- Saunders, D.Andrew. 1980. *Adirondack Mammals*. Adirondack Wildlife Program, State University of New York (College of Environmental Science and forestry -Syracuse) 216 pp.
- Whitaker, J.O. Jr. and W.J. Hamilton Jr. 1998. *Mammals of the Eastern United States*. 3rd edition. Cornell University Press. Ithaca. NY. 583 pp.



Common Name: **Long-tailed or Rock Shrew**
Scientific Name: **Sorex dispar**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: S2

State Trend: unknown

Extirpated in VT? No

Regional SGCN? yes

Assessment Narrative:

The long-tailed shrew (rock shrew) is listed as a Regional Species of Greatest Conservation Need (RSGCN) among the 13 Northeastern states. It is further listed as an S2 species in Vermont. The long-tailed shrew is currently listed as a C-2 species by the USFWS indicating the species may be endangered or threatened but insufficient information is currently available to allow preparation of rules for listing the species. The total number of known occurrences in the state is 32 (Tumosa 2001, Chipman 1994, Kilpatrick and Benoit 2011, VT Natural Heritage database). Eleven specimens of the long-tailed shrew were obtained between 2008 and 2011 as part of the development of a small mammal atlas in Vermont. Two of the specimens collected during this effort provided the first records of existence in the Northeastern Highlands Biophysical Region and in Orange County. Eleven specimens were collected prior to 1940, one specimen was taken on Camels Hump in 1968, six specimens were obtained between 1972 and 1989, and three long-tailed shrews were captured at sites in the Northern Green Mountains biophysical region in 1994.

It is believed that the long-tail shrew occurs in limited, localized, at-risk populations however current data is to limited to fully assess the species' status. In Vermont, the species is also believed to be primarily associated with talus slopes and is only occasionally found in association with mountain streams and never in large forest openings such as clearcuts (Kilpatrick and Benoit 2011).

Distribution

The long-tailed shrew was historically found in the prominent talus habitat located along the western slopes of the Southern Green Mountains in Mendon and Wallingford (Rutland County) and was further documented to occur on Camel's Hump (Chittenden and Washington Counties) however no recent records of their occurrence in these locations exist. More recently, the species has been reported from the talus dominated habitats found on Mount Mansfield (Chittenden and Lamoille Counties), Wheeler Mountain (Caledonia County), West Mountain (Essex County) and Mount Ascutney (Windsor County). The long-tailed shrew has also been reported from the Champlain Hills since 1990 and was documented in the towns of Jericho and Vershire.

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Not Probable
Northern Green Mtns	Confident	Southern Green Mtns	Historic Records Only
Northern VT Piedmont	Confident	Taconic Mtns	Not Probable
Northeastern Highlands	Confident		

Distribution by Watershed:



Common Name: **Long-tailed or Rock Shrew**
Scientific Name: **Sorex dispar**
Species Group: **Mammal**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☐

The long-tailed shrew prefers cold, mesic forests and is typically found in close proximity to streams having undercut banks. The species often inhabits cool, talus slopes and moss covered boulders and logs. It is believed that moss covered rocks and logs provide critical shade and protective cover. Similarly, forested talus slopes are also believed to be an important habitat feature where long-tailed shrews spend most of their time in the labyrinth of spaces between rocks about a foot beneath the surface (Kirkland 1981). Although generally associated with coniferous forests, the long-tailed shrew may also be found in deciduous and mixed forest types. May be associated with rock vole.

Habitat Types:

Cliffs and Talus

Northern Hardwood

Spruce Fir Northern Hardwood

Current Threats

Habitat Threats:

Conversion of Habitat

Energy Infrastructure and Development

Climate Change

Incompatible Recreation

Description of habitat threat(s): Ski trails and associated structures could impact the habitat of the long-tailed shrew. Conversion of habitat as a result of quarrying activities could also destroy critical rocky, talus habitat.

Climate Change may significantly warm and dry the moist talus slopes favored by these shrews. In addition, the upper elevation development of wind energy facilities could result in the conversion of suitable long-tailed shrew habitat to a more open, bare rock and/or grass dominated habitat.

Non-Habitat Threats:

Genetics

Pollution

Loss of Prey Base

Description of non-habitat threat(s): Change in prey base due to acid rain deposition at high elevations. Shrews feed on invertebrates and therefore may accumulate pesticides and heavy metals in body tissue (Tumosa 2001). Mining, mercury deposition, and the application of sewage sludge can all negatively affect long-tailed shrews due to a build up of toxins in the body. Furthermore, because the species is believed to occur in limited, localized, at-risk populations, there is a risk of reduced survival and fecundity due to inbreeding depression.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Long-tailed or Rock Shrew**
 Scientific Name: **Sorex dispar**
 Species Group: **Mammal**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	Identify habitats critical for the perpetuation of the species
Research	Basic Life History	High	Determine home range and other life history needs.
Research	Distribution and Abundance	High	1) Determine distribution and abundance in a multi year monitoring effort. 2) Re-census historical habitats and survey in other likely habitats. 3) Map confirmed habitats.
Research	Threats and Their Significance	High	Determine significance of environmental toxicity on preferred prey base and survival.
Research	Population Genetics	High	Determine the isolation of existing populations and the need for the protection of movement corridors.
Monitoring	Population Change	Medium	Determine current status of the population and monitor changes to this population through the future.
Monitoring	Habitat Change	High	Monitor populations near or adjacent to high elevation development to determine long range changes.
Monitoring	Range Shifts	High	Monitor current populations to determine any change due to climate change.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Standards	Medium	Ensure that ski trail development and maintainance and energy development follows best management practices.	Number of habitats protected	VFPR, GMP, Ski Areas	SWG
Privately-Owned Protected Areas	High	Minimize fragmentation (the permanent conversion of habitats as a result of development) between populations in core habitats	Number of travel corridors identified and protected.	TNC, VLT, Coverts, VHCB, VFPR	SWG, VHCB
Habitat Restoration	High	Determine appropriate management strategies to improve and conserve habitat.	Number of Habitats identified and protected	TNC, University of Vermont, Middlebury College, VFPR	SWG



Common Name: **Long-tailed or Rock Shrew**
Scientific Name: **Sorex dispar**
Species Group: **Mammal**

Bibliography

- Chipman, R. B. 1994. Distribution, relative abundance, and habitat use by small mammals in Vermont. Unpublished M.S. thesis, University of Vermont, 168 pp.
- Degraaf, R.M. And Mariko Yamasake. 2001. New England Wildlife: Habitat, Natural History, and Distribution. University Press of New England, Hanover.
- French, T.W., and K.L.Crowell.1985. Distribution and status of the uellow-nosed bole and rock shrew in New York. New York Game and Fish Journal, 32:26-40.
- Kilpatrick, C. W., and J. Benoit. 2011. Small mammal project. University of Vermont/NorthWoods Stewardship Center, final report submitted to Vermont Fish and Wildlife Department
- Kirk, George L. 1916. The mammals of Vermont, Joint Bulletin No. 2 Vermont botanical and Bird Club 2: 28-34.
- Kirkland, Gordon L. 1981.Sorex dispar and Sores gaspensis. Mammalian Species No. 155 American Society Mammalogy 4pp.
- Osgood, Frederick L. Jr., 1938. The mammals of Vermont . J.Mammalogy 19(4): 435-441.
- Richmond, N.D., and W.C. Grimm. 1950. Ecology and distribution of the shrew Sorex dispar in Pennsylvania. Ecology, 31: 279-282.
- Tumosa, J. 2001. United States Forest Service species data collection form. Sorex dispar. 15pp.
- Wihou, D.E. and Sue Ruft ed. 1999. The Smithsonian Book of North American Mammals. Smithsonian Institute Press Washington.
- Whiticker, J.O., Jr. and W.J. Hamilton, Jr. 1998. Mammals of the Eastern U. S. Comstock Publishing, Cornell Univ. Press. Ithaca.



Common Name: **Pygmy Shrew**
Scientific Name: **Sorex hoyi**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2

State Trend: Unknown

Extirpated in VT? no

Regional SGCN? No

Assessment Narrative:

The pygmy shrew is listed as an S2 species in Vermont. The species appears to be rare in the state based on the scarcity of occurrence records though this could be due to survey methods. The pygmy shrew is hard to catch and difficult to identify so it may be more abundant than records would suggest. Very little is known about the historic population of this species in Vermont.

Distribution

Very few records of occurrence of pygmy shrews exist in Vermont. Furthermore, the species was not detected during a state wide small mammal survey conducted between 2008 and 2010 (Kilpatrick and Benoit, 2011).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Probable
Champlain Hills	Confident	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Habitat requirements of the pygmy shrew are relatively unknown. It is believed that mesic forests and fields are used but it has also been recorded in swamps and marshes. Critical habitat is often listed as boreal forests where wet and dry areas occur together. Disturbed sites and cultivated areas with leaf litter and downed logs may also be important habitats for pygmy shrews. The species is believed to require moist leaf mold near water (DeGraff and Yamasaki, 2001) and is typically found within 100 yards of water. There is no evidence in the literature that it prefers any particular forest age class. It was found in 4 different age classes of cove hardwood stands in southern Appalachia (Ford et al. 1996 in Tumosa 2001).



Common Name: **Pygmy Shrew**
Scientific Name: **Sorex hoyi**
Species Group: **Mammal**

Habitat Types:

Northern Hardwood
Hardwood Swamps
Marshes and Sedge Meadows
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Early Succession Boreal Hardwoods

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration

Description of habitat threat(s): The distribution and abundance of pygmy shrews in Vermont, as well as the species' specific habitat requirements, are poorly understood. Therefore, additional information is required before a comprehensive threat assessment can be completed.

Non-Habitat Threats:

Loss of Prey Base

Description of non-habitat threat(s): It is speculated that changes to habitat resulting from succession, alteration and/or conversion could result in the diminishment of the species' prey base.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Collect baseline data on habitat use and identify critical habitats
Research	Distribution and Abundance	High	Collect baseline data on distribution and abundance



Common Name: **Pygmy Shrew**
Scientific Name: **Sorex hoyi**
Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Standards	Medium	Protect, through voluntary management practices, habitat from wetland, to grassland to forest.	Monitoring and demonstrating use of these habitats	NRCS, VLT, Coverts, Consulting Foresters	EQIP, SWG,
Habitat Restoration	Medium	Restore any missing habitats identified above.	Shrew restoration and use of restored habitats.	UVM, Middlebury, Johnson State College	SWG
Compatible Resource Use	Medium	Identify and maintain a mosaic of habitats	Number of habitats identified and maintained	VFPR., USFS, Coverts	SWG
Research	High	Determine habitat requirements and distribution	Development and adoption of habitat guidelines for the species	UVM, VFWD	SWG

Bibliography

- Brannon, M.P. 2000. Niche relationships of two syntopic species of shrews, *Sorex fumeus* and *S. cinereus*, in the southern Appalachian mountains. *Journal of Mammalogy* 81(4): 1053-1061.
- Degraaf, R.M. and M. Yamasaki. 2001. *New England Wildlife*. Univ. Press of New England. Hanover. N.H.
- Kilpatrick, C. W., and J. Benoit. 2011. Small mammal project. University of Vermont/NorthWoods Stewardship Center, final report submitted to Vermont Fish and Wildlife Department
- Kirk, George L. 1916. The mammals of Vermont, Joint Bulletin No. 2 Vermont botanical and Bird Club 2: 28-34.
- Long, Charles A. 1974. *Microsorex hoyi* and *microsorex thompsoni*. *Mammalian Species* No. 33: 1-4.
- Wilson D.E. and S. Ruff. 1999. *The Smithsonian Book of North American Mammals*. Smithsonian Institution Press. Washington . D.C.
- Osgood, F.L. 1938. The mammals of Vermont *J. of Mammalogy*. 19(4): 435-441.
- Whitaker, J.O., Jr. and William J. Hamilton.1998. *Mammals of the Eastern United States*. Comstock Publishing, Ithaca.
- Saunders, D.A. *Adirondack Mammals* State Univ. of N.Y.
- Tumosa, J. 2001. United States Forest Service species data collection form for *Microsorex hoyi*. 17 pp.



Common Name: **Hairy-tailed Mole**
 Scientific Name: **Parascalops breweri**
 Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S3S4

State Trend: unknown

Extirpated in VT? no

Regional SGCN? Yes

Assessment Narrative:

The hairy-tailed mole is listed as a Regional Species of Greatest Conservation Need (RSGCN) among the 13 Northeastern states.

Believed to be relatively common, but population status and trends are unknown. The Vermont Small Mammal Atlas verified distributional records in Orleans, Essex, Chittenden, Caledonia, Addison, Washington, Windsor, and Windham counties based on results of surveys (for herps), incidental pick up or photographs of dead specimens, and from voucher specimens at the Zadock Thomson Natural History Collection at the UCMM (Kilpatrick and Benoit 2011). Little is known about this species' status and habitat requirements. Loss of habitat with sandy and sandy loam soils is a concern

Distribution

The Vermont Small Mammal Atlas verified distributional records in Orleans, Essex, Chittenden, Caledonia, Addison, Washington, Windsor, and Windham counties based on results of surveys (for herps), incidental pick up or photographs of dead specimens, and from voucher specimens at the Zadock Thomson Natural History Collection at the University of Vermont (Kilpatrick and Benoit 2011).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Historic Records Only
Northern Green Mtns	Probable	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Historic Records Only
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

Found in all places with well-drained sandy loam soils (e.g. agricultural fields and older forests). Open deciduous woodlands with thick humus are preferred. Hairy tailed moles are also adapted to second growth stands, old fields, and hedgerows. They prefer well-drained, light, moist soil with well-mixed organic matter and minerals and avoid soils that are hard, dry, or with a large clay content. Species is not restricted to any one habitat type or successional stage (Hallett 1978).



Common Name: **Hairy-tailed Mole**
 Scientific Name: **Parascalops breweri**
 Species Group: **Mammal**

Habitat Types:

Northern Hardwood
 Spruce Fir Northern Hardwood
 Oak-Pine Northern Hardwood
 Grasslands, Hedgerows, Old Field, Shrub, or Orchard
 Lawns, Gardens, and Row Crops

Current Threats

Habitat Threats:

Conversion of Habitat
 Unknown Habitat Threats

Description of habitat threat(s): Conversion of required habitats to houses, roads or other development may negatively impact the species..

Non-Habitat Threats:

Trampling or Direct Impacts

Description of non-habitat threat(s): Because of human/mole conflicts proximity to humans can result in decline. The application of pesticides/rodenticides may also cause localized population declines, particularly in orchards. The status of the species in forested habitats is unknown.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Collect baseline data on habitat requirements.
Research	Distribution and Abundance	Medium	Collect baseline data on distribution and abundance.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Standards	Medium	Develop guidelines for pest control professionals for the non-lethal control of the species	Number of trained pest control professionals	Agricultural Extension, Pest Control Professionals	SWG
Research	Medium	Monitor distribution and abundance of species	Distribution map	Agricultural Extension, UVM, Pest Control Professionals	SWG



Common Name: **Hairy-tailed Mole**
Scientific Name: **Parascalops breweri**
Species Group: **Mammal**

Bibliography

DeGraaf, R. M., and M. Yamasaki,. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover, New Hampshire, USA.

Eadie, W.R. 1939. A contribution to the biology of *Parascalops breweri*. *Journal of Mammalogy* 20:150-173.

Godin, A. J. 1977. Wild mammals of New England. Johns Hopkins University Press, Baltimore, Maryland, USA.

Gorman, M. L., and R. D. Stone. 1990. The natural history of moles. Cornell University Press, Ithaca, New York, USA.

Hallett, J.G. 1978. *Parascalops breweri*. *Mammalian Species* 98:1-4.

Kilpatrick, C. W., and J. Benoit. 2011. Small mammal project. University of Vermont/NorthWoods Stewardship Center, final report submitted to Vermont Fish and Wildlife Department



Common Name: **Little Brown Bat/Myotis**
Scientific Name: **Myotis lucifugus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G3

Global Trend:

State Rank: S1

State Trend: Stable

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The Little Brown Bat/Myotis has been identified as a Very High Concern Regional Species of Greatest Conservation Need. Before White-nose Syndrome, the little brown bat was considered to be relatively common and was one of the most frequently captured bat species in state-wide surveys. The Little Brown Bat/Myotis relies heavily on human dwellings as maternity sites and less frequently uses trees. Both maternity colony habitat and winter hibernacula are vulnerable and at risk. Every year bats lose hundreds of possible building roosts as a result of exclusion and eviction or the actual removal of old barns and other structures. Although Little Brown Bats/Myotis are known to hibernate in slightly greater than 20 sites in Vermont, the vast majority of the population hibernates in a single cave. This species is also impacted by the removal or killing of bats in structures, as well as recreational spelunking in hibernacula. Little brown myotis have experienced population declines of 90% in Vermont due to White-nose Syndrome (Darling and Smith 2011) and have experienced similar or greater mortality rates region-wide (Turner et al. 2011). The state-wide population is a fraction of what it once was and concentrated gatherings of bats at maternity colonies are particularly vulnerable to incidental mortality as evidenced by citizen reports of up to 23 bats found dead in a furnace in one summer after they flew down the chimney, probably in search of a warm roost. Trend information is needed on this species in the years following White-nose Syndrome to determine whether populations will recover from the disease.

Distribution

Distribution, including both maternity colonies and dispersed males, was historically statewide from spring through early fall before the massive population declines caused by White-nose Syndrome (WNS). Little Brown Bats/Myotis migrate to their winter hibernacula both in Vermont and in neighboring states such as New York. This species has been historically documented at nearly every known bat hibernacula in the state. In the years following WNS, maternity colonies appear to be concentrated in the greater Champlain Valley and northern Taconic Mountains and a few in the Southern Vermont Piedmont, though males and non-reproductive females likely still exist state-wide according to acoustic survey data.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:



Common Name: **Little Brown Bat/Myotis**
Scientific Name: **Myotis lucifugus**
Species Group: **Mammal**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

During the winter Little Brown Bats/Myotis hibernate in caves with a constant temperature of 40 degrees F and a relative humidity of 80% (Banfield 1974: 42 in DeGraff and Yamasaki, 2001). Little Brown Bats/Myotis often hibernate in large clusters. To prevent dehydration they awaken every ten to fourteen days to consume water. This is thought to act as a buffer against water loss, enabling longer hibernation between arousals (Sanders 2004). During the summer the Little Brown Bat/Myotis often inhabits attics where the temperature may average 100 degrees (Chenger 2004). Females form large nursery colonies that numbered in the hundreds or even thousands of individuals before White-nose Syndrome. Capture data and citizen reports indicate that males spend the summer months scattered around the state, either solitary or in small bachelor groups in buildings or trees. Colonies usually exist close to water because little brown bats seem to prefer to forage over water. When foraging, the bats may repeat a set hunting pattern within a few miles of the roost (Chenger 2004). Little brown bats eat moths, wasps, gnats, crane flies, and beetles. Young are born in May, June, or early July. Average litter size is one (Davis and Hitchcock 1965).

Habitat Types:

Northern Hardwood
Hardwood Swamps
Spruce Fir Northern Hardwood
Marshes and Sedge Meadows
Subterranean
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops
Wet Swales and Ditches
Shrub Swamps
Building or Structure
Mine
Powerlines/RR/Roadsides
Aquatic: Fluvial
Aquatic: Lacustrine
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Incompatible Recreation



Common Name: Little Brown Bat/Myotis
Scientific Name: *Myotis lucifugus*
Species Group: Mammal

Description of habitat threat(s): Disturbance in hibernacula reduces fat reserves and negatively affects reproduction and survivability. In addition, every year Little Brown Bats/Myotis lose hundreds of possible building roosts due to exclusion or the actual destruction of buildings. Direct killing of bats is common due to human fears about rabies, bat bites and histoplasmosis.

Non-Habitat Threats:

Genetics

Disease

Description of non-habitat threat(s): The Little Brown Bat/Myotis has suffered population declines upwards of 90 in Vermont and the Northeast (Turner et al. 2014) and White Nose Syndrome continues to be a threat. Pesticides and environmental poisons have had negative impacts on, and increased the mortality rates of, bat populations. Bats store some lipophilic pesticides in brown adipose fat tissues. These stores are released as bats use their fat reserves during hibernation. Depending upon tissue levels of the pesticide, as well as the amount of fat used over a given time period, bats can be exposed to both chronic and acute poisoning which can result in death. At lower levels, chronic poisoning may raise a bat's metabolism, burning the limited fat resources more quickly and possibly causing them to starve to death. In addition, broad spectrum insecticides can deplete insect diversity and limit the food sources available for bats.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	Continue the maternity colony location and monitoring program to plot changes in distribution, abundance, and population size in the years following White-nose Syndrome.
Research	Threats and Their Significance	Medium	Research and quantify the effect of evicting from or incidentally taking maternity colonies in structures on reproductive success.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	Medium	Research the effectiveness of bat houses for maternity colonies evicted and excluded from buildings.
Monitoring	Population Change	High	Monitor changes in population size in the years following White-nose Syndrome.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	Medium	Investigate observed post-WNS range shifts from state-wide occurrences of maternity colonies to concentrations in the Champlain Valley.
Monitoring	Monitor Threats	High	Monitor the continued population effects of White-nose Syndrome and cooperate on research about individual survivors.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Little Brown Bat/Myotis**
 Scientific Name: **Myotis lucifugus**
 Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Technical Assistance, Training, Learning Networks	High	Train nuisance wildlife control operators (NWCOS) in proper bat exclusion techniques. Work with homeowners/landowners to safely exclude bats and erect bat houses for displaced colonies.	Implement Best Management Practices and train nuisance wildlife control operators.	Wildlife Rehabilitators, NWCOS, Homeowners associations	USFWS, PR
Habitat Restoration	High	Maintain at least 20 maternity colony sites and a minimum of 10,000 adult females.	Number of maternity sites and bats protected	Coverts, Vermont Woodlands Magazine, NWF, UVM, Mammals subcommittee of ESA,	SWG, PR USFWS, WNS
Habitat Restoration	High	Protect hibernacula containing 100 or more little brown bats	Number of hibernacula protected	Vermont Cavers Assoc., UVM, TNC, VLT, Coverts	USFWS, TNC, VLT

Bibliography

- Banfield, A.W.F. The mammals of Canada. Toronto, ON: University of Toronto Press; 1974. 438p. In R.M. Degraaf and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover.
- Chenger, J. 2003. Bat Inventory for Project Lands of the Upper Connecticut River Basin. Prepared for the U.S. Army Corp of Engineers New England District by Bat Conservation and Management, Inc. 102 pp
- Davis, W.H. and H.B. Hitchcock. 1965. Biology and migration of the bat, *Myotis lucifugus* in New England Journal of Mammalogy, 46: 296-313.
- Fenton, M.B., and R.M.R. Barclay, 1980. *Myotis lucifugus*. Mammalian species, 142:1-8.
- Fenton, M.B. 1969. Summer activity of *Myotis lucifugus* (Chiroptera vespertilionidas) at hibernacula in Ontario and Quebec. Canadian Journal of Zoology, 47:597-602.
- Godin, A. 1977. Wild mammals of New England. Baltimore: The John Hopkeins University Press.
- Griffin, D.R. 1940. Notes on the life histories of New England cave bats. Journal of Mammalogy, 21:181-187.
- Sanders, Chris. 2004. Vermont Multi-Species Bat Conservation and Recovery Plan. 55pp.
- Trombulak, S.C., P.E. Higuera, and M. DesMeules. 2001. Population trends of wintering bats in Vermont. Northeastern Naturalists, 8:51-62.
- Turner, GG, DM Reader, and JTH Coleman. 2011. A five-year assessment of mortality and geographic spread of white-nose syndrome in North America and a look to the future. Bat Research News, 52:13-27.
- Van Zyll de Jong, C.G. 1985. Handbook of Canadian Mammals. Volume 2. Bats. National Museums of Canada, Ottawa, Ontario, Canada. 212 pp.



Common Name: **Indiana Bat**
Scientific Name: **Myotis sodalis**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G2

Global Trend:

State Rank: S1

State Trend: Stable

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The Indiana bat has been identified as a Very High Concern Regional Species of Greatest Conservation Need. Indiana bats are listed as endangered in Vermont. They have declined range wide by 60% since monitoring began in the 1960's. Historic numbers were estimated at 800,000 in the late 1960's and by 1997 the range wide population was down to 350,000 (USFWS, 1999 in Sanders, 2004). Surveys in the 2000's indicated that regionally the population was rebounding and may have been increasing until the deadly fungal disease, White-nose Syndrome (WNS), was found in the state. Nationally, declines could have been related to disturbance in hibernacula and more recently in the northeast this species has suffered mass mortality from WNS (Turner 2011). Limited dispersal may be a problem for pregnant females. Vermont is the only New England state known to harbor maternity colonies. Radio-transmitted Indiana bats roosting in the Champlain Valley come from hibernacula in New York (Sanders, 2004). Because the majority of Vermont's summer population is believed to hibernate in a single abandoned mine in Essex County, New York, they are especially vulnerable to disturbance and disease transmission. In Vermont in the 1940-50s, Indiana bats were reported in the 1000s in hibernacula. Historic hibernacula included the Plymouth Caves, Nickwacket Cave, Dorset Cave, and the Ely Copper Mine. Currently, Vermont has two hibernacula used by Indiana bats: Brandon Silver Mine (3 bats in 2011) and Little Skinner Hollow (53 bats in 2013). This species has high interannual fidelity to roost sites and is vulnerable to habitat fragmentation.

Distribution

The Indiana bat is distributed throughout the lower Champlain valley and northern Taconic Mountains during the summer, with populations concentrated around roost trees and maternity colony sites. Small numbers of this species hibernate in Vermont, though most of the Champlain Valley population is known to hibernate across the Lake in a large abandoned mine in NY state. Hibernacula: Brandon Silver Mine, owned by The Nature Conservancy (TNC); Dorset Cave, gated and owned by TNC. Skinner Hollow, unprotected and privately owned. private ownership. Nickwackett Cave, gated and privately owned.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Historic Record(s) Only
Northern VT Piedmont	Historic Record(s) Only	Taconic Mtns	Confident
Northeastern Highlands	Not Probable		

Distribution by Watershed:



Common Name: **Indiana Bat**
Scientific Name: **Myotis sodalis**
Species Group: **Mammal**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Maternity roosts are in large diameter shaggy hardwoods with sloughing bark (maple, shagbark hickory, poplar) or snags. Indiana bats roost under loose or peeling bark or in crevices and require nearby water (within a few hundred meters) to forage over. Colonies typically select one or more primary roost tree that receives direct sunlight for most of the day. Additional alternate roost trees may be shaded or in the open. During the winter months Indiana bats hibernate and require caves with a specific microclimate. Cave conditions that include cool, stable temperatures are preferred. Roost sites that are below 10 degrees Celsius when they arrive and 3-6 degrees in mid-winter allow for population increases (Tuttle and Kennedy, 2002 in Tumosa, 2003). Relative humidity above 78% but below saturation is also important. It appears that there is fidelity to the hibernaculum. Indiana bats in Kentucky travel over 300 miles to maternity areas in Michigan (Kurta and Murray, 2000 in the Vermont bat conservation plan). They have also been documented flying over 20 miles in one hour during migration (Sanders and Cheng, 2001 in the Vermont bat conservation plan). Indiana bats are insectivorous, eating mostly flies, moths, beetles, and caddis flies. Mosquitoes, midges, bees and other flying insects are also consumed (USFWS, 1999 in Tumosa, 2001). During the swarming period, the area within 0-2 miles of the hibernaculum is critical for foraging and night roosting; 2-5 miles is important, and 5-10 miles gets used but not as frequently. Connectivity between habitats may be important but is poorly understood.

Habitat Types:

Northern Hardwood
Seeps and Pools
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Open Peatlands
Marshes and Sedge Meadows
Subterranean
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Wet Shores
Shrub Swamps
Mine
Powerlines/RR/Roadsides
Aquatic: Fluvial
Aquatic: Lacustrine
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat



Common Name: **Indiana Bat**
Scientific Name: **Myotis sodalis**
Species Group: **Mammal**

Habitat Alteration

Climate Change

Incompatible Recreation

Description of habitat threat(s): Disturbance of winter hibernacula is a significant problem to Indiana bats. These bats have been documented to lose 15-20% of their body weight during hibernation in an undisturbed hibernaculum (Johnson et al. 1997, in Sanders, 2004). Disturbance of hibernating bats causes them to awaken and forces them to use additional limited energy reserves (Sanders, 2004). Arousal can use up enough fat to sustain a bat for 10-30 days (Thomas et al. 1990, Thomas 1995). Changes in temperature and light, as well as direct contact, can cause a bat to awaken and deplete stored fat reserves. Alterations to cave and mine openings can change the microclimate of a hibernacula and affect bat survival. Loss of maternity roosts may also be a problem to survivability of young. Maternity roosts can house several hundred individual bats. Felling of a maternity roost tree can impact the survival of both adults and young. Development within close proximity of hibernacula, particularly along travel corridors could also be detrimental to survival. Destruction/development of summer habitats are likely to negatively affect bats if potential roost sites and foraging areas are altered (Tumosa 2003).

Non-Habitat Threats:

Genetics

Disease

Loss of Prey Base

Description of non-habitat threat(s): Pesticides and environmental poisons have had negative impacts on, and increased the mortality rates of, bat populations. Bats store some lipophilic pesticides in brown adipose fat tissues. These stores are released as bats use their fat reserves during hibernation. Depending upon tissue levels of the pesticide, as well as the amount of fat used over a given time period, bats can be exposed to both chronic and acute poisoning which can result in death. At lower levels, chronic poisoning may raise a bat's metabolism, burning the limited fat resources more quickly and possibly causing them to starve to death. In addition, broad spectrum insecticides can deplete insect diversity and limit the food sources available for bats.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Indiana Bat**
 Scientific Name: **Myotis sodalis**
 Species Group: **Mammal**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	Refine knowledge of maternity roost tree characteristics.
Research	Basic Life History	Medium	1) Determine the summer range of bats that use VT hibernacula. 2) Assess the degree of local recruitment to determine if are Vermont populations reproducing.
Research	Distribution and Abundance	Medium	Monitor changes in distribution and abundance in the years following White-nose Syndrome.
Research	Threats and Their Significance	Low	Determine what other factors besides habitat loss and White-nose Syndrome influence population trends.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	High	Learn more about the role of Vermont hibernacula in the region
Monitoring	Population Change	High	Monitor population trends in the years following White-nose Syndrome to determine if the species continues to decline.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	High	Investigate range shifts associated with summer habitat and roost tree loss in the Champlain Valley.
Monitoring	Monitor Threats	High	Monitor the continued population effects of White-nose Syndrome and cooperate on research about individual survivors.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Privately-Owned Protected Areas	High	Protect at least four USFWS Level I or II hibernacula in Vermont or New York	Number of hibernacula protected.	UVM, USFS, Cavers Organization, TNC, NY DEC	SWG, USFWS, NYDEC
Protected Area Management	Medium	Maintain and protect all maternity roost trees that support over 100 adults. Conserve summer foraging habitat that supports 2500 adults.	Number of roost trees identified and protected. Acres of foraging habitat conserved.	UVM, Coverts, Cavers, TNC, NY DEC,	SWG, USFWS, NYDEC
Research	Medium	Collect distribution and abundance data through the Northamerican Bat Monitoring Project (NABat) to contribute to range-wide trend information over time.	Number of NABat randomized grid cells surveyed each year.	USGS, USFWS refuges, USFS, National Parks, FPR	ENDG, PR, USFWS



Common Name: **Indiana Bat**
Scientific Name: **Myotis sodalis**
Species Group: **Mammal**

Bibliography

Beverly, J., J.D. Kiser, and V. Brack, Jr., PhD. 2002. A survey for eastern forest, with emphasis on the federally endangered Indiana bat (*Myotis sodalis*). Environmental Solutions and Innovations, LLC. 56pp.

Chenger, J. 2003. Bat Inventory for Project Lands of the Upper Connecticut River Basin. Prepared for the U.S. Army Corp of Engineers New England District by Bat Conservation and Management, Inc. 102 pp

Degraaf, R.M, and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover.

Godin, A. (1977). Wild mammals of New England. Baltimore: The John Hopkeins University Press.

Kiser, J.D., J. Beverly, and V. Brack, Jr., PhD. 2002. A survey of eastern forest bat communities in the Lake Champlanin Valley, with emphasis on the federally endangered Indiana bat (*Myotis sodalis*). Environmental solutions and Innovations, LLC. 70pp.

Kurta, A., and S.W. Murray. 2002. Philopatry and migration of banded Indiana bats (*Myotis sodalis*) and effects of radio transmitters. Journal of Mammalogy, 83:585-589.

Kurta, A., K.J. Williams, and R. Mies. 1996. Ecological, behavioral, and thermal observations of a peripheral population of Indiana bats (*Myotis sodalis*). Pp.102-117, in Bats and Forests symposium (R.M.R. Barclay and R.M. Brigham, eds.). Research Branch, British Columbia Ministry of Forests, Victoria British Columbia, CA Working Paper. 23:1-292.

Kurta, A.k D. King, J.A. Teramino, J.M. Stribley, and K.J. Williams. 1993. Sumer roosts of the endangered Indiana bat (*Myotis sodalis*) on the northern edge of its range. American Midland Naturalists, 129:132-138.

Sanders, Chris. 2004. Vermont Multi-Species Bat Conservation and Recovery Plan. 55pp.

Tumosa, J. 2003. United States Forest Service Species Data Collection Form. 32pp.

Turner, GG, DM Reader, and JTH Coleman. 2011. A five-year assessment of mortality and geographic spread of white-nose syndrome in North America and a look to the future. Bat Research News, 52:13-27.

Van Zyll de Jong, C.G. 1985. Handbook of Canadian Mammals. Volume 2. Bats. National Museums of Canada, Ottawa, Ontario, Canada. 212 pp.



Common Name: **Small-footed Bat**
 Scientific Name: **Myotis leibii**
 Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G1G3

Global Trend:

State Rank: S1

State Trend: unknown

Extirpated in VT? No

Regional SGCN? yes

Assessment Narrative:

The small-footed bat occurs throughout southeast Canada and the eastern United States, but is found in very low numbers. Regionally it seems to be at risk. In New England, this bat is listed as threatened in Vermont, endangered in New Hampshire, and a species of concern in Maine, Massachusetts, and Connecticut as well as in New York State. The small-footed bat has been identified as a Very High Concern Regional Species of Greatest Conservation Need. There is a general lack of information about this species. It is found in small numbers (i.e., 2-50 bats) in the major caves in Vermont, with one larger population of 110 individuals observed at Pike Hill Mine in 2013. This bat may be particularly susceptible to disturbance and is known to be associated with dams, exposed cliff faces, and talus during the summer. Three small-footed bats were caught in mist nets in 2003 at the Union Village Dam, North Hartland dam, and Townshend dam (a female, male, and female respectively) by a contractor for the US Army Corp of Engineers (Chenger 2003). Though infrequently captured in the summer and noted for their ability to detect and avoid mist nets, records exist from over a dozen towns around the state. Small-footed bats are susceptible to White-nose Syndrome, but have demonstrated relatively low population declines, as evidenced by regional hibernacula data (Turner et al. 2011). The USFWS was petitioned to evaluate this species for federal listing but concluded in 2013 that the listing was not warranted (USFWS 2013).

Distribution

The small-footed bat is widely, though sparsely, distributed throughout the state, as evidenced by mist net captures from around the state, and is likely associated with dams, exposed cliff faces, and talus. This species is documented at hibernacula in Brandon, Sudbury, Manchester, Stockbridge, Vershire, and Corinth.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

In both winter and summer the small-footed bat is closely associated with rocky habitat such as caves, cliffs, talus piles, quarry faces, and rock outcrops. It hibernates in very cold sites, often in the entrance areas of caves and mines sometimes using small cracks or piles of breakdown on cave and mine floors. Hibernacula surveys probably undercount the species. They may also hibernate in talus piles and cliffs that have deep crevices;



Common Name: Small-footed Bat
Scientific Name: *Myotis leibii*
Species Group: Mammal

however, the extent of this behavior in Vermont is unknown. No maternity sites have been found in Vermont, however, in other states they use barns and buildings, cliffs and bridges, but are primarily found under exfoliating tree bark (Sanders 2004). Changer (2004) documented small-footed bats using crevices in rocks and large rip-rap on a manmade dam face in New Hampshire. A radio-transmitted small-footed bat was found to use power line corridors (Kilpatrick, pers com). Areas that promote an abundance of insects are crucial to small-footed bat survival (Tomosa, 2003). Beaver ponds with abundant snags may provide roosting and foraging sites. Micro Habitat: outcrops

Habitat Types:

Cliffs and Talus
Northern Hardwood
Oak-Pine Northern Hardwood
Subterranean
Wet Shores
Building or Structure
Mine
Powerlines/RR/Roadsides
Aquatic: Fluvial
Other Cultural
Aquatic: Lacustrine

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Impacts of Roads or Transportation Systems
Climate Change
Incompatible Recreation

Description of habitat threat(s): These bats may be more susceptible to disturbance in the hibernacula. Disturbance of hibernating bats causes them to awaken and forces them to use additional limited energy reserves (Sanders, 2004). Arousal can use up enough fat to sustain a bat for 10-30 days (Thomas et al., 1990; Thomas, 1995; Martin et al, 1966). Changes in temperature and light, as well as, direct contact can cause a bat to awaken and deplete stored fat reserves. Alterations to cave mine openings can change the microclimate of a mine and affect bat survival. Loss of maternity roosts may also be a problem to survivability of young. Little is known about the summer habitat requirements of this bat but destruction/development of summer habitats are likely to negatively affect bats if potential roost sites and foraging areas are altered (Tumosa 2003) Maternity roosts may be present in rock outcroppings along roadsides and could therefore be susceptible to habitat disturbance or alteration during highway work. Warm winters and drought conditions are likely to increase bat body temperatures and corresponding metabolic demands which may influence survivability and reproduction.



Common Name: **Small-footed Bat**
 Scientific Name: **Myotis leibii**
 Species Group: **Mammal**

Non-Habitat Threats:

Genetics
 Disease
 Reproductive Traits
 Loss of Prey Base
 Loss of Metapopulation Structure
 Unknown Non-Habitat Threats

Description of non-habitat threat(s): Insecticides and pesticides have been implicated in the decline of several bat species (Belwood 1998 in Tumosa 2003). Environmental poisons have had negative impacts on, and increased the mortality rates of, bat populations. Bats store some lipophilic pesticides in brown adipose fat tissues. These stores are released as bats use their fat reserves during hibernation. Depending upon tissue levels of the pesticide, as well as the amount of fat used over a given time period, bats can be exposed to both chronic and acute poisoning which can result in death. At lower levels, chronic poisoning may raise a bat's metabolism, burning the limited fat resources more quickly and possibly causing them to starve to death. In addition, broad spectrum insecticides can deplete insect diversity and limit the food sources available for bats.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	1) Determine summer habitat utilization from known hibernacula in NY and Vershire in a telemetry study and 2) research the use of transportation corridor rock outcroppings
Research	Basic Life History	Low	
Research	Distribution and Abundance	High	1) estimate the statewide population by evaluating population densities in summer and winter habitat. 2) Document estimated populations of reproductive females.
Research	Threats and Their Significance	Medium	Research the impacts of transportation corridor maintenance activities on rock outcroppings used by this species.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	Low	
Monitoring	Population Change	Medium	1) Monitor distribution and abundance to determine critical summer and winter habitats as well as population status. 2) Develop a monitoring plan to document the number of reproductive females.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	Medium	Monitor changes in summer and winter habitat use in the aftermath of White-nose Syndrome and possible effects from population declines of other Myotis species.
Monitoring	Monitor Threats	Medium	Monitor changes in hibernating populations in sites that are gated to limit human entry versus open to visitation.



Common Name: **Small-footed Bat**
 Scientific Name: **Myotis leibii**
 Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Standards	High	Protect all VT hibernacula with 5 or more small-footed bats.	Number of hibernacula protected	TNC, VLT, Coverts	TNC, VLT, Forest Legacy, VHCB, USFWS
Research	Medium	Locate summer maternity roost sites and define roost characteristics.	Number of summer roost sites located.	GMNF, USFWS Refuges, VTRANS, Private quarry owners, CRAG Vermont	PR, USFWS

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Small-footed Bat**
Scientific Name: **Myotis leibii**
Species Group: **Mammal**

Bibliography

- Barbour, R.W. and W.H. Davis. 1969. Bats of America. The University of Kentucky Press, Lexington, Kentucky.
- Belwood, J.J. 1998. In Ohio's Backyard: Bats. Ohio Biological Survey Backyard Series No 1. 196pp in (Tumosa, J., J.Belwood, S. Lemieux, T. Gokee, J.Smith. 2003. U.S. Forest Service species data collection form for Myotis leibii, Eastern small-footed bat 35 pp.)
- Best, T.L. and J.B. Jennings. 1997. Myotis leibii. Mammalian species, 547:1-6.
- Chenger, J. 2003. Bat Inventory for Project Lands of the Upper Connecticut River Basin. Prepared for the U.S. Army Corp of Engineers New England District by Bat Conservation and Management, Inc. 102pp.
- Degraaf, R.M. and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover.
- Godin, A. (1977). Wild mammals of New England. Baltimore: The John Hopkeins University Press.
- Martin, R.L., J.T. Pawluk, and T. B. Clancy. 1966. Observations on hibernation of Myotis subulatus. Journal of Mammalogy, 47:348-349.
- NatureServe: On online encyclopedia of life [web application]. 2001. Version 1.3 Arlington, Virginia, U.S.A.: Association for Biodiversity Information. Available: <http://www.natureserve.org/>. In (Tumosa, J., J.Belwood, S. Lemieux, T. Gokee, J.Smith. 2003. U.S. Forest Service species data collection form for Myotis leibii, Eastern small-footed bat 35 pp.)
- Sanders, Chris. 2004. Vermont Multi-Species Bat Conservation and Recovery Plan. 55pp.
- Thomas, D.W. 1995. Hibernating bats are sensitive to nontactile human disturbance. Journal of Mammalogy. 76:940-946.
- Thomas, D.W. 1992. Status of the eastern small-footed bat (Myotis leibii) in Vermont. Final Report Vermont Fish and Wildlife , Nongame and Natural Heritage Program. 82pp.
- Thomas, D.W., M. Dorais, and J.M. Bergeron. 1990. Winter energy budgets and cost of arousals for hibernating little brown bats, Myotis lucifugus. Journal of Mammalogy 71:475-479.
- Trombulak, S.C., P. E. Higuera, and M. DesMueles. 2001. Population trends of wintering bats in Vermont. Northeastern Naturalist, 8:51-62.
- Tumosa, J., J.Belwood, S. Lemieux, T. Gokee, J.Smith. 2003. U.S. Forest Service species data collection form for Myotis leibii, Eastern small-footed bat 35 pp.
- Van Zyll de Jong, C.G. 1985. Handbook of Canadian Mammals. Volume 2. Bats. National Museums of Canada, Ottawa, Ontario, Canada. 212 pp.



Common Name: **Northern Long-eared Bat**
Scientific Name: **Myotis septentrionalis**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G2G3

Global Trend:

State Rank: S1

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The Northern Long-eared Bat has been identified as a Very High Concern Regional Species of Greatest Conservation Need and was listed as Federally Threatened on April 2, 2015. This species is extremely vulnerable to White-nose Syndrome, with state-wide surveys indicating declines of 93-100% from this disease (Darling and Smith 2011) and regional hibernacula declines of 98% (Turner et al. 2011). This species is in serious danger of extirpation in Vermont and extinction across its range as the disease continues to spread each winter (Frick et al. 2012 and USFWS 2013). Loss of maternity roosts could be a concern. Little information exists regarding summer roosting needs in VT, although neighboring NH has documented northern long-eared bats using a variety of tree species in close proximity, switching roosts frequently, and using trees that have a larger DBH than the average size in the stand (Sasse 1996). Recreational spelunking could also affect winter survivability. Information is needed on population trends and recruitment to determine if Vermont still has a reproductively viable population of this species.

Distribution

Distribution is statewide as recorded through captures, and hibernacula and acoustic surveys. However, this species has suffered drastic population declines due to White-nose Syndrome and their current distribution is not well known.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Northern Long-eared Bats hibernate in parts of caves and mines that are relatively cool and moist where the air is still. Hibernation may begin in August and may last for 8-9 months in northern latitudes. In the summer Northern Long-eared Bats roost by day in buildings and under tree bark, shutters, bat houses and bridges. At night they use caves to roost. They tend to be more solitary than other bats (Chenger 2004). They are gleaners and Northern long-eared bats forage in forested hillsides rather than in stream associated woodlands and consume a variety of night flying insects. They are well suited to forest interior habitats. Micro Habitat: roost sites



Common Name: Northern Long-eared Bat
Scientific Name: *Myotis septentrionalis*
Species Group: Mammal

Habitat Types:

Northern Hardwood
Hardwood Swamps
Softwood Swamps
Seeps and Pools
Oak-Pine Northern Hardwood
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Early Succession Upland Oak
Subterranean
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Wet Shores
Shrub Swamps
Early Succession Northern Hardwoods
Mine
Aquatic: Fluvial
Aquatic: Lacustrine
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Incompatible Recreation

Description of habitat threat(s): Disturbance of hibernating bats causes them to awaken and forces them to use additional limited energy reserves (Sanders, 2004). Arousal can use up enough fat to sustain a bat for 10-30 days (Thomas et al. 1990, Thomas 1995). Changes in temperature and light, as well as, direct contact can cause a bat to awaken and deplete stored fat reserves. Alterations to cave mine openings can change the microclimate of a mine and affect bat survival. Loss of maternity roosts may also be a problem to survivability of young. Felling of a maternity roost tree can impact the survival of both adults and young. Development/roads within close proximity of hibernacula, particularly along travel corridors could also be detrimental to survival. Destruction/development of summer habitats are likely to negatively affect bats if potential roost sites and foraging areas are altered (Tumosa 2003).

Non-Habitat Threats:

Genetics
Disease



Common Name: **Northern Long-eared Bat**
 Scientific Name: **Myotis septentrionalis**
 Species Group: **Mammal**

Description of non-habitat threat(s): The Northern Long-eared Bat has suffered population declines of 90-99% in Vermont and the northeast (Turner et al. 2014) and White Nose Syndrome continues to be a threat. Pesticides and environmental poisons have had negative impacts on, and increased the mortality rates of, bat populations. Bats store some lipophilic pesticides in brown adipose fat tissues. These stores are released as bats use their fat reserves during hibernation. Depending upon tissue levels of the pesticide, as well as the amount of fat used over a given time period, bats can be exposed to both chronic and acute poisoning which can result in death. At lower levels, chronic poisoning may raise a bat's metabolism, burning the limited fat resources more quickly and possibly causing them to starve to death. In addition, broad spectrum insecticides can deplete insect diversity and limit the food sources available for bats.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Determine summer habitat and roost tree characteristics.
Research	Basic Life History	High	Determine the spring migratory distance of bats emerging from hibernation and traveling to their summer range.
Research	Distribution and Abundance	High	Investigate the current distribution and abundance of this species in the years following White-nose Syndrome.
Research	Threats and Their Significance	High	Research what threats secondary to White-nose Syndrome are the most detrimental to the small remaining population.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	Low	
Monitoring	Population Change	High	Monitor population trends in the years following White-nose Syndrome.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	High	Investigate possible range shifts to prime summer or winter habitat in the years following White-nose Syndrome when population size and competition for habitat are extremely low.
Monitoring	Monitor Threats	High	Monitor the continued population effects of White-nose Syndrome and cooperate on research about individual survivors.

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Northern Long-eared Bat**
Scientific Name: **Myotis septentrionalis**
Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Privately-Owned Protected Areas	High	Protect all hibernacula containing northern long-eared bats in line with Federal threatened listing.	Number of hibernacula protected	UVM, Middlebury College, Vt. Cavers Assoc, VLT, TNC	SWG, TNC, USFWS
Creating Privately-Owned Protected Areas	High	Protect all roost trees documented as used by northern long-eared bats in line with Federal threatened listing.	Number of roost trees protected	UVM, Middlebury College, Vt., VLT, TNC, Woodland Owners Association	SWG, ENDG, WNS, TNC, USFWS

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Northern Long-eared Bat
Scientific Name: *Myotis septentrionalis*
Species Group: Mammal

Bibliography

- Barbour, R.W. and W.H. Davis. 1969. Bats of America. The University of Kentucky Press, Lexington, Kentucky.
- Beverly, J., J.D. Kiser, and V. Brack, Jr., PhD. 2002. A survey for eastern forest, with emphasis on the federally endangered Indiana bat (*Myotis sodalis*). Environmental Solutions and Innovations, LLC. 56pp.
- Chenger, J. 2003. Bat Inventory for Project Lands of the Upper Connecticut River Basin. Prepared for the U.S. Army Corp of Engineers New England District by Bat Conservation and Management, Inc. 102pp.
- Cryan, PM. 2003. Seasonal distribution of migratory tree bats (*Lasiurus* and *Lasionycteris*) in North America. Journal of Mammalogy 84:579-593
- Decker J. and C.W. Kilpatrick. 2002. Small mammals of the Guthrie-Bancroft farm, Year two, Colby Hill Ecological Project, Lincoln and Bristol, VT. 2001 Final Report 22pp.
- Decker J. and C.W. Kilpatrick. 2003. Small mammals of the Guthrie-Bancroft farm, Year three, Colby Hill Ecological Project, Lincoln and Bristol, VT. 2002 Final Report 22pp.
- Degraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover.
- Godin, A. (1977). Wild mammals of New England. Baltimore: The John Hopkins University Press
- Fitch, J. H., and K. A. Shump, jr. 1979. *Myotis keeni*. Mammalian Species, 121:1-3.
- Foster, R.W., and A. Kurth. 1999. Roosting Ecology of the northern bat (*Myotis septentrionalis*) and comparisons with the endangered Indiana bat (*Myotis sodalis*). Journal of Mammalogy, 80:659-672.
- Degraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover.
- Kilpatrick, C. W. 2001. Small mammal survey of the Nulhegan Basin Division of the Silvio O. Conte NFWR and the State of Vermont's West Mountain Wildlife Management Area, Essex County, Vermont. Final Report March 15, 2001 submitted to the Nature Conservancy.
- Kiser, J.D., R.R. Kiser, V. Brack, Jr., PhD., and E.R. Britzke. 2001. A survey for eastern forest bats on Green Mountain and Finger Lakes National Forests, with emphasis on the federally endangered Indiana bat (*Myotis sodalis*). Environmental Solutions and Innovations, LLC, 91pp.
- Kiser, J.D., J. Beverly, and V. Brack, Jr., PhD. 2002. A survey of eastern forest bat communities in the Lake Champlain Valley, with emphasis on the federally endangered Indiana bat (*Myotis sodalis*). Environmental solutions and Innovations, LLC. 70pp.
- Krusie, R.A., M. Yamasaki, C.D. Neefus, and P.J. Pekins. 1996. Bat habitat use in white Mountain National forest. Journal of Wildlife Management, 60:625-631.
- NatureServe: On online encyclopedia of life [web application]. 2001. Version 1.3 Arlington, Virginia, U.S.A.: Association for Biodiversity Information. Available: <http://www.natureserve.org/>. In (Tumosa, J., J. Belwood, S. Lemieux, T. Gokee, J. Smith. 2003. U.S. Forest Service species data collection form for *Myotis leibii*, Eastern small-footed bat 35 pp.)
- Owen, S.F., M. A. Menzel, W.M. Ford, B. R. Chapman, K.V. Miller, J. W. Edwards, and P.B. Wood. 2003. Home-range size and habitat used by northern *Myotis* (*Myotis septentrionalis*). American Midland Naturalists, 150:352-359.
- Reynolds, D.S. 2000. Woodland bat survey, Green Mountain National Forest, July 03-12, 2000. North east Ecological Services. 28pp.
- Reynolds, D.S. 2000. Woodland bat survey, Green Mountain National Forest, July 20-29, 2000. North east Ecological Services. 28pp.
- Sasse, D. B., and P. J. Pekins. 1996. Summer roosting ecology of northern long-eared bats (*Myotis septentrionalis*) in the White Mountain National Forest. Pp. 91-101, in Bats and Forest Symposium (R. M. R. Barclay and R. M. Brigham, eds.). British Columbia Ministry of Forest, Victoria, BC.

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: Northern Long-eared Bat
Scientific Name: *Myotis septentrionalis*
Species Group: Mammal

Sanders, Chris. 2004. Vermont Multi-Species Bat Conservation and Recovery Plan. 55pp.

Thomas, D.W. 1995. Hibernating bats are sensitive to nontactile human disturbance. *Journal of Mammalogy* 76:940-946.

Thomas, D.W., M.Dorais, and J.M. Bergeron. 1990. Winter energy budgets and cost of arousals for hibernating little brown bats, *Myotis lucifugus*. *Journal of Mammalogy*: 45:148-149.

Thomas, D.W. 1992. Status of the Eastern Small-footed bat (*Myotis leibii*) in Vermont. Final report for contract VOTHER. Nongame and Natural Heritage Program, Vermont fish and Wildlife Department, Waterbury, Vermont in (Sanders, Chris. 2004. Vermont Multi-Species Bat Conservation and Recovery Plan. 55pp.)

Toth, E. 1999. Woodland bat survey and core monitoring team training, August 3-11, 1999. Green Mountain National Forest. 23pp.

Tumosa, J. 2003. Green Mountain National Forest species data collection form. *Myotis septentrionalis*.

Turner, GG, DM Reader, and JTH Coleman. 2011. A five-year assessment of mortality and geographic spread of white-nose syndrome in North America and a look to the future. *Bat Research News*, 52:13-27.

Van Zyll de Jong, C.G. 1985. Handbook of Canadian Mammals. Volume 2. Bats. National Museums of Canada, Ottawa, Ontario, Canada. 212 pp.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Silver-haired Bat**
Scientific Name: **Lasionycteris noctivagans**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S2B

State Trend: Unknown

Extirpated in VT? no

Regional SGCN? Yes

Assessment Narrative:

Status of this bat is unknown but presumed to be much lower than historical levels. Many factors could be influencing the decline. Silver-haired bats migrate along the eastern seaboard in winter and could encounter factors that affect its survival. In some parts of the country it is associated with late successional forests with a snag density of more than 21 snags/ hectare. Loss of forest habitat throughout the 1800's probably contributed to the decline of this bat in New England. Other factors such as pesticides, availability of prey, and loss of maternity roosts could also be influencing the status of this bat. The silver-haired bat has been identified as a Very High Concern Regional Species of Greatest Conservation Need. Once the most common bat in the region in the 1800s it has experienced a significant decline throughout the Northeast. This species is currently considered rare and difficult to monitor. The silver-haired bat is documented as the shortest-lived (i.e., average 2 years, maximum 12 years), possibly indicating that this species is more sensitive to changes than other bat species. Silver-haired bats are the second most-commonly reported species found dead below turbines in Vermont's operating wind facilities (Sheffield, Lowell, and Georgia) during surveys conducted between April and October. Wind energy development is an increasing threat to this species, especially during fall migration (Leclair et al. 2009).

Distribution

One capture record exists for this species in Springfield and a maternity roost was found in a building in Chittenden. Data from mortality surveys below operating wind turbines in Sheffield, Lowell, and Georgia have provided new occurrence data for this migratory species, as well as acoustic data around the state from 2010 to 2014. Silver-haired bats are assumed to be widespread but very little is known about how abundant or how evenly distributed they are in Vermont. This species migrates south for the winter.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Silver-haired bats will range up to 5km from roost tree to forage areas. In summer, they roost under the bark of late-successional and old-growth boreal forests and perhaps along woodland edges. They forage in forest openings, including clear cuts, and over water and sometimes roost in buildings. In other parts of the country



Common Name: Silver-haired Bat
Scientific Name: *Lasionycteris noctivagans*
Species Group: Mammal

they are associated with late successional forests with snag densities of 21 snags/hectare. They form maternity colonies almost exclusively in tree cavities and will periodically switch roosts throughout the maternity season. Like big brown bats, the silver-haired bats feed on many insect pest species such as flies, midges, leafhoppers, moths, mosquitoes, beetles, crane flies, lacewings caddis flies, ants, crickets, and spiders.

Habitat Types:

Spruce Fir Northern Hardwood
Northern Hardwood
Oak-Pine Northern Hardwood
Open Peatlands
Marshes and Sedge Meadows
Early Succession Boreal Conifers
Early Succession Pine and Hemlock
Early Succession Upland Oak
Early Succession Other Types
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Wet Shores
Shrub Swamps
Early Succession Boreal Hardwoods
Early Succession Spruce-Fir
Early Succession Northern Hardwoods
Building or Structure
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lower CT River
Aquatic: Lacustrine
Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development
Habitat Alteration

Description of habitat threat(s): Conversion of forest habitat as a result of rural development that leads to loss of mature and older forests used as roosting habitat. Because silver-haired bats are migratory, they could be limited by wind and radio towers as well as powerlines. Wind energy development causes significant direct mortality to this species through collisions with turbine blades and barotrauma. Predators include several kinds of birds including blue jays therefore increased suburbanization could increase loss to



Common Name: **Silver-haired Bat**
Scientific Name: **Lasionycteris noctivagans**
Species Group: **Mammal**

predation.

Non-Habitat Threats:

Genetics

Disease

Description of non-habitat threat(s): Pesticides and environmental poisons have had negative impacts on, and increased the mortality rates of, bat populations. Bats store some lipophilic pesticides in brown adipose fat tissues. These stores are released as bats use their fat reserves during hibernation. Depending upon tissue levels of the pesticide, as well as the amount of fat used over a given time period, bats can be exposed to both chronic and acute poisoning which can result in death. At lower levels, chronic poisoning may raise a bat's metabolism, burning the limited fat resources more quickly and possibly causing them to starve to death. In addition, broad spectrum insecticides can deplete insect diversity and limit the food sources available for bats. Silver-haired bats are also susceptible to a virulent strain of rabies. This normally solitary species is more vulnerable to population impacts when concentrated in the spring and fall migration.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Determine habitat requirements in Vermont.
Research	Basic Life History	Low	Research the possibility of food competition and partitioning between red, hoary, silver-haired and eastern pipistrelle bats.
Research	Distribution and Abundance	High	Collect baseline data on distribution, abundance in Vermont.
Research	Threats and Their Significance	High	Collect mortality data from wind energy facilities.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	Low	Research the effects of pesticides on mortality and reproductive success.
Monitoring	Population Change	High	Monitor changes in abundance after the onset of operating wind turbines in the Northeast.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	Medium	Monitor changes in migratory patterns after habitat conversion of ridgelines and direct mortality due to wind development.
Monitoring	Monitor Threats	High	Research migratory patterns and impacts from power lines, wind towers, and road mortality

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Silver-haired Bat**
Scientific Name: **Lasionycteris noctivagans**
Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Policy & Regulations	High	Design, standardize, and implement mitigation guidelines, such as curtailment regimes, to decrease the threat of and direct take from wind energy development in Vermont.	Percentage of operating wind turbines that meet minimum mitigation guidelines.	Bat Wind Energy Cooperative, USFWS, USFS, Wind energy companies	Wind industry, USFWS, ENDG



Common Name: **Silver-haired Bat**
Scientific Name: **Lasionycteris noctivagans**
Species Group: **Mammal**

Bibliography

- Banfield, A.W.F. 1974. The mammals of Canada. University of Toronto Press, Toronto. 438 pp.
- Barbour, R.W. and W.H. Davis. 1969. Bats of America. The University of Kentucky Press, Lexington, Kentucky.
- Barclay, R.M.R. 1985. Ong- versus short-range foraging strategies of hoary (*Lasiurus cinereus*) and silver-haired (*Lasionycteris noctivagans*) bats and the consequences for prey selection. Canadian Journal of Zoology, 64: 2700-2705.
- Chenger, J. 2003. Bat Inventory for Project Lands of the Upper Connecticut River Basin. Prepared for the U.S. Army Corp of Engineers New England District by Bat Conservation and Management, Inc. 102pp.
- Cryan, P. 2000. Seasonal Distribution of Male and Female Hoary Bats *Lasiurus cinereus* in Continental North America. Bat Research News. 41(4): 114.
- Cryan, P.M. 2003. Seasonal distribution of migratory tree bats (*Lasiurus* and *Lasionycteris*) in North America. Journal of Mammalogy 84:579-593
- Godin, A. (1977). Wild mammals of New England. Baltimore: The John Hopkins University Press.
- Krusic, R.A. 1995. Habitat use and identification of bats in the White Mountain National Forest. M.S. Thesis, University of New Hampshire, Durham, New Hampshire. 86pp. (Cited in R.M. Degraaf and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover).
- Krusic, R.A. and C.D. Neefus. 1996. Habitat associations of bat species in the White Mountain National Forest. Pages 185-198 in R.M.R. Barclay and R.M. Brigham, editors. Bats and forest symposium, 19-21 October 1995, Victoria British Columbia, Canada. Research Branch, British Columbia Ministry of Forestry, Victoria British Columbia. Working paper 23. (Cited in R.M. Degraaf and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover).
- Kunz, T.H. 1982. *Lasionycteris noctivagans*. American Society of Mammalogists. Mammalian Species No. 172:1-5.
- Owen, S.F., M.A. Menzel, J.W. Edwards, W.M. Ford, J.M. Menzel, B.R. Chapman, P.B. Wood, and K.V. Miller. 2004. Bat activity in harvested and intact forest stands in the Allegheny Mountains. North. J. Appl. For. 21(3):154-159.
- Parsons, H.J., D.A. Smith, and R.F. Whittam. 1986. Maternity colonies of silver-haired bats, *Lasionycteris noctivagans*, in Ontario and Saskatchewan. Journal of Mammalogy 67: 598-600.
- Sanders, Chris. 2004. Vermont Multi-Species Bat Conservation and Recovery Plan. 55pp.
- Sasse, D.B. 1995. Summer roosting ecology of cavity-dwelling bats in the White Mountain National Forest. M.S. Thesis, University of New Hampshire, Durham, New Hampshire. 85pp. (Cited in R.M. Degraaf and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover).
- Thomas, D.W. 1988. The distribution of bats in different ages of Douglas fir forests. Journal of Wildlife Management, 52(4):619-626
- Van Zyll de Jong, C.G. 1985. Handbook of Canadian Mammals. Volume 2. Bats. National Museums of Canada, Ottawa, Ontario, Canada. 212 pp.
- Vonhoff, M.J. 1999. Patterns of tree use, group composition, and group stability in silver-haired bats: implications for forest management. Bat Research News 40 (4): 199



Common Name: **Tri-colored bat**
Scientific Name: **Perimyotis subflavus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G3

Global Trend:

State Rank: S1

State Trend: Declining

Extirpated in VT? No

Regional SGCN? Yes

Assessment Narrative:

The tri-colored bat is one of six species that overwinter in Vermont. This species occurs in small numbers in Vermont hibernacula and is only infrequently captured in mist net surveys. Its small size and multiple young (i.e., two, versus one for most bats) makes it more vulnerable. The tri-colored bat has been identified as a Very High Concern Regional Species of Greatest Conservation Need. This species was formerly known as the eastern pipistrelle, *Pipistrellus subflavus* (Menu 1984). Little is known about the tri-colored bat in Vermont, where this species is near the northern edge of its summer range (Whitaker 1998). It appears uncommon based on available survey data. Historic summer captures total less than 10 and many hibernacula surveys reveal 2-6 individuals roosting solitarily. By 2011, White-nose Syndrome had caused regional population declines for this species upwards of 75% (Turner et al. 2011) in hibernacula surveys. Tri-colored bats are extremely vulnerable to WNS. Fungal infection rates during late hibernation reach nearly 100% and fungal loads in this species are among the highest documented (Langwig et al. 2014). The tricolored bat was added to Vermont's endangered species list in 2012. The concentration of this species in caves and mines to hibernate makes them particularly vulnerable to human disturbance.

Distribution

Distribution is probably statewide but sparse. Capture records, hibernacula survey records, or acoustic recordings exist from most regions, though little is known about how evenly distributed they are in Vermont. Tri-colored bats have been documented in small numbers in the majority of known hibernacula. However, seven of the hibernacula with tri-colored bats documented before White-nose Syndrome (WNS) revealed none of this species in post-WNS surveys.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Probable
Champlain Hills	Probable	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The Tri-colored bat forages over wetlands, riparian areas, and forest edges, ingesting ants, moths, small beetles, mosquitoes and other insects. Possibly uses trees for maternity roosts, although in Vermont, the Tri-colored bat has not been found with other tree roosting bats. In Indiana they are found in sugar maple and American elms, as well as tulip and sycamore trees. Tri-colored bat is also found in the dead foliage of oaks.



Common Name: **Tri-colored bat**
Scientific Name: **Perimyotis subflavus**
Species Group: **Mammal**

They hibernate in caves mines and rock crevices where humidity is high and temperatures are around 10 to 15 degrees centigrade.

Habitat Types:

Northern Hardwood
Hardwood Swamps
Oak-Pine Northern Hardwood
Floodplain Forests
Early Succession Upland Oak
Subterranean
Wet Shores
Early Succession Boreal Hardwoods
Early Succession Northern Hardwoods
Mine
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lower CT River
Aquatic: Lacustrine
Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Incompatible Recreation

Description of habitat threat(s): Hibernating bats are limited by degradation, destruction and disturbance of hibernacula (caves and mines). Bats disturbed within the hibernacula use significant stores of fat each time they are awakened. If awakened enough times, bats can deplete their fat reserves and not have enough energy resources to complete spring migration, survive post emergence periods of bad weather or initiate and successfully complete gestation. In some cases, awakening hibernating bats can directly lead to their death. Closure of mines or caves in winter, when bats are present, would lead to the destruction of the entire colony. Slight alterations in cave/mine microclimate as a result of modifications to the opening etc. could also negatively impact hibernating bats. Removal of trees which serve as bat roosts, especially those serving as maternity roosts can directly kill entire colonies of bats. Wind energy turbines located on ridge tops have been found to directly kill bats as well..

Non-Habitat Threats:

Genetics
Disease



Common Name: **Tri-colored bat**
 Scientific Name: **Perimyotis subflavus**
 Species Group: **Mammal**

Loss of Metapopulation Structure

Description of non-habitat threat(s): Pesticides and environmental poisons have had negative impacts on, and increased the mortality rates of, bat populations. Bats store some lipophilic pesticides in brown adipose fat tissues. These stores are released as bats use their fat reserves during hibernation. Depending upon levels of the pesticide in the tissue, as well as, the amount of fat used over a given time period, bats can be exposed to both chronic and acute poisoning which can result in death. At lower levels, chronic poisoning may raise a bat's metabolism, burning the limited fat resources more quickly and possibly causing them to starve to death. In addition, broad spectrum insecticides can deplete insect diversity and limit the food sources available for bats.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Collect baseline data on habitat requirements.
Research	Basic Life History	Low	Research possibility of food competition and partitioning between red, hoary, silver-haired and eastern pipistrelle bats.
Research	Distribution and Abundance	High	Collect baseline data on distribution and abundance
Research	Threats and Their Significance	High	Research what threats secondary to White-nose Syndrome are the most detrimental to the small remaining population.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	Low	
Monitoring	Population Change	High	Monitor population trends in the years following White-nose Syndrome to determine if the species continues to decline.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	Medium	Investigate the characteristics of hibernacula still occupied by this species post-White-nose Syndrome versus abandoned sites.
Monitoring	Monitor Threats	High	Monitor the continued population effects of White-nose Syndrome and cooperate on research about individual survivors.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Creating Privately-Owned Protected Areas	High	Protect all hibernacula containing more than 5 tri-colored bats.	Number of hibernacula protected	UVM, USFS, Cavers Organization, TNC, Private landowners	ENDG, USFWS



Common Name: **Tri-colored bat**
Scientific Name: **Perimyotis subflavus**
Species Group: **Mammal**

Bibliography

- Banfield, A.W.F. 1974. The mammals of Canada. University of Canada. University of Toronto Press, Toronto. 438pp.
- Davis, W.H. , and R. E. Mumford. 1962. Ecological notes on the bat *Pipistrellus subflavus*. American midland Naturalists, 68:394-398.
- Fujita, M.S., and T.H. Kunz. 1984. *Pipistrellus subflavus*. Mammalian Species, 228:1-6.
- Godin, A. (1977). Wild mammals of New England. Baltimore: The John Hopkeins University Press.
- Saunders, D. Andrew. 1980. Adirondack Mammals. Adirondack Wildlife Program, State University of New York (College of Environmental Science and Forestry - Syracuse). 216 pp.
- Chenger, J. 2003. Bat Inventory for Project Lands of the Upper Connecticut River Basin. Prepared for the U.S. Army Corp of Engineers New England District by Bat Conservation and Management, Inc. 102 pp
- DeGraaf, Richard M. and Yamasaki, Mariko. New England Wildlife: Habitat, Natural History, and Distribution. University Press of New England, Hanover, NH, 2001.
- Turner, GG, DM Reader, and JTH Coleman. 2011. A five-year assessment of mortality and geographic spread of white-nose syndrome in North America and a look to the future. Bat Research News, 52:13-27.
- Veilleux, J.P., J.O. Whitaker, Jr., and S.L. Veilleux. 2003. Tree-roosting ecology of reproductive female eastern pipistrelles, *Pipistrellus subflavus*, in Indiana. Journal of Mammalogy, 84:1068-1075.



Common Name: **Big Brown Bat**
Scientific Name: **Eptesicus fuscus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S4

State Trend: Stable

Extirpated in VT? no

Regional SGCN? Yes

Assessment Narrative:

The Big Brown Bat has been identified as a High Concern Regional Species of Greatest Conservation Need. Because Big Brown Bat maternity sites are most often in human structures such as barns, sheds, houses, and churches, they are frequently in conflict with people. At best, they may be excluded from these structures, forcing a split into several smaller maternity sites. At worst, they may be exterminated by pest control agents or homeowners. This species is also vulnerable to the effects of White-nose Syndrome (WNS), though it has not seen the same drastic declines in Vermont or the northeast as some other hibernating species (Turner et al. 2011). Big Brown Bats are among the first bats to give birth and often have 2 offspring. Before WNS, little brown bat colonies were nearly twice as common as Big Brown Bat colonies in New England (D.S. Reynolds and T. H. Kunz, unpub. data, 1999). Because of the 90% declines of the little brown bat (another structure-dwelling species) experienced from WNS, the VFWD conducted extensive surveys of citizen-reported bat colonies in buildings between 2011 and 2014. By 2014 that ratio had reversed and Big Brown Bats made up 249 of the 283 structure-dwelling maternity colonies identified by the VFWD. Big Brown Bats are commonly captured during mist net surveys and are among the most commonly detected species in acoustic surveys done by neighboring states (Carl Herzog, NYDEC and Kate Moran, CTDEEP, personal communication). However, Big Brown Bats are difficult to survey during the winter because not only do they overwinter in caves and mines, but they also hibernate in structures, cliffs, and wood piles, where they are more difficult to detect. The actual population size of this species in VT is unknown but their long-term conservation is important as they may be one of the few species able to thrive after WNS has spread across the rest of North America and threatened the viability of many other hibernating species.

Distribution

Distribution is statewide as recorded through captures, citizen reports of bats in buildings, wind turbine mortality data, and acoustic survey data. During the winter, the Big Brown Bat is found in small numbers in about half the known bat hibernacula in Vermont, but also hibernates in structures around the state.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:



Common Name: **Big Brown Bat**
Scientific Name: **Eptesicus fuscus**
Species Group: **Mammal**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

In summer, Big Brown Bats roost in the attics of churches, houses, and old abandoned structures and deciduous tree cavities. In winter they hibernate in very cold areas (cave entrances and cliff faces) often with temperatures very close to and sometimes below freezing. This is the only bat species in VT known to hibernate in buildings. These low temperatures allow them to drastically slow their metabolism (Sanders 2004). Right now, Big Brown Bats hibernate in fewer than 20 sites in Vermont. Big Brown Bats consume beetles, ants, flies, mosquitoes, mayflies, stoneflies, and other insects. They emerge from their summer roost at dusk and fly a steady, nearly straight course to foraging areas (Chenger, 2004). There may be fidelity to the feeding grounds and some bats use the same grounds night after night. Little is known about where the majority of these bat winter, though reports of 1-6 bats hibernating in buildings are increasingly frequent as citizen reporting of bat activity has increased since White-nose Syndrome.

Habitat Types:

Cliffs and Talus
Hardwood Swamps
Softwood Swamps
Seeps and Pools
Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Early Succession Pine and Hemlock
Early Succession Upland Oak
Subterranean
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops
Wet Swales and Ditches
Wet Shores
Shrub Swamps
Building or Structure
Mine
Powerlines/RR/Roadsides
Aquatic: Fluvial
Aquatic: Lacustrine



Common Name: **Big Brown Bat**
Scientific Name: **Eptesicus fuscus**
Species Group: **Mammal**

Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Alteration

Description of habitat threat(s): Every year Big Brown Bats lose hundreds of possible building roosts due to exclusion or the actual destruction of buildings. Direct killing of bats is common due to human fears about rabies, bat bites and histoplasmosis. In addition, alterations or impacts to winter hibernacula also limits the future of this bat.

Non-Habitat Threats:

Genetics

Disease

Description of non-habitat threat(s): Big Brown Bats are one of 6 species in Vermont that are susceptible to White-nose Syndrome, though direct mortality for this species has been lower than for other susceptible species (Turner et al. 2011). The long-term and reproductive effects of this disease are unknown. Citizen reports of abandoned, dying, and dead young found below bat houses, and building roosts increased in VT and other Northeast states in 2012-2014 and is yet unexplained. Big Brown Bats are threatened by direct take when roosting in buildings during the summer and winter due to human fears of bats and rabies, as well as routine building maintenance (e.g., roof replacement), weatherization, and pest control activities. Pesticides and environmental poisons have had negative impacts on, and increased the mortality rates of, bat populations. Bats store some lipophilic pesticides in brown adipose fat tissues. These stores are released as bats use their fat reserves during hibernation. Depending upon levels of the pesticide in the tissue, as well as, the amount of fat used over a given time period, bats can be exposed to both chronic and acute poisoning which can result in death. At low levels, chronic poisoning may raise a bat's metabolism, burning the limited fat resources more quickly and possibly causing them to starve to death. In addition, broad spectrum insecticides can deplete insect diversity and limit the food sources available for bats.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Big Brown Bat**
 Scientific Name: **Eptesicus fuscus**
 Species Group: **Mammal**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Investigate the importance of human-made structures for hibernation.
Research	Basic Life History	Low	
Research	Distribution and Abundance	Medium	Document and map summer maternity colonies and human-made structures used for hibernation.
Research	Threats and Their Significance	Medium	Research long-term population effects of White-nose Syndrome
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	High	Track changes in abundance, distribution, and colony size over time by locating and monitoring summer maternity colonies
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	Medium	Investigate the increased use of colony locations (structures) or foraging areas previously dominated by little brown bats and other species in the genus Myotis that have declines drastically since White-nose Syndrome.
Monitoring	Monitor Threats	Medium	Monitor long-term population changes in the years following White-nose Syndrome

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Habitat Restoration		Protect hibernacula that are found to contain 30 or more big brown bats	Number of hibernacula protected	Vermont Cavers Association, TNC, VLT, USFS, ACE, Coverts	SWG, USFWS
Habitat Restoration		Maintain at least 50 maternity sites and a minimum of 5,000 adult female individuals in Vermont	Number of maternity sites protected	NRCS, Coverts, USFWS, ACE, VLT, TNC	USFWS, NRCS, EQIP,
Technical Assistance, Training, Learning Networks		Train nuisance wildlife control operators (NWCOs) in proper bat exclusion techniques. Work with homeowners/landowners to safely exclude bats and erect bat houses for displaced colonies.	Implement Best Management Practices and train nuisance wildlife control operators.	Wildlife Rehabilitators, NWCO's, Homeowners associations	USFWS, PR



Common Name: **Big Brown Bat**
Scientific Name: ***Eptesicus fuscus***
Species Group: **Mammal**

Bibliography

- Banfield, A.W.F. 1974. The mammals of Canada. University of Canada. University of Toronto Press, Toronto. 438pp.
- Godin, A. (1977). Wild mammals of New England. Baltimore: The John Hopkins University Press.
- Banfield, A.W.F. 1974. The mammals of Canada. University of Canada. University of Toronto Press, Toronto. 438pp.
- Sanders, Chris. 2004. Vermont Multi-species Bat Conservation and Recovery Plan. Prepared for the Vermont Bat Conservation and Recovery Plan Team. 55pp.
- Saunders, D. Andrew. 1980. Adirondack Mammals. Adirondack Wildlife Program, State University of New York (College of Environmental Science and Forestry - Syracuse). 216 pp.
- Chenger, JJ 2003. Bat Inventory for Project Lands of the Upper Connecticut River Basin. Prepared for the U.S. Army Corp of Engineers New England District by Bat Conservation and Management, Inc. 102 pp
- DeGraaf, Richard M. and Yamasaki, Mariko. New England Wildlife: Habitat, Natural History, and Distribution. University Press of New England, Hanover, NH, 2001.
- Kurta, A. & Baker, R.H. 1990. *Eptesicus fuscus*. Mammalian Species. 356:1-10



Common Name: **Eastern Red Bat**
 Scientific Name: **Lasiurus borealis**
 Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S4B

State Trend: unknown

Extirpated in VT? no

Regional SGCN? Yes

Assessment Narrative:

Eastern Red Bats spends the winter in the southern U.S. or Mexico. They migrate back and forth along the Eastern seaboard. A study in New York (Fisher 1896) reported red bats to be the second most common bat and reports from the late 1800's and early 1900's talk about "great flights of them during the whole day" (Mearns, 1898). This bat has a larger litter size than most other bats, ranging from one to five young. The Eastern Red Bat has been identified as a Very High Concern Regional Species of Greatest Conservation Need. Once one of the most abundant bats in many parts of their range, Eastern Red Bats appear to have declined dramatically over the last 100 years. Little was known about the Vermont population based on traditional survey methods. However, since the development of wind energy facilities in Vermont, data from fatality monitoring below operating turbines has added to population distribution information. Eastern Red Bats are the third most-commonly reported species found dead below turbines in Vermont's operating wind facilities (Sheffield, Lowell, and Georgia) during surveys conducted between April and October. Wind energy development is an increasing threat to this species, especially during fall migration. In addition, this species may be vulnerable to climate change as violent spring and autumn thunderstorms reportedly account for a large percentage of Eastern Red Bat deaths to migrating individuals and to females that are hesitant to separate from young during the birthing season (Leclaire et al. 2009).

Distribution

Capture records exist from around the state. Data from mortality surveys below operating wind turbines in Sheffield, Lowell, and Georgia have provided new occurrence data for this migratory species, as well as acoustic data from 2010 to 2014. The eastern red bat appears to be widespread throughout Vermont, though very little is known about how abundant or how evenly distributed they are in the state. This species migrates south for the winter.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Probable
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:



Common Name: **Eastern Red Bat**
Scientific Name: **Lasiurus borealis**
Species Group: **Mammal**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The eastern red bat is a solitary rooster which often hangs by one foot from branches in the foliage appearing as dead leaves in the crown of the tree. It prefers older forests with dense canopy foliage and open understory as well as hedgerows with elms and eastern red cedar stands. They are fast flyers that forage in open areas along hedgerows and field edges. Eastern red bats are also frequently observed foraging around lights in rural and suburban areas. The eastern red bats migrate south to Gulf states to hibernate. Tree bats such as the red, silver-haired, and hoary are the least studied of the bats and little is known about their status or habitat needs in Vermont. Eastern red bats feed on moths, crickets, flies, mosquitoes, beetles, cicadas, and other insects. Micro Habitat: red cedar

Habitat Types:

Hardwood Swamps
Seeps and Pools
Spruce Fir Northern Hardwood
Northern Hardwood
Oak-Pine Northern Hardwood
Early Succession Pine and Hemlock
Early Succession Upland Oak
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops
Shrub Swamps
Early Succession Northern Hardwoods
Powerlines/RR/Roadsides
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lower CT River
Aquatic: Lacustrine
Aquatic: Man-Made Water Bodies
Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development
Habitat Alteration



Common Name: **Eastern Red Bat**
Scientific Name: **Lasiurus borealis**
Species Group: **Mammal**

Climate Change

Description of habitat threat(s): Problems include conversion/degradation of forest habitat, as well as, rural development leading to loss of mature forest. Loss of American elms, a major roost tree, may be a continuing factor in the decline of the red bat. Because red bats are migratory, they could be limited by wind and radio towers as well as powerlines. Wind energy development causes significant direct mortality to this species through collisions with turbine blades and barotrauma. This species may be vulnerable to climate change as violent spring and autumn thunderstorms reportedly account for a large percentage of eastern red bat deaths to migrating individuals and to females that are hesitant to separate from young during the birthing season (LeCLaire et al. 2009). Predators include several kinds of birds including blue jays therefore increased suburbanization could increase loss to predation.

Non-Habitat Threats:

Genetics

Disease

Loss of Prey Base

Description of non-habitat threat(s): Pesticides and environmental poisons have had negative impacts on, and increased the mortality rates of, bat populations. Bats store some lipophilic pesticides in brown adipose fat tissues. These stores are released as bats use their fat reserves during hibernation. Depending upon tissue levels of the pesticide, as well as the amount of fat used over a given time period, bats can be exposed to both chronic and acute poisoning which can result in death. At lower levels, chronic poisoning may raise a bat's metabolism, burning the limited fat resources more quickly and possibly causing them to starve to death. In addition, broad spectrum insecticides can deplete insect diversity and limit the food sources available for bats. This normally solitary species is more vulnerable to population impacts when concentrated in the spring and fall migration.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Eastern Red Bat**
 Scientific Name: **Lasiurus borealis**
 Species Group: **Mammal**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Collect baseline data on habitat requirements
Research	Basic Life History	Low	Research possibility of food competition and partitioning between red, hoary, silver-haired and eastern pipistrelle bats.
Research	Distribution and Abundance	High	Collect baseline data on distribution and abundance in Vermont
Research	Threats and Their Significance	High	Collect mortality data from wind energy facilities.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	Low	Research the effects of pesticides on mortality and reproductive success.
Monitoring	Population Change	High	Monitor changes in abundance after the onset of operating wind turbines in the Northeast.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	Medium	Monitor changes in migratory patterns after habitat conversion of ridgelines and direct mortality due to wind development.
Monitoring	Monitor Threats	High	Research migratory patterns and impacts from power lines, wind towers, and road mortality

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Policy & Regulations	High	Design, standardize, and implement mitigation guidelines, such as curtailment regimes, to decrease the threat of and direct take from wind energy development in Vermont.	Percentage of operating wind turbines that meet minimum mitigation guidelines.	Bat Wind Energy Cooperative, USFWS, USFS, Wind energy companies	Wind industry, USFWS, ENDG



Common Name: Eastern Red Bat
Scientific Name: *Lasiurus borealis*
Species Group: Mammal

Bibliography

- Banfield, A.W.F. 1974. The mammals of Canada. University of Canada. University of Toronto Press, Toronto. 438pp.
- Barbour, R.W., and W.H. Davis. 1969. Bats of America. The University of Kentucky Press, Lexington, Kentucky.
- Chenger, J. 2003. Bat Inventory for Project Lands of the Upper Connecticut River Basin. Prepared for the U.S. Army Corp of Engineers New England District by Bat Conservation and Management, Inc. 102 pp
- Cryan, P.M. 2003. Seasonal distribution of migratory tree bats (*Lasiurus* and *Lasionycteris*) in North America. *Journal of Mammalogy* 84: 579-593.
- DeGraaf, Richard M. and Yamasaki, Mariko. New England Wildlife: Habitat, Natural History, and Distribution. University Press of New England, Hanover, NH, 2001.
- Godin, A. (1977). Wild mammals of New England. Baltimore: The John Hopkins University Press.
- Hart, J.A., G.L. Kirkland, Jr. and S.C. Grossman. 1993. Relative abundance and habitat use by tree bats, *Lasiurus* spp., in southcentral Pennsylvania. *Canadian field-Naturalist* 107: 208-212.
- Hickey, M.B.C. and M.B. Fenton. 1990. Foraging by red bats (*Lasiurus borealis*): do intraspecific chases mean territoriality? *Canadian Field Journal of Zoology* 68: 2477-2482.
- Krusic, R.A. and C.D. Neefus. 1996. Habitat associations of bat species in the White Mountain National Forest. Pages 185-198 in R.M.R. Barclay and R.M. Brigham, editors. Bats and forests symposium, 19-21 October 1995, Victoria British Columbia, Canada. Research Branch, British Columbia Ministry of Forestry, Victoria British Columbia. Working paper 23. (Cited in R.M. Degraaf and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover.)
- Laval, R.K. and M.L. Laval. 1979. Notes on reproductive, behavior, and abundance of the red bat, *Lasiurus borealis*. *Journal of Mammalogy* 60(1): 209-212.
- Owen, S.F., M.A. Menzel, J.W. Edwards, W.M. Ford, J.M. Menzel, B.R. Chapman, P.B. Wood, and K.V. Miller. 2004. Bat activity in harvested and intact forest stands in the Allegheny Mountains. *North. J. Appl. For.* 21(3): 154-159.
- Sasse, D.B. 1995. Summer roosting ecology of cavity-dwelling bats in the White Mountain National Forest. M.S. Thesis, University of New Hampshire, Durham, New Hampshire. 85pp. (Cited in R.M. Degraaf and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover.)
- van Zyll de Jong, C.G. 1985. Handbook of Canadian Mammals. Volume 2. Bats. National Museums of Canada, Ottawa, Ontario, Canada. 212pp.
- Saunders, D. Andrew. 1980. Adirondack Mammals. Adirondack Wildlife Program, State University of New York (College of Environmental Science and Forestry - Syracuse). 216 pp.
- Sanders, Chris. 2004. Vermont Multi-Species Bat Conservation and Recovery Plan. 55pp.
- Shump, K. A., Jr., and A. U. Shump. 1982. *Lasiurus borealis*. *Mammalian Species*, 183:1-6.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Hoary Bat**
Scientific Name: **Lasiurus cinereus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S3B

State Trend: unknown

Extirpated in VT? no

Regional SGCN? Yes

Assessment Narrative:

Hoary Bats are the largest bats of northeastern North America. The range-wide population has declined significantly since 1900. Historically, few records existed for this species in Vermont, due largely to the difficulty in capturing this fast, high-flying species in nets. Due to their solitary nature, we know the least about the three tree bat species in Vermont (red, hoary, and silver-haired). However, capture records, combined with more recent acoustic survey and wind mortality data indicate that this species is wide-spread throughout Vermont. The Hoary Bat has been identified as a Very High Concern Regional Species of Greatest Conservation Need. Add statements about wind mortality. Hoary Bats are the most commonly reported species found dead below turbines in Vermont's operating wind facilities (Sheffield, Lowell, and Georgia) during surveys conducted between April and October. Wind energy development is an increasing threat to this species, especially during fall migration (Leclaire et al. 2009).

Distribution

Capture records from Thetford, Springfield, Orwell, Brandon, Salisbury, and historic record from Rutland. Data from mortality surveys below operating wind turbines in Sheffield, Lowell, and Georgia have provided new occurrence data for this migratory species, as well as acoustic data around the state from 2010 to 2014. The Hoary Bat appears to be widespread throughout Vermont, though very little is known about how abundant or how evenly distributed they are in the state. This species migrates south for the winter.

Distribution by Biophysical Region:

Champlain Valley	Historic Record(s) Only	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

In the summer, during the day, Hoary Bats may stay concealed in the foliage of trees, well-concealed but with an open understory, generally 10 to 17 feet above the ground and often on the edge of a clearing. They emerge after dark to feed and may make round trips of up to 24 miles to forage. They forage over wetlands, openings, lakes and edges. They are fast flyers. Northern populations make long seasonal migrations to and from warmer winter habitats in the southern United States or Mexico. The sexes are segregated throughout most of the summer range. Foods include moths, true bugs, mosquitoes, and other insects. Hoary Bats have two young in mid-May through June or July. Females are solitary roosters and roost exclusively in trees. They may roost in



Common Name: **Hoary Bat**
Scientific Name: **Lasiurus cinereus**
Species Group: **Mammal**

the same tree in subsequent years.

Habitat Types:

Hardwood Swamps
Softwood Swamps
Seeps and Pools
Spruce Fir Northern Hardwood
Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests
Early Succession Boreal Conifers
Early Succession Pine and Hemlock
Early Succession Upland Oak
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Early Succession Boreal Hardwoods
Early Succession Spruce-Fir
Early Succession Northern Hardwoods
Powerlines/RR/Roadsides
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lower CT River
Aquatic: Lacustrine
Aquatic: Man-Made Water Bodies
Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development
Habitat Alteration

Description of habitat threat(s): Problems include conversion/degradation of forest habitat, as well as rural development leading to loss of mature forest. Because Hoary Bats are migratory, they could be impacted by wind and radio towers as well as powerlines. Wind energy development causes significant direct mortality to this species through collisions with turbine blades and barotrauma. Predators include several kinds of birds including blue jays therefore increased suburbanization could increase loss to predation

Non-Habitat Threats:



Common Name: **Hoary Bat**
 Scientific Name: **Lasiurus cinereus**
 Species Group: **Mammal**

Genetics

Disease

Description of non-habitat threat(s): Pesticides and environmental poisons have had negative impacts on, and increased the mortality rates of, bat populations. Bats store some lipophilic pesticides in brown adipose fat tissues. These stores are released as bats use their fat reserves during hibernation. Depending upon tissue levels of the pesticide, as well as the amount of fat used over a given time period, bats can be exposed to both chronic and acute poisoning which can result in death. At lower levels, chronic poisoning may raise a bat's metabolism, burning the limited fat resources more quickly and possibly causing them to starve to death. In addition, broad spectrum insecticides can deplete insect diversity and limit the food sources available for bats. This normally solitary species is more vulnerable to population impacts when concentrated in the spring and fall migration.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Collect baseline data on habitat requirements.
Research	Basic Life History	Low	Research the possibility of food competition and partitioning between red, hoary, silver-haired and eastern pipistrelle bats.
Research	Distribution and Abundance	High	Collect baseline data on distribution and abundance in Vermont
Research	Threats and Their Significance	High	Collect mortality data from wind energy facilities.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Research	Other Research	Low	Research the effects of pesticides on mortality and reproductive success.
Monitoring	Population Change	High	Monitor changes in abundance after the onset of operating wind turbines in the Northeast.
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	Medium	Monitor changes in migratory patterns after habitat conversion of ridgelines and direct mortality due to wind development.
Monitoring	Monitor Threats	High	Research migratory patterns and impacts from power lines, wind towers, and road mortality



Common Name: **Hoary Bat**
 Scientific Name: **Lasiurus cinereus**
 Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Policy & Regulations	High	Design, standardize, and implement mitigation guidelines, such as curtailment regimes, to decrease the threat of and direct take from wind energy development in Vermont.	Percentage of operating wind turbines that meet minimum mitigation guidelines.	Bat Wind Energy Cooperative, USFWS, USFS, Wind energy companies	Wind industry, USFWS, ENDG

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Hoary Bat**
Scientific Name: **Lasiurus cinereus**
Species Group: **Mammal**

Bibliography

- Banfield, A.W.F. 1974. The Mammals of Canada. University of Toronto Press, Toronto. 438pp.
- Barbour, R.W. and W.H. Davis. 1969. Bats of America. The University of Kentucky Press, Lexington, Kentucky.
- Barclay, R.M.R. 1985. Ong- versus short-range foraging strategies of hoary (*Lasiurus cinereus*) and silver-haired (*Lasionycteris noctivagans*) bats and the consequences for prey selection. *Canadian Journal of Zoology*, 64: 2700-2705.
- Chenger, J. 2003. Bat Inventory for Project Lands of the Upper Connecticut River Basin. Prepared for the U.S. Army Corp of Engineers New England District by Bat Conservation and Management, Inc. 102pp.
- Cryan, P. 2000. Seasonal Distribution of Male and Female Hoary Bats *Lasiurus cinereus* in Continental North America. *Bat Research News*. 41(4): 114.
- Decker J. and C.W. Kilpatrick. 2002. Small mammals of the Guthrie-Bancroft farm—Year three, Colby Hill Ecological Project, Lincoln and Bristol, VT. Final Report 22pp.
- Findley, J.S. and C. Jones. 1964. Seasonal distribution of the hoary bat. *Journal of Mammalogy*. 45: 461-470.
- Furlonger, C.L., H.J. Dewar, and M.B. Fenton. 1987. Habitat use by foraging insectivorous bats. *Canadian Journal of Zoology* 65:284-288.
- Godin, A. (1977). Wild mammals of New England. Baltimore: The John Hopkins University Press.
- Hart, J.A., G.L. Kirkland, Jr. and S.C. Grossman. 1993. Relative abundance and habitat use by tree bats, *Lasiurus* spp., in southcentral Pennsylvania. *Canadian Field Naturalist* 107: 208-212.
- Krusic, R.A. 1995. Habitat use and identification of bats in the White Mountain National Forest. M.S. Thesis, University of New Hampshire, Durham, New Hampshire. 86pp. (Cited in R.M. Degraaf and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover).
- Krusic, R.A. and C.D. Neefus. 1996. Habitat associations of bat species in the White Mountain National Forest. Pages 185-198 in R.M.R. Barclay and R.M. Brigham, editors. Bats and forest symposium, 19-21 October 1995, Victoria British Columbia, Canada. Research Branch, British Columbia Ministry of Forestry, Victoria British Columbia. Working paper 23. (Cited in R.M. Degraaf and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover).
- Owen, S.F., M.A. Menzel, J.W. Edwards, W.M. Ford, J.M. Menzel, B.R. Chapman, P.B. Wood, and K.V. Miller. 2004. Bat activity in harvested and intact forest stands in the Allegheny Mountains. *North. J. Appl. For.* 21(3):154-159.
- Rolseth, S.L., C.E. Koehler, and R.M.R. Barclay. 1994. Differences in the diets of juvenile and adult hoary bats, *Lasiurus cinereus*. *Journal of Mammalogy* 75 (2): 394-398. [Manitoba]
- Sanders, Chris. 2004. Vermont Multi-Species Bat Conservation and Recovery Plan. 55pp.
- Sasse, D.B. 1995. Summer roosting ecology of cavity-dwelling bats in the White Mountain National Forest. M.S. Thesis, University of New Hampshire, Durham, New Hampshire. 85pp. (Cited in R.M. Degraaf and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover).
- Shump, K.A. and A.U. Shump. 1982. *Lasiurus cinereus*. American Society of Mammalogists. Mammalian Species No. 185: 1-5.
- Van Zyll de Jong, C.G. 1985. Handbook of Canadian Mammals. Volume 2. Bats. National Museums of Canada, Ottawa, Ontario, Canada. 212 pp.

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **New England Cottontail**
Scientific Name: **Sylvilagus transitionalis**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: SU

State Trend: Unknown

Extirpated in VT? no

Regional SGCN? yes

Assessment Narrative:

The New England cottontail is rare, possibly extirpated in Vermont. The New England Cottontail is the only rabbit native to the northeastern United States east of the Hudson River Valley of New York including New England. It's range has contracted by an estimated 86% since 1960. Outside of Vermont, only five smaller populations occupy its historic New England range. The cottontail is recognized as a SGCN in the Wildlife Action Plans of all New England States and New York. In 2006 it was designated a candidate for listing under the federal Endangered Species Act

The New England Cottontail is listed as a Regional Species of Greatest Conservation Need among the 13 Northeastern states. A regional effort has been mounted to restore the New England Cottontail (<http://www.newenglandcottontail.org/>).

The New England cottontail was abundant in Vermont prior to the 1940s, however, the species was last documented in the state in 1946. Widespread introductions of the eastern cottontail (*Sylvilagus floridanus*) and habitat changes have resulted in apparent competition and possibly hybridization with eastern cottontails. Despite concerted trapping efforts in Vermont, no evidence of New England cottontails has been found since 1991.

Distribution

Distribution by Biophysical Region:

Champlain Valley	Historic Records Only	Southern VT Piedmont	Historic Records Only
Champlain Hills	Not Probable	Vermont Valley	Historic Records Only
Northern Green Mtns	Not Probable	Southern Green Mtns	Historic Records Only
Northern VT Piedmont	Historic Records Only	Taconic Mtns	Historic Records Only
Northeastern Highlands	Not Probable		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

New England cottontails are associated with many types of vegetation but are most often found in early successional old fields, 10-25 years post-disturbance with high stem density (9000-10,000 stems/hectare). It is critical that patches of dense hardwood and softwood shrubs, seedlings and saplings at least .5 meters tall and less than 7.5 meters in diameter be closely spaced to facilitate usage. Connectivity between patches is also important. Isolated patches are much less frequently used (Tumosa 2001). New England cottontails seldom



Common Name: New England Cottontail
Scientific Name: *Sylvilagus transitionalis*
Species Group: Mammal

venture far from dense cover and in winter will inhabit larger patches (greater than 10 ha) (DeGraff and Yamasaki, 2001). They cannot colonize areas already inhabited by Eastern cottontail. Home ranges can be linear along riparian areas, roadsides etc.

Habitat Types:

Early Succession Pine and Hemlock
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Early Succession Boreal Hardwoods

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Habitat Fragmentation
Invasion by Exotic Species

Description of habitat threat(s): Fragmentation and isolation of patches results in lower survival rates and skewed sex ratios and increases vulnerability to extirpation due to chance events (natureserve.org). Habitat patches less than 3 acres in size increases the risk of predation. Decline in patch size (less than 15-75 ha) and increase in juxtaposition (greater than 500m) reduces survivability of New England cottontails. Loss of 10-25 year post-disturbance habitat due to conversion, succession and fragmentation also negatively influences New England cottontail recovery. Competition from eastern cottontail is also a problem. The eastern cottontail will occupy a habitat first and exclude NE Cottontail.

Non-Habitat Threats:

Competition

Description of non-habitat threat(s): Competition with eastern cottontail is widely recognized as a limiting factor for New England cottontail populations.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Determine habitat requirements in Vermont.
Research	Distribution and Abundance	Medium	Continue to monitor for occurrence in likely Vermont habitats.
Research	Taxonomy	Medium	Genetically test trapped rabbits to determine distribution of floridanus vs. transitionalis
Monitoring	Monitor Threats	Medium	Monitor changes in early successional habitats in regards to size, age, and juxtaposition

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **New England Cottontail**
 Scientific Name: ***Sylvilagus transitionalis***
 Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Species Restoration	High	Support the implementation of the Conservation Strategy for the New England Cottontail (Fuller and Tur 2012)		Other New England states, VLT, TNC, USFWS	SWG, PR
Ex-Situ Conservation		Identify regional refugia until habitat can be developed w/in a state. Maintain isolated populations until a long-term plan is developed.	Number of isolated populations conserved. Number of regional refugia conserved.	Other New England states, VLT, TNC, USFWS	SWG, PR

Bibliography

Chapman, Joseph A., Kenneth L. Cramer, Nico J. Dppenaar, and Terence J. Robinson. 1992. Systematics and biogeography of the New England cottontail. SYLVILAGUS TRANSITIONALIS (Bangs, 1895), with the description of a new species from the Appalachian Mountains. Proc. Biol. Soc. Wash. 105 (4): 841-865.

Feldhamer, George A., Bruce C. Thompson, and Joseph A. Chapman: 2003: Wild Mammals of North America, 2nd Edition. The Johns Hopkins University Press, Baltimore and London.

Fuller, S. and A. Tur. 2012. Conservation Strategy for the New England Cottontail (*Sylvilagus transitionalis*) http://www.newenglandcottontail.org/sites/default/files/research_documents/conservation_strategy_final_12-3-12.pdf

Kirk, George L. 1916. The mammals of Vermont, Joint Bulletin No. 2 Vermont botanical and Bird Club 2: 28-34.

Litvaitis, John A. 1993 Status of the New england cottontail in the Lake Champlain drainage of Vermont. Nongame and Natural Heritage program. Vermont Department of Fish and Wildlife, Waterbury, VT.

Litvaitis, John A., and B. Johnson, W. Jakubus, and K. Morris. 2003a. Distribution and Habitat features associated with remnant populations of New England cottontails in Maine. Can. J. Zool. L81: 877-887.

Litvaitis, John A., and Michael N. Marchand, Jeffery P. Tash, Mathew Oberkrieser, Vanessa Johnson, and Marian K Litvaitis. 2003b. Intrim progress report II: A regional inventory of New Engalnd cottontails. Department of Natural Resources and Zoology Department, University of New Hampshire.

Litvaitis, J.A., J P. Tash, M.K. Litvaitis, M. N.Marchand,A. I. Kovach, R. Innes.2006. A range-wide survey to determine the current distribution of New England cottontails. Wildlife Society Bulletin. jh34(4): 1190-1197.

Osgood, Frederick L. Jr., 1938. The mammals of Vermont . J.Mammalogy 19(4): 435-441.

Probert, Brenda L. and John A. Litvaitis. 1996. Behavioral interactions between invading and endemic lagomorphs: Implications for conserving a declining species. Biol. Conserv. 76: 289-295.

Smith, Donald F. and John A. Litvaitis. 2000. Foraging strategies of sympatirc lagomorphs: Implications for differential success in fragmented landscapes. Can.J. Zool. 78 (12): 2134-2141.

Tumosa, Judy. 2001. United States Forest Service species data collection form for *Sylvilagus transitionalis*. 19pp.

Wilson, Don E. and Sue Ruff. 1999. The Smithsonian Book of North American Mammals. Smithsonian Institution Press, Washington and London.

Whitaker, John O. and W.J. Hamilton, Jr. 1998. Mammals of the Eastern United States. Comstock Publ. Co., Ithaca, N.Y.



Common Name: **Snowshoe Hare**
 Scientific Name: **Lepus americanus**
 Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5

State Trend: Declining

Extirpated in VT? No

Regional SGCN? no

Assessment Narrative:

The Snowshoe Hare experiences cyclical population changes on a 10-year period mostly in the northern parts of its range. Populations near the southern limits of its range, including Vermont, are believed to be less cyclical. Early successional softwood and mixed softwood- hardwood patches are critical habitats. Dense softwood and hardwood understory cover is highly important as it provides feeding, escape, and thermal cover for hares (Carreker 1985, Litvaitis et al. 1985). Forest succession and an overall decrease in active forest management practices in recent decades (Morin et al. 2014) has led to a reduction in suitable habitat and a decline in the state and regional Snowshoe Hare populations. Furthermore, changes in the climate that produce anomalously warm temperatures and decreased snowfall may diminish the hares' competitive advantages leading to higher predation rates and chronic declines in hare abundance (Schmitz et al. 2003). Consequently, lower hare populations may affect other wildlife species that rely on abundant hare populations as a source of prey (Chapman and Feldhamer 1982). The Snowshoe Hare is a keystone species in the northern transitional and boreal forest. If it should disappear, many species of predators would go with it and the structure of the plant community would be altered substantially (Krebs et al. 2001).

Distribution

The Snowshoe Hare was formerly found throughout Vermont with highest populations found in the mountains and lowland swamps (Foote 1946). Clearing of the land for agriculture and the introduction of the eastern cottontail have reduced hare habitat and populations especially in the Champlain and Connecticut River Valleys where hares currently exist only in larger forested blocks away from agriculture and development. The species is currently more commonly found in the Green Mountains, Taconic Mountains, and the northeastern part of the state. Hare populations generally increase from south to north in the state.

Distribution by Biophysical Region:

Champlain Valley	Probable	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Prefers large expanses of forest habitat, with low brushy cover and needs diverse forest size/age classes for feeding and cover.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Snowshoe Hare**
Scientific Name: **Lepus americanus**
Species Group: **Mammal**

Habitat Types:

Hardwood Swamps
Softwood Swamps
Spruce Fir Northern Hardwood
Early Succession Boreal Conifers
Early Succession Pine and Hemlock
Shrub Swamps
Early Succession Spruce-Fir
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Habitat Succession
Climate Change

Description of habitat threat(s): The natural succession of forests, particularly with respect to an observed decrease in active forest management across the region, is believed to be the leading cause of snowshoe hare population declines (DeGraaf & Yamasaki 2001). The availability of suitable cover and sufficient quantities of preferred browse plays an important role in hare productivity and survival (DeGraaf & Yamasaki 2001). Although the potential effects of climate change on this species are poorly understood, it is widely speculated that a warming climate could impact the species' ability to persist due to decreased snowfall and commensurate shifts in predator communities.

Description of non-habitat threat(s):

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	High	Collect Snowshoe Hare baseline data on the distribution and abundance in Vermont.
Research	Threats and Their Significance	Medium	Evaluate current forest management trends in Vermont and assess implications for Snowshoe Hare.
Monitoring	Population Change	Medium	Develop and implement a hare monitoring protocol in the state for evaluating population trends over time
Monitoring	Habitat Change	High	Periodically perform quantitative assessments of hare habitat in Vermont in order to detect trends and evaluate effectiveness of conservation strategies.
Monitoring	Range Shifts	Low	Develop and implement a protocol for monitoring range shifts in carnivores as a result of a changing climate.
Monitoring	Monitor Threats	Low	

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Snowshoe Hare**
 Scientific Name: **Lepus americanus**
 Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Policy & Regulations	High	Support and cooperate with regional efforts to curb the effects of climate change via the development and implementation of appropriate policy and regulations.			
Publically-Owned Protected Areas	High	Use even age management methods to increase young softwood and young mixed softwood/hardwood forests on state and federal lands.	Number of acres of early successional habitat in VT Forest Inventory Analysis (USFS).	ANR, USFS, USFWS, Coverts, RGS, WMI	PR,SWG
Conservation Payments/Financial Incentives	High	Encourage private landowners to use even age management methods to increase young softwood and young mixed softwood/hardwood forests through incentive programs (e.g., Current Use USDA Wildlife Habitat programs).	Number of acres of early successional habitat in VT Forest Inventory Analysis (USFS).	ANR, USFS, USFWS, Coverts, RGS, WMI, private landowners	PR,SWG

Bibliography

- Carreker, R.G. 1985. Habitat suitability index models: snowshoe hare. USFWS Biol. Rep. 82 (10:101). 21pp.
- Chapman, J.A., and G.A. Feldhamer. 1982. Snowshoe Hare in: Wild Mammals of North America. Johns Hopkins University Press, Baltimore, MD. 1147pp.
- DeGraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover, NH. 482 pp.
- Foote, L.E. 1946. A history of wild game in Vermont. Federal Aid in Wildlife Restoration Project No. 1-R. 51pp.
- Krebs CJ, Boonstra R, Boutin S, Sinclair ARE. 2001. What drives the 10-year cycle of snowshoe hares? BioScience 51: 25–35.
- Litvaitis, J.A., J.A. Sherburne, and J.A. Bissonette. 1985. Influence of understory characteristics on snowshoe hare habitat use and density. J. Wildl. Manage.: 49(4):866-873.
- Morin, Randall S.; Pugh, Scott A. 2014. Forests of Vermont, 2013. Resource Update FS-30. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 4 p.
- Schmitz, Oswald J., et al. "Ecosystem responses to global climate change: moving beyond color mapping." BioScience 53.12 (2003): 1199-1205

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Southern Flying Squirrel**
Scientific Name: **Glaucomys volans**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S4

State Trend: stable

Extirpated in VT? no

Regional SGCN? no

Assessment Narrative:

Species is listed as S4 and is apparently secure but little is known about this species or population trends in the state. Generally less is known about flying squirrels than other squirrels because of their nocturnal habits. Southern flying squirrels are expanding their range northward and have recently been documented from the Northeastern Highlands (Kilpatrick and Benoit 2011). Although the number of sites where the northern and southern flying squirrels occur in sympatry in the state are limited (Kilpatrick and Benoit 2011), increase competition for nest sites (tree cavities) may occur. While it is expected that the southern flying squirrel will dominate in these situations (Wells-Gosling and Heaney 1984), limited empirical data from field studies are available.

Distribution

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Probable
Northern Green Mtns	Probable	Southern Green Mtns	Confident
Northern VT Piedmont	Probable	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

In the eastern United States the southern flying squirrel is usually found at lower elevations in deciduous forests (Dolan and Carter 1977). In the northern areas of its range it also inhabits mixed woodlands of hardwoods and conifers, particularly where hardwoods predominate (Dolan and Carter 1977). Individuals and family groups require several nests; a primary nest (usually a tree cavity) that is used more or less continuously and several secondary nests (often stick nest) that serve as sheltered stations for feeding and defecating (Muul 1968). The availability of nest sites may be a limiting factor of population size (Muul 1968).

Habitat Types:

Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests



Common Name: **Southern Flying Squirrel**
 Scientific Name: **Glaucomys volans**
 Species Group: **Mammal**

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Alteration

Climate Change

Description of habitat threat(s): Although little is known about the potential habitat related threats to this species, it is believed southern flying squirrels are vulnerable to the degradation of preferred habitats resulting from climate change, forest pests, and forestry practices. Habitat threats of this nature are of particular concern with respect to the availability of sufficient quantities of large diameter deciduous trees with cavities suitable for use as nesting sites.

Non-Habitat Threats:

Trampling or Direct Impacts

Description of non-habitat threat(s): Southern flying squirrels are known to occupy residential structures with some frequency. Pest control professionals often respond to complaints of nuisance squirrel behavior via lethal control measures.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	High	Determine distribution and abundance by conducting targeted surveys and through collaboration with pest control professionals
Monitoring	Population Change	Medium	Monitor for changes in population by periodically assessing the species distribution and abundance

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Standards	Medium	Develop guidelines for pest control professionals for the non-lethal control of the species	Change in the number of trained pest control professionals	Pest Control Professionals	SWG
Standards	Medium	Develop guidelines for retention of suitable cavity trees on public and private forest land	Change in the number of suitable cavity trees retained	VFPR	SWG
Research	High	Monitor distribution and abundance of species	Distribution map	UVM, Pest Control Professionals	SWG



Common Name: **Southern Flying Squirrel**
Scientific Name: ***Glaucomys volans***
Species Group: **Mammal**

Bibliography

Degraaf, R. M. and Rudis, D.D. 1986. New England wildlife: habitat, natural history, and distribution. Gen. Tech. Rep. NE-108. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station; 491 pp.

Dolan P.G., and D.C. Carter. 1977. *Glaucomys volans*. Mammalian Species, 78 1-6.

Godin, A. J. 1977. Wild Mammals of New England. John Hopkins University Press, Baltimore, 304 pp.

Kilpatrick, C. W., and J. Benoit. 2011. Small Mammal Project. Report to Vermont Fish and Wildlife Department, 92 pp.

Muul, I. 1968. Behavior and physiological influences on the distribution of the flying squirrel, *Glaucomys volans*. Miscellaneous Publications of the Museum of Zoology, University of Michigan, 134:1-66.

Wells-Gosling, N., and L. R. Heaney. 1984. *Glaucomys sabrinus*. Mammalian Species, 220:1-8.



Common Name: **Northern Flying Squirrel**
Scientific Name: **Glaucomys sabrinus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S4

State Trend: stable

Extirpated in VT? no

Regional SGCN? no

Assessment Narrative:

The Northern Flying Squirrel has a state rarity rank of apparently secure (S4) but little is known about its biology, population and threats within the state. Northern flying squirrels are cavity nesters that frequently nest in woodpecker holes (Wells-Gosling and Heaney 1984). Although they also use non-cavity stick nests, these exposed structures are unsuitable as winter nests, requiring the utilization of cavities during winter months (Cowan 1936). Experimental studies (Weigl 1977) have shown that the smaller southern flying squirrel (*Glaucomys volans*) is dominant and more aggressive sometimes displacing northern flying squirrels from nest boxes. The northern range expansion of the southern flying squirrel may have several negative impacts on populations of northern flying squirrels. The southern flying squirrel may have a greater ability to locate and dominate tree cavities thus displacing northern flying squirrels from hardwood forest (Wells-Gosling and Heaney 1984). Furthermore, in areas where the two species are sympatric, the earlier breeding southern flying squirrel may have an advantage by being the first to occupy tree cavities as nest sites for their young (Wells-Gosling and Heaney 1984). Additionally, southern flying squirrels have a parasite (*Strongyloides* sp.) that appears to be debilitating or lethal to northern flying squirrels (Weigl 1977). Finally, the dietary requirements of northern flying squirrels are not understood. These squirrels cannot be maintained on a diet of spruce seed (Brink and Dean 1966). Fungi and lichens may be the predominant or only foods eaten at certain times of the year (Cowan 1936, Connor 1960, McKeever 1960, Wrigley 1969, Maser et al. 1978, Mowrey et al. 1981, Maser et al. 1985, Mayer et al. 2005)

Distribution

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Historic Records
Northern Green Mtns	Confident	Southern Green Mtns	Probable
Northern VT Piedmont	Confident	Taconic Mtns	Historic Records
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Known to inhabit a wide variety of woodland habitats including spruce-fir and mixed hemlocks and adjacent mature hardwoods (Weigl 1978). Under experimental conditions, Weigl (1978) found that northern flying squirrels would select for either deciduous or coniferous habitat whereas the southern flying squirrel strongly selected deciduous habitat. Given the southern flying squirrels ability to displace northern flying squirrels from



Common Name: **Northern Flying Squirrel**
Scientific Name: **Glaucomys sabrinus**
Species Group: **Mammal**

tree cavities, northern flying squirrels are likely displaced from hardwood forest in areas where the two species are sympatric. Northern flying squirrels require mature trees with cavities for winter nest sites (Cowan 1936). The species feeds on hypogeous fungi in the summer and arboreal lichens and hypogeous fungi in the winter (DeGraff et al, 1986, Rosentreter et al. 1997, Curran et al. 2000, Vernes et al. 2004).

Current Threats

Habitat Threats:

Conversion of Habitat

Habitat Alteration

Climate Change

Description of habitat threat(s):

Non-Habitat Threats:

Competition

Description of non-habitat threat(s): Predicted changes in the climate may allow the southern flying squirrel population to shift northward thereby increasing competition with northern flying squirrels. Increased competition for suitable nesting cavities may be amongst the most significant impact resulting from climate change particularly with respect to the outright loss of nest cavities and/or the displacement from the use of nest cavities. Climate change may also limit northern flying squirrel populations by influencing the abundance of key dietary requirements such as lichen and fungi.

Increased sympatry with southern flying squirrels may result in the spread of parasites to northern flying squirrels. Although the intestinal nematode, *Strongylorides robustus*, infects both southern and northern flying squirrels, it is more prevalent in southern flying squirrels (Wetzel and Weigl 1994) and appears to be more deleterious to northern flying squirrels (Pauli et al. 2004).

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	High	Monitor distribution and abundance of this species.
Research	Threats and Their Significance	Medium	Determine the prevalence of the <i>Strongylorides robustus</i> parasite in flying squirrel populations in Vermont
Monitoring	Range Shifts	High	Monitor changes in the distribution of flying squirrels to determine the degree of sympatry.



Common Name: **Northern Flying Squirrel**
Scientific Name: **Glaucomys sabrinus**
Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Standards	Medium	Develop guidelines for retention of suitable cavity trees on public and private forest land.	Change in the number of suitable cavity trees retained	VFPR	SWG, PR
Research	High	Monitor distribution and abundance of species	Distribution maps	UVM	SWG, PR



Common Name: Northern Flying Squirrel
Scientific Name: *Glaucomys sabrinus*
Species Group: Mammal

Bibliography

- Brink, C. H., and F. C. Dean. 1966. Spruce seed as a food of red squirrels and flying squirrels in interior Alaska. *Journal of Wildlife Management*, 30:503-512.
- Connor, P. F. 1960. The small mammals of Otsego and Schohaire counties, New York. *New York State Museum Science Service Bulletin*, 382:1-84.
- Cowan, I. McT. 1936. Nesting habits of the flying squirrel, *Glaucomys sabrinus*. *Journal of Mammalogy*, 17:58-60.
- Curran, R. S., E. A. Smreciu, T. Lehesvirta, and K. W. Larson. 2000. Fungi in the winter diet of northern flying squirrels and red squirrels in the boreal mixedwood forest of northeastern Alberta. *Canadian Journal of Botany*, 78:1514-1520.
- Degraaf, R.M. and Rudis, D.D. 1986. New England wildlife: habitat, natural history, and distribution. Gen.Tech. Rep. NE-108. Broomall, PA: United States Department of Agriculture, Forest Service, Northeast Forest Experiment Station. 491 pp
- Godin, A. J. 1977. *Wild Mammals of New England*. John Hopkins University Press, Baltimore, 304 pp.
- Maser, C., J. M. Trappe, and R. A. Nausbaum. 1978. Fungal-small mammal interrelationships with emphasis on Oregon coniferous forest. *Ecology*, 59:799-809.
- Maser, Z., C. Maser, and J. M. Trappe. 1985. Food habits of the northern flying squirrel (*Glaucomys sabrinus*) in Oregon. *Canadian Journal of Zoology*, 63:1084-1088.
- McKeever, S. 1960. Food of the northern flying squirrel in California. *Journal of Mammalogy*, 41:270-271.
- Meyer, M. D., M. P. North, and D. A. Kelt. 2005. Fungi in the diet of northern flying squirrels and lodgepole chipmunks in the Sierra Nevada. *Canadian Journal of Zoology*, 83:1581-1589.
- Mowrey, R. A., G. A. Laursen, and T. A. Moore. 1981. Hypogeous fungi and small mammal mycophagy in Alaska taiga. *Proceedings of Alaska Science Conference*, 32:120-121.
- Pauli, J. N., S. A. Dubay, E. M. Anderson, and S. J. Taft. 2004. *Strongylorides robustus* and the northern sympatric populations of northern (*Glaucomys sabrinus*) and southern (*G. volans*) flying squirrels. *Journal of Wildlife Diseases*, 40:579-582.
- Rosentreter, R., G. D. Hayward, and M. Wicklow-Howard. 1997. Northern flying squirrel seasonal food habits in the interior conifer forest of Central Idaho, USA. *Northwest Science*, 71:97-102.
- Saunders, D. A. 1988. *Adirondack Mammals*. College of Environmental Science and Forestry, State University of New York, Syracuse, 216 pp.
- Vernes, K., S. Blois, and F. Barlocher. 2004. Seasonal and yearly changes in the consumption of hypogenous fungi by northern flying squirrels and red squirrels in old-growth forest, New Brunswick. *Canadian Journal of Zoology* 82:110-117.
- Weigl, P. D. 1977. Status of the northern flying squirrel, *Glaucomys sabrinus coloratus*, in North Carolina. Pp. 398-400, in *Endangered and Threatened Plants and Animals of North Carolina* (J. E. Cooper, S. S. Robinson, and J. B. Funderburg, eds.). North Carolina State Museum of Natural History, Raleigh, 444 pp.
- Wells-Gosling, N., and L. R. Heaney. 1984. *Glaucomys sabrinus*. *Mammalian Species*, 220:1-8.
- Wetzel, E. J., and P. D. Weigl. 1994. Ecological implications for flying squirrels (*Glaucomys* spp.) of effects of temperature on the vitro development and behavior of *Strongylorides robustus*. *American Midland Naturalist*, 131:43-54.
- Wrigley, R. E. 1969. Ecological notes on the mammals of southeastern Quebec. *Canadian Field-Naturalists*, 83:201-211

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Rock Vole**
Scientific Name: **Microtus chrotorrhinus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: S2

State Trend: unknown

Extirpated in VT? no

Regional SGCN? No

Assessment Narrative:

Ranked as S2 in Vermont and considered a species of special concern. Talus slopes are the species' refugal habitat. In some locations, rock voles may be found in early successional forest habitat (Kirkland 1977; Martell and Radvanyi 1977) and krumholtz. There is uncertainty as to why the population fluctuates so much and there are relatively few known populations. The relatively narrow habitat requirements of this species make it vulnerable to habitat alterations. Furthermore, because rock voles occur in disjunct populations, it is dependent upon movement corridors. It is also speculated that these disjunct populations could be negatively impacted by landscape changes that favor the meadow vole (*Microtus pennsylvanicus*) which is a suspected competitor of the rock vole.

Distribution

There are a number of historic records indicating the species' existence and distribution in the state. These records include: 20 specimens from Island Pond at 1400' elevation (1937-1940); two specimens from Brighton on the talus slopes of NW Bluff Mountain (1953); one specimen from near Smugglers cave, Mt. Mansfield (1954); four specimens from Nebraska Notch, Mt. Mansfield (1958-1959); and two specimens (one male and one female) from Nebraska Notch, Mt Mansfield (1966). More contemporary records of the rock vole in Vermont include: Whenlock WMA (Chipman, 1994); West Mountain WMA (Kilpatrick, 2001); East Mountain, East Haven (Kilpatrick, 2005), and East Charleston (Kilpatrick, pers. comm.). The Vermont Small Mammal Atlas also recorded six specimens from four sites in Essex and Caledonia counties between 2008 and 2010 (Kilpatrick and Benoit 2011). Unknown populations were verified inhabiting talus slopes on Brousseau Mountain (Averill, Essex Co.), Umpire Mountain (Victory, Essex Co.) and Wheeler Mountain (Sutton, Caledonia Co.) and the population from West Mountain WMA was verified to still exist (Kilpatrick and Benoit 2011).

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Not Probable
Champlain Hills	Historic Records Only	Vermont Valley	Not Probable
Northern Green Mtns	Confident	Southern Green Mtns	Not Probable
Northern VT Piedmont	Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Confident		

Distribution by Watershed:



Common Name: **Rock Vole**
Scientific Name: **Microtus chrotorrhinus**
Species Group: **Mammal**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Occurs in disjunct populations that are not genetically differentiated so movement corridors may be important. This species is very habitat selective. They use moist talus habitats among mossy rocks and logs in spruce/ fir and northern hardwood forests, cedar swamps, and krummholz. May be naturally rare due to habitat specificity. Rock vole has been reported in three-five year old clearcuts with slash however, not in Vermont. Critical habitat includes cool, moist talus and mossy rocks usually with a stream or other surface water in the immediate vicinity.

Habitat Types:

Cliffs and Talus

Softwood Swamps

Spruce Fir Northern Hardwood

Current Threats

Habitat Threats:

Habitat Alteration

Description of habitat threat(s): Mesic aspect of habitat is important so the loss of forest cover may dry out the site. Loss of connectivity may be a problem. Habitat is isolated and local populations may go extinct. Repopulation may require habitat corridors of coniferous forests that connect optimal habitats. Activities that destroy or degrade talus habitat would impact rock vole populations.

Non-Habitat Threats:

Competition

Description of non-habitat threat(s): Competition from meadow mouse as a result of habitat conversion, particularly near talus areas, could limit the rock vole. Metapopulation structure is not clearly understood but local populations appear to go extinct and then are repopulated. In Massachusetts and West Virginia populations were negatively affected by high levels of deer over the long term (Healey and Brooks 1988).

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Rock Vole**
Scientific Name: **Microtus chrotorrhinus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: S2

State Trend: unknown

Extirpated in VT? no

Regional SGCN? No

Assessment Narrative:

Ranked as S2 in Vermont and considered a species of special concern. Talus slopes are the species' refugal habitat. In some locations, rock voles may be found in early successional forest habitat (Kirkland 1977; Martell and Radvanyi 1977) and krumholtz. There is uncertainty as to why the population fluctuates so much and there are relatively few known populations. The relatively narrow habitat requirements of this species make it vulnerable to habitat alterations. Furthermore, because rock voles occur in disjunct populations, it is dependent upon movement corridors. It is also speculated that these disjunct populations could be negatively impacted by landscape changes that favor the meadow vole (*Microtus pennsylvanicus*) which is a suspected competitor of the rock vole.

Distribution

There are a number of historic records indicating the species' existence and distribution in the state. These records include: 20 specimens from Island Pond at 1400' elevation (1937-1940); two specimens from Brighton on the talus slopes of NW Bluff Mountain (1953); one specimen from near Smugglers cave, Mt. Mansfield (1954); four specimens from Nebraska Notch, Mt. Mansfield (1958-1959); and two specimens (one male and one female) from Nebraska Notch, Mt Mansfield (1966). More contemporary records of the rock vole in Vermont include: Whenlock WMA (Chipman, 1994); West Mountain WMA (Kilpatrick, 2001); East Mountain, East Haven (Kilpatrick, 2005), and East Charleston (Kilpatrick, pers. comm.). The Vermont Small Mammal Atlas also recorded six specimens from four sites in Essex and Caledonia counties between 2008 and 2010 (Kilpatrick and Benoit 2011). Unknown populations were verified inhabiting talus slopes on Brousseau Mountain (Averill, Essex Co.), Umpire Mountain (Victory, Essex Co.) and Wheeler Mountain (Sutton, Caledonia Co.) and the population from West Mountain WMA was verified to still exist (Kilpatrick and Benoit 2011).

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Not Probable
Champlain Hills	Historic Records Only	Vermont Valley	Not Probable
Northern Green Mtns	Confident	Southern Green Mtns	Not Probable
Northern VT Piedmont	Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Confident		

Distribution by Watershed:



Common Name: **Rock Vole**
Scientific Name: **Microtus chrotorrhinus**
Species Group: **Mammal**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Occurs in disjunct populations that are not genetically differentiated so movement corridors may be important. This species is very habitat selective. They use moist talus habitats among mossy rocks and logs in spruce/ fir and northern hardwood forests, cedar swamps, and krummholz. May be naturally rare due to habitat specificity. Rock vole has been reported in three-five year old clearcuts with slash however, not in Vermont. Critical habitat includes cool, moist talus and mossy rocks usually with a stream or other surface water in the immediate vicinity.

Habitat Types:

Cliffs and Talus

Softwood Swamps

Spruce Fir Northern Hardwood

Current Threats

Habitat Threats:

Habitat Alteration

Description of habitat threat(s): Mesic aspect of habitat is important so the loss of forest cover may dry out the site. Loss of connectivity may be a problem. Habitat is isolated and local populations may go extinct. Repopulation may require habitat corridors of coniferous forests that connect optimal habitats. Activities that destroy or degrade talus habitat would impact rock vole populations.

Non-Habitat Threats:

Competition

Description of non-habitat threat(s): Competition from meadow mouse as a result of habitat conversion, particularly near talus areas, could limit the rock vole. Metapopulation structure is not clearly understood but local populations appear to go extinct and then are repopulated. In Massachusetts and West Virginia populations were negatively affected by high levels of deer over the long term (Healey and Brooks 1988).

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Rock Vole**
 Scientific Name: **Microtus chrotorrhinus**
 Species Group: **Mammal**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Basic Life History	Medium	Telemeter to determine home range movements
Research	Distribution and Abundance	High	Determine distribution and abundance as well as corridor needs
Research	Population Genetics	Medium	Research genetics to determine changes in population structure and size.
Research	Other Research	High	Determine appropriate management strategies to improve and conserve habitat.
Monitoring	Population Change	Medium	In a multi year monitoring effort, re-census historical habitats and survey in other likely habitats and map confirmed habitats.
Monitoring	Monitor Threats	Medium	Monitor encroachment by meadow mice.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Compatible Resource Use	Medium	Minimize permanent fragmentation between populations.	Amount of habitat between populations protected or conserved.	UVM	SWG

Bibliography

Degraaf, R.M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution, University Press of New England, Hanover, New Hampshire, USA.

Chipman, R. B. 1994. Distribution, relative abundance, and habitat use by small mammals in Vermont. M.S. Thesis, University of Vermont, Burlington, 168 pp.

Healy, W. M., and R. T. Brooks. 1988. Small mammal abundance in northern hardwood stands in West Virginia. *Journal of Wildlife Management*. 52:491-496.

Kilpatrick, C. W. 2001. Small mammal survey of the Nulhegan Basin Division of the Silvio O. Conte NFWR and the State of Vermont's West Mountain Wildlife Management Area, Essex County, Vermont. Final Report March 15, 2001 submitted to the Nature Conservancy.

Kilpatrick, C. W. 2005. East Mountain Mammal Survey August 2004- December 2004. Final report May 15, 2005, Prepared for Kingdom Commons Group.

Kilpatrick, C. W., and J. Benoit. 2011. Small mammal project. University of Vermont/NorthWoods Stewardship Center, final report submitted to Vermont Fish and Wildlife Department.

Kirkland, G. L., Jr. 1977. The rock vole, *Microtus chrotorrhinus* (Miller) (Mammalia: Rodentia) in West Virginia. *Annals of Carnegie Museum* 46:45-53.

Kirkland, G.L., Jr., and F.J. Jannett. 1982. *Mammalian Species* 180:1-5.

Martell, A.M., and A. Radvanyi. 1977. Changes in small mammal populations after clear cutting of northern Ontario black spruce forest. *Canadian Field Naturalist* 91:41-46.

Osgood, F. L., Jr. 1938. First Vermont record of the rock vole. *Journal of Mammalogy* 19:108.

Tumosa, J. 2001. United States Forest Service species data collection form for *Microtus chrotorrhinus*. 19 pp



Common Name: **Woodland Vole**
 Scientific Name: **Microtus pinetorum**
 Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S3

State Trend: unknown

Extirpated in VT? no

Regional SGCN? no

Assessment Narrative:

The woodland vole is frequently considered a pest in agricultural settings (especially in apple orchards) though much of the reported damage is the result of meadow voles (*Microtus pennsylvanicus*). Despite appearing to do well in agricultural landscapes, little is known about this species outside this setting or in its native habitat. Fewer than 50 specimens have been collected in the state and is known historically from very few localities.

Distribution

Known historically from very few localities including the flanks of Ide Mountain, West Lyndon Center (Miller, 1964); Island Pond (Miller, 1964); Sherburne (Osgood, 1936); and from Colchester and Duxbury (Kilpatrick, pers. com). Woodland voles occur in orchards in Putney, Mendon, and Bennington (Kilpatrick, 1979). The Vermont Small Mammal Atlas obtained two specimens from two localities in Orleans and Windsor counties from 2008 to 2010; one was trapped in the Skitchewaug WMA (species verified by DNA sequencing) and another collected from a garden in Charleston (Kilpatrick and Benoit 2011). Records were also verified from Addison County and Chittenden County (Kilpatrick and Benoit 2011).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Probable
Northern Green Mtns	Probable	Southern Green Mtns	Probable
Northern VT Piedmont	Confident	Taconic Mtns	Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

Defining habitat characteristic of the woodland vole is well-drained sandy loam soils. Found in all places with these soils (e.g. agricultural fields and older forests). Favors well-drained upland forests, grasslands, meadows, or orchards but can be found in marshes and swamps (DeGraff and Yamasaki, 2001). May require a ground cover of leaves or duff or grass. Forages primarily below ground digging tunnel systems 3 inches to 12 inches below ground. Nests are found under dead and down material, rocks, or in burrows. They are active throughout the year and eat tubers, roots and bulbs, seeds, nuts fruits, bark and leaves (DeGraff and Yamasaki, 2001). Can be a problem in orchards.. Prefers large expanses of forest and grassland habitats,



Common Name: **Woodland Vole**
 Scientific Name: **Microtus pinetorum**
 Species Group: **Mammal**

Habitat Types:

Northern Hardwood
 Spruce Fir Northern Hardwood
 Oak-Pine Northern Hardwood
 Grasslands, Hedgerows, Old Field, Shrub, or Orchard
 Lawns, Gardens, and Row Crops

Current Threats

Habitat Threats:

Unknown Habitat Threats

Description of habitat threat(s): Habitat requirements unknown.

Non-Habitat Threats:

Pollution

Trampling or Direct Impacts

Description of non-habitat threat(s): Because of human/vole conflicts, the application of rodenticides may cause a decline of this species in orchards and other developed lands. The status of the woodland vole in forested habitats is unknown.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Develop baseline data on habitat requirements outside of agricultural areas.
Research	Distribution and Abundance	Medium	Develop baseline data on distribution and abundance outside of agricultural areas.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Standards	Medium	Develop guidelines for pest control professionals for the non-lethal control of the species.	Number of trained pest control professionals	Agricultural extension, Pest Control Professionals	SWG



Common Name: **Woodland Vole**
Scientific Name: ***Microtus pinetorum***
Species Group: **Mammal**

Bibliography

- Benton, A.H. 1955. Observations on the life history of the northern pine mouse. *Journal of Mammalogy*, 36: 52-62.
- DeGraaf, R. M., and M. Yamasaki. 2001. *New England Wildlife: Habitat, Natural History, and Distribution*. University Press of New England, Hanover. 482 pp.
- Hamilton, W.J. 1938. Life history notes on the northern pine mouse. *Journal of Mammalogy*, 19:163-170.
- Kilpatrick, C.W. 1979. Habitat, climatological and management variables associated with vole populations in Vermont orchards, Unpublished Report. 27pp. For U.S. Fish and Wildlife Service.
- Kilpatrick, C. W., and J. Benoit. 2011. Small Mammal Project. Report to Vermont Fish and Wildlife Department, 92 pp.
- Miller, D.H. 1964. Northern records of the pine mouse in Vermont. *Journal of Mammalogy*, 45:627-628.
- Miller, D.H. and L.L. Getz. 1969. Life-history notes on *Microtus pinetorum* in central Connecticut. *Journal of Mammalogy*, 50: 777-784.
- Osgood, F.L. 1936. Pine mice in Vermont. *Journal of Mammalogy*. 17: 291-292.
- Smolen, M.J. 1981. *Microtus pinetorum*. *Mammalian species*, 147: 1-7.



Common Name: **Muskrat**
 Scientific Name: **Ondatra zibethicus**
 Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5

State Trend: Declining

Extirpated in VT? no

Regional SGCN? no

Assessment Narrative:

The muskrat has traditionally been one of the most heavily exploited furbearers in North America owing to its abundance, relative ease of capture and highly prized fur (Boutin and Birkenholz 1987). Across its range today, most jurisdictions, including Vermont, maintain regulated trapping and hunting seasons for the species. The muskrat plays an important ecological role serving as a significant prey source for a variety of predators including raptors, river otter and American mink (Holmengen et al 2009). In recent decades, anecdotal evidence indicates a nationwide decline in muskrat populations. Such noted declines have been most prominent in the northeast. Despite much knowledge regarding the biology and management of muskrats, little empirical evidence exists indicating either the magnitude of such declines and/or any possible contributing factors (Roberts and Crimmins 2010).

Distribution

Muskrat harvest records in Vermont indicate well established populations in all major watersheds.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Known Watersheds

Middle Connecticut
 West
 Waits
 Upper Connecticut-Mascoma
 Black-Ottawquechee
 Deerfield
 Hudson-Hoosic
 Mettawee River
 Lake Champlain
 Lamoille River



Common Name: **Muskrat**
Scientific Name: **Ondatra zibethicus**
Species Group: **Mammal**

Missisquoi River
Otter Creek
Passumpsic
St. Francois River
Upper Connecticut
White
Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Muskrats occupy almost every type of freshwater aquatic habitat in eastern North America (Boutin and Birkenholz 1987). Muskrat have flexible habitat requirements as long as there is permanent water and protection through burrows and vegetated lodges. Highest population densities exist where emergent vegetation is at a 1:1 ratio to open water.

Habitat Types:

Marshes and Sedge Meadows
Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lower CT River
Aquatic: Lacustrine
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Invasion by Exotic Species
Unknown Habitat Threats

Description of habitat threat(s): Although the specific effects of habitat alteration on muskrats are poorly understood, the anthropogenic degradation of muskrat habitat is widely recognized as a potential contributing factor to the decline of populations throughout the region. Increased sedimentation and stream flashiness resulting from poorly planned land management and/or excessive development could, for example, alter the ratio of open water to emergent vegetation within watersheds to the detriment of muskrats. Similarly, human activities resulting in the spread of invasive plant species, such as phragmites, can cause a reduction in the abundance and diversity of native taxa, including muskrats, by creating monotypic stands.

Non-Habitat Threats:

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Muskrat**
 Scientific Name: **Ondatra zibethicus**
 Species Group: **Mammal**

Genetics

Predation or Herbivory

Loss of Relationship with Other Species

Description of non-habitat threat(s): Previous studies of contaminant levels in muskrats have shown that muskrats bioaccumulate heavy metals (Halbrook et al. 1993, Stevens et al. 1997). While the direct effects of such contaminants on muskrats remain uncertain, there is continued concern that the long-term persistence of such contaminants in the environment could limit muskrat populations. While the significance and magnitude of other non-habitat threats are poorly understood, it is speculated that changes in predatory communities, diseases and alterations of natural water cycles all potentially contribute to observed declines in muskrat populations regionally.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	High	Continue closely monitoring the distribution and abundance in Vermont
Research	Threats and Their Significance	High	Determine what factors may be influencing population declines, focusing in particular on pollution and habitat degradation.
Research	Other Research	Medium	Conduct a cause specific mortality study to aid in the identification of significant mortality factors in Vermont.
Monitoring	Monitor Threats	Medium	Monitor the accumulation of contaminants such as heavy metals and PCBs in the tissues of muskrats throughout all watersheds in Vermont.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	High	Determine causes of observed declines in regional muskrat populations	Number of hypotheses evaluated	UVM, AFWA	SWG, PR
Policy & Regulations	Medium	Support and cooperate with regional efforts to curb pollution via the development and implementation of appropriate policy and regulations	Reduction in the prevalence of contaminants in Vermont's water bodies	DEC, EPA	
Compliance & Enforcement	High	Enforce existing laws with respect to water quality protection	Increased compliance with existing laws	DEC, EPA	
Compliance & Enforcement	High	Enforce existing laws with respect to riparian and wetland habitat protection	Area and/or linear distance of riparian and wetland habitat protected	DEC, EPA, USACE	
Invasive Species Control & Prevention	Medium	Identify and restore muskrat habitat impaired by invasive plants, and develop and implement measures aimed at preventing further introduction of such species	Acreage of habitat restored and number of preventative measures adopted		



Common Name: **Muskrat**
Scientific Name: **Ondatra zibethicus**
Species Group: **Mammal**

Bibliography

Boutin, S. and Birkenholz, D.E., .1987. Muskrat. In M. Novak, J.A. Baker, M.E. Obbard, and B. Malloch, eds. Wild furbearer management and Conservation in North America. Ontario Trappers Assoc., North Bay.

Everett, J.J. and R.G. Anthony.1976. Heavy metal accumulation in muskrats in relation to water quality. Trans. Northeastern Fish and Wildlife Conf., 33:105-116.

Stevens, R.T., T.L. Ashwood, and J.M. Sleeman 1997. Mercury in hair of muskrats (*Ondatra zibethicus*) and mink (*Mustela vison*) from the U.S. Department of Energy, Oak Ridge Reservation. Bulletin of Environmental Contamination and Toxicology, 58: 720-725.

Willner, G.R., G.A. Feldhammer, E.E. Zucker, and J.A. Champmand, 1980. *Ondatra zibethicus*. Mammalian Species, 141: 1-8.



Common Name: **Southern Bog Lemming**
Scientific Name: **Synaptomys cooperi**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S3

State Trend: Fluctuating

Extirpated in VT? no

Regional SGCN? yes

Assessment Narrative:

Although the southern bog lemming is relatively rare in collections, it is by no means an uncommon animal (Whitaker and Hamilton 1998). A number of historical records are available primarily for southern Vermont (Kirk 1916, Osgood 1938, Godin 1977). When combined with recent records (Brooks et al. 1998, Kilpatrick 2003, Decher and Kilpatrick 2005, Kilpatrick and Benoit 2011) some 268 specimens of the southern bog lemming confirm the occurrence at over 35 different localities throughout the state (see Kilpatrick and Benoit 2011). The species is believed to exist in scattered colonies that often inhabit only a small portion of the suitable habitat. Although little is known about potential threats to this species in Vermont, it is believed southern bog lemmings are vulnerable to changes in habitat, competition with meadow voles and to a variety of disease and parasites.

The southern bog lemming is listed as a Regional Species of Greatest Conservation Need (RSGCN) among the 13 Northeastern states.

Distribution

The southern bog lemming is known from throughout the state with the exception of Grand Isle, Franklin, and Orange Counties

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Historical Records
Champlain Hills	Confident	Vermont Valley	Historical Records
Northern Green Mtns	Confident	Southern Green Mtns	Historical Records
Northern VT Piedmont	Probable	Taconic Mtns	Historical Records
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

The southern bog lemming uses a wide variety of habitats in addition to sphagnum bogs, including wet meadows and marshes, grassy openings in woods, and among mossy boulders in spruce forests (Linzey 1983). In Southern Canada, New York and New England most captures are associated with sphagnum bogs or heavily forested areas (Coventry 1942, Goodwin 1932, Hamilton 1941). The southern bog lemming will use clearcuts and other small forest openings with adequate ground cover (Kirkland 1977). Recent small mammal surveys in Vermont (Kilpatrick and Benoit 2011) found southern bog lemming among small rock outcrop in a mesic spruce forest and in a red pine plantation. Doult et al. (1973) suggested that the major feature common to *Synaptomys* habitats was that they were marginal for *Microtus* and Linzey (1981, 1984) documented



Common Name: Southern Bog Lemming
Scientific Name: *Synaptomys cooperi*
Species Group: Mammal

competitive exclusion of *Synaptomys* by *Microtus* in southwestern Virginia. Southern bog lemmings have been collected from hairy-tailed mole burrows (Eadie 1939).

Habitat Types:

Outcrops and Alpine
Northern Hardwood
Softwood Swamps
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Open Peatlands
Marshes and Sedge Meadows
Grasslands, Hedgerows, Old Field, Shrub, or Orchard

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Climate Change

Description of habitat threat(s): Although little has been documented about the potential habitat related threats to this species, it is believed southern bog lemmings are vulnerable to the degradation of preferred habitats resulting from climate change, forest succession, and/or direct human impacts. Habitat threats are of particular concern with respect to a potentially drying climate and the direct loss of sphagnum bogs.

Non-Habitat Threats:

Competition
Disease

Description of non-habitat threat(s): Competition from *Microtus* (meadow vole) in sites where habitat has been altered and/or forest succession has favored this species. Southern bog lemmings carry a heavy ectoparasite parasite load (Wassel et al. 1978) and several endoparasites have been confirmed (Erickson 1938, Whitaker and Adalis 1971).

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Determine baseline information
Research	Distribution and Abundance	High	Determine baseline information
Monitoring	Population Change	High	1) Begin low-level monitoring in appropriate habitats to determine distribution, abundance, and population status and trends. 2) Better understand distribution, abundance and changes in population.



Common Name: **Southern Bog Lemming**
Scientific Name: **Synaptomys cooperi**
Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	High	Monitor distribution and abundance of species	Distribution maps	UVM	SWG, PR



Common Name: Southern Bog Lemming
Scientific Name: *Synaptomys cooperi*
Species Group: Mammal

Bibliography

- Buckner, C.H. 1957. Home range of *Synaptomys cooperi*. *Journal of Mammalogy* 38:132.
- Beasley, L.E. and L.L. Getz. 1986. Comparison of demography of sympatric populations of *Microtus ochrogaster* and *Synaptomys cooperi*. *Acta Theriologica* 31:395-400.
- Brooks, R. T., H. R. Smith, and W. M. Healy. 1998. Small mammal abundance at three elevations on a mountain in central Vermont, USA: a 16 year record. *Forest Ecology and Management*, 110:181-193.
- Connor, P.F. 1959. The bog lemming *Synaptomys cooperi* in Southern New Jersey. Publication of the Museum, Michigan State University, Biological Series 1:161-248.
- Coventry, A. F. 1942. *Synaptomys cooperi* in forested regions. *Journal of Mammalogy*, 23:450-451.
- Decher, J., and C. W. Kilpatrick. 2005. Small mammals of the Guthrie-Bancroft Farm – Year 4 Colby Hill Ecological Project, Lincoln and Bristol, Vermont.
- DeGraaf, R. M., and M. Yamasaki. 2001. *New England Wildlife: Habitat, Natural History, and Distribution*. University Press of New England, Hanover, NH.
- Doutt, J. K., C. A. Heppenstall, and J. E. Guilday. 1973. *Mammals of Pennsylvania*. Third ed. Pennsylvania Game Commission, Harrisburg, 283 pp.
- Eadie, W. R. 1939. A contribution to the biology of *Parascalops breweri*. *Journal of Mammalogy*, 20:150-173.
- Erickson, A. B. 1938. Parasites of some Minnesota Cricetidae and Zapodidae, and a host catalogue of helminth parasites of native American mice. *American Midland Naturalists*, 20:575-589.
- Godin, A.J. 1977. *Wild Mammals of New England*. Johns Hopkins University Press. Baltimore, MD, 304 pp.
- Goodwin, G. G. 1932. New records and some observations on Connecticut mammals. *Journal of Mammalogy*, 13:36-40.
- Hamilton, W. J., Jr. 1941. On the occurrence of *Synaptomys cooperi* in forested regions. *Journal of Mammalogy*, 22:195.
- Kilpatrick, C. W. 2003. Small mammal survey of the Vermont Ecosystem Management Project Plots, Steenville Brooke Research Area, Mount Mansfield State Forest.
- Kilpatrick, C. W., and J. Benoit. 2011. Small Mammal Project. Report submitted Vermont Fish and Wildlife Department, 92 pp.
- Kirk, G. L. 1916. The mammals of Vermont. *Joint Bulletin, Vermont Botanical and Bird Club*, 2:28-34.
- Kirkland, G. L., Jr. 1977. Responses of small mammals to the clearcutting of northern Appalachian forest. *Journal of Mammalogy*, 58:600-609.
- Krupa, J.J. 1996. Invasion of the meadow vole (*Microtus pennsylvanicus*) in southeastern Kentucky and its possible impact on the Southern bog Lemming (*Synaptomys cooperi*). *American Midland Naturalist* 135:14-22.
- Linsey, A.V. 1981. Patterns of coexistence in *Microtus pennsylvanicus* and *Synaptomys cooperi*. Unpublished Ph.D. dissertation, Virginia Polytechnic Institute and State University, Blacksburg, 97 pp.
- Linsey, A.V. 1983. *Synaptomys cooperi*. *Mammalian Species*, 210:1-5.
- Linzey, A.V. 1984. Patterns of coexistence in *Synaptomys cooperi* and *Microtus pennsylvanicus*. *Ecology* 65:382-393.
- Linzey, D.W. 1995. *Mammals of Great Smoky Mountains National Park*. McDonald and Woodward Publishing Company, Blacksburg, VA.
- Osgood, F. L., Jr. 1938. The mammals of Vermont. *Journal of Mammalogy*, 19:435-441.
- Wassel, M. E., G. L. Tieben, and J. O. Whitaker, Jr. 1978. The ectoparasites of the southern bog lemming, *Synaptomys cooperi*, in Indiana. *Proceedings of the Indiana Academy of Science*, 87:446-449.



Common Name: **Southern Bog Lemming**
Scientific Name: **Synaptomys cooperi**
Species Group: **Mammal**

Whitaker, J. O., Jr., and D. Adalis. 1971. Trematodes and cestodes from the digestive tracts of *Synaptomys cooperi* and three species of *Microtus* from Indiana. *Proceedings of the Indiana Academy of Science*, 80:489-494.

Whitaker, J. O., Jr., and W. J. Hamilton, Jr. 1998. *Mammals of the Eastern United States*. Third ed. Cornell University Press, Ithaca, NY, 583 pp.

Wilson, G.H.. 1997. Taxonomic status and biogeography of the Southern bog lemming, *Synaptomys cooperi*, on the Central Great Plains. *Journal of Mammalogy* 78: 444-458



Common Name: **Northern bog lemming**
Scientific Name: **Synaptomys borealis**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G4

Global Trend:

State Rank: SU

State Trend: Unknown

Extirpated in VT? no

Regional SGCN? No

Assessment Narrative:

Although there are no historical or recent records of the northern bog lemming in Vermont, records are known from surrounding states including a recent specimen from Whiteface Mountain, NY (Sanderson 1988), and three specimens from New Hampshire including two from Coos Co., from Fabyans near the base of Mt Washington (Preble 1899) and Bean's Purchase (Yamasaki 1997) and one from Mt Moosilauke, Grafton Co. (Clough and Albright 1987). Five specimens have been verified from Maine from two localities in Piscataquis Co., one being Mt Katahdin and the other a low elevation site near the western border of Baxter State Park (Clough and Albright 1987). Additional specimens are known from Quebec (Cross 1938, Banfield 1974). The northern bog lemming is among the rarest mammals in New England and eastern Canada and is likely vulnerable to local extirpation (Banfield 1974). The subterranean habits of bog lemmings (Banfield 1974, Godin 1977, Degraff and Yamasaki 2001) likely results in infrequent captures of these rodents by traditional collecting methods. This, combined with the difficulty in identification (Clough and Albright 1987), probably contributes substantially to the rarity of northern bog lemmings in surveys and collections of small mammals. A recent small mammal survey in New Hampshire (Yamasaki 1997) employing methods to increase the captures of several rare small mammal species captured a single northern bog lemming at one of the 108 sites surveyed. No northern bog lemmings were captured at the 51 sites recently surveyed in Vermont and none were identified among the southern bog lemming specimens examined (Kilpatrick and Benoit 2011). Despite a lack of evidence of the species in the state, Vermont appears to have viable habitat.

Distribution

Distribution by Biophysical Region:

Champlain Valley	Unknown	Southern VT Piedmont	Unknown
Champlain Hills	Unknown	Vermont Valley	Unknown
Northern Green Mtns	Unknown	Southern Green Mtns	Unknown
Northern VT Piedmont	Unknown	Taconic Mtns	Unknown
Northeastern Highlands	Unknown		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The northern bog lemming has been taken at high elevation sites (3700 - 4500 ft.) in spruce-fir forest with dense herbaceous and mossy understory and in alpine sedge meadows containing sphagnum and surrounded by dense spruce-fir Krummholtz (Clough and Albright 1987). At least two records are known from relatively low elevations (1300 - 1600 ft.) in New Hampshire from habitats ranging from a stand of spruce-budworm



Common Name: Northern bog lemming
Scientific Name: *Synaptomys borealis*
Species Group: Mammal

killed spruce-fir forest with a shrub and ground layer consisting of a dense covering of raspberry, ferns, and sedges, and having sphagnum moss in scattered damp places (Clough and Albright 1987) to a wet meadow and mossy streamside (Preble 1898). Habitat requirements included moist loose soils of leaf mold with sphagnum present (Banfield 1974, DeGraff and Yamasaki, 2001). Northern bog lemmings feed on grasses and sedges and use burrows several inches below the ground (Banfield 1974). They are active year round, in summer constructing spherical nest of dried grasses in burrows and in winter nesting on the ground (Banfield 1974).

Habitat Types:

Outcrops and Alpine
Softwood Swamps
Spruce Fir Northern Hardwood
Open Peatlands

Current Threats

Habitat Threats:

Conversion of Habitat
Impacts of Roads or Transportation Systems
Climate Change

Description of habitat threat(s): Two hypotheses for the rarity of this species have been proposed by Clough and Albright (1987); northern bog lemmings require a habitat that is scarce and/or the species cannot coexist with other species of small mammals. Neither hypothesis is strongly supported by the available data. However, habitat conversion that results in the elimination of peat lands, sphagnum bogs and moist wooded areas with a solid floor of thick sphagnum could be a problem for the northern bog lemming. Climate change that results in increasing temperatures, could result in dryer habitats that would allow the meadow vole population to increase and thereby compete and displace northern bog lemmings. Development of roads, trails and powerlines could also provide access for meadow vole populations and result in increased competition with the northern bog lemming.

Non-Habitat Threats:

Competition
Description of non-habitat threat(s): Habitat changes that benefit the meadow vole could result in increased competition that negatively affects the northern bog lemming.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Determine habitat requirements. Map appropriate habitat.
Research	Distribution and Abundance	High	Conduct a dedicated search for northern bog lemming using species specific methods (pit fall traps and drift fences) in sphagnum-dominated vegetative communities.
Monitoring	Range Shifts	Medium	



Common Name: **Northern bog lemming**
 Scientific Name: **Synaptomys borealis**
 Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Research	Medium	Conduct targeted surveys for northern bog lemmings.	Distribution map	UVM	SWG
Research	High	Determine habitat requirements	Map appropriate habitats	UVM	SWG

Bibliography

- Banfield, A. W. F. 1974. The Mammals of Canada. University of Toronto Press, Toronto, Ontario, 438 pp.
- Clough, G.C. and J. J. Albright. 1987. Occurrence of the Northern bog lemming, *Synaptomys borealis*, in the northeastern United States. Canadian Field Naturalist 101:611-613.
- Cross, E.C. 1938. *Synaptomys borealis* from Godbout, Quebec. Journal of Mammalogy 19:378.
- DeGraaf, R. M., and M. Yamaski. 2001. New England Wildlife: habitat, natural history, and distribution. University Press of New England, Hanover, NH.
- Dutcher, B.H. 1903. Mammals of Mt. Katahdin, Maine. Proceedings of the Biological Society of Washington. 16:63-72.
- Godin, A.J. 1977. Wild mammals of New England. Johns Hopkins University Press. Baltimore, MD. 304 pp.
- Hall, E.R. and E.L. Cockrum. 1953. A synopsis of the North American Microtine rodents. University of Kansas Publications, Museum of Natural History. 5:373-498.
- Kilpatrick, C. W., and J. Benoit. 2011. Small Mammal Project. Report submitted to Vermont Fish and Wildlife Department. 92 pp
- Pearson, D.E. 1991. The northern bog lemming in Montana and the contiguous United States: Distribution, Ecology and relic species theory. Unpublished Senior Theses, University of Montana (Missoula) 33 pp.
- Preble, E.A. 1899. Description of a new lemming mouse from the White Mountains, New Hampshire. Proceedings of the Biological Society of Washington 13: 43-45.
- Reichel, J.D. and S.G. Beckstrom. 1994. Northern bog lemming survey: 1993. Montana Natural Heritage Program, Helena, MT. 87pp
- Saunders, D. A. 1988. Adirondack Mammals. State University of New York (College of Environmental Science and Forestry, Syracuse), 216 pp.
- Yamaski, M. 1977. White Mountain National Forest small mammal identification and collection report. Northeast Forest Experimental Station RWU-4155, Durham, NH.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Wolf**
Scientific Name: **Canis sp?**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G4

Global Trend:

State Rank: SX

State Trend: N/A

Extirpated in VT? Yes

Regional SGCN? No

Assessment Narrative:

Believed to be extirpated in Vermont and the rest of New England. Based on bounty records, wolves were historically common in Vermont but were eliminated from the state by the mid to late 1800's as the result of a \$20.00 bounty and habitat changes. There is uncertainty regarding the genetic ancestry of the wolves that inhabited the northeastern United States historically, including Vermont (Wilson et al. 2003, Koblmuller et al. 2009, Kays et al. 2009). Rigorous DNA analysis of additional historic samples from Vermont and the northeastern United States may help clarify this issue.

The wolf is currently considered extirpated in the Northeast but populations exist in southern Canada with potential for migrants to arrive in Vermont within next 20 years. However, the St. Lawrence River and adjacent agricultural/urban/suburban environments in southern Quebec and Ontario may pose substantial barriers. Additionally, dispersal rates for wolves in southern Ontario and Quebec appear to be relatively low and canids are harvested heavily in these regions, which will likely reduce the number of wolves successfully dispersing into New England (Wydeven et al. 1998). The ability of coyote hunters in the northeast to effectively discern wolves from coyotes in the field may also influence the likelihood of natural wolf recolonization. Recovery/reintroduction efforts are complicated by taxonomic uncertainty about the wolf or wolves that historically occupied the region, by public attitudes towards wolves, and by potential interactions with the eastern coyote. Regardless, populations of gray wolves, eastern wolves, and wolves of mixed ancestry in Ontario and Quebec are within plausible dispersal distance of Vermont (Wydeven et al. 1998, Fuller et al. 2003). Thus, it is possible that eastern and/or gray wolves enter Vermont periodically and the potential for natural recolonization of the state exists

Distribution

It is believed that wolves existed throughout Vermont prior to European settlement. This belief is supported by bounty records which clearly indicate the existence of wolves in nearly all biophysical regions of the state.

Distribution by Biophysical Region:

Champlain Valley	Historic Records Only	Southern VT Piedmont	Historic Records Only
Champlain Hills	Historic Records Only	Vermont Valley	Historic Records Only
Northern Green Mtns	Historic Records Only	Southern Green Mtns	Historic Records Only
Northern VT Piedmont	Historic Records Only	Taconic Mtns	Historic Records Only
Northeastern Highlands	Historic Records Only		

Distribution by Watershed:



Common Name: **Wolf**
Scientific Name: **Canis sp?**
Species Group: **Mammal**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Wolves are considered to be habitat generalists and usually select habitat to maximize predation success rather than for specific vegetation characteristics per se (e.g., Mech et al. 2003). Much of the suitable habitat for wolves in Vermont is likely forested, however wolves would be likely to occupy any patches of undeveloped terrestrial habitat that support adequate prey densities and where they are protected from human-caused mortality. Although wolves use a variety of habitat types across their range, they tend to occupy relatively contiguous patches of forests in remote areas with relatively low human and road densities (Mladenoff et al. 1995, Benson et al. 2012). Wolves requires an adequate prey base to persist. Deer, moose, and beaver would likely be the main prey for wolves in Vermont Mladenoff and Sickley (1998) and Harrison and Chapin (1998) estimated that approximately 53, 500 to 58, 500 km² of suitable habitat remains in northern New England. Mladenoff and Sickley (1998) further suggested that this habitat could support 702 to 1439 wolves. Harrison and Chapin (1998) suggested that 2470 km² and 1430 km² of suitable “core” and “dispersal” habitat, respectively, existed in Vermont based on road densities, human densities, and available forested habitat. Fuller et al. (2003) recommended that the smallest demographically viable wolf population might include 2-3 adjacent packs of 4 wolves each that were 40-60 km from other wolves. Thus, the 950 km² of suitable core habitat estimated to exist in Vermont might support approximately 8 packs of 4 wolves at average ungulate densities (8 deer/ km²) with wolf territories of approximately 300 km² (Fuller et al. 2003). Some of the core habitat identified by Harrison and Chapin (1998) is somewhat isolated in the central and southwestern portions of the state which might limit connectivity between patches. However, there is considerable evidence of wolves crossing highways and areas used intensively by humans in both Europe and North America (Merrill and Mech 2000, reviewed by Boitani 2003) suggesting that wolves might be able to successfully navigate the fragmented New England landscape. Mech (2006) found that Mladenoff and Sickely’s predictive model for wolf recolonization in Wisconsin (and potentially for the Northeast) failed to account for the wolf’s adaptability and capacity to colonize areas deemed <50% probable, including 22% of colonized areas with low probability. Additionally, Harrison and Chapin (1998) noted that much of the core habitat in Vermont is in the northeastern portion of the state and is contiguous with an expansive area of suitable habitat in New Hampshire, Maine, and Quebec meaning that wolves in Vermont could be connected with a larger regional population should recolonization occur. Territory size and density of wolves are strongly influenced by the availability of prey. Mean territory size is larger (>1000 km²) and smaller (< 200 km²) in areas with lower and greater prey densities, respectively (Mech and Boitani 2003, Fuller et al. 2003). Thus, the estimates for wolf numbers and territory sizes would likely shift depending on the local densities of deer and moose in areas of suitable habitat within Vermont. Regional corridors and habitat linkages are critical to maintaining wolves in potentially fragmented landscapes. Three important elements to wolf population viability are adequate prey, absence of excessive human exploitation, and relatively undeveloped blocks of habitat (Fritts and Carbyn 1995; Fuller 1997; Haight et al. 1998 in Parson 2003).



Common Name: **Wolf**
Scientific Name: **Canis sp?**
Species Group: **Mammal**

Habitat Types:

Upland Shores
Outcrops and Alpine
Cliffs and Talus
Hardwood Swamps
Softwood Swamps
Spruce Fir Northern Hardwood
Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Early Succession Boreal Conifers
Early Succession Boreal Hardwoods
Early Succession Pine and Hemlock
Early Succession Upland Oak
Early Succession Other Types
Wet Shores
Shrub Swamps
Early Succession Spruce-Fir
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Habitat Fragmentation
Impacts of Roads or Transportation Systems

Description of habitat threat(s): Human activity associated with roads, vehicles, and houses seem to negatively influence use of an area by wolves. Conversion of forest and other natural habitat to non-forest (development and agriculture) also negatively affects wolf densities. Wolves cannot survive without adequate prey, adequate protection, and adequate public support (Theberge et al, 1996 in Tumosa 2003). Connectivity with other wolf packs in the region is important to recovery of wolves in the northeast. Potential core habitat in southern Vermont appears to be disconnected from core habitat in northeastern Vermont (Harrison and Chapin 1998).

Non-Habitat Threats:

Genetics



Common Name: **Wolf**
Scientific Name: **Canis sp?**
Species Group: **Mammal**

Competition

Parasites

Harvest or Collection

Loss of Prey Base

Trampling or Direct Impacts

Description of non-habitat threat(s): Competition/hybridization with eastern coyotes may influence the probability of successful wolf recolonization of Vermont. Eastern wolves readily hybridize with eastern coyotes where they come into contact (e.g., Rutledge et al. 2010, Benson et al. 2012). Hybridization would likely be rampant in Vermont between recolonizing eastern wolves (which would be at low density) and coyotes (which would be much more abundant). Conversely, gray wolves and admixed gray wolves such as those inhabiting Minnesota, Wisconsin, and Michigan have not been documented to hybridize with coyotes in the wild (e.g., Wheeldon et al. 2010). Thus, dispersing gray wolves from Quebec and Ontario may have a higher probability of avoiding genetic swamping from eastern coyotes and establishing a viable population in Vermont. The eastern coyote is now the dominant large canid predator in the Northeast and it is not clear how the existing coyote population would respond to the establishment of a wolf population. A better understanding of the ecological role of the eastern coyote in Vermont would help clarify the extent to which these smaller canids are able to occupy the ecological niche of wolves.

Thiel (1985) found that when wolves were persecuted by humans in Wisconsin populations did not persist where road densities exceeded approximately 1km/km². However, with sufficient protection from human-caused mortality wolves have been documented persisting at road densities greater than 1km/km² as public attitudes about wolves shifted (Mech 1989, Fuller et al. 1992, reviewed in Fuller et al. 2003). Thus, protection from hunting and trapping mortality may facilitate viable wolf populations in fragmented habitat with higher human population and road densities.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	Medium	Document and map the distribution of large wild canids based on DNA analysis.
Research	Population Genetics	High	Determine the genetics of large wild canids in Vermont and monitor wolf colonization events.
Research	Taxonomy	High	Determine the species of wolf historically found in Vermont.
Research	Other Research	High	Determine public attitudes towards wolves in Vermont and New England
Monitoring	Other Monitoring Needs	High	Monitor wolf colonization events

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Wolf**
Scientific Name: **Canis sp?**
Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications	High	Determine public attitudes towards wolf recovery possibly by partnering with University researchers to conduct a rigorous evaluation of public opinions.		NWF, Keeping Track, Sportsmen's Federation, University researcher	USFWS, SWG
Policy & Regulations	Medium	Develop statewide protocol to guide state/federal wildlife management actions in response to wolf immigration. Results of the species restoration strategy may provide information that can be used to reevaluate the rank for this strategy in the future.	Degree to which partners adopt the protocols	USFWS, USFS, NWF, VTFSC, Agency of Agriculture, NRCS, Farm Bureau, RPCs, Law Enforcement	USFWS, SWG
Species Restoration	High	Evaluate VT large canid ancestry via DNA analysis/ morphology to monitor possible recolonization. Obtain tissue samples and morphological measurements from large canids trapped, shot, hit by cars, or otherwise observed in VT.		NWF, Keeping Track, Sportsmen's Federation, VTA,	NWF, USFWS, SWG



Common Name: **Wolf**
Scientific Name: **Canis sp?**
Species Group: **Mammal**

Bibliography

- Benson, J.F., B.R. Patterson, and T.J. Wheeldon. 2012. Spatial genetic and morphologic structure of wolves and coyotes in relation to environmental heterogeneity in a *Canis* hybrid zone. *Molecular Ecology* 21:5934-5954.
- Boitani, L. 2003. Wolf conservation and recovery. In: Mech, L.D., and Boitani, L. (ed.), *Wolves: Ecology, Behavior, and Conservation*. University of Chicago Press, pp. 317-340.
- Fascione, N., L.G.L. Osborn, S.R. Kendrot, and P.C. Paquet. 2001. *Canis soupus*: Eastern wolf genetics and its implications for wolf recovery in the northeast United States. *Endangered Species UPDATE* 18(4):159-163
- Fritts, S. H. and L. N. Carbyn. 1995. Population viability, nature reserves, and the outlook for gray wolf conservation in North America. *Restoration Ecology* 3(1):26-38.
- Fuller, T.K., W.E. Berg, G.L. Radde, M.S. Lenarz, and G.B. Joselyn. 1992. A history and current estimate of wolf distribution and numbers in Minnesota. *Wildlife Society Bulletin* 20:42-55.
- Fuller, T. K. 1997. Guidelines for gray wolf management in the northern Great Lakes Region . Second Edition. Ely, MN: International Wolf Center.
- Fuller, T.K., L.D. Mech, and J.F. Cochrane. 2003. Wolf population dynamics. In: Mech, L.D., and Boitani, L. (ed.), *Wolves: Ecology, Behavior, and Conservation*. University of Chicago Press, pp. 161-191.
- Haight, R. G., D. J. Mladenoff, and A. P. Wydeven. 1998. Modeling disjunct gray wolf populations in semi-wild landscapes. *Conservation Biology* 12(4):879-888
- Harrison, D.J. and Chapin, t.G. 1998. Extent and connectivity of habitat of wolves in eastern North America. *Wildlife Society Bulletin* 26: 767-775.
- Kobl Müller, S., M. Nord, R.K. Wayne, and J.A. Leonard. 2009. Status and origin of the Great Lakes wolf. *Moelcular Ecology* 18:2313-2326.
- Mech, L.D. 1995a. The challenge and opportunity of recovering wolf populations. *Conservation Biology*. 9(2):270-278 in Parsons, D.R. 2003. Natural History Characteristics of Focal Species in the New Mexico Highlands Wildlands Network. *Wildlands Project*. 69 pp.
- Mech, L.D. 1974. Mammalian species, *Canis lupus*. The American Society of Mammalogists 37: 1-6 in (Tumosa, J. 2003. Green Mountain National Forest species data collection form for *Canis lupus*.38 pp.)
- Mech, L.D. 1989. Wolf population survival in an areas of high road density. *American Midland Naturalist* 121:387-389.
- Mech, L.D., and Boitani, L. 2003. Wolf social Ecology. In: Mech, L.D., and Boitani, L. (ed.), *Wolves: Ecology, Behavior, and Conservation*. University of Chicago Press, pp. 1-34.
- Mech, L.D. 2006. Prediction Failure of a Wolf Landscape Model. *Wildlife Society Bulletin* 34(3): 874-877.
- Merrill, S.B., and L.D. Mech. 2000. Details of extensive movements by Minnesota wolves. *American Midland Naturalist* 144: 428-33.
- Mladenoff, D.J., T.A. Sickely, R.G. Haight, and A.P. Wydeven. 1995. A regional landscape analysis and prediction of favorable gray wolf habitat in the northern Great Lakes region. *Conservation Biology* 9:279-294.
- Mladenoff, D.J., and Sickley, T.A. 1998. Assessing potential gray wolf restoration in the northeastern United States: a spatial prediction of favorable habitat and potential population levels. *Journal of Wildlife Management* 62: 1-10.
- Rutledge, L.Y., C.J. Garroway, K.M. Loveless, and B.R. Patterson. 2010. Genetic differentiation of eastern wolves in Algonquin Park despite bridging gene flow between coyotes and grey wolves. *Heredity* 105: 520-531.
- Therberge, J. B., Therberge, M.T., and Forbes, G. 1996. What Algonquin Park wolf research has to instruct about recovery in northeastern United States. In: *Proceedings Defenders of Wildlife's wolves of America Conference*. 14-16 November 1996, Albany, New York.in (Tumosa, J. 2003. Green Mountain National Forest species data collection form for *Canis lupus*.38 pp.)
- Thiel, D. 1985. Relationship between road density and wolf habitat suitability in Wisconsin. *American Midland Naturalist* 113:404-407.
- Wheeldon, T. J., B.R. Patterson, and B.N. White. 2010. Sympatric wolf and coyote populations of the western Great Lakes region are reproductively isolated. *Molecular Ecology* 10/2010: 4428-4440.
- Wilson P.J., Grewal S., Lawford I.D., Heal J.N.M., Granacki A.G., Pennock D., Theberge J.D., Theberge M.T., Voigt D.R., Waddell W., Chambers R.E., Paquet P.C., Goulet G., Cluff D., and White B.N. 2000 DNA profiles of the eastern Canadian wolf and the red wolf provide evidence for a common evolutionary history independent of the gray wolf *Can. J. Zool.* 78: 2156–2166
- Woods, C. A. 1973. *Erethizon dorsatum*. *Mammalian Species*, 29:1-6.
- Wilson, P.J., S. Grewal, T. McFadden, R.C. Chambers, and B.N. White. 2003. Mitochondrial DNA extracted from eastern North American wolves killed in the 1800s is not of gray wolf origin. *Canadian Journal of Zoology* 81:936-940.
- Wydeven, A.P., T.K. Fuller, W. Weber, and K. MacDonald. 1998. The potential for wolf recovery in the northeastern United States via dispersal from southeastern Canada. *Wildlife Society Bulletin* 26:776-784



Common Name: **Gray Fox**
Scientific Name: **Urocyon cinereoargenteus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5

State Trend: Stable

Extirpated in VT? No

Regional SGCN? no

Assessment Narrative:

Gray foxes are widespread throughout Vermont and occupy most major habitat types including forests, shrublands, agricultural areas, and the margins of urban environments. Despite being relatively common, little is known about basic characteristics of the species in the state, including distribution, demographics, diet and space use behavior, and interactions with other species. Similarly, little is known about threats facing the species. Gray foxes elsewhere are negatively impacted by competition from larger carnivores such as red foxes, coyotes, and bobcats, and diseases such as rabies and canine distemper. Gray foxes also appear to be expanding their range northward into Quebec.

Distribution

Gray Fox harvest records indicate a widespread distribution of the species in Vermont with records of occurrence in all biophysical regions.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Few studies have been undertaken on gray foxes regionally in New England and even range-wide, despite their widespread distribution and perception as a common species (Fuller and Cypher 2004). Studies elsewhere indicate that foxes occur in densities that range from 0.4/km² (California) to 1.5/km² (Florida), and that foxes occupy home ranges that vary in size from 75 ha (West Virginia) to 653 ha (Alabama) (Fritzell and Haroldson 1982, Fuller and Cypher 2004). In Vermont, a radio-telemetry study indicated that average gray fox home range size was 4.43 km² in the Champlain Valley (n=5, 2 females/3 males, Ingle 1990). Gray foxes in this study occurred primarily in hardwood forested areas and avoided open habitats. Basic demographic estimates, such as density and population size, and home range/habitat use characteristics have not been adequately quantified in Vermont. Gray foxes elsewhere associate mainly with deciduous forest, but use other forest types, shrublands, agricultural lands, fields, and farmlands, and the margins of urban environments (Fritzell and Haroldson 1982). They typically use successional forests, habitat mosaics and managed woodlands.



Common Name: **Gray Fox**
Scientific Name: **Urocyon cinereoargenteus**
Species Group: **Mammal**

Habitat Types:

Northern Hardwood
Hardwood Swamps
Softwood Swamps
Seeps and Pools
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Early Succession Boreal Conifers
Early Succession Pine and Hemlock
Early Succession Upland Oak
Early Succession Other Types
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Wet Shores
Shrub Swamps
Early Succession Boreal Hardwoods
Early Succession Spruce-Fir
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Habitat Alteration

Unknown Habitat Threats

Description of habitat threat(s): Unknown, but distribution and abundance appears to be linked to forest habitats. Changes in forest cover, especially deciduous forest, due to development (e.g., residential housing, roads, urban expansion) may impact the species in Vermont.

Non-Habitat Threats:

Competition

Disease

Description of non-habitat threat(s): Competition and mortality from coyotes, bobcats, and red foxes represent potential threats. These three species have been shown to compete with gray foxes elsewhere (Chamberlain and Leopold 2005, Farias et al. 2005), and may negatively impact the species in Vermont. Diseases such as rabies and distemper represents another potential concern (Fuller and Cypher 2004).



Common Name: **Gray Fox**
Scientific Name: **Urocyon cinereoargenteus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5

State Trend: Stable

Extirpated in VT? No

Regional SGCN? no

Assessment Narrative:

Gray foxes are widespread throughout Vermont and occupy most major habitat types including forests, shrublands, agricultural areas, and the margins of urban environments. Despite being relatively common, little is known about basic characteristics of the species in the state, including distribution, demographics, diet and space use behavior, and interactions with other species. Similarly, little is known about threats facing the species. Gray foxes elsewhere are negatively impacted by competition from larger carnivores such as red foxes, coyotes, and bobcats, and diseases such as rabies and canine distemper. Gray foxes also appear to be expanding their range northward into Quebec.

Distribution

Gray Fox harvest records indicate a widespread distribution of the species in Vermont with records of occurrence in all biophysical regions.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Few studies have been undertaken on gray foxes regionally in New England and even range-wide, despite their widespread distribution and perception as a common species (Fuller and Cypher 2004). Studies elsewhere indicate that foxes occur in densities that range from 0.4/km² (California) to 1.5/km² (Florida), and that foxes occupy home ranges that vary in size from 75 ha (West Virginia) to 653 ha (Alabama) (Fritzell and Haroldson 1982, Fuller and Cypher 2004). In Vermont, a radio-telemetry study indicated that average gray fox home range size was 4.43 km² in the Champlain Valley (n=5, 2 females/3 males, Ingle 1990). Gray foxes in this study occurred primarily in hardwood forested areas and avoided open habitats. Basic demographic estimates, such as density and population size, and home range/habitat use characteristics have not been adequately quantified in Vermont. Gray foxes elsewhere associate mainly with deciduous forest, but use other forest types, shrublands, agricultural lands, fields, and farmlands, and the margins of urban environments (Fritzell and Haroldson 1982). They typically use successional forests, habitat mosaics and managed woodlands.



Common Name: **Gray Fox**
Scientific Name: **Urocyon cinereoargenteus**
Species Group: **Mammal**

Habitat Types:

Northern Hardwood
Hardwood Swamps
Softwood Swamps
Seeps and Pools
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Early Succession Boreal Conifers
Early Succession Pine and Hemlock
Early Succession Upland Oak
Early Succession Other Types
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Wet Shores
Shrub Swamps
Early Succession Boreal Hardwoods
Early Succession Spruce-Fir
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Habitat Alteration

Unknown Habitat Threats

Description of habitat threat(s): Unknown, but distribution and abundance appears to be linked to forest habitats. Changes in forest cover, especially deciduous forest, due to development (e.g., residential housing, roads, urban expansion) may impact the species in Vermont.

Non-Habitat Threats:

Competition

Disease

Description of non-habitat threat(s): Competition and mortality from coyotes, bobcats, and red foxes represent potential threats. These three species have been shown to compete with gray foxes elsewhere (Chamberlain and Leopold 2005, Farias et al. 2005), and may negatively impact the species in Vermont. Diseases such as rabies and distemper represents another potential concern (Fuller and Cypher 2004).

**Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report**



Common Name: **Gray Fox**
Scientific Name: **Urocyon cinereoargenteus**
Species Group: **Mammal**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Medium	Identify important habitats and quantify patterns of habitat selection.
Research	Basic Life History	Medium	Estimate home range characteristics.
Research	Distribution and Abundance	High	Refine distribution and abundance data.
Research	Threats and Their Significance	Medium	1) Examine how habitat alteration impacts distribution and abundance. 2) Determine the effects of zoonotic diseases (distemper and rabies). 3) Determine effects of competition with coyotes and other sympatric carnivores such as fisher.
Monitoring	Range Shifts	Medium	Determine possible range shifts and population changes due to climate change.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Compatible Resource Use	High	Promote less development of high quality habitats.	Amount of high quality habitat protected or conserved	VTrans, Town Planning Commission, Town and Regional Cons Comms, VLT, Keeping Track	SWG, Vtrans



Common Name: **Gray Fox**
Scientific Name: ***Urocyon cinereoargenteus***
Species Group: **Mammal**

Bibliography

Bozarth, C. A., S. L. Lance, D. J. Civitello, J. L. Glenn, and J. E. Maldonado. 2011. Phylogeography of the gray fox (*Urocyon cinereoargenteus*) in the eastern United States. *Journal of Mammalogy* 92:283-294.

Chamberlain, M. J., and B. D. Leopold. 2005. Overlap in space use among bobcats (*Lynx rufus*), coyotes (*Canis latrans*) and gray foxes (*Urocyon cinereoargenteus*). *American Midland Naturalist*:171-179.

Cypher, B. L. 2003. Foxes (*Vulpes* species, *Urocyon* species, and *Alopex lagopus*). Pages 511-546 in G. A. Feldhamer, B. C. Thompson, and J. A. Chapman (eds.) *Wild mammals of North America biology, management, and conservation*, 2nd edition. Johns Hopkins University Press, Baltimore, Maryland, USA.

Farias, V., T. K. Fuller, R. K. Wayne, and R. M. Sauvajot. 2005. Survival and cause-specific mortality of gray foxes (*Urocyon cinereoargenteus*) in southern California. *Journal of Zoology* 266:249-254.

Frizzell, E. K., and K. J. Haroldson. 1982. *Urocyon cinereoargenteus*. *Mammalian Species* 189:1-8.

Fuller, T. K., and B. L. Cypher. 2004. Gray fox *Urocyon cinereoargenteus* (Schreber, 1775). Pages 92-97 in C. Sillero-Zubiri, M. Hoffmann, and D. W. Macdonald (eds.), *Canids: foxes, wolves, jackals and dogs. Status survey and conservation action plan*, IUCN/SSC Canid Specialist Group, Gland, Switzerland and Cambridge, United Kingdom.

Ingle, M. A. 1990. Ecology of red foxes and gray foxes and spatial relationships with coyotes in an agricultural region of Vermont. Master of Science thesis, University of Vermont, Burlington, Vermont, USA.

Mcalpine, D., J. D. Martin, and C. Libby. 2008. First occurrence of the grey fox, *Urocyon cinereoargenteus*, in New Brunswick: a climate-change mediate range expansion? *Canadian Field-Naturalist* 122-169-171.



Common Name: **Black Bear**
 Scientific Name: **Ursus americanus**
 Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5

State Trend: Stable

Extirpated in VT? no

Regional SGCN? no

Assessment Narrative:

Currently, the black bear population is stable and found through out Vermont. Certain habitat types are critical to the sustainability of Vermont's black bear population.

Critical habitat (including mast stands, forested wetlands, and travel corridors) could be vulnerable due to habitat loss and conversion from development and fragmentation. Black bear depends on large expanses of forested blocks with connectivity that could be lost due to development and fragmentation. It is already possible that Vermont has two distinct genetic populations due to fragmentation with the southern population below critical levels.

Distribution

Distribution is statewide. Higher densities are found along the Green Mountains and in the Northeast Highlands and Northern VT Piedmont.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills		Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Eastern deciduous forest, mast producing hardwood forests, (beech, birch maple and oak hickory). Large blocks of forest with large stands of mast producing trees (.e.g. beech, oak and cherry). Uses forested wetlands, released apple orchards. Requires connectivity of habitat. Micro Habitat: mast stands, wetlands, rich wet sites



Common Name: **Black Bear**
Scientific Name: **Ursus americanus**
Species Group: **Mammal**

Habitat Types:

Outcrops and Alpine
Northern Hardwood
Softwood Swamps
Seeps and Pools
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests
Hardwood Swamps
Open Peatlands
Marshes and Sedge Meadows
Early Succession Boreal Conifers
Early Succession Pine and Hemlock
Early Succession Upland Oak
Early Succession Other Types
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Lawns, Gardens, and Row Crops
Wet Shores
Shrub Swamps
Early Succession Boreal Hardwoods
Early Succession Spruce-Fir
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Conversion of Habitat
Energy Infrastructure and Development
Impacts of Roads or Transportation Systems
Habitat Fragmentation

Description of habitat threat(s): Food is the most important resource for black bears. Nuts of American beech trees are an important primary fall food source in Northern New England. Nutritional plane of female bears and their reproductive rates would decline substantially if American beech trees diminish significantly in the forests of Vermont. Forest management practices, disease events and weather events may affect stands of mast producing trees.

Description of non-habitat threat(s):

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Black Bear**
 Scientific Name: **Ursus americanus**
 Species Group: **Mammal**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Conduct comprehensive critical habitat inventory including location, and condition of mast stands and monitor changes through time. Develop a universal wildlife habitat management unit meaningful for basic life history requirements of black bears.
Research	Threats and Their Significance	Medium	Examine impacts and avoidance of commercial industrial wind energy generation in high elevation areas, and effects to denning behavior and sounds associated with operations of these industrial facilities.
Monitoring	Habitat Change	High	Examine long term effects of ice storm damage to high elevation beech stands, examine tree disease impacts on health and vigor of mast producing beech and oak stands. Examine trends in beech and oak stand health and distribution.
Monitoring	Monitor Threats	High	Examine highway development and improvements related to bear genetics statewide.
Monitoring	Other Monitoring Needs	Medium	Monitor other elements as identified in the statewide management plan for this species (e.g. sale of black bear parts, specific harvest parameters, monitor trends in large forest block patterns and connectivity).

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Policy & Regulations	High	Maintain viable bear population through regulating harvest and managing bear habitat.			
Technical Assistance, Training, Learning Networks	Medium	Collaborate with Agency of Transportation regarding highway placement and passage of wildlife.			
Technical Assistance, Training, Learning Networks	High	Develop information for private land owners for managing necessary habitat.			
Technical Assistance, Training, Learning Networks	Medium	Provide information to towns and regional planning commissions regarding habitat requirements.		Agency of Transportation	PR, FHWA, SWG
Compatible Resource Use	High	Conserve contiguous blocks of remote interior habitat and the linkages between them.			
Standards	High	Manage necessary habitats on publicly owned land.			



Common Name: **Black Bear**
Scientific Name: **Ursus americanus**
Species Group: **Mammal**

Bibliography

Goodrich, J.M., and J. Berger. 1994. Winter recreation and hibernation black bear *Ursus americanus*. *Biological Conservation*, 67:105-110.

Hammond, F.M. 2002. The Effects of Resort and Residential Development on Black Bears in Vermont. Final Report. Vermont Department of Fish and Wildlife. 114pp.

Larriere, S. 2001. *Ursus americanus*. *Mammalian Species*, 647:1-11.

McLaughlin, C.R. 1998. Modeling effects of food and harvests on female black bear populations. PhD Dissertation. Univ. of Maine. 263pp.

Kolenosky, G.B. and S.M. Strathearn 1987. Black Bear. In M. Novak, J.A. Baker, M.E. Obbard, and B. Malloch, eds. *Wild furbearer management and conservation in North America*. Ontario Trappers Assoc., North Bay.

Litvaitis, John A. 2001 Importance of Early Successional Habitat to Mammals in Eastern Forests. *Wildlife Society Bulletin* 29(2):466-473.

Elowe, K.D. 1984. Home range, movements, and habitat preferences of black bear in western Massachusetts M.S. Thesis. University of Mass. Amherst. 112pp.

Elowe, K.D. 1987 Factors affecting black bear reproductive success and cub survival in Massachusetts. Ph.D. Dissertation. University of Mass. Amherst, 71pp.

Elowe, K.D. and W. E. Dodge. 1989 Factors affecting black bear reproductive success and cub survival. *J. Wildl. Manage.* 53:962-968.

Hugie, R.D. Black bear ecology and management in northern conifer-deciduous forests of Maine. Ph.D. Thesis Univer. Montana, Missoula, 202pp.

Eiler, J.H. 1981. Reproductive biology of black bears in the Great Smoky Mountains of Tennessee. M.S. Thesis., Univ. Tennessee, Knoxville. 128pp.

Eiler, J.H., W.G. Wathen and M.R. Pelton. 1989. Reproduction in Black Bears in the Southern Appalachian Mountains. *J. Wildl. Manage.* 53:353-360.

Beecham, J.J. 1980. Population characteristics, denning and growth patterns of black bears in Idaho. Ph.D. Thesis, Univ. Montana Missoula. 101pp.

Johnkel, C.J. and I.M. Cowan. 1971 The black bear in the spruce-fir forest. *Wildl. Monograph*. 27.57pp.

Pelton, M.R. 1982. Black Bear, Pages 504-514 in J.A. Champan and G.A. Feldhamer, eds. *Wild Mammals of North America*. The Johns Hopkins Univ. Press, Baltimore, Md.

Schooley, R.L. 1990. Habitat use, fall movements, and denning ecology of female black bears in Maine. M.S. Thesis, Univ. Maine, Orono. 115pp.

Schooley, R.L., C.R. McLaughlin, G.J. Matula, Jr., and W.B. Krohn. 1994. Denning Chronology of female black bears: effects of food, weather, and reproduction. *Journal of Mammalogy* 75: 466-477.

Wiley, C.H. 1978. The Vermont black bear. Vermont Fish and Wildlife Dept., Montpelier, Vt. 73pp.



Common Name: **American Marten**
 Scientific Name: **Martes americana**
 Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1

State Trend: Increasing

Extirpated in VT? Unknown

Regional SGCN? Yes

Assessment Narrative:

Despite having been previously extirpated from Vermont, recent evidence indicates the presence of two distinct populations of American marten in the state (VFWD unpublished data). Although little is known regarding their full extent and distribution, these populations are likely at risk due to their presumed small size and limited distribution. Relative to most other forest-associated mammals, marten have large spatial requirements, low population densities and specific habitat needs (Buskirk and Ruggerio 1994) making populations particularly vulnerable to factors influencing habitat suitability. Forest management practices that fail to consider marten habitat requirements, for example, may result in a decrease in marten density and productivity over the landscape (Gosse et al. 2005, Payer and Harrison 1999, Johnson et al. 2009, Fuller and Harrison 2005). Furthermore, interspecific relations with sympatric carnivores such as fisher and red fox are widely hypothesized to be limiting factors for marten population recovery and expansion (Krohn et al 2004, Siren 2009). Vermont furbearer harvest data indicate widespread and abundant populations of many competing carnivores throughout the state (VFWD unpublished data). Last, the strong correlation between marten occurrence and the annual accumulation of suitable snow depths makes the persistence of this species in Vermont vulnerable to changes in the climate (Krohn 2012, Kelly 2005, Siren 2009, and Carroll 2007).

Distribution

Although believed to have occurred throughout the state prior to European contact, American marten were extirpated from Vermont in the 1800's due to excessive land clearing and unregulated trapping. Since 2000, a total of 25 marten occurrences have been confirmed in Vermont (VFWD unpublished data). The majority of these were reported from the northeast corner of the state in Essex (13), Caledonia (4) and Orleans (1) counties. The remaining marten were reported from the high elevation towns of the southern Green Mountains in Bennington (4) and Windham (3) counties. Additionally, remote camera surveys conducted in 2012 documented the occurrence of two individual marten in the town of Sunderland (Bennington County).

Distribution by Biophysical Region:

Champlain Valley	Not Probable	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Probable	Southern Green Mtns	Confident
Northern VT Piedmont	Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Confident		

Distribution by Watershed:



Common Name: **American Marten**
Scientific Name: **Martes americana**
Species Group: **Mammal**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

In general, American marten are associated with forested habitats that provide overhead cover and complex physical structure near the forest floor (Payer and Harrison 2003, Andruskiw 2008, Godbout and Ouellet 2010). Although these forest characteristics are most closely associated with older seral stages, the use of younger, managed forests by marten has also been well documented where previous harvesting practices have favored the retention of coarse woody debris, and have maintained residual basal areas greater than 18m²/ha and at least a 30% canopy closure in winter (Thompson et al 2012, Payer and Harrison 2003, Fuller and Harrison 2005). In the northeast, suitable marten habitat is provided by a wide range of forest types including mixed coniferous-deciduous forests and forests dominated by deciduous trees (Kelly 2005, Payer and Harrison 1999). Marten avoid open areas such as those occurring naturally on the landscape (e.g. wetland meadows and stands recently disturbed by fire, Gosse et al. 2005) and those resulting from human activities (e.g. clearcutting and infrastructure development; Payer and Harrison 1999, Siren 2009). Jensen et al (2012) documented a significant demographic response of the marten population to fluctuations in annual mast crop production indicating the importance of mast producing trees as a component of suitable marten habitat. Several studies have documented a close association of annual snow fall rates and occupied marten habitat suggesting a strong preference for deep snow where certain morphological adaptations may give marten competitive advantages over sympatric carnivores (Krohn 2004, Kelly 2005, Carroll 2007).

Habitat Types:

Softwood Swamps
Spruce Fir Northern Hardwood
Northern Hardwood

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Climate Change
Habitat Fragmentation

Description of habitat threat(s): Because American marten life history is strongly influenced by adult survival (Buskirk et al. 2012), the recovery and growth of Vermont's marten populations will require favorable environmental conditions over long periods of time. Thus, habitat stochasticity resulting from anticipated changes in the climate (Carroll 2007, Krohn 2012, Kelly 2005), the broadscale implementation of forest management practices that do not adequately account for marten habitat requirements (Thompson et al. 2012, Fuller and Harrison 2005, Payer and Harrison 2003), and further fragmentation of the landscape (Siren 2009) jeopardizes the persistence of marten in the state.

Non-Habitat Threats:

Competition
Disease



Common Name: **American Marten**
 Scientific Name: **Martes americana**
 Species Group: **Mammal**

Harvest or Collection

Predation or Herbivory

Loss of Prey Base

Description of non-habitat threat(s): Competition with, and predation by, sympatric carnivores such as fisher and red fox could negatively influence the distribution and persistence of marten in Vermont (Krohn et al 1995, Kelly 2005, Siren 2009). The effects of climate change will likely exacerbate the adverse impact of interspecific completion on marten as carnivore communities shift northward into marten range and the species' competitive advantages are diminished as a result of lower snowfall accumulations (Carroll 2007, Krohn et al. 2005). Although the incidental take of marten in fisher traps has been documented in Vermont (VFWD unpublished data), it is not currently known to be a limiting factor of the marten population. In fact, the continued harvest and management of competing carnivores could prove to be an overall benefit to marten despite this infrequent take. Although difficult to assess, the impacts of unregulated take and interspecific competition need to be considered where the maintenance of marten populations is a priority.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Perform a habitat suitability analysis of Vermont in order to identify key marten habitats, to help predict distribution of the speices in the state and to facilitate the developemnt of appropriate conservation actions.
Research	Distribution and Abundance	High	Collect baseline data on marten distribution and abundance in Vermont in order to assess the status of the population and develop appropriate conservation strategies.
Research	Threats and Their Significance	Medium	Examine the affects of interspecific competition with fisher and assess how certain habitat features, fisher harvests and snow depths influence this relationship.
Research	Population Genetics	High	Conduct a genetic analysis of marten in Vermont in order to determine the source of the species in the state, particularly of the southern population.
Research	Other Research	High	Assess the effectiveness and practicality of various trap configurations and trapping techniques for minimizing the incidental take of marten.
Monitoring	Population Change	Medium	Develop and implement a plan for monitoring the marten population in Vermont.
Monitoring	Habitat Change	Medium	Develop and implement a plan for monitoring changes in suitable marten habitat resulting from habitat conversions, forest management practices and climate change.
Monitoring	Range Shifts	Low	Monitor range shifts of competing carnivore populations resulting from climate change.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Marten**
 Scientific Name: **Martes americana**
 Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Policy & Regulations	Medium	Support and cooperate with larger efforts to curb global climate change.			
Compatible Resource Use	High	Promote forest management practices that provide for the life history requirements of marten	Number of acres of forest land positively influenced	Coverts, UVA, USFS, Industrial forest landowners, VFPR, VLT landowners, Forest Legacy landowners	
Standards	Medium	Develop best management practices for forest management within key marten habitats	The successful development and subsequent dissemination of best management practices	UVM, VFPR	SWG
Standards	High	Develop best management practices for fisher trapping in order to minimize incidental take of marten	Number of trappers employing best management practices and the number of marten taken	Vermont Trappers Association, AFWA, NHFG, MDIFW, NY DEC	Vermont Trappers Association, AFWA, SWG
Compatible Resource Use	High	Continue managing competing carnivores within key marten habitats, particularly fisher, via regulated trapping	Maintenance of healthy furbearer populations	Vermont Trappers Association	
Standards	Medium	Develop and implement guidelines and mitigation strategies for minimizing impacts to key marten habitats from regulated land use activities such as the development of energy infrastructure	Number of acres protected from conversion		

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **American Marten**
Scientific Name: **Martes americana**
Species Group: **Mammal**

Bibliography

Bissonette, J.A. , D.J. Harrison, , C.D. Hargis, T.G. Chapin. 1997. The influence of spatial scale and scale-sensitive properties on habitat selection by American marten. In *wildlife and Landscape Ecology*, ed. J.A. Bissonette. New York, NY: Springer.

Chapin, T.G., D.J. Harrison, D.D. Katnik. 1996. Influence of Landscape Pattern, Forest Type, and Forest Structure on Use of Habitat by Marten in Maine. 78pp.

DiStefano, J.J., K. Royar, D. Pence, J.E. Denoncour. 1990. Marten Recovery Plan for Vermont. 19pp.

Krohn, W. B., K. D. Elowe, and R. B. Boone. 1995. Relations between fishers, snowfall, and martens. *Forestry Chronical*, 71:97-105.

Novak M., J.A. Baker, M.E. Obbard, and B. Malloch, eds. *Wild furbearer management and conservation in North America*. Ontario Trappers Assoc., North Bay.

Thompson, Zadock. 1853. *Natural History of Vermont*. Charles E. Tuttle. Rutland, Vermont. 286 pp.

Trombulak, S.C. and K. Royar. 2000. "Restoring the Wild: Species Recovery and Reintroduction," in *Wilderness Comes Home: Rewilding the Northeast*, edited by Christopher McGrory Klyza, Hanover, N.H. University Press of New England.



Common Name: **Long-tailed Weasel**
Scientific Name: **Mustela frenata**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S3S4

State Trend: unknown

Extirpated in VT? no

Regional SGCN? no

Assessment Narrative:

The distribution and abundance of long-tailed weasels in Vermont are poorly understood and no records of their occurrence were collected during a statewide small mammal survey between 2008 and 2010 (Kilpatrick and Benoit, 2011). Although the extent to which these factors influence the population is poorly understood, the species is vulnerable to current pest control practices and could be potentially impacted by the application of pesticides.

Distribution

Only 22 verified records of the long-tailed weasel are available for Vermont but these confirm a wide spread distribution of this species in Orleans, Essex, Chittenden, Caledonia, Addison, Rutland, Windsor and Bennington counties. No additional records of their occurrence were collected during a state wide small mammal survey between 2008 and 2010 (Kilpatrick and Benoit, 2011)

Distribution by Biophysical Region:

Champlain Valley	Probable	Southern VT Piedmont	Confident
Champlain Hills	Probable	Vermont Valley	Probable
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Probable	Taconic Mtns	Probable
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

The long-tailed weasel inhabits the broadest range of any of the weasels from low elevations to above treeline across the continent (Novak et al, 1987). They occupy a variety of habitats from forest and shrubs adjacent to stone walls to fields, wetlands and standing water. Where it overlaps with the short-tailed weasel, it may occupy more open habitats while the short-tailed weasel is more common in forested or wetland areas. Areas with high prey density are important. The long-tailed weasel feeds on small mammals such as mice, rabbits, voles and ground nesting birds. Water seems to be a critical factor. Hamilton (1933) reported that they can drink 25cc of water per day and therefore, it may be restricted to habitats in close proximity to standing water. The long-tailed weasel is more of a food generalist than the short-tailed weasel. On average, long-tailed weasels will take 1.5 voles per day (Powell 1973 in Wild Furbearer Mgt 1987). The weasel uses excavated burrows or holes and/or crevices for den sites (DeGraff and Yamasaki, 2001).



Common Name: Long-tailed Weasel
Scientific Name: *Mustela frenata*
Species Group: Mammal

Habitat Types:

Northern Hardwood
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Marshes and Sedge Meadows
Early Succession Upland Oak
Grasslands, Hedgerows, Old Field, Shrub, or Orchard
Wet Shores
Early Succession Boreal Hardwoods
Early Succession Northern Hardwoods
Aquatic: Man-Made Water Bodies

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration

Description of habitat threat(s): Although the full extent and nature of impacts are poorly understood, it is suspected that the conversion of habitat via natural succession or anthropogenic degradation could negatively affect weasel populations.

Non-Habitat Threats:

Predation or Herbivory

Description of non-habitat threat(s): Predation on long-tailed weasels by domestic pets, foxes and raptors could be a factor limiting the distribution and abundance of this species. Similarly, when existing in close proximity to humans, exposure to pest control practices and potential for road kill may be a problem. Weasels could be affected directly and/or indirectly by pesticide use.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	High	Determine abundance, distribution, and status of the Vermont population.
Research	Threats and Their Significance	Medium	Examine how current pest control practices, including the use of pesticides, influence long-tailed weasel populations.
Research	Other Research	Low	Examine how predation, particularly by domestic pets, influences long-tailed weasel populations.
Monitoring	Population Change	Medium	Develop and implement a plan for monitoring the long-tailed weasel population in Vermont.
Monitoring	Habitat Change	Medium	Examine how forest succession and anthropogenic changes of the landscape influence long-tailed weasel populations.



Common Name: **Long-tailed Weasel**
 Scientific Name: **Mustela frenata**
 Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Awareness Raising and Communications	Medium	Develop outreach materials informing the public of the importance of keeping domestic pets under control	Development and dissemination of outreach materials		
Standards	Medium	Develop Best Management Practices for pest control professionals and landowners to follow for minimizing damage by and lethal control of long-tailed weasels	Development and dissemination of BMPs	NWCOs, Pest Control Professionals	SWG

Bibliography

- Degraaf, R.M., and Yamasaki, M. 2001. New England Wildlife: Habitat, Natural History, and Distribution, University Press of New England, Hanover and London.
- Hamilton, W.J., Jr. 1933. The weasels of New York: their natural history and economic status, American Midland Naturalist, 14:289-344.
- Novak, M., J.A. Baker, M.E. Obbard, and B. Malloch. Wild Furbearer Management and Conservation in North America. Ontario Trappers Association. 1150pp.
- Sheffield, S.R., and H.H. Thomas. 1997. *Mustela frenata*. Mammalian species, 570: 1-9
- Simm, D.A. 1979. North American weasels: resource utilization and distribution. Canadian Journal of Zoology, 57:504-520.
- Whitacker, J.O. and W.J. Hamilton, Jr. 1998. Mammals of the Eastern United States. Comstock Publishing, Ithaca, N.Y.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Northern River Otter**
Scientific Name: **Lontra canadensis**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5

State Trend: stable

Extirpated in VT? no

Regional SGCN? no

Assessment Narrative:

The Northern River Otter's piscivorous diet and high trophic position make it a noteworthy indicator of pollution in aquatic systems (Melquist and Dronkert 1987). Of 20 otter tested for mercury in Vermont in 2001, for example, two had levels higher than that recommended by the EPA (K. Royar, pers. Com). Prey may also be susceptible to pollution and acid rain. Because of their strict aquatic nature, otter populations are susceptible to changes in riverine and lacustrine habitats which alter the physical character of these habitats and/or impact the prey upon which they depend.

Distribution

Otter are annually harvested in every watershed in Vermont during a regulated trapping season.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Known Watersheds

Middle Connecticut

West

Waits

Upper Connecticut-Mascoma

Black-Ottawquechee

Deerfield

Hudson-Hoosic

Mettawee River

Lake Champlain

Lamoille River

Missiquoi River

Otter Creek



Common Name: **Northern River Otter**
Scientific Name: **Lontra canadensis**
Species Group: **Mammal**

Passumpsic
St. Francois River
Upper Connecticut
White
Winooski River

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☐

Otter are adaptable to many different wetland habitats including beaver created wetlands, lakes, streams and ponds. Intact vegetation along the perimeter of streams, lakes and wetlands is an important habitat feature of otter habitat. Beaver bank dens and lodges are also used by otter. Beaver created wetlands provide critical foraging and denning habitat. Log jams resulting from fallen trees also provide shelter and foraging habitat. Otter also require healthy aquatic systems that provide an adequate prey base.

Habitat Types:

Aquatic: Fluvial
Aquatic: Large Lake Champlain Tribs Below Falls
Aquatic: Lower CT River
Aquatic: Lacustrine
Aquatic: Lake Champlain

Current Threats

Habitat Threats:

Habitat Alteration
Impacts of Roads or Transportation Systems
Sedimentation

Description of habitat threat(s): Forested riparian buffers are key components of otter habitat. Loss and/or degradation could influence otter habitat selection and productivity. Historically, otter were limited by human encroachment, habitat destruction, and unregulated harvest. In Vermont, the extirpation of beaver, loss of habitat, and pollution resulted in a much reduced population throughout the 1800's and early 1900's. Otter populations have rebounded with the return of the beaver. Although not strongly supported in recent literature, it is expected that increasing development pressure and pollutants such as mercury could negatively affect future population levels. Despite this potential vulnerability, contemporary harvest records in Vermont indicate a well distributed, abundant population of otter in recent decades. Furthermore, should Vermont's otter population begin experiencing the effects of development, pollution and/or climate stressors, the mechanisms for detecting and addressing such population trends are currently in place.

Non-Habitat Threats:

Pollution

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Northern River Otter**
 Scientific Name: **Lontra canadensis**
 Species Group: **Mammal**

Loss of Prey Base

Description of non-habitat threat(s): Although the effects of pollutants are not believed to be a limiting factor for the otter population in Vermont, contaminants such as PCB's, mercury, and other heavy metals are known to accumulate in the tissue of otter and negatively affect reproduction and survival.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	Medium	Monitor distribution and abundance
Research	Threats and Their Significance	High	Determine the impact of heavy metals and contaminants on otter populations in each watershed.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Privately-Owned Protected Areas		Maintain riparian buffer strips along streams, rivers, lakes, ponds, and wetland habitats.	Number of linear miles of vegetated riparian buffers	Trout Unlimited, NRCS, USFWS, NWTF, DEC, Vt. F&P	SWG, USFWS
Species Restoration		Provide a suitable prey base.		Trout unlimited, DEC	TU, DEC, USFWS, SWG
Policy & Regulations		Eliminate acid rain and the input of mercury into otter habitat.	Decrease acid, mercury, and heavy metal deposition into Vermont lakes, rivers, and streams	DEC, EPA,	DEC, EPA
Compliance & Enforcement		Enforce the Clean Water Act	Increase the number of bodies of water that meet class A designation	Trout Unlimited, NRCS, USFWS, USFS, Wild Turkey Federation, DEC, Vt. Forests & Parks	EQIP, SWG, EPA, NWTF

Bibliography

Lariviere S, Walton LR (1998) *Lontra canadensis*. Mammalian Species, 587, 1 -8.

Melquist, W.E. and A.E. Dronkert. 1987. River Otter. In M. Novak, J.A. Baker, M.E. Obbard, and B. Malloch, eds. Wild furbearer management and Conservation in North America. Ontario Trappers Assoc., North Bay.

Parsons, D.R. 2003. Natural History Characteristics of Focal Species in the New Mexico Highlands Wildlands Network. Wildlands Project. 69 pp.



Common Name: **Canada Lynx**
 Scientific Name: **Lynx canadensis**
 Species Group: **Mammal**

Conservation Assessment

Final Assessment: High Priority

Global Rank: G5

Global Trend:

State Rank: S1

State Trend: Fluctuating

Extirpated in VT? yes

Regional SGCN? no

Assessment Narrative:

Recovery of lynx in Vermont may be limited by global climate change (Carroll 2007, Hoving et al. 2005). Although the influence of competition from coyote, fisher, and bobcat, which could also be exacerbated by global climate change (Peers et al. 2013), may not be clearly understood (Ray et al. 2002), there is some indication that lynx populations existing at the margins of their range may be limited by these sympatric carnivores (Peers et al. 2013, Vashon et al. 2012). Harvest records for fisher, bobcat and coyote in northeast Vermont (VFWD unpublished data) and track surveys conducted within Vermont's two largest blocks of unfragmented suitable lynx habitat (Farrell 2012) indicate well-established populations of these competing carnivores. Suitable lynx habitat in Vermont is limited and occurs in relatively small patches distributed over the northeastern portions of the state. As a result of this habitat condition, the effects of fragmentation could result in the isolation of Vermont's lynx population from populations to the north further jeopardizing its ability to persist in the state (Koehler et al. 2008, Murray et al. 2008). Also, because Canada lynx exhibit strong selection for habitats where snowshoe hares are abundant (Fuller et al. 2007, Vashon et al. 2008, Squires et al. 2010), the suitability of Vermont's currently occupied lynx habitat could change markedly with future changes in landscape-level hare densities and changing habitat associated with forest management; thus, successful conservation of lynx populations in Vermont will require the protection and management of large tracts of snowshoe hare habitat (Simons-Legaard et al. 2013, Murray et al. 2008).

Distribution

Historical records of Canada lynx in Vermont are scarce. Prior to this century, lynx were documented in the state on only four occasions (Windham 1928, St. Albans 1968, Calais 1797 and Addison County 1937: Vermont archived bounty records). Since 2003, nine lynx sightings have been confirmed in Vermont. Eight of the sightings were recorded in Essex County and one in Orleans County (unpublished data, VFWD). Since 2012, Intensive snow track and remote camera surveys have successfully detected lynx in the Nulhegan Basin (Bernier 2011 & 2013). Reproduction was first documented in 2012 in the Nulhegan Basin when the tracks of three lynx, a presumed family group, were observed travelling together in late February (Bernier 2011).

Distribution by Biophysical Region:

Champlain Valley	Historic Records Only	Southern VT Piedmont	Not Probable
Champlain Hills	Not Probable	Vermont Valley	Not Probable
Northern Green Mtns	Not Probable	Southern Green Mtns	Not Probable
Northern VT Piedmont	Probable	Taconic Mtns	Not Probable
Northeastern Highlands	Confident		

Distribution by Watershed:



Common Name: **Canada Lynx**
Scientific Name: **Lynx canadensis**
Species Group: **Mammal**

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☐

Along the southern periphery of their range, lynx prefer a variety of habitat types including mid-successional coniferous forests and edge habitat with moderate to abundant understory cover (Koehler et al. 2008, Maletzke et al. 2008, Vashon et al. 2008b). Lynx tend to avoid open areas and mature forests having little horizontal cover (Vashon et al. 2008b). Lynx select for stands where snowshoe hare are abundant (2.4 hares/ha, Vashon et al. 2008) such as areas of dense softwood in association with 11 - 21 year old regenerating clear-cuts or similarly aged partially harvested stands (Fuller et al. 2007, Simons-Legaard 2013). Organ et al. (2008) identified the "tip up mounds" of blown down trees as features commonly used as natal dens and further found that the presence of within stand structure capable of providing visual obscurity at 5 meters from the den was a significant predictor of den site selection by lynx. Hoving (2005) determined that lynx populations in this region are unlikely to occur in areas of low annual snowfall (<270cm) or areas dominated by deciduous forests.

Habitat Types:

Spruce Fir Northern Hardwood
Early Succession Boreal Conifers
Early Succession Spruce-Fir

Current Threats

Habitat Threats:

Habitat Succession
Habitat Alteration
Impacts of Roads or Transportation Systems
Climate Change
Habitat Fragmentation

Description of habitat threat(s): Changes in the climate that result in the reduction of annual snowfall could greatly influence the distribution of lynx in the northeast (Hoving 2005). Decreased snowfall can affect lynx through decreased prey vulnerability and decreased competitive advantage over sympatric carnivores (Carroll 2007). Furthermore, although there is evidence that the degree of diet specialization of lynx in the southern parts of their range is less than in their northern counterparts, the long-term persistence of lynx in Vermont could be limited by the availability of suitable snowshoe hare densities (Roth et al. 2007, Simons-Legaard et al. 2013). Thus, the loss of suitable hare habitat from both natural (i.e. forest succession) and human caused disturbances (i.e. forest management favoring deciduous forest composition) could adversely affect lynx in Vermont. In addition, because the viability of lynx populations in the southern part of their range is suspect in the absence of ingress from northern populations (Murray et al. 2008), the maintenance of landscape connectivity with these northern areas of occupancy is of critical importance. Although Farrell (2012) concluded that lynx connectivity across the northeast is expected to remain stable in the coming decades, the long-term persistence of lynx in Vermont remains dependent upon interstate and international commitments to maintaining these connective habitats (Murray et al. 2007).

Non-Habitat Threats:



Common Name: **Canada Lynx**
Scientific Name: **Lynx canadensis**
Species Group: **Mammal**

Competition

Predation or Herbivory

Loss of Prey Base

Loss of Metapopulation Structure

Description of non-habitat threat(s): Peers et al. (2013) determined that lynx are subjected to niche displacement in areas of overlap with bobcat. In Vermont, bobcat harvest data (VFWD unpublished data) and the results of extensive snow track surveys conducted since 2012 (Bernier 2012 & 2013) indicate a well-established, sympatric bobcat population. Furthermore, the effects of climate change could increase the competitive pressure on lynx by altering the distribution and abundance of competing carnivores populations and by decreasing their competitive advantages over these sympatric species (Carroll 2007). In addition, the primary source of mortality of lynx in Maine was predation, especially by fisher, accounting for nearly 42% of lynx deaths (Vashon et al. 2012). Similar to bobcats, harvest data and track survey results also indicate an abundance of fisher within Vermont's most suitable lynx habitats.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Distribution and Abundance	High	Collect baseline data on lynx distribution and abundance in Vermont in order to assess the status of the population and develop appropriate conservation strategies.
Research	Threats and Their Significance	High	Examine the affects of competition with sympatric carnivores and assess how certain habitat features such as snow depth, managing furbearer populations, and a changing climate may influence this relationship.
Monitoring	Population Change	High	Continue monitoring for the presence of the species in the state.
Monitoring	Habitat Change	Medium	Develop and implement a plan for monitoring changes in suitable lynx habitat resulting from habitat conversions, forest management practices and climate change.
Monitoring	Range Shifts	Low	Monitor range shifts of competing carnivore populations resulting from climate change.
Monitoring	Monitor Threats	Medium	Identify and monitor impacts to key connective corridors serving to link Vermont's lynx population with core populations to the north.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Canada Lynx**
 Scientific Name: **Lynx canadensis**
 Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Compatible Resource Use	High	Promote forest management practices that provide for the life history requirements of lynx	# of acres of snowshoe hare habitat available within potential lynx range	Vt. Forest and Parks Dept, Industrial forest landowners, Coverts	EQIP, SWG, USFWS
Compatible Resource Use	High	Maintain connectivity of habitat between Maine, New Hampshire, Quebec and Vermont.	# of acres of corridor habitat conserved	TNC, VLT, NHF&G, Conservation Fund, NWF, Keeping Track, Coverts	TNC, VLT, Conservation Fund, USFWS, Forest Legacy
Compatible Resource Use	High	Continue managing competing carnivores within key lynx habitats, particularly fisher, via regulated trapping	Maintenance of healthy furbearer populations	Vermont Trappers Association	
Policy & Regulations	Medium	Support and cooperate with larger efforts to curb global climate change.			



Common Name: **Canada Lynx**
Scientific Name: **Lynx canadensis**
Species Group: **Mammal**

Bibliography

- Degraaf, R.M., and Yamasaki, M. 2001. New England Wildlife: Habitat, Natural History, and Distribution, University Press of New England, Hanover and London.
- Fuller, A., and Homyack . 1999. University of Maine (Find Citation)
- Hamilton, J., and J. Whitaker. 1979. Mammals of the eastern United States. Cornell University Press, Ithaca, N.Y.
- Hoving, C.L. 2001. Historical occurrence and habitat ecology of Canada lynx (*lynx canadensis*) in eastern North America. M.S. Thesis, University of Maine, Orono.
- McCord , C. M. and J. E Cardoza. 1982. Bobcat and Lynx Pages 728-766 in J.A. Chapman and G.A. Feldhamer, eds. Wildlife Mammals of North America: biology, management,, and economics. Johns Hopkins Univ. Press, Baltimore, Md.
- Novak, M., J. Baker, M. Obbard, and B. Mallock, editors. 1987. Wild furbearer management and conservation in North America. Ontario Ministry of Natural Resources, Toronto
- Osgood, Frederick L. Jr., 1938. The mammals of Vermont . J.Mammalogy 19(4): 435-441.
- Ray, J.C., J.F. Organ, and M.S. O'Brien. 2002. Canada lynx (*lynx Canadensis*) in the Northern Appalachians: Current Knowledge, Research Priorities, and a Call for Regional Cooperation and Action. Report of a Meeting held in Portland Maine, April 2002. 22pgs.
- Ruediger, B., J. Claar, S. Gniadek, B. Holt, L.Lewis, S. Mighton, B. Naney, G. Patton, T. Rinaldi, J. Trick, A. Vandehay, F. Wahl, N. Warren, D. Wenger, and A. Williamson. 2000. Canada lynx conservation assessment and strategy. USDA Forest Service, USDI fish and Wildlife Service, USDI Bureau of Land Management, USDI National Park Service. Missoula, MT.
- Tumlison, R. 1987. *Felis lynx*. Mammalian Species 269:1-8.
- Tumosa, J. and D. Batchelder. U. S Forest Service. Species Data Collection Form
- Vashon, J., A. Vashon, and S. Crowley. 2003. Partnership for lynx conservation in Maine, Dec. 2001 - Dec. 2002 field report. Unpublished Report, Maine Dept. Inland Fish and Wildlife.
- Vashon, J., J. Holloway, A Winters, S. Crowley, and C. Todd. 2003. Snow track survey of Canada lynx in the Boundary Plateau and St John Upland ecoregions of Maine. Unpublished report. Maine Dept. Inland Fish and Wildlife.



Common Name: **Bobcat**
Scientific Name: **Lynx rufus**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S4

State Trend: Unknown

Extirpated in VT? no

Regional SGCN? Yes

Assessment Narrative:

The bobcat is apparently common and well distributed throughout Vermont although higher densities appear to exist in the Champlain Valley and the Taconics possibly due to higher prey densities. Bobcats have declined since the middle of the 20th century due to land use changes affecting prey densities and to increasing competition from other carnivores such as fisher and coyote. Statewide population estimates are unknown, but carrying capacity has been estimated.

The bobcat uses a variety of habitats and the relative suitability of habitats in the Vermont landscape have been quantified (see below). Bobcat occurrence appears to be positively related to the amount of mixed forest and forested wetland habitats. Critical habitats, such as those used for denning remain largely unquantified.

Landscape change represents a primary threat to bobcats, especially as they appear to depend on connected expanses of undeveloped habitat. Conversion of natural habitat to housing and other forms of development will most likely affect the distribution and abundance of the species in Vermont. Similarly, the impacts of climate change, particularly with respect to changes in prey and sympatric carnivore distribution and abundance, may present significant challenges to bobcats through the future.

Distribution

Bobcats occupy home ranges that include a variety of habitats. Average home range size for bobcats based on a study in the Champlain Valley was 57.3 km² (Donovan et al. 2011) Male home ranges (n=10) averaged 70.9 km² while female home ranges (n=4) averaged 22.9 km². Based on patterns of use in home ranges, bobcats respond positively to shrub, deciduous forest, coniferous forest, and wetland cover types within 1 km of a location and negatively to roads and mixed forest cover within 1 km of a location. Similar results have been found in New Hampshire with bobcats preferring areas with few roads, limited human development, high stream densities, and steep topography (Broman et al. 2014). Another study conducted repeated surveys throughout Vermont and concluded that bobcat probability of occupancy was positively related to the percentage of both mixed forest and forested wetland habitat within 1 km of survey sites (Long et al. 2011). In Vermont, steep, rocky cliffs may be important as winter refuges and breeding habitat.

The size of the bobcat population is uncertain in Vermont. Donovan et al. (2012) estimated the maximum carrying capacity of females in northwestern Vermont (WMU 1, 1,153 km²) as 42. Using a similar approach, carrying capacity across Vermont has been estimated as 1,150 (835 females, 316 males) (J. Murdoch, pers. comm.).

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconnic Mtns	Confident
Northeastern Highlands	Confident		



Common Name: **Bobcat**
Scientific Name: **Lynx rufus**
Species Group: **Mammal**

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☒ General Literature ☒

Bobcats occupy home ranges that include a variety of habitats. Average home range size for bobcats based on a study in the Champlain Valley was 57.3 km² (Donovan et al. 2011) Male home ranges (n=10) averaged 70.9 km² while female home ranges (n=4) averaged 22.9 km². Based on patterns of use in home ranges, bobcats respond positively to shrub, deciduous forest, coniferous forest, and wetland cover types within 1 km of a location and negatively to roads and mixed forest cover within 1 km of a location. Similar results have been found in New Hampshire with bobcats preferring areas with few roads, limited human development, high stream densities, and steep topography (Broman et al. 2014). Another study conducted repeated surveys throughout Vermont and concluded that bobcat probability of occupancy was positively related to the percentage of both mixed forest and forested wetland habitat within 1 km of survey sites (Long et al. 2011). In Vermont, steep, rocky cliffs may be important as winter refuges and breeding habitat.

The size of the bobcat population is uncertain in Vermont. Donovan et al. (2012) estimated the maximum carrying capacity of females in northwestern Vermont (WMU 1, 1,153 km²) as 42. Using a similar approach, carrying capacity across Vermont has been estimated as 1,150 (835 females, 316 males) (J. Murdoch, pers. Comm.).

Habitat Types:

Cliffs and Talus
Hardwood Swamps
Softwood Swamps
Spruce Fir Northern Hardwood
Northern Hardwood
Oak-Pine Northern Hardwood
Floodplain Forests
Open Peatlands
Marshes and Sedge Meadows
Early Succession Boreal Conifers
Early Succession Pine and Hemlock
Early Succession Upland Oak
Early Succession Other Types
Wet Shores
Shrub Swamps
Early Succession Boreal Hardwoods



Common Name: **Bobcat**
Scientific Name: **Lynx rufus**
Species Group: **Mammal**

Early Succession Spruce-Fir
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Succession
Habitat Alteration
Habitat Fragmentation

Description of habitat threat(s): Bobcats distribution appears to relate mainly to forest cover and forest wetland habitat, both of which positively influence probability of occurrence in the landscape. Changes to these two habitats and others that offer important resources like rocky ledges for denning represent a primary threat to the species. Conversion of habitats due to development like residential housing and roads or even climate change will most likely affect bobcat distribution and abundance (Bettigole et al. 2014).

Non-Habitat Threats:

Competition
Loss of Prey Base

Description of non-habitat threat(s): Bobcat numbers have declined since coyotes became established in Vermont. The specific impacts of coyotes and other carnivores such as fisher remain largely unstudied in the Northern Forest. Prey species have also declined in some areas due to loss of early successional habitat and have presumably impacted bobcat numbers.

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	High	Identify and quantify critical habitats for reproduction, such as rocky, ledge areas.
Research	Distribution and Abundance	High	Determine the location of source and sink populations and identify the habitat parameters associated with these populations.
Research	Threats and Their Significance	Medium	1) Examine how habitat loss, conversion, and fragmentation impacts distribution and abundance. 2) Determine competition effects with coyotes and other sympatric carnivores such as fisher.
Monitoring	Range Shifts	Medium	Assess possible range shifts and population changes due to climate change.



Common Name: **Bobcat**
 Scientific Name: **Lynx rufus**
 Species Group: **Mammal**

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Compatible Resource Use	High	Promote less development of high quality habitats.	Amount of high quality habitat protected or conserved	VTrans, Town Planning Commission, VLT, Regional and Town Cons Comms, Keeping Track	SWG, AOT
Species Restoration	Medium	Provide important prey base	Number of acres of rabbit and hare habitat protected	Coverts, USFS, VWA, Northern, USFS, VFPR, Ruffed Grouse Society	USFWS, Ruffed Grouse Society, EQIP
Species Restoration	Medium	Identify necessary habitats and develop actions for protection	Number of necessary habitats mapped and protected	Coverts, USFS, VWA, VLT, UVM	UVM, VLT, USFS, USFWS

Bibliography

- Anderson, E. M., and M. J. Lovallo. 2003. Bobcat and lynx (*Lynx rufus* and *Lynx canadensis*). Pages 758-788 in G. A. Feldhamer, B. C. Thompson, and J. A. Chapman (eds.) *Wild mammals of North America biology, management, and conservation*, 2nd edition. Johns Hopkins University Press, Baltimore, Maryland, USA.
- Bettigole, C. A., T. M. Donovan, R. Manning, J. Austin, and R. Long. 2014. Acceptability of residential development in a regional landscape: potential effects on wildlife occupancy patterns. *Biological Conservation* 169:401-409.
- Broman, D. J. A., J. A. Litvaitis, M. Ellingwood, P. Tate, and G. C. Reed. 2014. Modeling bobcat *Lynx rufus* habitat associations using telemetry locations and citizen-scientist observations: are the results comparable? *Wildlife Biology* 20:229-237.
- Donovan, T. M., G. S. Warrington, W. S. Schwenk, and J. H. Dinitz. 2012. Estimating landscape carrying capacity through maximum clique analysis. *Ecological Applications* 22:2265-2276.
- Donovan, T. M., M. Freeman, H. Abouelezz, K. Royar, A. Howard, and R. Mickey. 2011. Quantifying home range habitat requirements for bobcats (*Lynx rufus*). *Biological Conservation* 144:2799-2809.
- Lavoie, M., P-Y Collin, F. Lemieux, H. Jolicoeur, P. Cana-Marquis, and S. Larivière. 2009. Understanding fluctuations in bobcat harvest at the northern limit of their range. *Journal of Wildlife Management* 73:870-875.
- Larivière, S., and L. R. Walton. 1997. *Lynx rufus*. *Mammalian Species* 563:1-8.
- Litvaitis, J. A., J. P. Tash, and C. L. Stevens. 2006. The rise and fall of bobcat populations in New Hampshire: relevance of historical harvests to understanding current patterns of abundance and distribution. *Biological Conservation* 128:517-528.
- Long, R. A., T. M. Donovan, P. MacKay, W. J. Zielinski, and J. S. Buzas. 2011. Predicting carnivore occurrence with noninvasive surveys and occupancy modeling. *Landscape Ecology* 26:327-340.



Common Name: **Eastern Mountain Lion**
Scientific Name: **Puma concolor cougar**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: SH

State Trend: N/A

Extirpated in VT? yes

Regional SGCN? No

Assessment Narrative:

The Mountain Lion, also known as Puma, Cougar and Catamount is listed as endangered in Vermont. It is believed to be extirpated in the East (except in southern Florida). The USFWS declared the Eastern cougar (*Puma concolor cougar*) extinct in 2011 though it remains federally endangered pending delisting. Anecdotal reports of field sightings are fairly frequent; however, both field and incidental evidence is absent. Even in lowest densities, Mountain Lions are hit, shot, snared, wander into towns and cities, and are photographed on cell phones, point & shoot cameras, and random remote wildlife cams. A Black Hills, SD male left field and incidental evidence in four states across 1500 miles before being hit by a car in Milford, CT, June 2011. All North American Mountain Lions are one subspecies genetically, though the taxonomy remains disputed (Culver et al. 2000); which suggests that conservation efforts should be focused on the entire puma Genus. Confirmations of Mountain Lions with both North and South American DNA (former captives or descendants) have been documented in Ontario (Rosatte, 2011), Quebec and New Brunswick (Lang, et al. 2013), There is no evidence of breeding in eastern Canada. The closest breeding colonies to Vermont remain southwest Florida, the Dakotas and Nebraska. Recent research show mountain lions are keystone species for ecosystem functioning (Ripple et al. 2014).

Distribution

Distribution by Biophysical Region:

Champlain Valley	Historic Records Only	Southern VT Piedmont	Historic Records Only
Champlain Hills	Historic Records Only	Vermont Valley	Historic Records Only
Northern Green Mtns	Historic Records Only	Southern Green Mtns	Historic Records Only
Northern VT Piedmont	Historic Records Only	Taconic Mtns	Historic Records Only
Northeastern Highlands	Historic Records Only		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☒ Extensive Local Knowledge ☐ Regional Literature ☐ General Literature ☒

Mountain Lions are no longer understood to be wilderness obligates, with the widest range across more habitats, including urban landscapes, of any terrestrial mammal in the western hemisphere. Beier (1993), using simulated population dynamics, estimated that an area of 1,000 to 2,200 square kilometers (372 to 818 square miles, depending on the demographics of a particular population) was needed for a population of 15-20 adult cougars to have a very low risk (<98%) of extinction within 100 years. Area of 600 - 1600 km², and smaller (Beier. 1993), might suffice where adequate dispersal corridors allow movement among populations. Smallest documented home range is 39 km² (Laundre and Loxterman 2006). Mountain Lions are breeding in suburban-



Common Name: Eastern Mountain Lion
Scientific Name: *Puma concolor couguar*
Species Group: Mammal

exurban-wildland matrix habitat throughout the western US, and have recovered range east to the Dakotas/Nebraska without assistance. Space-use patterns differ little between wildland and residential environments (Kertson et al, 2011), though reproductive behaviors (communication/denning) require greater buffers from development than non-reproductive behaviors (movement/feeding) within the suburban/exurban/wildland matrix (Wilmers et al, 2013). Specific dispersal barriers include roads and nighttime illumination (Beier 1993, 1995); identifying and protecting wildlife corridors can mitigate dispersal mortalities. Male dispersal and settlement patterns based on mating opportunities; female patterns based on avoiding other Mountain Lions (Stoner et al. 2013). Mountain Lions are the epitome of a generalist predator (Knopf and Boyce 2014), though they favor and are adapted for medium-sized ungulates. Deer/ Elk wintering habitat is seasonally favored. (Lindzey 1987).

Adirondack Park, an area roughly comparable to the state of Vermont, could support as many as 350 Mountain Lions (Laundre, 2013). Glick (2014) found that the Northeast region east of the Hudson River could support from 322 - 2,535 Mountain Lions.

Habitat Types:

Outcrops and Alpine
Cliffs and Talus
Northern Hardwood
Spruce Fir Northern Hardwood
Oak-Pine Northern Hardwood
Open Peatlands
Marshes and Sedge Meadows
Early Succession Boreal Conifers
Early Succession Pine and Hemlock
Early Succession Upland Oak
Early Succession Other Types
Wet Shores
Shrub Swamps
Early Succession Boreal Hardwoods
Early Succession Spruce-Fir
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Conversion of Habitat
Habitat Alteration
Impacts of Roads or Transportation Systems
Habitat Fragmentation



Common Name: **Eastern Mountain Lion**
Scientific Name: **Puma concolor cougar**
Species Group: **Mammal**

Description of habitat threat(s): Where they still exist, pumas can be found in a multitude of habitats, ranging from closed forest to semi-open shrublands. Human development/disturbance appears to affect little the use of areas by pumas as they are found in suburban to exurban environments. Human intolerance to their presence in these areas is the main negative impact on their survival. Prey availability and habitat characteristics can affect puma distribution and survival. Loss of habitat connectivity between source populations limits dispersal, range expansion, and genetic variability (Ernest et al. 2003).

Non-Habitat Threats:

Harvest or Collection

Loss of Prey Base

Trampling or Direct Impacts

Description of non-habitat threat(s): Negative human attitudes among certain demographics towards pumas in regards to human safety and perceived impacts on deer populations can impact successful establishment/ maintenance of puma populations in the East Florida public attitude surveys found broad public support for puma recovery, including residents of a proposed relocation region and among sportsmen (Duda and Young. 1995; Cramer. 1995). However, a successful test-release of Texas pumas to southern Georgia/north Florida concluded that resistance from just a handful of individuals can impede recovery efforts (Belden and McCown. 1996). Pending federal delisting could jeopardize any potential for recolonization if eastern state protections are not established, maintained and enforced,

Research and Monitoring Needs

Type	Need	Priority	Description
Monitoring	Population Change	Medium	Continue current low-level monitoring and incidental Mountain Lion evidence documentation (track, scat, kills, photographs, etc.). Consider active pheromone station monitoring (e.g Lang et al. 2013) to detect VT presence. Collect genetic material for testing.

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Policy & Regulations	High	Pending federal delisting, maintain and enforce state protections of entire puma Genus.			
Research	Medium	Identify areas within state that could support viable populations (Glick 2014) and develop a state recovery plan.			
Awareness Raising and Communications	High	Determine public attitudes towards puma recovery efforts in VT (e.g. McGovern and Kretser 2014); Provide interpretive and public education material about pumas.			

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: Eastern Mountain Lion
Scientific Name: *Puma concolor cougar*
Species Group: Mammal

Bibliography

- Anderson, A.E. 1983. A Critical review of literature on Puma (*Felis concolor*). Colorado Division of Wildlife, Special Report no. 54. 91pp.
- Belden, R.C. and J.W. McCown. 1996. Florida panther reintroduction feasibility study. Florida Game and Fresh Water Fish Commission, Bureau of Wildlife Research. Final Report.
- Beier, P. 1993. Determining minimum habitat areas and habitat corridors for cougars. *Conservation Biology* 7:94-108.
- Beier, P. 1995. Dispersal of juvenile cougars in fragmented habitat. *Journal of Wildlife Management* 59:228-237.
- Beier, P. 1996. Predator Defense correspondence. http://www.predatordefense.org/docs/cougars_Beier_sightings.pdf.
- Cramer, P.C. 1995. The northeast Florida panther education program. The University of Florida Final Report to Florida Panther Reintroduction Feasibility Study. Florida Advisory Council on Environmental Education, Tallahassee.
- Culver, M., W.E. Johnson, J. Pecon-Slaterry, S.J. O'Brien. 2000. Genomic Ancestry of the American Puma (*Puma concolor*). *Journal of Heredity*, 91. 186-197.
- Currier, MJP 1983. *Felis concolor*. *Mammalian Species* 200:1-7.
- Degraaf, R.M. and M. Yamasaki. 2001. *New England Wildlife*. Univ. Press of New England. Hanover. N.H.
- Duda, M. and K.C. Young. 1995. Floridian's knowledge, opinions, and attitudes toward panther habitat and panther-related issues: public opinion survey results report. Report for the Advisory Council on Environmental Education. . Responsive Management, Mark Damian Duda and Associates, Inc, Harrison, Virginia, USA.
- Ernest, H.B., W.M. Boyce, V.C. Bleich, Bernie May, S.J. Stiver and G. Torres. 2003. Genetic Structure of Mountain Lion (*Puma concolor*) populations In California. *Conservation Genetics* 4:353-356.
- Kitchell, J.A. 1999. Statement of Purpose and Reason Draft Species Data Records, *Felis concolor*. USFS Region 9.
- Knopff, K., and Mark Boyce. 2014. Prey Specialization in Multiprey Systems. Pages 194-210. Transactions of the 72nd North American Wildlife Natural Resources Conference. Wildlife Management Institute.
- Lindzey, F. 1987. Mountain lion. Pages 656-668. In Novak, M.; Baker, J.A.; Obbard, M.E.; Malloch, B. (editors). *Wild Furbearer management and conservation in North America*. Toronto: Ontario Ministry of Natural Resources and Ontario Trappers Association.
- Lang, L., N. Tessier, M. Gauthier, R. Wissink, H. Jolicoeur, and F.-J. Lapointe. 2013. Genetic Confirmation of Cougars (*Puma concolor*) in Eastern Canada. *Northeastern Naturalist* 20(3): 383-396.
- Laundre, J.W. 2013. The Feasibility of the Northeastern USA Supporting the Return of the Cougar (*Puma concolor*). *Oryx*/ Volume 47/ Issue 1/ January 2013, pp 96-104.
- Laundre, J.W. and J. Loxterman. 2006. Impact of Edge Habitat on Summer Home Range Size in Female Pumas. *American Midland Naturalist*. 157:221-229.
- Maehr, D.S., 1997. *The Florida Panther: life and death of a vanishing carnivore*. Washington D.C., Island Press.
- McGovern, E.B. and H.E. Kretser. 2014. *Puma concolor cougar in the Adirondack Park: Resident and Visitor Perspectives*. Wildlife Conservation Society, Adirondack Program Technical Paper #5.
- Ripple, W.J., J. Estes, R.L. Beschta, C.C. Wilmers, E.G. Ritchie, M. Hebblewhite, J. Berger, B. Elmhagen, M. Letnic, M.P. Nelson, O.J. Schmitz, D.W. Smith, A.D. Wallach. and A.J. Wirsing. 2014. Status and Ecological Effects of the World's Largest Carnivores. *Science* 343, 1241484.
- Rosatte, R. 2011. Evidence to Support the Presence of Cougars (*Puma concolor*) in Ontario, Canada. *Canadian Field-Naturalist* 25: 116-125.
- Russell, K.R. 1978. Mountain Lion. Pages 207-225. In *Big Game of North America Ecology and Management*. Stackpole Company, Harrisburg, Pennsylvania.

***Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report***



Common Name: **Eastern Mountain Lion**
Scientific Name: **Puma concolor couguar**
Species Group: **Mammal**

U.S. Fish and Wildlife Service (USFWS) Northeast Region. 2011. Press release: U.S. Fish and Wildlife Service concludes eastern cougar extinct. 2 March 2011. USFWS Northeast Region Page.
<<http://www.fws.gov/northeast/ECougar/newsreleasefinal.html>>.

Wilmers, C., Y. Wang, B. Nickel, P. Houghtaling, Y. Shakeri, M. L. Allen, J. Kermish-Wells, V. Yovovich, T. Williams. 2013. Scale Dependant Behavioral Responses to Human Development by a Large Predator, the Puma. PLOS One 8(4): e60590.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Moose**
Scientific Name: **Alces alces**
Species Group: **Mammal**

Conservation Assessment

Final Assessment: Medium Priority

Global Rank: G5

Global Trend:

State Rank: S5

State Trend: Declining

Extirpated in VT? yes

Regional SGCN? no

Assessment Narrative:

Moose were extirpated from Vermont by the early 19th century due to forest clearing and no legal protection. Following regrowth of the forest, restoration of beavers, and nearly a century of protection, moose immigrated from New Hampshire in the 1970's and their numbers and distribution in Vermont grew rapidly in the 1980's and 90's. By the time the former WAP was written in 2005, moose numbered over 5,000 animals and were reproducing throughout the state. Moose were recognized in the 2005 WAP as a "special category" species, along with beaver and white-tailed deer, due to their socioeconomic value and potential of having a significant ecological effect on the landscape. SWG funds were not intended to be directed at these three species at that time.

Currently, the statewide moose population is about half of what it was in 2005. Most of this reduction was by design in order to bring numbers in northeastern Vermont down below ecological carrying capacity and allow for adequate regeneration of trees in managed stands. The current population estimate of 2500 moose is below the minimum target of 3,000 as called for in Vermont's 10-year Big Game Management Plan. Moose health and nutrition as reflected by body weight and ovulation rate has declined, and warmer weather from spring through autumn has likely contributed to higher incidence of parasites, most notably the winter tick and brainworm, and abnormally high levels of heat stress.

Distribution

Highest densities in the Northeastern Highlands and Northern Vermont Piedmont.

Distribution by Biophysical Region:

Champlain Valley	Confident	Southern VT Piedmont	Confident
Champlain Hills	Confident	Vermont Valley	Confident
Northern Green Mtns	Confident	Southern Green Mtns	Confident
Northern VT Piedmont	Confident	Taconic Mtns	Confident
Northeastern Highlands	Confident		

Distribution by Watershed:

Habitat Description

Habitat Information is based on the following:

Limited Local Knowledge ☐ Extensive Local Knowledge ☒ Regional Literature ☒ General Literature ☒



Common Name: **Moose**
Scientific Name: **Alces alces**
Species Group: **Mammal**

Habitat Types:

Hardwood Swamps
Softwood Swamps
Spruce Fir Northern Hardwood
Northern Hardwood
Marshes and Sedge Meadows
Early Succession Boreal Conifers
Early Succession Boreal Hardwoods
Early Succession Pine and Hemlock
Shrub Swamps
Early Succession Spruce-Fir
Early Succession Northern Hardwoods

Current Threats

Habitat Threats:

Habitat Fragmentation
Climate Change

Description of habitat threat(s): Fragmentation from ski area and recreational trail expansions; ridgetop windfarms. Heat stress from warming climate.

Non-Habitat Threats:

Parasites

Description of non-habitat threat(s): Increased levels of parasites, most notably *Dermacentor albipictus* and *Paralaphostrongylus tenuis*.

Vermont Department of Fish and Wildlife
Wildlife Action Plan - Revision 2015 Draft
Species Assessment Report



Common Name: **Moose**
 Scientific Name: **Alces alces**
 Species Group: **Mammal**

Research and Monitoring Needs

Type	Need	Priority	Description
Research	Habitat Requirements	Low	
Research	Basic Life History	Low	
Research	Distribution and Abundance	Low	
Research	Threats and Their Significance	High	Health condition and effects from parasites and disease.
Research	Population Genetics	Low	
Research	Taxonomy	Low	
Monitoring	Population Change	High	
Monitoring	Habitat Change	Low	
Monitoring	Range Shifts	Low	
Monitoring	Monitor Threats	High	

Species Strategies

Strategy Type	Strategy Priority	Strategy Description	Performance Measure	Potential Partners	Potential Funding Sources
Species Restoration	Medium	Keep moose densities below 0.75/sq km. and densities below 10/2.6sq km in order to reduce winter tick and brainworm infection rates.	Reduced levels of winter tick infestation. Reduced incidence of brainworm cases.	USFWS	SWG, PR
Habitat Restoration	Medium	Increase amounts of early successional habitat, especially in the Central and Southern Green Mountains.	Improved Moose body weights and ovulation rates	USFWS	SWG, PR

Bibliography

DeGraaf, R. M., and M. Yamasaki. 2001. New England wildlife: habitat, natural history, and distribution. University Press of New England, Hanover, NH. 482 pp.

Vermont's Wildlife Action Plan

Habitat & Community Summaries

Appendix B

Draft • 10/1/2015

Conservation at Multiple Scales	2

Terrestrial Habitats

Northern Hardwood Forest Summary	9
Spruce-Fir-Northern Hardwood Forest Summary	16
Oak-Pine-Northern Hardwood Forest Summary	23
Young Forest-Old Forest Summary	32
Floodplain Forest Summary	33
Hardwood Swamp Summary	38
Softwood Swamps Summary	43
Vernal Pools & Seeps Summary	48
Open Peatlands Summary	53
Marshes and Sedge Meadows Summary	58
Wet Shores Summary	62
Shrub Swamps Summary	66
Upland Shores Summary	70
Outcrops & Upland Meadows Summary	73
Cliff & Talus Summary	76
Grassland & Hedgerow Summary	79
Mines and Quarries Summary	84
Subterranean Summary	87
Buildings & Other Structures Summary	90

Aquatic Habitats

Riparian Summary	93
Streams & Rivers Summary	98
Lake Champlain Tributaries Summary	106
Lower Connecticut River Summary	112
Lake Champlain Summary	117
Lakes Summary (excluding Lake Champlain)	122

Conservation at Multiple Scales

Introduction

This chapter explains how conservation is organized in this Wildlife Action Plan.

Vermont's list of Species of Greatest Conservation Need (SGCN) comprises 133 vertebrate species 200 invertebrate species (such as the Tawny Emperor Butterfly, Cobblestone Tiger Beetle, and Giant Floater mussel) and 813 plants (vascular and bryophytes). Developing individual conservation plans for each of these species would have been exhausting and impractical. Moreover, implementing so many individual plans would be impossible due to insufficient staffing, resources and funds. In short, it would be monumentally inefficient.

Fortunately an easier and more efficient approach exists. It consists of designing and implementing conservation at multiple scales. This is commonly referred to as the “coarse filter-fine filter” approach and is widely accepted by scientists, wildlife managers and planners. The underlying concept is that if examples of all coarse-filter features are conserved at the scale at which they naturally occur, most of the species they contain—from the largest trees and mammals to the smallest insects—will also be conserved (Hunter 1991; NCASI 2004; Schulte et al. 2006). The coarse-filter approach is well documented in the scientific literature (Jenkins 1985; Noss 1987; Hunter et al. 1988; Hunter 1991; Noss and Cooperrider 1994; Haufler et al. 1996; Jenkins 1996; Poiani et al. 2000; USDA 2004). Habitat management historically practiced by Fish and Wildlife agencies to create young forests and shrublands that benefit dozens of “shrub and early-successional species” including Moose, New England Cottontail, American Woodcock and Ruffed Grouse is an example of a ‘habitat-scale’ coarse filter.

To best and most efficiently conserve all our SGCN, this Wildlife Action Plan focuses on three scales of conservation:

1. **Landscapes:** Include the features that contribute to ecological function at the state and regional levels, including a network of large, connected habitat blocks and another of aquatic habitats and riparian areas. Species requiring large habitat block, mixes of forest, wetlands and waters and connections between them will benefit most from landscape-level conservation but most other SGCN can also benefit.
2. **Habitats and Natural Communities:** Include the range of naturally occurring and anthropogenic habitats (such as young forest and grasslands). Terrestrial natural communities follow the classification system developed by Sorenson and Thompson (2005) which ties in with the ecological systems classification developed for the Northeast Association of Fish & Wildlife Agencies (Gawler 2008) for the 13 northeastern states. Aquatic communities follow the classification developed by Langdon et.al. (1998).
3. **Species and Groups of Species:** these are the SGCN for which we have identified specific conservation needs that would not be covered by conservation efforts at the other two scales.

Not all species, however, are best conserved by coarse-filters alone. For example, species dependent on multiple habitats at different times during their life cycles, those that occur in small geographic areas, those with highly specialized needs, those that travel across large geographic areas and those that are particularly rare often require focused attention. To ensure that the needs of these species are also addressed, fine filter conservation strategies are also needed. Species-specific conservation summaries can be found in Appendix A.

Efficiency in conservation effort can be realized by first identifying landscape conservation priorities that will effectively capture many natural communities, habitats, and species found within them. Natural community and habitat level conservation can effectively capture many of the remaining species. And finally, species-specific conservation action will be required for those species that are not captured at landscape or habitat/natural community scales. Typically these are species that are very rare, are declining across their range, aggregate for breeding, and/or require large home ranges.

Given the species focus of the congressional requirements for Wildlife Action Plan development, we began at the species level by assessing SGCN individually (Appendix A). Then SGCN were organized by taxonomic group and by the habitats they use. This resulted in conservation strategies at the three levels listed above (and in table 4.1).

Table B.1 Organization of Conservation Information in this Report

Level	Organization	Location in this Action Plan
1-Species	6 group summaries (amphibians & reptiles, birds, fishes, invertebrates, mammals and plants)	Chapter 4
	133 individual species and 15 invertebrates group summaries	Appendix A
2-Habitats & Natural Communities	125 communities & cultural habitats grouped into 24 summaries	Appendix B
3-Landscapes	Statewide and regional conservation strategies	Chapter 1
	Landscapes Landscape Report	Chapter 5 Appendix F

Selection of Classification Systems

We delineated landscapes based on the following elements: Interior Forest Blocks, Connectivity Blocks, Surface Waters and Riparian Areas, Riparian Areas for Connectivity, Physical Landscape Diversity Blocks, and Wildlife Road Crossings. Landscape conservation is discussed in chapter 5 and Appendix F of this Wildlife Action Plan.

Though great strides have been made in developing vegetation classification systems that function at the site, landscape, region and national scales (Barnes 1979, Allen and Starr 1982, Forman and Godron 1986, Cleland et. al 1997, Grossman et. al 1998), they are incomplete. In particular, no system satisfactorily integrates aquatic and terrestrial communities and cultural habitats¹ used by wildlife nationwide.

In lieu of a unified habitat classification system, Vermont's Action Plan technical teams selected the best features of five peer-reviewed vegetation classification systems that can be crosswalked with those used in other states to support broader scale conservation efforts—regionally, nationally, and internationally. Forest Cover Types (Eyre 1980) and U.S Forest Service Forest Inventory & Analysis Types (USDA 2003) were used for early successional stage forests. Natural Communities (Thompson and Sorenson 2000) were the basis for most terrestrial vegetation. "A Classification of the Aquatic Communities of Vermont" by Langdon et al. (1998) was adapted for aquatic community designations and cultural habitats¹ were adapted from Reschke (1990). Landscape scale communities were adapted from Poiani et.al. (2000).

One hundred twenty five aquatic and natural community types, cultural habitats and land cover types, capturing most of the habitat required by SGCN were selected from the five systems (table

¹ Cultural habitats are communities and sites that are either created and/or maintained by human activities or are modified by human influence to such a degree that the physical condition is substantially different from what existed prior to human influence (adapted from Reschke 1990).

4.2). Each was assigned to one of 22 categories. Because Lake Champlain and the Connecticut River harbor most of the fish diversity in Vermont, these two waterbodies were broken out from the taxonomy to provide for a more targeted assessment. Technical teams then developed assessment summaries for each that include descriptions and general locations; current conditions; desired conditions based on the needs of associated SGCN; priority problems; conservation strategies to address problems (along with the identification of potential conservation partners and funding sources); and a listing of relevant plans and planning processes pertinent to a habitat type.

Our terrestrial classification is designed to roll up to the Northeast Terrestrial Habitat Classification System (Gawler 2008) with standardized terminology and compatible habitat classifications. It allows the Action Plan to describe the aspects of conservation which are particular to Vermont, while facilitating conservation at a broader regional level. A Companion to the Terrestrial and Aquatic Maps has been published by TNC (Anderson et al. 2013). It includes profiles of each habitat type in the Northeast, distribution maps, state acreage figures, SGCN identification concern, and an assessment of overall conditions in the region. Habitat conservation summaries can be found in Appendix B.

Habitat Succession, Species of Greatest Conservation Need & the Action Plan

Plant succession produces cumulative change in the types of plant species occupying a given area through time. Succession is complicated by factors such as disturbance (large and small), local conditions, seed banks and soil legacies (Oliver 1981). A highly simplified timeline begins when land is cleared. Pioneer species typically return first followed by other species generally better adapted to the new and changing conditions created by the previous suite of species. Given sufficient time and appropriate conditions the area moves roughly through early, middle, and late successional stages—often referred to as mature or old growth. A disturbance, if sufficiently large, can re-set the clock anytime and succession begins again. The best known examples are forest succession but it occurs in virtually all vegetated areas. For example, lichen communities on granite mountaintops experience successional changes (Wessels 2002).

Succession can significantly impact habitat for Species of Greatest Conservation Need and other wildlife as in the edge habitat example noted earlier. Generally as succession moves from early (young forests) to late stages some wildlife will lose out (e.g., Spruce Grouse, American Woodcock, Cottontail Rabbit) and others will benefit (e.g., American Marten, Northern Goshawk). Others still prefer a mix of successional stages in appropriate configurations (e.g., Black Bear, Canada Lynx).

Over the past two centuries the mix of successional stages available to Vermont's wildlife has changed dramatically in both distribution and abundance. Though precise estimates (current and historic) are unavailable, prior to 1800 a significant percentage of Vermont's forests were in late-successional stages (>150-300 years and older). One-hundred years later young forests (early-successional stages of 1-15 years) dominated the state and today mid-successional forests (60-100 years) are most abundant. Wildlife populations have responded in turn. Vermont's SGCN list contains relatively few species requiring mid-successional forests and more that thrive in early and late-successional representations.

Because the loss of late-successional forests in the eastern U.S. occurred prior to the advent of modern wildlife biology and the current scarcity of later-successional stages (particularly northern hardwood forest types) our understanding how wildlife utilized these stages is not as advanced as our knowledge of wildlife in early successional stages. Historic records and research in late-successional areas elsewhere indicate that the distribution and abundance of some wildlife species was much greater when late-successional forests were in greater abundance—even if these species

can survive without them. Given the lack of this condition on the landscape it is advisable to increase its availability to wildlife.

The habitat, community and landscape summaries that follow here and in Chapter 6 and Appendix B address the habitat the needs of Species of Greatest Conservation Need that use that vegetation types in one or more successional stages. Conservation strategies address these particular successional stage needs as well as those of species that prefer a mosaic of successional stages.

Table 4.2: Landscape, Community, Habitat & Cover Type Categories

* Categories marked with an asterisk "*" are considered major categories for the purposes of organizing this report (24 in all). Conservation summaries were developed addressing characteristics and location, current and desired condition, SGCN using this habitat category, priority problems impacting this category, conservation strategies to address the problems and a list of other plans and planning entities with significant interest in this area.

***Landscapes**

Interior Forest Blocks
Connectivity Blocks
Surface Waters and Riparian Areas
Riparian Areas for Connectivity
Physical Landscape Diversity Blocks
Wildlife Road Crossing

Aquatic Communities

***Riparian Areas**

***Riverine** (Langdon et.al. 1998)

Brook trout
Brook trout-slimy sculpin
Blacknose dace-slimy sculpin
Blacknose dace-bluntnose minnow
Blacknose dace creek chub
Tessellated darter-fallfish
Blacknose dace-slimy sculpin
White sucker-tessellated darter

***Lower Connecticut River**

(Atlantic salmon-American shad community)

***Lower Lake Champlain Tributaries**

(Redhorse-lake sturgeon community)

Cultural Habitats

(Reschle 1990)

***Building & structures**

***Mine & Gravel Pits**

***Grassland & Hedgerows**

Grasslands
Hedgerow
Old field/shrub
Orchard

***Lakes & Ponds**

Dystrophic lakes
Meso-eutrophic lakes
Oligotrophic lakes
High elevation acidic lakes

***Lake Champlain**

***Young Forests**

(Successional Stages, Forest Cover Types, Eyre 1980, US Dept of Agriculture 2003)

Stages: Seedling/Sapling Sapling/Pole Timber, Pole Timber

Cover types

Boreal Conifers
Balsam fir
Black spruce
White spruce
Boreal Hardwoods
Aspen
Pin cherry
Paper birch
Spruce-Fir
Red spruce
Red spruce-balsam fir
Paper birch-red spruce-balsam fir
Pine and Hemlock
Eastern white pine

Table 4.2 continued: Terrestrial Natural Communities (Thompson & Sorenson 2005)

Open or Shrub Wetlands

***Open Peatlands**

Alpine peatland
Dwarf shrub bog
Black spruce woodland bog
Pitch pine woodland bog
Poor fen
Rich fen
Intermediate fen

***Marshes & Sedge Meadows**

Deep bulrush marsh
Deep broadleaf marsh
Shallow emergent marsh
Sedge meadow
Cattail marsh
Wild rice marsh

***Wet Shores**

Calcareous riverside seep
River cobble shore
Lakeshore grassland
Riverside sand or gravel shore
Outwash plain pondshore
River mud shore
Rivershore grassland

***Shrub Swamps**

Buttonbush basin swamp
Alder swamp
Alluvial shrub swamp
Sweet gale shoreline swamp
Buttonbush swamp

Forested Wetlands

***Floodplain Forests**

Silver maple-ostrich fern riverine floodplain forest
Lakeside floodplain forest
Silver maple-sensitive fern riverine floodplain forest
Sugar maple-ostrich fern riverine floodplain forest

***Hardwood Swamps**

Red maple-black ash swamp
Red maple-northern white cedar swamp
Calcareous red maple-tamarack swamp
Red or silver maple-green ash swamp
Red maple-black gum swamp
Red maple-white pine-huckleberry swamp

***Softwood Swamps**

Northern white cedar swamp
Spruce-fir-tamarack swamp
Black spruce swamp
Hemlock swamp

***Seeps & Vernal Pools**

Vernal pools
Seeps

Open Upland Communities

***Upland shores**

Riverside outcrop
Lake sand beach
Lake shale or cobble beach
Erosional river bluff
Sand dune

***Outcrops & Upland Meadows**

Alpine meadow
Boreal outcrop
Serpentine outcrop
Temperate acidic outcrop
Temperate calcareous outcrop

***Cliffs & Talus**

Boreal acidic cliff
Boreal calcareous cliff
Temperate acidic cliff
Temperate calcareous cliff
Open talus

Upland Forests & Woodlands

***Spruce-Fir Northern Hardwood Forest**

Subalpine krummholz
Montane spruce-fir forest
Lowland spruce-fir forest
Montane yellow birch-red spruce forest
Boreal talus woodland
Cold-air talus woodland
Red spruce-northern hardwood forest
Red Spruce-Heath Rocky Ridge Forest

***Northern Hardwood Forest**

Northern hardwood forest
Rich northern hardwood forest
Mesic red oak-northern hardwood forest
Hemlock forest
Hemlock-northern hardwood forest
Northern hardwood talus woodland

***Oak-Pine-Northern Hardwood Forest**

Limestone bluff cedar-pine forest
Mesic maple-ash-hickory-oak forest
Mesic Clayplain Forest
White pine-red oak-black oak forest
Dry oak forest
Dry Red Oak-White Pine Forest
Pine-oak-heath sandplain forest
Dry oak-hickory-hophornbeam forest
Red cedar woodland
Red pine forest or woodland
Pitch pine-oak-heath rocky summit
Dry oak woodland
Sand-Over-Clay Forest
Temperate Hemlock Forest
Temperate Hemlock-Hardwood Forest
Transition Hardwoods Limestone Forest

Literature Cited

- Allen, T., F.H.; Starr, 1982. Hierarchy: perspectives for ecological complexity. Chicago: The University of Chicago Press.
- Anderson, M.G., M. Clark, C.E. Ferree, A. Jospe, A. Olivero Sheldon, and K.J. Weaver. 2013. Northeast Habitat Guides: A companion to the terrestrial and aquatic habitat maps. Submitted to the Regional Conservation Needs Grants Program of the Northeast Association of Fish and Wildlife Agencies. The Nature Conservancy, Eastern Conservation Science, Eastern Regional Office. Boston, MA.
<http://rcngrants.org/content/guide-terrestrial-habitat-map>.
- Association of Fish and Wildlife Agencies (AFWA), Teaming With Wildlife Committee, State Wildlife Action Plan Best Practices Working Group. 2012. Best Practices for State Wildlife Action Plans—Voluntary Guidance to States for Revision and Implementation. Washington, DC: Association of Fish and Wildlife Agencies. www.fishwildlife.org/files/SWAPBestPractices.pdf.
- Barnes, R.B. 1979. Wildlife habitat from a forest resource inventory: is it possible? Transactions of the Northeast Section of the Wildlife Society 36:151-159.
- Cleland D.T., P.E. Avers, W.H McNab, M.E. Jensen, R.G. Bailey, T. King, and W.E. Russell 1997. National Hierarchical Framework of Ecological Units. Ecosystem Management: Applications for Sustainable Forest and Wildlife Resources., Mark S. Boyce and Alan Haney, editors. Yale University Press, New Haven.
- Eyre, F. H. ed. 1980. Forest Cover Types of the United States. Society of American Foresters. Washington, DC.
- Forman, R.T.T.; Godron, M. 1986. Landscape Ecology. New York: Wiley and Sons
- Gawler, S. C. 2008. Northeastern Terrestrial Wildlife Habitat Classification. Report to the Virginia Department of Game and Inland Fisheries on behalf of the Northeast Association of Fish and Wildlife Agencies and the National Fish and Wildlife Foundation. NatureServe, Boston, Massachusetts.
- Grossman, D.H., D. Faber-Langendoen, A.W. Weakley, M. Anderson, P. Bourgeron, R. Crawford, K. Goodin, S. Landaal, K. Metzler, K.D. Patterson, M. Pyne, M. Reid and L. Sneddon. 1998. International Classification of Ecological Communities: Terrestrial Vegetation of the United States. Volume I: The National Vegetation Classification Standard. The Nature Conservancy.
- Langdon, R., J. Andrews, K. Cox, S. Fiske, N. Kamman, and S. Warren. 1998. A classification of the aquatic communities of Vermont. The Nature Conservancy and the Vermont Biodiversity Project, Montpelier, VT.
- Oliver, C.D. 1981. Forest Development in North America Following Major Disturbances. Forest Ecology and Management 3:153-168
- Poiani, K.P., B.D. Richter, M.G. Anderson, and H.E. Richter. 2000. Conservation at multiple scales: functional sites, landscapes, and networks. BioScience 50:133-146
- Smith, R.L. 1980 Ecology and Field Biology, 3th edition. Harper & Row. New York.
- Reschke, Carol. 1990. Ecological Communities of New York State. Latham, NY: New York Natural Heritage Program, NYSDEC, www.dec.state.ny.us/website/dfwmr/heritage/EcolComm.htm
- Simberloff D. 1998. Flagships, umbrellas, and keystones: Is single-species management passe in the landscape era? Biological Conservation 83:247–257.
- Thompson, E. H., and E. R. Sorenson. 2005. Wetland, Woodland, Wildland—A guide to the natural communities of Vermont. University Press of New England, Hanover.
- U.S. Department of Agriculture, Forest Service. 2003. Forest inventory and analysis national core field guide, volume 1. U.S. Department of Agriculture, Forest Service, Forest Inventory and Analysis, 201 14th St., Washington, D.C., 20250.

Wessels, T. 2002. *The Granite Landscape: A Natural History of America's Mountain Domes, From Acadia to Yosemite*. The Countryman Press. Woodstock, VT

Northern Hardwood Forest Summary

Characteristics and Location

The Northern Hardwood Forest is best developed at Vermont's middle elevations and these are widespread in the state. Beech, sugar maple, and yellow birch are the predominant tree species, but hemlock, red oak, red maple, white ash, basswood, and white pine can be common as well, and red spruce makes an occasional appearance.

These are the dominant communities in nearly all biophysical regions, excepting the higher elevations of the Green Mountains and the warmer regions of the Champlain Valley, Taconic Mountains, and Southern Vermont Piedmont. Where the natural communities serve as landscape level habitat (i.e., matrix), they should be represented in large blocks of contiguous forest (1,000 acre to 20,000 acre blocks or larger) of various successional stages, elevations, and soils.

The natural communities that comprise Northern Hardwood forest formation habitat are found in every biophysical region of the state.

Natural communities of the Northern Hardwood Forest

Northern Hardwood Forest: A variable community, generally dominated by beech, sugar maple, and yellow birch. This community occurs as a landscape natural community type (i.e., matrix) throughout the state.

Rich Northern Hardwood Forest: High diversity hardwood-dominated forests of sugar maple, white ash, basswood, and hophornbeam, with excellent productivity and high herb diversity. These forests are closely associated with limestone and other calcium-rich bedrock types. Maidenhair fern, blue cohosh and wood nettle are characteristic herbs. This community occurs as a landscape natural community type (i.e., matrix) in the Taconic Mountains.

Mesic Red Oak-Northern Hardwood Forest: Northern hardwood species and red oak co-dominate. Mostly on south-facing slopes in the northern parts of Vermont, but also on cooler settings in the warmer regions of Vermont.

Hemlock Forest: Small forest patches dominated by hemlock, often on shallow soils and cool sites. Found throughout Vermont.

Hemlock-Northern Hardwood Forest: Mixed forest of hemlock and northern hardwoods. This community occurs as a landscape natural community type (i.e., matrix) in at least the Southern Vermont Piedmont and the Taconic Mountains.

Northern Hardwood Talus Woodland: A small patch community with characteristic species including yellow birch, mountain maple, red berried elder, rock polypody, and Virginia creeper.

Northern Hardwood Forest Condition

Historical Perspective: Northern Hardwood Forests have dominated the Vermont landscape for at least the last 4,500 years, a period over which there was a gradual cooling of the climate. These past forests are believed to have fairly closely resembled the composition of forests of today. Notable differences in the presettlement northern hardwood forests were the predominance of beech, making up over 40% of the trees (Siccama 1971) and the lower abundance of sugar maple. Although red spruce has decreased in abundance since presettlement times at mid-elevations, it has increased in abundance in valleys due to regeneration in old fields (Hamburg and Cogbill 1988). Similarly, white pine is now more abundant due to its regrowth in abandoned fields (Cogbill 2000). Presettlement forests also likely had much less red maple, white birch, and poplars than the forests of today, as these species are associated with younger forests (Cogbill 2000).

Current Condition: Vermont's Northern Hardwood Forest has become more widespread as farmland on the slopes and in the valleys has reverted to forest. However, human population growth and economic development result in forestland conversion and fragmentation that yield smaller blocks of contiguous Northern Hardwood Forest. While much of the Northern Hardwood Forest has been cleared or logged at one time, current land management trends will likely yield less early successional habitat in the future.

Desired Condition (SGCN Needs): Northern Hardwood Forest should be represented in both large blocks of contiguous forestland that contribute to the full complement of landscape level habitat for wide-ranging species and interior forest dwelling species, as well as in the natural community types that serve specific SGCN associated with that type. The large, contiguous forest blocks of Northern Hardwood Forest should exist in 1,000 to 20,000 acre blocks and should include representation of all successional stages, elevations, and soils should be well represented within each biophysical region. Prey wildlife species supported by northern hardwoods are an important component to maintaining several of the wide-ranging wildlife. In addition, the value of hard mast as wildlife food (i.e., nuts and acorns) from northern hardwoods is important for many SGCN with stands of bear-scarred American beech being a classic example. Interior forest conditions that occur in larger unfragmented forest blocks are critical for many species.

Implementing the 2005 Wildlife Action Plan

Actions by the Vermont Fish & Wildlife Department and partners to implement the Wildlife Action Plan in Northern Hardwood Forests since 2005 include:

Contiguous forest/habitat blocks and associated linkages were identified and prioritized as part of the "habitat block project" conducted from 2007 to 2014. Using GIS analysis of existing data, this project identified 4,055 unfragmented forest blocks in Vermont and ranked each block for its biological and physical landscape diversity values. The project also identified a modeling tool for identifying likely wildlife corridors in Vermont. Partners included Vermont Land Trust, the Forests, Parks & Recreation Department, The Nature Conservancy, Audubon Vermont, and Green Mountain National Forest. The project results are now used extensively in VFWD technical assistance to towns. The project report is "Vermont Habitat Blocks and Habitat Connectivity: An Analysis using Geographic Information Systems."

VFWD has acquired in fee and through conservation easements many high priority sites that include landscape scale Northern Hardwood Forests and provide critical landscape connectivity. These include Bird Mountain in Rutland County and Athens Dome in Windham County. From 2005-2013, the Department acquired 41 separate parcels (excluding fishing access areas) in fee totaling more than 4,100 acres to be added to WMAs or to create new WMAs. VFWD also acquired more than 2,300 acres under conservation easement during the same period. All of these projects either directly or indirectly benefit SGCN. Partner organizations including the Forests, Parks & Recreation Department, The Nature Conservancy, The Trust for Public Land, Vermont Land Trust and many local land trusts acquired and managed lands similarly benefitting SGCN.

VFWD provided technical assistance to private landowners, user groups and forest managers to manage for SGCN including, species associated with early successional and late successional habitat through the Natural Resources Conservation Service-funded WHIP and EQIP programs. Over the period from 2003-2013, the Department has worked on approximately 986 WHIP and 220 EQIP projects representing a total of 1,206 new wildlife habitat enhancement projects with as many private landowners throughout Vermont. Within each of these projects the following practices are the most common: Early Successional Habitat Development (Patch Cuts), Upland Wildlife Habitat Development (Mast and Apple Tree Release), and Invasive Species Control (in the form of Herbaceous weed control, and Brush Management).

VFWD provided technical assistance to every Vermont Regional Planning Commission and nearly every town on a variety of wildlife and land planning related issues, including SGCN conservation, habitat blocks, and wildlife corridors. [Conserving Vermont's Natural Heritage](#) (Austin et.al. 2004) was reprinted and distribution of this planning document continues.

The [Vermont Forest Roundtable](#) was first convened in 2006 as a venue for information exchange on keeping Vermont's forests as forests. Organized by the Vermont Natural Resources Council, the Roundtable regularly hosts consulting foresters, professional planners, state agency officials (including VFWD and VFPR), landowners, sportsmen, forest products industry representatives, conservation groups, biomass energy organizations and academics. The Roundtable formed with an initial focus on parcelization and forest fragmentation issues. It's since facilitated discussions on trends in Vermont's real estate market and rising forestland values, property tax policy, land use and conservation planning, estate planning, landowner incentive programs such as the Current Use Program, and the long-term sustainability of the forest products industry.

VFWD and the Vermont Agency of Transportation (VTrans) established a joint Wildlife-Transportation Steering Committee in 2007 to guide and support interagency cooperation to make Vermont's transportation system safer for both people and wildlife. VTrans published its [Vermont Transportation & Habitat Connectivity Guidance Document](#) in 2012. Together they currently support three wildlife camera and road tracking projects to advance our understanding of wildlife's use of transportation infrastructure. These studies are providing VTrans with improved infrastructure design criteria and VFWD with an enhanced understanding of wildlife movement at key locations in the state.

The [Staying Connected Initiative](#) was established in 2008 to maintain and improve landscape connectivity across the Northern Appalachian/Acadian region of the eastern U.S. and

Canada (NY, VT, NH, ME, MA and the eastern provinces) through research, land use planning, land management, land protection and road barrier mitigation. The comprehensive approach of the partnership allows the targeting of specific wildlife movement pinch points and coordinated action and affords some assurance that expensive state investment in wildlife-friendly transportation infrastructure is not undone by conflicting land uses in the near vicinity beyond the transportation right-of-way. Partners include VFWD, TNC, VNRC, VTrans, NWF, Wildlife Conservation Society, and the fish and wildlife and transportation agencies of partner states). VFWD has also worked closely with the North Atlantic Landscape Conservation Cooperative on a pilot conservation design for the Connecticut River watershed.

Audubon Vermont's [Foresters for the Birds](#) program, developed in partnership with the Vermont Parks & Recreation Department in 2008, provides foresters and landowners with education and technical assistance to manage forest lands for bird habitats. The program is proving to be an excellent mechanism to bring forest landowners with an interest in bird conservation into being active forest stewards.

In 2014-2015 VFWD and partners including Vermont Land Trust, Vermont Forests, Parks & Recreation, The Nature Conservancy, and the Northwoods Stewardship Center produced "Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape" (Sorenson et al. 2015). This report identifies coarse-filter conservation targets for landscape scale features including forest blocks, riparian areas, wildlife and landscape connectivity, and physical landscape diversity that are necessary to effectively conserve many finer scale conservation elements in the face of climate change and habitat loss, including natural communities, rare species, and SGCN.

Species of Greatest Conservation Need in Northern Hardwood Forest

High Priority

Canada Warbler (*Cardellina canadensis*)
 Jefferson Salamander (*Ambystoma jeffersonianum*)
 Fowler's Toad (*Anaxyrus fowleri*)
 Spotted Turtle (*Clemmys guttata*)
 Wood Turtle (*Glyptemys insculpta*)
 Butterflies-Hardwood Forest Group
 Silver-haired Bat (*Lasionycteris noctivigans*)
 Eastern Red Bat (*Lasiurus borealis*)
 Hoary Bat (*Lasiurus cinereus*)
 Woodland Vole (*Microtus pinetorum*)
 Long-tailed or Rock Shrew (*Sorex dispar*)
 Pygmy Shrew (*Sorex hoyi*)
 Water Shrew (*Sorex palustris*)
 Southern Bog Lemming (*Synaptomys cooperi*)

Medium Priority

Red-shouldered Hawk (*Buteo lineatus*)
 Chimney Swift (*Chaetura pelagica*)
 Black-billed Cuckoo (*Coccyzus erythrophthalmus*)
 Black-throated Blue Warbler (*Dendroica caerulescens*)
 Wood Thrush (*Hylocichla mustelina*)
 American Woodcock (*Scolopax minor*)
 Chestnut-sided Warbler (*Dendroica pensylvanica*)
 Ruffed Grouse (*Bonasa umbellus*)
 Blue-spotted Salamander (*Ambystoma laterale*)
 Spotted Salamander (*Ambystoma maculatum*)
 Four-toed Salamander (*Hemidactylium scutatum*)
 DeKay's Brownsnake (*Storeria dekayi*)
 Long-tailed Weasel (*Mustela frenata*)
 Hairy-tailed Mole (*Parascalops breweri*)
 Cinereus or Masked Shrew (*Sorex cinereus*)
 Smoky Shrew (*Sorex fumeus*)
 Common Gray Fox (*Urocyon cinereoargenteus*)
 Black Bear (*Ursus americanus*)

SGCN Note: Plant SGCN not listed here: 59 species. The SGCN invertebrate group listed here contains numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/ Information Need Category	Problem/ Information Need Detail	Rank
Habitat Conversion	Permanent conversion of large blocks of forestland to housing development, commercial development, and roads	High
Habitat Fragmentation	Break up of large forest blocks, riparian corridors, and migration paths	High
Impacts of Roads and Trails	Human and motorized disturbance from new roads and trails in sensitive habitats (e.g., denning sites, breeding sites, feeding areas)	High
Distribution of successional stages	Lack of appropriate landscape level approach to management resulting in a lack of either late or early successional habitat in appropriate size and juxtaposition.	High
Climate Change	May affect species composition	Low
Pollution	Acid rain, sulfur and mercury deposition	High
Invasive Exotic Species	Introduction of exotics species such as sudden oak death, hemlock wooly adelgid, beech bark disease, emerald ash borer, and garlic mustard could affect survival of species such as marten, black bear, Edwards hairstreak, West Virginia white, small mammals songbirds, etc.	High
Incompatible Recreation	Inappropriate location of ski, hiking, snowmobile trails, illegal ATV use, rock climbing.	Medium
Habitat Degradation	Loss of key feeding areas (beech stands, riparian areas, snags, cavity trees, etc.). Loss of dead and down material, fragmentation of contiguous forests..	High
Herbivory	Excessive deer and moose browsing alters tree regeneration, composition, and ability to compete with invasive exotics	Medium

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Develop a landscape level planning effort for public/private lands that address SGCN needs, including early and late successional species and integrates habitat for prey species (e.g., deer, moose, beaver, snowshoe hare) according to population objectives of species management plans	Adoption of goals and objectives for public and private lands by private and public land entities	USFWS, USFS, ANR, NRCS	PR, SWG, NRCS, USFWS
Expand the landscape-scale conservation design (Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape) completed in 2015 to address fine scale elements of the "coarse-fine filter" approach, including similar analyses and mapping to set conservation goals for all natural community types, habitats, SGCN and other species for which these serve as coarse filters, and to identify the SGCN and other rare species not "captured" by a coarse filter and need specific conservation attention.	Conservation targets (numeric and distributional goals) for natural community and habitat scale elements.	TNC, VLT, FPR, DEC, CHC	SWG

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Encourage long-term conservation efforts to keep forests forested including supporting Vermont's Use Value Appraisal, Forest Legacy, State Lands Acquisition and Management Programs and Vermont Housing and Conservation Board projects to protect intact forests.	Number of acres conserved	ANR, FPR, VLT, TNC, TPL, VHCB, Local Land Trusts	VHCB, VLT, Forest Legacy
Provide technical assistance to private landowners, user groups and forest managers to manage for SGCN including, SGCN associated with early successional and late successional habitat. Distribute <i>Landowners Guide - Wildlife Habitat Management for Lands in Vermont</i> (VFWD 2015)	Number landowners managing for SGCN.	NRCS, TNC, ANR, SAF, VWA, Covert	SWG
Financial incentives for private landowners to reduce problems and fragmentation to habitats for SGCN and to restore and enhance degraded habitats	Number of acres affected/restored	VFWD, NRCS	WHIP
Technical assistance to town and regional planning organizations, distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns contacted; No. towns incorporating wide-ranging species into planning	VFWD, RPCs, AVCC, VFS	VFWD
Technical assistance to state and federal land management agencies	Number of state and federal land management plans that include SGCN conservation.	ANR, USFWS, USFS	ANR, USFWS, NRCS
Work with VTrans to identify and maintain wildlife highway/road crossings and recreational user groups to avoid road and trail placement in sensitive habitat types.	Number functional linkages across highways/roads	VFWD, VTrans, VAST, GMHA	SWG, PR, VTrans
Manage deer and moose populations at levels that provide suitable harvest opportunities, but do not impair forest regeneration	Number of deer and moose/square mile.	VFWD	PR
Investigate the impact of invasive earthworms on Vermont forest habitats: survey the extent of infestations, and develop education and technical assistance programs, best management practices and rules as needed.		VFWD, VFPR, UVM	SWG

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
Bat Conservation Plan	Bat habitat conservation	VFWD
Partners in Flight	Bird conservation plan	PIF, VCE, VFWD, Audubon, USFWS
Vermont Forest Resources Plan (2015 Update Draft)	Conservation of healthy forests and the sustainable use and management of Vermont's Forests	VFPR
Vermont Transportation & Habitat Connectivity Guidance Document.	Informs transportation planning, design, construction, operations and maintenance activities and related wildlife and ecological systems monitoring	VTrans

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Cogbill, Charles V. 2000. Vegetation of the presettlement forests of northern New England and New York. *Rhodora*, Volume 102, No. 911, 250-276.
- Hamburg, S.P. and C.V. Cogbill. 1988. Historical decline of red spruce populations and climatic warming. *Nature*, Volume 331, 428-431.
- Horton, J., M. Langlais, T. Morton, D. Paganelli, N. Patch, S. Wilmot. May 2015. [Creating and Maintaining Resilient Forests in Vermont: Adapting Forests to Climate Change](#). Vermont Forests, Parks & Recreation Department.
- Siccama, Thomas G. 1971. Presettlement and present forest vegetation in northern Vermont with special reference to Chittenden County. *American Midland Naturalist*. Volume 85, No. 1, 153-172.
- Thompson, E.H. and E.R. Sorenson. 2005. Wetland, Woodland, Wildland, A Guide to the Natural Communities of Vermont.
- Vermont Agency of Transportation. 2012. [Vermont Transportation & Habitat Connectivity Guidance Document](#). Montpelier. VT
- Vermont Fish & Wildlife Department. 2015 Landowners Guide - Wildlife Habitat Management for Lands in Vermont.
- Sorenson, E., R. Zaino, J. Hilke and E. Thompson. 2015. Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape. Vermont Fish & Wildlife Department and Vermont Land Trust. Montpelier, VT.
- Vermont Forests, Parks & Recreation Department. 2015. Vermont Forest Resources Plan (2015 Update Draft).
- Vermont Forests, Parks & Recreation Department. April, 2015. 2015 [Vermont Forest Fragmentation Report: Report to the Vermont Legislature](#), Submitted to House and Senate Committees on Natural Resources and Energy and the House Committee on Fish, Wildlife, and Water Resources.

Spruce-Fir-Northern Hardwood Forest Summary

Characteristics and location

These forests characterize our coldest regions. At higher elevations and in low cold, moist areas, red spruce and balsam fir may dominate the canopy. Warmer or better drained sites have significant amounts of hardwoods (yellow birch, sugar maple, and beech) along with softwoods in the canopy. Human or natural disturbance can also lead to temporary dominance by hardwood species.

These forests occur where growing seasons are short, summers are cool, and winters are harsh. The conifer-dominated forests blanket our highest peaks above 2,500 feet as well as occurring in cold lowland pockets within large areas of Northern Hardwood Forest. The mixed forests of red spruce and northern hardwoods are more widely distributed.

Natural communities of the Spruce-Fir-Northern Hardwood Forest

Subalpine Krummholz: Low, dense thickets of balsam fir and black spruce at high elevations. Generally shallow to bedrock.

Montane Spruce-fir Forest: Dominated by red spruce and balsam fir, with occasional heartleaf birch, paper birch, and yellow birch. Higher elevations of the Green Mountains and other ranges generally above 2,500 feet.

Lowland Spruce-Fir Forest: Dominated by red spruce and balsam fir, with occasional white spruce, black spruce, paper birch, and yellow birch. Lowlands of Northeastern Highlands and cold valleys elsewhere.

Montane Yellow Birch-Red Spruce Forest: Mixed forest of mountain slopes at elevations typically from 2,000 to 2,900 feet, dominated by yellow birch and red spruce.

Red Spruce-Northern Hardwood Forest: Mixed forest of red spruce, yellow birch, sugar maple, beech, and balsam fir found on generally cooler and drier sites than Northern Hardwood Forest, generally below 2,400 feet elevation.

Red Spruce-Heath Rocky Ridge Forest: A forest of red spruce and heath shrubs (low blueberries) that occurs on ridgelines, low summits, and exposed ledges where there are thin, well-drained soils over acidic bedrock. It is uncommon, but forms small to large forest patches at 1,500' to 2,500' elevations in all but Vermont's lowest elevations.

Boreal Talus Woodlands: Rockfall slopes in cold settings dominated by heart-leaved paper birch with occasional red spruce. Appalachian polypody, skunk currant, and mountain maple are often abundant.

Cold-Air Talus Woodland: Rare. Found where cold air drains at the bases of large talus areas. Characteristic plants are black spruce, abundant mosses and liverworts, foliose lichens, and Labrador tea.

Spruce-Fir-Northern Hardwood Forest Condition

Historical Perspective: In recent geologic time, forests dominated by spruce and fir became established in eastern North America only as recently as 8,000 years ago (Webb 1987). A warming trend, known as the hypsithermal interval, occurred from about 6,000 to 4,000 years ago, at which time spruce and fir dominated forests were greatly reduced in distribution. There has been a general expansion of spruce and fir since this time associated with a general cooling of climate (Klyza and Trombulak 1999).

Balsam fir has increased substantially when compared to presettlement forests, likely the result of its competitive advantage over spruce after heavy cutting (Whitney 1994). Red spruce has decreased in abundance at mid-elevation as a result of natural climate warming after the "little ice age" and forest harvesting, whereas it has increased in abundance in valley settings as a result of regeneration in old fields (Hamburg and Cogbill 1988).

Current Condition: Many of the natural communities within the spruce–fir–northern hardwood formation exist at high elevations and are often on shallow, acidic, infertile soils. They are, therefore, particularly susceptible to global climate change and acid rain. Montane Spruce-Fir Forest is commonly considered one of the landscape forest types most vulnerable to expected climatic warming. In addition, fragmentation through permanent conversion of forest blocks to roads, houses, ridgeline development, and ski trails pose the most significant problems to this forest type and the species that depend on it.

Desired Condition (SGCN Needs): Many of the below listed SGCN depend upon large, contiguous, interconnected, forest blocks. Where they exist within a biophysical region, examples of large, intact blocks of appropriate natural communities should be conserved to ensure the long-term viability of the associated SGCN (i.e. Montane Spruce-Fir Forest: Blackpoll Warbler, Olive-sided Flycatcher, Bicknell's Thrush, Bay-breasted Warbler; Lowland Spruce Fir Forest: Black-backed Woodpecker, Gray Jay, Bay-breasted Warbler), and Spruce Grouse. Contiguous forest blocks will ideally exist in 1,000-20,000 acre blocks at various elevations and of various soil types. Conservation of these blocks should incorporate SGCN distribution and habitat needs.

Implementing the 2005 Wildlife Action Plan

Actions by the Vermont Fish & Wildlife Department and partners to implement the Wildlife Action Plan in Spruce-Fir-Northern Hardwood Forest's since 2005 include:

Contiguous forest/habitat blocks and associated linkages were identified and prioritized as part of the "habitat block project" conducted from 2007 to 2014. Using GIS analysis of existing data, this project identified 4,055 unfragmented forest blocks in Vermont and ranked each block for its biological and physical landscape diversity values. The project also identified a modeling tool for identifying likely wildlife corridors in Vermont. Partners included Vermont Land Trust, the Forests, Parks & Recreation Department, The Nature Conservancy, Audubon Vermont, and Green Mountain National Forest. The project results are now used extensively in VFWD technical assistance to towns. The project report is "Vermont Habitat Blocks and Habitat Connectivity: An Analysis using Geographic Information Systems."

VFWD has acquired in fee and through conservation easements many high priority sites that include landscape scale Spruce-Fir-Northern Hardwood Forests and provide critical landscape connectivity. These include Bird Mountain in Rutland County and Athens Dome in Windham County. All of these projects either directly or indirectly benefit SGCN. Partner organizations including the Forests, Parks & Recreation Department, The Nature Conservancy, The Trust for Public Land, Vermont Land Trust and many local land trusts acquired and managed lands similarly benefitting SGCN.

VFWD provided technical assistance to private landowners, user groups and forest managers to manage for SGCN including, species associated with early successional and late successional habitat through the Natural Resources Conservation Service-funded WHIP and EQIP programs. Over the period from 2003-2013, the Department has worked on approximately 986 WHIP and 220 EQIP projects representing a total of 1,206 new wildlife habitat enhancement projects with as many private landowners throughout Vermont. Within each of these projects the following practices are the most common: Early Successional Habitat Development (Patch Cuts), Upland Wildlife Habitat Development (Mast and Apple Tree Release), and Invasive Species Control (in the form of Herbaceous weed control, and Brush Management).

VFWD provided technical assistance to every Vermont Regional Planning Commission and nearly every town on a variety of wildlife and land planning related issues, including SGCN conservation, habitat blocks, and wildlife corridors. [Conserving Vermont's Natural Heritage](#) (Austin et.al. 2004) was reprinted and distribution of this planning document continues.

The [Vermont Forest Roundtable](#) was first convened in 2006 as a venue for information exchange on keeping Vermont's forests as forests. Organized by the Vermont Natural Resources Council, the Roundtable regularly hosts consulting foresters, professional planners, state agency officials (including VFWD and VFPR), landowners, sportsmen, forest products industry representatives, conservation groups, biomass energy organizations and academics. The Roundtable formed with an initial focus on parcelization and forest fragmentation issues. It's since facilitated discussions on trends in Vermont's real estate market and rising forestland values, property tax policy, land use and conservation planning, estate planning, landowner incentive programs such as the Current Use Program, and the long-term sustainability of the forest products industry.

VFWD and the Vermont Agency of Transportation (VTrans) established a joint Wildlife-Transportation Steering Committee in 2007 to guide and support interagency cooperation to make Vermont's transportation system safer for both people and wildlife. VTrans published its [Vermont Transportation & Habitat Connectivity Guidance Document](#) in 2012. Together they currently support three wildlife camera and road tracking projects to advance our understanding of wildlife's use of transportation infrastructure. These studies are providing VTrans with improved infrastructure design criteria and VFWD with an enhanced understanding of wildlife movement at key locations in the state.

The [Staying Connected Initiative](#) was established in 2008 to maintain and improve landscape connectivity across the Northern Appalachian/Acadian region of the eastern U.S. and Canada (NY, VT, NH, ME, MA and the eastern provinces) through research, land use planning, land management, land protection and road barrier mitigation. The comprehensive approach of the partnership allows the targeting of specific wildlife movement pinch points

and coordinated action and affords some assurance that expensive state investment in wildlife-friendly transportation infrastructure is not undone by conflicting land uses in the near vicinity beyond the transportation right-of-way. Partners include VFWD, TNC, VNRC, VTrans, NWF, Wildlife Conservation Society, and the fish and wildlife and transportation agencies of partner states). VFWD has also worked closely with the North Atlantic Landscape Conservation Cooperative on a pilot conservation design for the Connecticut River watershed.

Audubon Vermont's [Foresters for the Birds](#) program, developed in partnership with the Vermont Parks & Recreation Department in 2008, provides foresters and landowners with education and technical assistance to manage forest lands for bird habitats. The program is proving to be an excellent mechanism to bring forest landowners with an interest in bird conservation into being active forest stewards.

In 2014-2015 VFWD and partners including Vermont Land Trust, Vermont Forests, Parks & Recreation, The Nature Conservancy, and the Northwoods Stewardship Center produced "Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape" (Sorenson et al. 2015). This report identifies coarse-filter conservation targets for landscape scale features including forest blocks, riparian areas, wildlife and landscape connectivity, and physical landscape diversity that are necessary to effectively conserve many finer scale conservation elements in the face of climate change and habitat loss, including natural communities, rare species, and SGCN.

SGCN in Spruce-Fir Northern Hardwood Forest

High Priority

Bicknell's Thrush (*Catharus bicknelli*)
 Spruce Grouse (*Falcapennis canadensis*)
 Canada Warbler (*Cardellina canadensis*)
 Jefferson Salamander (*Ambystoma jeffersonianum*)
 Wood Turtle (*Glyptemys insculpta*)
 Beetles-Tiger Beetle Group
 Eastern Red Bat (*Lasiurus borealis*)
 Hoary Bat (*Lasiurus cinereus*)
 Lynx (*Lynx canadensis*)
 American Marten (*Martes americana*)
 Rock Vole (*Microtus chrotorrhinus*)
 Woodland Vole (*Microtus pinetorum*)
 Long-tailed or Rock Shrew (*Sorex dispar*)
 Water Shrew (*Sorex palustris*)
 Northern bog lemming (*Synaptomys borealis*)
 Southern Bog Lemming (*Synaptomys*)

Medium Priority

Northern Goshawk (*Accipiter gentilis*)
 Chimney Swift (*Chaetura pelagica*)
 Black-billed Cuckoo (*Coccyzus erythrophthalmus*)
 Olive-sided Flycatcher (*Contopus cooperi*)
 Black-throated Blue Warbler (*Dendroica caerulescens*)
 Bay-breasted Warbler (*Dendroica castanea*)
 Blackpoll Warbler (*Setophaga striata*)
 Gray Jay (*Perisoreus canadensis*)
 Black-backed Woodpecker (*Picoides arcticus*)
 Blue-spotted Salamander (*Ambystoma laterale*)
 Spotted Salamander (*Ambystoma maculatum*)
 Wolf (*Canis* ?)
 Mountain Lion (*Puma concolor cougar*)
 Long-tailed Weasel (*Mustela frenata*)
 Hairy-tailed Mole (*Parascalops breweri*)
 Cinereus or Masked Shrew (*Sorex cinereus*)
 Smoky Shrew (*Sorex fumeus*)
 Black Bear (*Ursus americanus*)

SGCN Note: Plant SGCN not listed here: 21 species. The SGCN invertebrate group listed here contains numerous species. Wolf and mountain lion utilize this spruce-fir northern hardwood forests but are addressed in the Landscape Forest Summary. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A1-A5.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/ Information Need Category	Problem/ Information Need Detail	Rank
Habitat Conversion	Permanent conversion of large blocks of forestland to housing development, and commercial development including: quarries, wind farm, roads, and recreational development	High
Habitat Fragmentation	Break up of large forest blocks, riparian corridors, and migration paths	High
Impacts of Roads and Trails	Human and motorized disturbance from new roads and trails in sensitive habitats (e.g., denning sites, breeding sites, feeding areas) Conversion of habitat to roads and trails may interrupt movement corridors and provide habitat for competing species.	Medium
Distribution of successional stages	Lack of appropriate landscape level approach to management resulting in habitat degradation (lack of either late or early successional habitat in appropriate size and juxtaposition).	Medium
Climate Change	Expected to alter species composition of many Montane-Spruce-Fir Northern Hardwood Forest types and communities and stress sensitive SGCN associated with these forests.	High
Pollution	Acid rain, sulfur and mercury deposition may affect prey base and vernal pool chemistry	High
Habitat Degradation	Loss of concentrated food, cover, breeding habitats (deer wintering areas, vernal pools, conifer wetlands, coarse woody debris etc.).	High
Incompatible recreation	Inappropriate location of ski, hiking, snowmobile trails, illegal ATV use, rock climbing.	Medium
Herbivory	Excessive deer and moose browsing alters native tree regeneration, composition, and resistance to invasive exotics.	Medium

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Expand the landscape-scale conservation design (Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape) completed in 2015 to address fine scale elements of the "coarse-fine filter" approach, including similar analyses and mapping to set conservation goals for all natural community types, habitats, SGCN and other species for which these serve as coarse filters, and to identify the SGCN and other rare species not "captured" by a coarse filter and need specific conservation attention. Identify and establish habitat for climate adaptation refugia	Numeric and distributional goals for landscape and natural community scale elements.	TNC, VLT, FPR, DEC	SWG

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Encourage long-term conservation efforts to keep forests forested including supporting Vermont's Forest Use Value Appraisal, Forest Legacy, State Lands Acquisition and Management Programs and Vermont Housing and Conservation Board projects to protect intact forests.	Number of acres conserved	ANR, FPR, VLT, TNC, TPL, VHCB, Local Land Trusts	VHCB, VLT, Forest Legacy
Technical assistance to private landowners, user groups and forest managers to maintain and enhance SGCN habitat in Spruce-Fir NHF.	Number landowners/user groups/forest managers managing for Spruce-Fir SGCN.	NRCS, TNC, VFWD, FPR, Coverts, SAF, VWA, Keeping Track	SWG
Financial incentives for private landowners to maintain and enhance SGCN habitat in Spruce-Fir NHF	Number of acres affected/restored	VFWD, NRCS	WHIP
Technical assistance to town and regional planning organizations to maintain and enhance SGCN habitat in Spruce-Fir NHF. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns contacted; Number of towns incorporating the needs of SGCN in Spruce-Fir NHF into planning	VFWD, RPCs, AVCC, VFS	VFWD
Technical assistance to state and federal land management agencies to maintain and enhance SGCN habitat in Spruce-Fir NHF	Number of state and federal land management plans for Spruce-Fir NHF providing for lynx and marten habitat. Number of state and federal land management plans for Spruce-Fir NHF that include SGCN in their management objectives.	ANR, USFWS, USFS, SAF	ANR
Maintain forested buffers along stream and rivers (See ANR buffer policy)	Number of miles of streams with intact buffers	ANR, VLT, TNC, NWF, Coverts	SWG, WHIP, Trout Unlimited, NRCS
Work with VTrans to identify and maintain wildlife highway/road crossings	Number functional linkages across highways/roads	VFWD, VTrans	SWG, PR, VTrans
Work with recreational groups to reduce the number of trails in sensitive habitats	Number of sensitive habitats with limited disturbance	GMC, VAST, VT Ski Area Association	
Increase cooperation/coordination between adjacent states and provinces to support and encourage trans-jurisdictional actions to address issues such as global climate change, acid rain and other pollutants.	Implementation of trans-jurisdictional actions.	USFWS, USFS, ANR, other states, TNC, Quebec,	USFWS, IAFWA
Manage moose populations at levels that provide suitable prey, but do not impair forest regeneration	Number of moose/square mile	ANR	PR
Investigate the impact of invasive earthworms on Vermont forest habitats: survey the extent of infestations, and develop education and technical assistance programs, best management practices and rules as needed.		VFWD, VFPR, UVM	SWG

Coordination with other plans

Plan or planning entity	Goal/Scope of plan	Lead

Bat Conservation Plan	Bat habitat conservation	VFWD
Spruce Grouse Recovery Plan	Spruce grouse reintroduction	VFWD
Partners in Flight	Bird conservation plan	PIF, VFWD, Audubon, USFWS
Riparian Buffer Guidance, and Riparian Buffers and Corridors Technical Papers (http://www.anr.state.vt.us/site/html/buff/anrbuffer2005.htm)	Helps in the development of recommendations and designs for Act 250-regulated projects that incorporate appropriate buffer zone widths for protecting riparian functions	ANR
2015 Update Vermont Forest Resources Plan (Draft)	Conservation and Management of VT Forests	VFPR
Vermont Transportation & Habitat Connectivity Guidance Document.	Informs transportation planning, design, construction, operations and maintenance activities and related wildlife and ecological systems monitoring	VTrans

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Hamburg, S.P. and C.V. Cogbill. 1988. Historical decline of red spruce populations and climatic warming. *Nature*, Volume 331, 428-431.
- Horton, J., M. Langlais, T. Morton, D. Paganelli, N. Patch, S. Wilmot. May 2015. [Creating and Maintaining Resilient Forests in Vermont: Adapting Forests to Climate Change](#). Vermont Forests, Parks & Recreation Department.
- Klyza, Christopher M. and Steven T. Trombulak. 1999. *The Story of Vermont: A Natural and Cultural History*. Middlebury College Press and University Press of New England.
- Sorenson, E., R. Zaino, J. Hilke and E. Thompson. 2015. *Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape*. Vermont Fish & Wildlife Department and Vermont Land Trust. Montpelier, VT.
- Thompson, E.H. and E.R. Sorenson. 2005. *Wetland, Woodland, Wildland, A Guide to the Natural Communities of Vermont*.
- Vermont Agency of Transportation. 2012. [Vermont Transportation & Habitat Connectivity Guidance Document](#). Montpelier. VT
- Vermont Forests, Parks & Recreation Department. 2015. *Vermont Forest Resources Plan (2015 Update Draft)*.
- Vermont Forests, Parks & Recreation Department. April, 2015. 2015 [Vermont Forest Fragmentation Report: Report to the Vermont Legislature](#), Submitted to House and Senate Committees on Natural Resources and Energy and the House Committee on Fish, Wildlife, and Water Resources.
- Webb, T., III. 1987. The appearance and disappearance of major vegetational assemblages: Long-term vegetational dynamics in eastern North America. *Vegetatio*, Volume 69, 177-187.
- Whitney, Gordon G. 1994. *From Coastal Wilderness to Fruited Plain: A History of Environmental Change in Temperate North America from 1500 to Present*. Cambridge University Press.

Oak-Pine-Northern Hardwood Forest Summary

Characteristics and Location

The Oak-Pine-Northern Hardwood Forest is best developed in the warmer regions of Vermont—the Southern Vermont Piedmont, Champlain Valley, and the lower elevations in the Taconic Mountains. Forest communities in this formation generally occur as large patches or locally as small patches within Northern Hardwood Forests and on dry, south-facing slopes and ridgetops. An exception to this is the Clayplain Forest of the Champlain Valley, which prior to European settlement occurred as a landscape scale (matrix) forest, but now has been reduced to forest fragments due to extensive agricultural use of the valley's clay soils. In the Oak-Northern Hardwood Forest Formation, hardwoods such as sugar maple, beech and white ash may be present, but warmer climate species such as red oak, shagbark hickory, and white oak are dominants of the forest canopy. White pine is also a prominent part of these forests.

The natural communities that comprise the Oak-Pine-Northern Hardwood forest type are diverse in their species composition, but all have species that occur in warmer climates, or on dryer sites such as south-facing slopes and ridges.

Natural communities of the Oak-Pine-Northern Hardwood Forest

Red Pine Forest or Woodland: Maintained by fire, these small areas are dominated by red pine, have very shallow soils, and have blueberries and huckleberries in the understory. They are widespread, and often surrounded by Northern Hardwood Forests.

Pitch Pine-Oak-Heath Rocky Summit: These are fire-adapted communities on dry, acidic ridgetops where red oak, white oak, pitch pine, scrub oak, and white pine are characteristic trees. Heath shrubs (blueberries and huckleberries) are abundant.

Limestone Bluff Cedar-Pine Forest: Northern white cedar dominates these areas of shallow soils over calcareous bedrock. Red pine, white pine, hemlock, and hardwoods are also present. Characteristic herbs are ebony sedge and rock polypody. This community has suffered high degree of loss from historic levels due to shoreline development.

Red Cedar Woodland: These are open glade-like communities on ledge crests, where red cedar is native and persistent, and grasses and sedges dominate the ground layer.

Dry Oak Woodland: These are very open areas with trees of low stature on dry, south facing hilltops. Grasses and Pennsylvania sedge are dominant on the forest floor.

Dry Oak Forest: These forests occur on rocky hilltops with very shallow, infertile soils. Red oak, chestnut oak and white oak can all be present; usually other tree species are absent. Heath shrubs dominate the understory.

Dry Oak-Hickory-Hophornbeam Forest: These forests occur on till-derived soils, but they are often found on hilltops and bedrock exposures are common. Soils are well drained, but are more fertile than in Dry Oak Forests. Red oak, sugar maple,

hophornbeam, and shagbark hickory are variously dominant. Sometimes sugar maple is the dominant tree, sometimes it is oak and hickory. Pennsylvania sedge forms lawns.

Mesic Maple-Ash-Hickory-Oak Forest: Sugar maple, white ash, hickories and red and white oak are present in varying abundances. This community needs better documentation.

Transition Hardwoods Limestone Forest: occurs in warm climate regions of Vermont where calcareous bedrock is close to the soil surface. Trees may be stunted and typical include sugar maple, white ash, shagbark hickory, basswood, hophornbeam, butternut, white oak, yellow oak (*Quercus muehlenbergii*), and bladdernut (*Staphylea trifolia*). A diverse community with a carpet of herbs reflecting calcium-rich conditions.

Mesic Clayplain Forest: Found on the Vergennes clay soils of the Champlain Valley, this forest is typically dominated by white oak, red maple, bur oak, swamp white oak, hemlock, and shagbark hickory. Maple-leaved viburnum is a typical shrub. Clayplain forests in Vermont have declined by 87.9% since pre-European settlement (Lapin 2003) due primarily to agricultural land use.

Sand-Over-Clay Forest: This large patch forest type occurs on specific soil types of the Champlain Valley where there is a sandy layer overlying clay. Hemlock, red maple, red oak, white oak, and black birch are all typical tree species and witch-hazel is a common shrub.

White Pine-Red Oak-Black Oak Forest: These forests are found on coarse-textured soils. Red and black oak co-dominate along with white pine. Beech and hemlock are also common. Heath shrubs are common in the understory.

Pine-Oak-Heath Sandplain Forest: This is a rare community type, found on dry sandy soils in warmer areas. Characteristic species are white pine, pitch pine, black oak, and red oak with an understory dominated by heath shrubs. Due to high development pressure, only 5% of the original 15,000 acres of sandplain forest in Chittenden County remain (Engstrom 1991).

Temperate Hemlock Forest: Similar to Hemlock Forest, but these dark, hemlock-dominated, small patch forests of warmer regions of the state have white oak, red oak, black birch as canopy associates, instead of northern hardwood species.

Temperate Hemlock-Hardwood Forest: Found in warmer climatic regions of Vermont, this mixed forest forest is co-dominated by hemlock, white oak, red oak, and black birch.

Transition Hardwood Talus Woodland: These talus woodlands are found in warmer areas, often on limestone but occasionally on slate, schist, granite, gneiss, or other rock. Some characteristic species are red oak, basswood, white ash, sweet birch, bitternut hickory, northern white cedar, hackberry, bulblet fern, and American yew.

Oak-Pine-Northern Hardwood Forest Condition

Historical Perspective: The natural communities that we recognize now are not static – they have changed dramatically over time as component species have migrated across the landscape in response to climatic change. The Oak-Pine-Northern Hardwood Forest Formation (and its characteristic species: pines, oaks, and hickories) provides a good example of how species migrations are independent of each other. After the retreat of the glaciers to the north, pine became well established in the northeastern United States by about 12,000 years ago, while oak was not well established until about 8,000 years ago, and hickory arrived in New England 2,000 to 3,000 years after the first increase in oak populations (Jacobson et al. 1987; Prentice et al. 1991).

It is often thought that white pine dominated the presettlement landscape of Vermont, but evidence from early land surveys indicates that it had a variable and restricted distribution (Cogbill 2000). Pine was abundant only in scattered areas of the Champlain and Connecticut River valleys, and was generally uncommon elsewhere. White pine has more than doubled in frequency since presettlement times, apparently due to its establishment and growth in abandoned agricultural fields (Cogbill 2000).

Current Condition: Of the three landscape level forests in Vermont, the Oak-Pine-Northern Hardwood Forest has been the most altered by human activities. The primary reason is that this forest type is most closely associated with the Champlain and Connecticut River Valleys – Vermont's most populated and prized agricultural regions. The Oak-Pine-Northern Hardwood Formation occurs in the warmest regions of the state that are generally the most desirable for settlement and agriculture. Human alteration of the landscape has most significantly altered two of natural community types of this formation: Mesic Clayplain Forest and Pine-Oak-Heath Sandplain Forest are now both considered rare forest types. In fact, in the southern Champlain Valley 87.9% of the Clay Plain Forest has been lost or degraded (Lapin 2003), primarily as a result of conversion to agricultural uses. One of Vermont's rarest and most threatened natural communities is the Pine-Oak-Heath Sandplain Forest of the northern Champlain Valley. As a consequence of its high value for residential development, it has been estimated that only 5% of the original 15,000 acres of sandplain forest now remain in Chittenden County (Engstrom 1991). Many of the rarest SGCN are directly associated with these communities.

Many of the other natural communities of this forest formation are small in size and often found in isolated settings. Several are found along drier ridgetops that make them less vulnerable to forestland conversion. However, fire suppression over the past 200 years or more has taken away one of the more important natural disturbances vital to regenerating some of the oak-pine forest types. Without fire, regenerating oak following timber removal is difficult, particularly when under the influence of herbivory (i.e., deer browsing, hare and rabbit girdling). Invasive plants (e.g., honeysuckle, buckthorn) and exotic insects (e.g., gypsy moth) can have significant effects on the quality of the wildlife habitat.

Desired Condition (SGCN Needs): Oak-Pine-Northern Hardwood Forest should be represented in both large blocks of contiguous forestland that contribute to the full complement of landscape level forest for wide-ranging species, as well as in the natural community types that serve specific SGCN associated with that type. Although contiguous forest blocks are limited in size and availability for the rarer forest types, where they exist,

large, contiguous forest blocks of Oak-Pine-Northern Hardwood Forest will ideally exist in 1,000 acre or more blocks of various elevations and soils. The oak component of this forest serves as important fall foods for numerous mammals, including some key prey species (e.g., deer, small mammals) for wide-ranging wildlife. Because much of the rarer Oak-Pine-Northern Hardwood Forest types have been converted to agriculture and development, the remaining fragmented blocks of these types will ideally be maintained, if not enlarged, as well as interconnected through forested or riparian corridors.

Implementing the 2005 Wildlife Action Plan

Actions by the Vermont Fish & Wildlife Department and partners to implement the Wildlife Action Plan in Oak-Pine Northern Hardwood Forest's since 2005 include:

Contiguous forest/habitat blocks and associated linkages were identified and prioritized as part of the "habitat block project" conducted from 2007 to 2014. Using GIS analysis of existing data, this project identified 4,055 unfragmented forest blocks in Vermont and ranked each block for its biological and physical landscape diversity values. The project also identified a modeling tool for identifying likely wildlife corridors in Vermont. Partners included Vermont Land Trust, the Forests, Parks & Recreation Department, The Nature Conservancy, Audubon Vermont, and Green Mountain National Forest. The project results are now used extensively in VFWD technical assistance to towns. The project report is "Vermont Habitat Blocks and Habitat Connectivity: An Analysis using Geographic Information Systems."

VFWD has been inventorying Oak-Pine-Northern Hardwood Forest types throughout Vermont since 2007 with the goal of identifying the most important forest blocks that are dominated by this forest formation. Approximately 100 sites have seen ecological and wildlife inventories so far.

In 2013, VFWD and The Nature Conservancy, and working with other partners, completed an inventory and prioritization of clayplain forest fragments in the Champlain Valley. The high priority examples of all clayplain forest types, including Wet Clayplain Forest and Wet Sand-Over-Clay Forest, were entered into the Department's Natural Heritage Database to be used for conservation planning.

VFWD has acquired in fee and through conservation easements many high priority sites that include landscape scale Oak-Pine-Northern Hardwood Forests and provide critical landscape connectivity. These include Bird Mountain and North Pawlet Hills, both in Rutland County). Partner organizations including the Forests, Parks & Recreation Department, The Nature Conservancy, The Trust for Public Land, Vermont Land Trust and many local land trusts acquired and managed lands similarly benefitting SGCN.

VFWD provided technical assistance to private landowners, user groups and forest managers to manage for SGCN including, species associated with early successional and late successional habitat through the Natural Resources Conservation Service-funded WHIP and EQIP programs. Over the period from 2003-2013, the Department has worked on approximately 986 WHIP and 220 EQIP projects representing a total of 1,206 new wildlife habitat enhancement projects with as many private landowners throughout Vermont. Within each of these projects the following practices are the most common: Early Successional

Habitat Development (Patch Cuts), Upland Wildlife Habitat Development (Mast and Apple Tree Release), and Invasive Species Control (in the form of Herbaceous weed control, and Brush Management).

VFWD provided technical assistance to every Vermont Regional Planning Commission and nearly every town on a variety of wildlife and land planning related issues, including SGCN conservation, habitat blocks, and wildlife corridors. [Conserving Vermont's Natural Heritage](#) (Austin et.al. 2004) was reprinted and distribution of this planning document continues.

The [Vermont Forest Roundtable](#) was first convened in 2006 as a venue for information exchange on keeping Vermont's forests as forests. Organized by the Vermont Natural Resources Council, the Roundtable regularly hosts consulting foresters, professional planners, state agency officials (including VFWD and VFPR), landowners, sportsmen, forest products industry representatives, conservation groups, biomass energy organizations and academics. The Roundtable formed with an initial focus on parcelization and forest fragmentation issues. It's since facilitated discussions on trends in Vermont's real estate market and rising forestland values, property tax policy, land use and conservation planning, estate planning, landowner incentive programs such as the Current Use Program, and the long-term sustainability of the forest products industry.

VFWD and the Vermont Agency of Transportation (VTrans) established a joint Wildlife-Transportation Steering Committee in 2007 to guide and support interagency cooperation to make Vermont's transportation system safer for both people and wildlife. VTrans published its [Vermont Transportation & Habitat Connectivity Guidance Document](#) in 2012. Together they currently support three wildlife camera and road tracking projects to advance our understanding of wildlife's use of transportation infrastructure. These studies are providing VTrans with improved infrastructure design criteria and VFWD with an enhanced understanding of wildlife movement at key locations in the state.

The [Staying Connected Initiative](#) was established in 2008 to maintain and improve landscape connectivity across the Northern Appalachian/Acadian region of the eastern U.S. and Canada (NY, VT, NH, ME, MA and the eastern provinces) through research, land use planning, land management, land protection and road barrier mitigation. The comprehensive approach of the partnership allows the targeting of specific wildlife movement pinch points and coordinated action and affords some assurance that expensive state investment in wildlife-friendly transportation infrastructure is not undone by conflicting land uses in the near vicinity beyond the transportation right-of-way. Partners include VFWD, TNC, VNRC, VTrans, NWF, Wildlife Conservation Society, and the fish and wildlife and transportation agencies of partner states). VFWD has also worked closely with the North Atlantic Landscape Conservation Cooperative on a pilot conservation design for the Connecticut River watershed.

Audubon Vermont's [Foresters for the Birds](#) program, developed in partnership with the Vermont Parks & Recreation Department in 2008, provides foresters and landowners with education and technical assistance to manage forest lands for bird habitats. The program is proving to be an excellent mechanism to bring forest landowners with an interest in bird conservation into being active forest stewards.

In 2014-2015 VFWD and partners including Vermont Land Trust, Vermont Forests, Parks & Recreation, The Nature Conservancy, and the Northwoods Stewardship Center produced “Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape” (Sorenson et al. 2015). This report identifies coarse-filter conservation targets for landscape scale features including forest blocks, riparian areas, wildlife and landscape connectivity, and physical landscape diversity that are necessary to effectively conserve many finer scale conservation elements in the face of climate change and habitat loss, including natural communities, rare species, and SGCN.

SGCN in Oak-Pine Northern Hardwood Forest

High Priority

Jefferson Salamander (*Ambystoma jeffersonianum*)
 Fowler’s Toad (*Anaxyrus fowleri*)
 Spotted Turtle (*Clemmys guttata*)
 Wood Turtle (*Glyptemys insculcata*)
 Timber Rattlesnake (*Crotalus horridus*)
 Eastern Ratsnake (*Pantherophis alleghaniensis*)
 Five-lined Skink (*Plestiodon fasciatus*)
 Butterflies-Hardwood Forest Group
 Beetles-Tiger Beetle Group
 Indiana Bat (*Myotis sodalis*)
 Silver-haired Bat (*Lasionycteris noctivagans*)
 Eastern Red Bat (*Lasiurus borealis*)
 Hoary Bat (*Lasiurus cinereus*)
 Woodland Vole (*Microtus pinetorum*)
 Southern Bog Lemming (*Synaptomys cooperi*)

Medium Priority

Northern Goshawk (*Accipiter gentilis*)
 Red-shouldered Hawk (*Buteo lineatus*)
 Chimney Swift (*Chaetura pelagica*)
 Black-throated Blue Warbler (*Dendroica caerulescens*)
 American Woodcock (*Scolopax minor*)
 Chestnut-sided Warbler (*Dendroica pensylvanica*)
 Ruffed Grouse (*Bonasa umbellus*)
 Blue-spotted Salamander (*Ambystoma laterale*)
 Spotted Salamander (*Ambystoma maculatum*)
 Four-toed Salamander (*Hemidactylium scutatum*)
 DeKay’s Brownsnake (*Storeria dekayi*)
 Long-tailed Weasel (*Mustela frenata*)
 Hairy-tailed Mole (*Parascalops breweri*)
 Cinereus or Masked Shrew (*Sorex cinereus*)

SGCN Note: Plant SGCN not listed here: 89 species. The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

Problem/ Information Need Category	Problem/ Information Need Detail	Rank
Habitat Conversion	Permanent conversion of forestland to housing development, commercial development, agriculture, and roads	High
Habitat Fragmentation	Break up of large forest blocks, riparian corridors, and migration paths. Wider ranging reptiles and birds depend upon contiguous habitat mosaics of 1000 ha or more.	High
Impacts of Roads and Trails	Human and motorized disturbance from new roads and trails in sensitive habitats (e.g., denning sites, breeding sites, feeding areas)	High
Inadequate Disturbance Regime	Fire Suppression: many habitats depend upon fire.	Medium
Climate Change	Alters water temperatures and levels for amphibians and reptiles.	Medium
Pollution	Acid rain affects on amphibians.	Medium
Habitat Degradation	Alteration of tree composition and loss of large, dead trees for cavities and roosts	Medium
Herbivory	Excessive deer browsing alters tree regeneration and composition	High
Invasive Exotic Species	Fragmented forest blocks encourage invasive plant species. Gypsy moth infestations affect oak productivity and survival.	High

Priority Conservation Strategies

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Develop a strategy and design that identifies the coarse-filter conservation values of landscape scale features (blocks of all forest types, connectivity, physical landscape diversity) and which SGCN and other rare species are expected to be captured by these coarse filters. Conduct a similar analysis and mapping to set conservation targets for all natural community types and the SGCN and other species for which these serve as coarse filters.	Conservation targets (numeric and distributional goals) for landscape and natural community scale elements.	TNC, VLT, FPR, DEC	SWG
Encourage long-term conservation efforts to keep forests forested including supporting Vermont's Forest Use Value Appraisal, Forest Legacy, State Lands Acquisition and Management Programs and Vermont Housing and Conservation Board projects to protect intact forests.	Number of acres conserved	ANR, FPR, VLT, TNC, TPL, VHCB, Local Land Trusts	VHCB, VLT, Forest Legacy
Technical assistance and/or financial incentives for private landowners, user groups and forest managers to maintain/enhance Oak-Pine NHF	Number of landowners managing land for SGCN	NRCS, TNC, ANR, SAF, VWA, VT Coverts	SWG
Financial incentives for private landowners to maintain and enhance SGCN habitat in Oak-Pine NHF	Number of acres affected/restored	VFWD, NRCS	WHIP
Technical assistance to town and regional planning organizations. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns & RPC's considering SGCN in their planning	VFWD, RPC's, AVCC, SAF, VWA, Coverts, VFS	VFWD
Technical assistance to state and federal land management agencies	No. state & federal land mgmt plans providing for SGCN, including use of prescribed fire	ANR, USFWS, USFS	ANR
Manage deer populations at levels that provide suitable harvest opportunities, but do not impair forest regeneration	Number of deer/square mile. Level of browse. Change in the # of wildlife road mortalities	ANR	PR
Continue working with VTrans & towns to identify and improve wildlife-highway/road crossings	Number of functional linkages across highways/roads	VFWD, VTrans	SWG, PR, VTrans
Increase cooperation/coordination between adjacent states and provinces. Develop trans-jurisdictional actions to address issues such as global climate change, acid rain & connectivity.	Implementation of trans-jurisdictional actions.	USFWS, USFS, ANR, other states, TNC, Quebec, VTA	USFWS, IAFWA
Investigate the impact of invasive earthworms on Vermont forest habitats: survey the extent of infestations, and develop education and technical assistance programs, best management practices and rules as needed.		VFWD, VFPR, UVM	SWG

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
Bat Conservation Plan	Bat habitat conservation	VFWD
ANR Long Range Management Plans	Management activities on ANR Lands	ANR
Green Mountain Forest Plan	Management activities on GMNF	USFS
Partners in Flight	Bird conservation plan	PIF, VFWD, Audubon, USFWS
The Nature Conservancy Champlain Valley Ecoregional Plan	Land conservation targets for the Champlain Valley Ecoregion	TNC
Champlain Basin Plan	Conservation of Champlain Basin resources	LCBP
Watershed Management Plans	Watershed plans for the Lake Champlain Basin	DEC
2015 Update Vermont Forest Resources Plan (Draft)	Conservation of healthy forests and the sustainable use and management of Vermont's Forests	FPR
Creating and Maintaining Resilient Forests in Vermont: Adapting Forests to Climate Change	Maintaining and improving forest resiliency	VFPR
Vermont Transportation & Habitat Connectivity Guidance Document.	Informs transportation planning, design, construction, operations and maintenance activities and related wildlife and ecological systems monitoring	VTrans

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Cogbill, Charles V. 2000. Vegetation of the presettlement forests of northern New England and New York. *Rhodora*, Volume 102, No. 911, 250-276.
- Engstrom, Brett. 1991. Sandplain Natural Communities of Chittenden County, Vermont: A report to the Vermont Department of Fish and Wildlife concerning the management and viability of a threatened habitat in Vermont. Unpublished report.
- Horton, J., M. Langlais, T. Morton, D. Paganelli, N. Patch, S. Wilmot. May 2015. [Creating and Maintaining Resilient Forests in Vermont: Adapting Forests to Climate Change](#). VT Forests, Parks & Recreation Department.
- Lapin, Marc F. 2003. Nature Conservation in an Agricultural Landscape: Forest Ecology, Fragmentation Analysis, and Systematic Site Prioritization, Southern Champlain Valley, Vermont, USA. PhD Dissertation, Cornell University.
- Jacobson, George L., Thompson Webb III, and Eric C. Grimm. 1987. Patterns and rates of vegetation change during the deglaciation of eastern North America. Pages 277-288 in W.F. Ruddiman and H.E. Wright, Jr., editors. *North America and Adjacent Oceans During the Last Deglaciation*. The Geology of North America, Volume K-3. Geological Society of America, Boulder, Colorado.
- Prentice, Colin., Patrick J. Bartlein, and Thompson Webb III. 1991. Vegetation and climatic change in eastern North America since the last glacial maximum. *Ecology*, Volume 72, No. 6, 2038-2056.
- Sorenson, E., R. Zaino, J. Hilke and E. Thompson. 2015. Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape. Vermont Fish & Wildlife Department and Vermont Land Trust. Montpelier, VT.
- Thompson, E.H. and E.R. Sorenson. 2005. *Wetland, Woodland, Wildland, A Guide to the Natural Communities of Vermont*.

Vermont Agency of Transportation. 2012. [Vermont Transportation & Habitat Connectivity Guidance Document](#). Montpelier. VT

Vermont Forests, Parks & Recreation Department. 2015. Vermont Forest Resources Plan (2015 Update Draft).

Vermont Forests, Parks & Recreation Department. April, 2015. 2015 [Vermont Forest Fragmentation Report: Report to the Vermont Legislature](#), Submitted to House and Senate Committees on Natural Resources and Energy and the House Committee on Fish, Wildlife, and Water Resources.

Young Forest-Old Forest Summary

Summary in development

Floodplain Forest Summary

Characteristics and Location

Floodplain forests are usually dominated by silver maple or occasionally sugar maple, with abundant ostrich fern or sensitive fern. They are closely associated with river and lake floodplains and have exposed mineral soils of alluvial origin.

Natural communities of the Floodplain Forest include: Silver Maple-Ostrich Fern Riverine Floodplain Forest, Silver Maple-Sensitive Fern Riverine Floodplain Forest, Sugar Maple-Ostrich Fern Riverine Floodplain Forest and Lakeside Floodplain Forest.

Natural communities of the Floodplain Forest:

Silver Maple-Ostrich Fern Riverine Floodplain Forest: This forest is found in the floodplains of moderate-gradient rivers. Silver maple and ostrich ferns are the dominant species and the soils are typically well drained sandy alluvium. Boxelder may be abundant in young forests.

Silver Maple-Sensitive Fern Riverine Floodplain Forest: These forests occur in the floodplains of large, low-gradient rivers. Silver maple is the dominant tree, but green ash and swamp white oak may be present. Sensitive fern and false nettle are characteristic. Soils are moist, typically mottled, silty alluvium.

Sugar Maple-Ostrich Fern Riverine Floodplain Forest: This uncommon floodplain forest type occurs along small to moderate sized high gradient rivers in areas of calcium-rich bedrock. Sugar maple, white ash, basswood, boxelder, and ostrich fern are common. There can be a diverse herbaceous layer. Soils are well drained sandy alluvium. Many examples of this community are uplands.

Northern Conifer Floodplain Forest: This rare floodplain forest occurs along small to moderate-sized rivers, primarily in northeastern Vermont. The silty alluvial soils typically support balsam fir, northern white cedar, white spruce, black cherry, and black ash.

Lakeside Floodplain Forest: These forests occur primarily within the flooding zone of Lake Champlain. Silver maple and green ash are the dominant trees. Herbs include sensitive fern, false nettle, marsh fern, white grass, and Tuckerman's sedge. Surface organic layers are present in the moist silty soils and there are mottles near the surface.

Floodplain Forest Condition

Historical Perspective: Although there is little specific information on distribution and composition of floodplain forests prior to European settlement in the region (Siccama 1971), it is expected that they covered large areas and were likely continuous bands of forest extending unbroken for miles along all of our major rivers. Forests of towering silver maple and American elm likely covered many of the active floodplains, with more diverse forests of sugar maple, red oak, and other species on higher terraces of former floodplains. (Sorenson et al. 1998). Although their total numbers were relatively small, evidence suggests that the Abenaki people that lived in Vermont concentrated their villages and agriculture on and

adjacent to the floodplains of the Connecticut River, other major rivers, and Lake Champlain (Klyza and Trombulak 1999).

Current Condition: High quality floodplain forests are now uncommon in Vermont because the vast majority of the floodplain forest in Vermont and the region has been converted to agricultural use. Floodplains have been prized as agricultural lands because of their high soil fertility associated with annual flooding and deposition and because of the absence of stones. As a result of their high value as agricultural lands, floodplain forests are now limited to fragments of their original size. The small percentage of riverine floodplains remaining in a forested condition is illustrated for Franklin County by a comparison made between the area of alluvial soils identified by the Natural Resources Conservation Service (USDA 1979) and the area of floodplain forests identified in a Vermont Fish and Wildlife Department floodplain forest inventory project (Sorenson et al. 1998). Although approximate, this comparison indicates that as little as 11% of the floodplains in Franklin County remain in a forested condition.

Significant changes to the flooding regimes of floodplain forests results from dam operation and the construction of roads, bridges, and culverts along rivers and in floodplains. Furthermore, the disturbed nature of many of the floodplain sites makes them vulnerable to invasive exotic plant species, such as goutweed (*Aegopodium podagraria*), garlic mustard (*Alliaria petiolata*), dame's-rocket (*Hesperis matronalis*), honeysuckle (*Lonicera* spp.), and Japanese knotweed (*Polygonum cuspidatum*) (Sorenson et al. 1998).

Desired Condition (SGCN Needs): Floodplain forest is essential to those SGCN that require habitat mosaics of aquatic and riparian areas and upland forest. Several of the species associated with floodplain forests require a riparian mosaic that depends upon functioning floodplain wetlands (e.g., pied-billed grebe, Odonata, American black duck); many of which are most abundant in the floodplains of larger river systems. Other species such as the water shrew and spotted salamander use floodplain forest directly adjacent to the stream or river. Lastly, there are some species that require large (10-1000ha) contiguous blocks of forested habitat along stream and rivers—these range from the bald eagle to the wood turtle. In all, floodplain forest provides habitat for a total of 49 wildlife SGCN and 28 plant SGCN. Desired conditions include functional floodplain forests in healthy examples (mature, unfragmented) distributed across their range. High water quality is also an essential element of floodplain forest quality. Focus should be give to the largest examples.

Implementing the 2005 Wildlife Action Plan

VFWD continues to focus conservation work on floodplain forests due to their high habitat functions, as well as being critical for river stability. An example is the acquisition of the Johnson Farm in the northeastern corner of Vermont, in the towns of Lemington and Canaan. Acquired in 2012, the Department now owns 283 acres and also manages public access on an additional 130 acres of eased land on the adjacent farm conserved by the Vermont Land Trust. The Johnson Farm Wildlife Management Area supports over eight miles of river and stream frontage, including 6.1 miles along the Connecticut River. Most of this shoreline area has well established riparian habitat or was subject to an extensive buffer restoration project in 2005-2006 through the USDA's Conservation Reserve Enhancement Program. The wetland natural community types found on the WMA includes floodplain

forest, alder swamp, sedge meadow, shallow emergent marsh, deep broadleaf marsh, cattail marsh, poor fen and river mud shore.

In 2013, the Department completed the Vermont BioFinder project, a map and database identifying Vermont's lands and waters supporting high priority ecosystems, natural communities, habitats, and species. A notable outcome of the project was a map of all aquatic features and the riparian areas/valley bottoms in which rivers and streams occur and the identification of these areas as critical conservation components for wildlife habitat, rare species, aquatic system health, and wildlife/landscape connectivity. The project mapping results for aquatic features, valley bottoms, and riparian connectivity together provide a tool for prioritizing restoration of riparian areas, including floodplain forests.

Species of Greatest Conservation Need in Floodplain Forests

High Priority

American Black Duck (*Anas rubripes*)
 Bald Eagle (*Haliaeetus leucocephalus*)
 Canada Warbler (*Cardellina canadensis*)
 Fowler's Toad (*Anaxyrus fowleri*)
 Wood Turtle (*Glyptemys insculpta*)
 Odonates-Bog/Fen/Swamp/Marshy Pond Group
 Freshwater Snails Group
 Butterflies-Hardwood Forest Group
 Lakes/Ponds Odonata Group
 Tri-colored bat (*Perimyotis subflavus*)
 Water Shrew (*Sorex palustris*)

Medium Priority

Great Blue Heron (*Ardea herodias*)
 Red-shouldered Hawk (*Buteo lineatus*)
 Chimney Swift (*Chaetura pelagica*)
 Cerulean Warbler (*Dendroica cerulea*)
 Black-crowned Night-heron (*Nycticorax nycticorax*)
 Pied-billed Grebe (*Podilymbus podiceps*)
 Spotted Salamander (*Ambystoma maculatum*)
 DeKay's Brownsnake (*Storeria dekayi*)
 Cinereus or Masked Shrew (*Sorex cinereus*)
 Black Bear (*Ursus americanus*)

SGCN Notes: Plant SGCN not listed here include 28 species. The SGCN invertebrate groups listed here contain numerous species Wolf and Black bear utilize floodplain forests but are addressed at the Landscape level. See that section for details. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need/Categories	Problem/Info Need Detail	Rank
Habitat Conversion	Agriculture and development	High
Habitat Fragmentation	Wider ranging birds, mammals, and reptiles require unfragmented habitat mosaics of 10-1000 ha or more	High
Inadequate Disturbance Regime	Dams, drainage ditching, filling, and runoff that affect flooding, erosion, and deposition	High
Habitat Degradation	Altering forest conditions along streams and rivers	High
Distribution of successional stages	Loss of mid-story forest cover due to lack of disturbance or active management.	Medium
Invasive Exotic Species	Loosestrife and common reed	High
Trampling or direct impacts	Human activity proximate to nesting birds	High
Inventory	Determine the location, distribution and condition of floodplain forests throughout their range.	Medium

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Locate additional floodplain forests not already mapped by FWD and assess management practices for these forests.	Number of sites located and assessed	ANR, FSA, UVM, VA	SWG
Identify riparian areas that are high priority for restoration of floodplain forest and other natural communities in order to increase river stability, water quality, wildlife habitat, and connectivity.	Number of sites located and on which restoration is successfully completed.	ANR, NRCS	SWG, WRP, EQUIP
Identify areas within the state with the largest matrix of floodplain forest for inclusion in conservation opportunity area.	Number of opportunity areas identified	ANR, UVM, NRCS	WRP, SWG
Consider protection of opportunity areas via acquisition of conservation easements, management leases and fee title acquisition	Number of sites conserved	ANR, VHCB, TNC, NRCS	VHCB, WRP, TNC
Manage exotic species on state owned sites and provide technical assistance to other landowners regarding control of exotics	Number of sites with control activities and/or invasive monitoring	ANR, NEPCoP, TNC, NRCS	WHIP, SWG
Technical assistance to private landowners, NGOs and government agencies to maintain and enhance floodplain forests for SGCN	Number of acres of floodplain forest managed for SGCN maintained, enhanced or restored. Number landowners incorporating SGCN into their land management.	NRCS, TNC, VFWD, FSA	WHIP, WRP, CREP, CRP, SWG
Technical assistance to towns and regional planning organizations to maintain and enhance floodplain forests for SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of acres of floodplain forest managed for SGCN maintained, enhanced or restored. Number landowners incorporating SGCN into their land management, Number of towns including SGCN in their planning.	NRCS, TNC, VFWD	SWG, WRP, CREP
Financial incentives for private landowners to maintain and enhance floodplain forests for SGCN	Number of acres conserved/restored	VFWD, NRCS	WHIP, WRP
Conservation easements on higher quality sites with greatest number of SGCN or T&E listed SGCN	Number of acres conserved for SGCN	ANR, VLT, TNC	VHCB, VLT, LIP
Manage or remove dams to restore more natural flooding regimes	Number sites with adequate flooding regimes	ANR, CT River Watershed Council	ACOE

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
Floodplain Forests of Vermont	Natural Community Inventory	ANR
Draft VT Bat Conservation Plan	Bat conservation	ANR

Bald eagle recovery plan	Bald eagle recovery	NWF, ANR
Partners in Flight	Bird conservation plan	ANR, VT Audubon, USFWS
2015 Update Vermont Forest Resources Plan (Draft)	Conservation of healthy forests and the sustainable use and management of Vermont's Forests	FPR
North American Waterfowl Plan	Waterfowl populations	USFWS, ANR, DU

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Horton, J., M. Langlais, T. Morton, D. Paganelli, N. Patch, S. Wilmot. May 2015. [Creating and Maintaining Resilient Forests in Vermont: Adapting Forests to Climate Change](#). Vermont Forests, Parks & Recreation Department.
- Klyza, Christopher M. and Steven T. Trombulak. 1999. The Story of Vermont: A Natural and Cultural History. Middlebury College Press and University Press of New England.
- Siccama, Thomas G. 1971. Presettlement and present forest vegetation in northern Vermont with special reference to Chittenden County. American Midland Naturalist. Volume 85, No. 1, 153-172.
- Sorenson, E., M. Lapin, B. Engstrom, and R. Popp. 1998. Floodplain forests of Vermont: some sites of ecological significance. NNHP, VT Fish and Wildlife Department.
- USDA, 1979. Soil Survey of Franklin County, Vermont. United States Department of Agriculture, Soil Conservation Service.
- Vermont Forests, Parks & Recreation Department. 2015. Vermont Forest Resources Plan (2015 Update Draft).
- Vermont Forests, Parks & Recreation Department. April, 2015. 2015 [Vermont Forest Fragmentation Report: Report to the Vermont Legislature](#), Submitted to House and Senate Committees on Natural Resources and Energy and the House Committee on Fish, Wildlife, and Water Resources.

Hardwood Swamp Summary

Characteristics and Location

Hardwood dominated swamps are the most common swamp type in the state. They are especially common in the warmer regions of the state on flatter topography and so reach their largest size and greatest abundance in the Champlain Valley and are least frequent in the Northeast Highlands. While two of the seven types are widely distributed, two others are restricted to a few biophysical regions, and the remaining three occur primarily in only a single biophysical region.

Hardwood swamps provide a number of functions, including flood storage, water quality protection, and fish, wildlife, and endangered species habitat. Because of their more open, deciduous canopy, hardwood swamps have more significant understory development than do softwood dominated swamps. This feature, along with their characteristic hummock and hollow topography, creates a landscape mosaic that provides an abundance of microhabitats.

Hardwood Swamp Natural Communities

The hardwood swamp formation includes the nine following natural community types:

Red Maple-Black Ash Seepage Swamp: This is the most common hardwood swamp type in the state. It occurs in all biophysical regions as either small or large patches. Although they occur in various settings, this natural community type is closely associated with groundwater seepage and does not typically experience surface flooding of long duration. While red maple is typically the dominant tree, black ash is very characteristic of this community. There are also other tree species present and well developed shrub and herbaceous layers.

Red Maple-Sphagnum Acidic Basin Swamp: This is another common swamp type that is widely distributed throughout the state. Typically it occurs in poorly drained basins with deep organic soils. Groundwater seepage is absent and the permanently saturated soils tend to be quite acidic. Since they occur in basins, most of these basin swamps are small and typically have no inlet or outlet streams. Red maple is the dominant tree, often with a co-dominance of yellow birch and various softwoods. The shrub layer is well developed, but the herb layer is less diverse, often with dominance by cinnamon fern. Moss cover typically approaches 100%.

Red Maple-Northern White Cedar Swamp: This uncommon community type exists as large patches mostly in the western part of the state. This community is limited to areas of calcareous bedrock and is often associated with floodplains, especially in the Champlain Valley. Although it can also occur in isolated basins, it can form huge wetland complexes where it is associated with larger rivers. Red maple, white cedar, and black ash typically dominate the canopy. Both the shrub and herbaceous layer tend to be sparse depending upon the degree of shading and the abundance of water-filled hollows.

Calcareous Red Maple-Tamarack Swamp: This is a rare community type that is restricted to areas of calcareous groundwater seepage. It is mostly restricted to the Vermont Valley with only a few examples in other biophysical regions. It typically

occupies small isolated basins, but may also occur as part of a large wetland complex. Red maple and tamarack dominate the canopy that can range from nearly closed to very open. In the latter situation, especially, a diversity of shrubs, herbaceous, and bryophyte species flourish.

Red or Silver Maple-Green Ash Swamp: This uncommon natural community type is largely restricted to the Champlain and Vermont Valleys. It occurs as large patches mostly associated with Lake Champlain. This swamp type typically undergoes extensive spring flooding that often results in saturated soils throughout the growing season. Although silver maple typically dominates, red maple and green ash may be very abundant. Both the shrub and herbaceous layer are well developed.

Red Maple-Black Gum Swamp: This rare community type occurs as small patches. It is mostly restricted to the southeastern part of the state with a few outliers in other regions. It occurs in small basins that are isolated from surface waters and that contain deep, saturated organic soils. Red maple and black gum co-dominate, but hemlock, yellow birch, and red spruce are also common.

Red Maple-White Pine-Huckleberry Swamp: This is a very rare natural community type that is restricted to the Champlain Valley. All known examples occur near the center of much larger wetland complexes. The canopy is dominated by red maple and white pine, but the most striking feature is the dense cover of huckleberry below. Typically, cinnamon fern dominates the herbaceous layer.

Wet Clayplain Forest: A wet forest occurring on the very poorly drained clay soil types of the Champlain Valley. These forests have a diversity of tree species, including swamp white oak, red maple, black ash, green ash, shagbark hickory, and hemlock. Due to their wetness, many Wet Clayplain Forests are the only forest fragments remaining in highly agricultural areas of the Champlain Valley.

Wet Sand-Over-Clay Forest: Similar to Wet Clayplain Forest, but occurring on wet soils with a sand layer overlying clay. Green ash, swamp white oak, and white pine are all common and there is typically a dense tall shrub layer.

Hardwood Swamps Condition

Current Condition: Although still relatively common in the state, hardwood swamps were formerly even more abundant. The primary activities resulting in loss of hardwood swamps were commercial and residential road development and road construction. Presently, agricultural conversion results in the greatest loss of swamps. Although protected by the Vermont Wetland Rules, many smaller examples are not mapped and therefore not protected under the regulations. Since many of these swamp types are most abundant in the lower, warmer regions of the state, they are subject to hydrologic impairment and incremental loss along the edges as the area around them is developed.

The primary problems to SGCN include agricultural conversion, invasion by exotics, altered hydrology, development and unrestricted logging.

Desired Condition: Forested wetlands provide habitat for a number of SGCN in the state. Hardwood dominated examples are especially diverse since they tend to be at lower

elevations and in warmer areas of the state than coniferous swamps. A total of 36 SGCN animals and 43 plant SGCN rely on one or more of these natural communities to provide habitat. Several of the species associated with hardwood swamps also require a habitat mosaic that depends on functioning swamps. Desired conditions include functional hardwood swamps in healthy examples (mature, unfragmented) across the distribution of their range. High water quality is also essential to habitat quality. Focus should be given to the largest examples.

Implementing the 2005 Wildlife Action Plan

In 2013, VFWD and The Nature Conservancy, and working with other partners, completed an inventory and prioritization of clayplain forest fragments in the Champlain Valley. The high priority examples of all clayplain forest types, including Wet Clayplain Forest and Wet Sand-Over-Clay Forest, were entered into the Department's Natural Heritage Database to be used for conservation planning.

Species of Greatest Conservation Need in Hardwood Swamps

High Priority

American Black Duck (*Anas rubripes*)
 Canada Warbler (*Cardellina canadensis*)
 Fowler's Toad (*Anaxyrus fowleri*)
 Spotted Turtle (*Clemmys guttata*)
 Wood Turtle (*Glyptemys insculpta*)
 Eastern Ratsnake (*Pantherophis alleghaniensis*)
 Odonates-Bog/Fen/Swamp/Marshy Pond Group
 Tri-colored bat (*Perimyotis subflavus*)
 Pygmy Shrew (*Sorex hoyi*)

Medium Priority

Great Blue Heron (*Ardea herodias*)
 Red-shouldered Hawk (*Buteo lineatus*)
 Chimney Swift (*Chaetura pelagica*)
 Rusty Blackbird (*Euphagus carolinus*)
 Black-crowned Night-heron (*Nycticorax nycticorax*)
 Pied-billed Grebe (*Podilymbus podiceps*)
 Blue-spotted Salamander (*Ambystoma laterale*)
 Spotted Salamander (*Ambystoma maculatum*)
 Four-toed Salamander (*Hemidactylium scutatum*)
 DeKay's Brownsnake (*Storeria dekayi*)
 Cinereus or Masked Shrew (*Sorex cinereus*)
 Black Bear (*Ursus americanus*)

SGCN Notes: Plant SGCN not listed here 43. The SGCN invertebrate group listed here contains numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need/Category	Problem/Info Need Detail	Rank
Habitat Conversion	Agriculture, road building, development	High
Hydrologic alteration	Sedimentation, development in watershed, road building, dams	High
Invasion by Exotic Species	Non-native species can spread and degrade the habitat for wildlife and eliminate some plant species	Medium
Habitat Degradation	Selective removal of cedar or black gum, logging on non-frozen ground, heavy cutting	High

Habitat Fragmentation	Roads, agriculture, and development break swamps into smaller patches	High
Inventory	Statewide inventory has been completed, but not all sites have been evaluated	Low

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Provide information to State Wetlands Office & EPA	Number of sites added to Natural Heritage Database	DEC, EPA	SWG, EPA
Locate hardwood swamps and assess their management practices.	Number of sites located and assessed	ANR, FSA, UVM, VA	SWG
Provide support and technical information to DEC wetlands office for designation of Class 1 wetlands	Number of wetlands reclassified to Class 1	ANR (FWD and DEC)	SWG, PR (technical assistance)
Identify areas within the state with the largest matrix of hardwood swamps for inclusion in conservation opportunity areas.	Number of opportunity areas identified	ANR, UVM	SWG
Consider protection of large hardwood swamps via acquisition of conservation easements, management leases and fee title acquisition	Number of sites conserved	ANR, VHCB, TNC	VHCB, TNC
Manage exotic species on state owned sites and provide technical assistance to other landowners regarding control of exotics	Number of sites with control activities and/or invasive monitoring. Number sites where invasive species are eliminated or controlled	NEPCoP, TNC, NRCS	SWG
Provide technical assistance to private landowners, NGOs and government organizations to plan and manage for SGCN in hardwood swamps. Distribute <i>Landowners Guide - Wildlife Habitat Management for Lands in Vermont</i> (VFWD 2015)	Number of acres maintained, enhanced or restored. Number landowners incorporating SGCN into their land management.	NRCS, TNC, VFWD, FSA	SWG, CREP, WHIP, CRP
Provide technical assistance to towns and regional planning organizations to plan and manage for SGCN in hardwood swamps. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of acres maintained, enhanced or restored. Number landowners incorporating SGCN into their land management, Number of towns including SGCN in their planning.	NRCS, TNC, VFWD	SWG, CREP
Financial incentives for private landowners	Number of acres conserved/restored	VFWD, NRCS	WHIP, WRP
Conservation easements on higher quality sites with greatest number of SGCN or T&E listed SGCN	Number of acres conserved for SGCN	ANR, VLT, TNC	VHCB, VLT, LIP

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Plan or planning entity	Goal/Scope of plan	Lead
New England Plant Conservation Program –	Recovery of various plant species in New England	ANR

various Conservation Plans		
North American Waterfowl Plan	Waterfowl conservation and management	ANR
Black Bear Plan	Bear conservation and management in Vermont	ANR

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Sorenson, E. R., R. Popp, M. Lew-Smith, B. Engstrom, M. Lapin, and M. Ferguson. 2004. Hardwood Swamps of Vermont: Distribution, Ecology, Classification, and Some Sites of Ecological Significance. NNHP, VT. Fish and Wildlife Department.
- Sorenson, E.R., B. Engstrom, M. Lapin, R. Popp and S. Parren. 1998. Northern White Cedar swamps and Red Maple-Northern White Cedar Swamps of Vermont: Some Sites of Ecological Significance. NNHP, VT. Fish and Wildlife Department.
- Thompson, E. H., and E. R. Sorenson. 2005. Wetland, Woodland, Wildland - A guide to the natural communities of Vermont. University Press of New England, Hanover and London,

Softwood Swamps Summary

Characteristics and Location

Most softwood swamps in Vermont are situated at higher elevations in the cooler regions of the state. The one exception are hemlock swamps which behave more like hardwood swamps and are located in the lower, warmer portions of the state. Because of the dense shading in softwood swamps, the understory shrub and herbaceous layers are generally quite sparse. Conversely due to these same moist shady conditions, bryophyte cover tends to be quite abundant. All of the natural communities in this formation occur as small patches on the landscape with the exception of spruce-fir tamarack swamps which occur as large patch communities.

Eight natural communities included within the softwood swamp group

Northern White Cedar Swamp: This is an uncommon natural community type that is associated with calcareous bedrock and groundwater seepage that makes the dissolved minerals available to the plants. Although it occurs in most of the state's biophysical regions, this community is more abundant in the northern half of the state since white cedar declines to the south. In addition to white cedar, balsam fir may be abundant, but the dense shading results in a sparse shrub and herb layers. Only bryophytes attain abundance in these swamps.

Northern White Cedar Sloping Seepage Forest: An uncommon type of cedar swamp known only from northeastern and northcentral Vermont and closely associated with calcium-rich bedrock. Northern white cedar dominated over a sparsely vegetated and gently sloping ground with mineral-enriched ground water flows just below the surface.

Boreal Acidic Northern White Cedar Swamp: An uncommon swamp of northeastern and north central Vermont found in landform settings with swamp watersheds and no inlet or outlet streams. Northern white cedar dominates the closed canopy over a carpet of wet Sphagnum mosses. **Spruce-Fir-Tamarack Swamp:** This uncommon natural community is totally absent from the warmer parts of the state. They typically occupy basins that are isolated from surface water movement and have deep organic soils. The canopy is dominated by red or black spruce, fir, and tamarack in varying abundance. Generally more tamarack is indicative of more mineral rich conditions while more black spruce is indicative of deeper peat and less enriched conditions. Despite the deep shade, a number of tall shrubs persist here, especially mountain holly and wild raisin. Herbs are sparse whereas bryophytes proliferate in the cool, moist conditions.

Red Spruce-Cinnamon Fern Swamp: This uncommon swamp type is most abundant in the southern Green Mountains, although it occurs throughout Vermont. Red spruce is dominant over a ground cover of cinnamon fern and Sphagnum mosses. Organic soils may be deep, but there is little mineral enrichment from groundwater.

Black Spruce Swamp: As the peaty soils become deeper and increasingly acidic and saturated, black spruce begins to replace the less tolerant red spruce. This community is restricted to only the coldest locations where they occupy basins that have gradually accumulated peat over the millennia. Black spruce dominates the canopy which is

generally rather low and sparse. These swamps have low shrub and herbaceous diversity due more to the cold, wet, acidic conditions than shading. In openings, low shrubs characteristic of bogs may be common, but bryophytes are ubiquitous throughout the community.

Hemlock-Balsam Fir-Black Ash Seepage Swamp: This uncommon swamp is found throughout Vermont at lower elevations. It is closely associated with mineral enrichment from groundwater seepage. Hemlock and/or balsam fir are dominant and black ash is typically present. Herbaceous plants and mosses are abundant and diverse, reflecting the mineral-enriched groundwater.

Hemlock-Sphagnum Acidic Basin Swamp: This rare swamp type occurs in the warmer regions of Vermont and only in landform settings with small watersheds. There are no inlet or outlet streams and peat accumulations are typically several feet or more. Hemlock is dominant over a moist swamp floor carpeted by Sphagnum mosses.

Softwood Swamps Conditions

Current Conditions: Softwoods swamps have been less impacted than either hardwood swamps or floodplain forest communities due to their location in the colder regions of the state and their generally saturated peat soils. As with the other two wetland types, softwood swamps also receive some protection from the Vermont Wetland Regulations. Nonetheless, they are still limited by habitat degradation and alteration, hydrologic impairment, and sedimentation from development on the fringes and in the watershed, road construction, and poorly planned logging. Exotic species, and herbivory, especially by moose, are also a concern. A potentially major problem for hemlock swamps is the presence in southern Vermont of the hemlock wooly adelgid, an introduced insect that could devastate the Vermont's hemlocks.

Desired Conditions: The eight natural communities in softwood swamp formation provide habitat for 26 SGCN animals. This includes a large number of birds, but also some turtles and salamanders. A total of 33 plant SGCN occur in softwood swamps; not surprisingly, the majority of which are bryophytes which thrive in the cool, moist, shady conditions. Only spruce-fir-tamarack swamps occur as large patches; however this community and northern white cedar swamps are often included within much larger wetland complexes. Three of the four community types exist as small patches, they are more easily protected; however protection would need to extend beyond the wetland boundary to include at least a portion of the watershed and should include connectivity to softwood swamps. In such situations protection would need to apply to the entire complex. Desired conditions include functional softwood swamps in healthy examples (mature, unfragmented) across the distribution of their range. High water quality is also essential to habitat quality.

Implementing the 2005 Wildlife Action Plan

In 2010, VFWD completed a statewide inventory of softwood swamps which included assessment of 162 sites. As a result of this project and data collected, the natural community classification was revised to include new types. Breeding bird and amphibian surveys were conducted so that animal species could be more closely associated with the natural

community types. Information was provided to landowners on the importance of their swamps for the habitat they provide and recommendations for management.

Species of Greatest Conservation Need in Softwood Swamps

High Priority

American Black Duck (*Anas rubripes*)
 Spruce Grouse (*Falcipennis canadensis*)
 Canada Warbler (*Cardellina canadensis*)
 Spotted Turtle (*Clemmys guttata*)
 Wood Turtle (*Glyptemys insculpta*)
 Odonates-Bog/Fen/Swamp/Marshy Pond Group
 American Marten (*Martes americana*)
 Rock Vole (*Microtus chrotorrhinus*)
 Northern bog lemming (*Synaptomys borealis*)
 Southern Bog Lemming (*Synaptomys cooperi*)

Medium Priority

Great Blue Heron (*Ardea herodias*)
 Red-shouldered Hawk (*Buteo lineatus*)
 Chimney Swift (*Chaetura pelagica*)
 Rusty Blackbird (*Euphagus carolinus*)
 Black-crowned Night-heron (*Nycticorax nycticorax*)
 Gray Jay (*Perisoreus canadensis*)
 Black-backed Woodpecker (*Picoides arcticus*)
 Pied-billed Grebe (*Podilymbus podiceps*)
 Blue-spotted Salamander (*Ambystoma laterale*)
 Spotted Salamander (*Ambystoma maculatum*)
 Four-toed Salamander (*Hemidactylium scutatum*)
 Cinereus or Masked Shrew (*Sorex cinereus*)
 Smoky Shrew (*Sorex fumeus*)
 Black Bear (*Ursus americanus*)

SGCN Notes: Plant SGCN not listed here 33. The SGCN invertebrate group listed here contains numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need/Category	Problem/Info Need Detail	Rank
Habitat Conversion	Development, road construction	High
Habitat Fragmentation	Roads and development fragment the habitat into smaller patches or from larger habitat mosaics for the wider-ranging species (e.g., wood turtle, American marten)	High
Hydrologic Alteration	Sedimentation, development in watershed, road building, dams	Medium
Invasion by Exotic Species	Non-native species (e.g., wooly adelgid) can spread and degrade the habitat for wildlife and eliminate some plant species	Medium
Habitat Degradation	Selective removal of cedar, logging on non-frozen ground, heavy cutting, lack of mature and overmature stands	High
Herbivory	Moose can eliminate regeneration in some community types	Medium
Inventory	Distribution, location and condition of these communities are not known. The ongoing statewide inventory needs to be completed to identify and protect the best examples	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Provide information to State Wetlands Office & EPA	Number of sites added to the Natural Heritage Database	DEC, EPA	SWG, EPA
Provide support and technical information to DEC wetlands office for designation of Class 1 wetlands	Number of wetlands reclassified to Class 1	ANR (FWD and DEC)	SWG, PR (technical assistance)
Locate additional softwood swamps of high significance and assess their management practices.	Number of sites located and assessed	ANR, FSA, UVM, VA	SWG
Identify areas within the state with the largest matrix of softwood swamps for inclusion in conservation opportunity areas.	Number of opportunity areas identified	ANR, UVM	SWG
Consider protection of large softwood swamps via acquisition of conservation easements, management leases and fee title acquisition	Number of sites conserved	ANR, VHCB, TNC	VHCB, TNC
Manage exotic species on state owned sites and provide technical assistance to other landowners regarding control of exotics	Number of sites with control activities and/or invasive monitoring. Number sites where invasive species are eliminated or controlled	ANR, NEPCoP, TNC, NRCS	SWG
Technical assistance and/or financial incentives to private landowners, NGOs and government organizations to maintain and enhance softwood swamps for SGCN,	Number landowners incorporating SGCN into their land management, Number of acres conserved/restored	NRCS, TNC, VFWD, FSA	SWG, WHIP, CREP, CRP, WRP
Technical assistance and/or financial incentives to towns and regional planning organizations to maintain and enhance softwood swamps for SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns considering SGCN in their planning for softwood swamps. Number of acres conserved/restored	NRCS, TNC, VFWD	SWG, WRP, WHIP, CRP, CREP
Conservation easements on higher quality sites with greatest number of SGCN or T&E listed SGCN	Number of acres conserved for SGCN	ANR, VLT, TNC, NRCS	VHCB, VLT, WRP

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Plan or planning entity	Goal/Scope of plan	Lead
New England Plant Conservation Program – various Conservation Plans	Recovery of various plant species in New England	ANR
Black bear plan	Bear population conservation and management	ANR
American Marten Recovery Plan	Recovery of American Marten in Vermont	ANR
North American Waterfowl Plan	Waterfowl conservation and management	ANR

References

Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. *Conserving Vermont's Natural Heritage. A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity*. Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.

Sorenson, E.R., B. Engstrom, M. Lapin, R. Popp and S. Parren. 1998. Northern White Cedar swamps and Red Maple-Northern White Cedar Swamps of Vermont: Some Sites of Ecological Significance. NNHP, VT. Fish and Wildlife Department.

- Sorenson, E., R. Popp, B.F. Engstrom, M. Lapin, and D. Farrell. 2010. Softwood Swamps of Vermont: Distribution, Ecology, Classification, and Some Sites of Ecological Significance. Natural Heritage Inventory, Vermont Fish and Wildlife Department,
- Thompson, E. H., and E. R. Sorenson. 2005. Wetland, Woodland, Wildland - A guide to the natural communities of Vermont. University Press of New England, Hanover and London,

Vernal Pools & Seeps Summary

Vernal pools are small, open-water wetlands that are filled by rain and snowmelt in spring or fall and are typically dry during the summer months. Such a pool is usually contained within a small forested basin, has no permanent inlet or outlet, and does not support fish. Forested swamps may also contain vernal pools in small depressions. During particular wet growing seasons, temporary pools may persist without drying completely. Years of filling and drying result in a unique type of set of conditions that supports a variety of wildlife specialized to take advantage of these conditions. Vernal pools are often rich in unique insects, molluscs, and other invertebrates, as well as amphibians. Vernal pools and adjacent forested uplands are critical breeding habitat for mole salamanders and wood frogs.

Seeps are small wetlands that occur on slopes or at the bases of slopes in upland forests. Groundwater discharge is evident at the seep margin. Scattered trees may be present but canopy closure is usually from the adjacent forest. Certain species are adapted to the living in these conditions, including some invertebrate and plant SGCN.

Vernal Pools & Seeps Condition

Current Condition: Vernal pools and the wildlife that use them face many problems, including direct loss of pools, degradation of pool quality, and alteration of the surrounding upland habitat that is critical for many amphibians non-breeding life stages. Hikers, their pets, and recreational vehicles that enter vernal pool risk destroying amphibian eggs and larvae and invertebrate SGCN. In addition, recreational vehicles that enter vernal pools can destroy the soil structure that is so important to maintaining these pools and the species that depend on them. Alterations within the forested basin that surrounds a pool can have significant impacts on the pool's hydrology and its species. Reduction in the volume of water that fills the pool means that drying will occur sooner. Loss of the adjacent canopy trees increases the solar energy reaching the pool, causing water temperature to rise more rapidly and drying the pool earlier in the warm season than usual. Premature drying has a negative impact on the invertebrates and young amphibians that require a minimum length of time (up to 4 months or more) to complete critical life stages. Removal of too many mature trees and downed logs in the surrounding upland habitat can impair the forest floor used by pool-breeding salamanders and frogs. Ditches and vehicle ruts in the surrounding forest often intercept spring migrating adults, luring them to lay eggs in spots that can dry well before the young can leave the water. Road construction or increased road traffic that bisects the upland amphibian habitat surrounding a vernal pool often results in the death of many of these animals as they make their annual migrations between the terrestrial and aquatic environments.

Seeps face problems similar to those of vernal pools. Activities that alter the hydrology of a seep to even a minor degree can eliminate the characteristics required by some wildlife species. The ecological significance of seeps (and vernal pools) is often not recognized during development planning, with the result being direct loss of these features.

Desired Condition (SGCN Needs): Functional vernal pools are those examples that are intact, well-buffered and interconnected to ensure productivity and movement of species associated with vernal pools. Spotted salamanders, blue-spotted salamanders, Jefferson salamanders, and wood frogs all use vernal pools for breeding. They spend almost their entire lives in the surrounding upland forests, moving up to 300 meters or more from the pool. The adults return for a brief period in the spring to leave their eggs. Water depth must

be great enough to cover the egg masses (generally 30cm or more) and provide continuous aquatic habitat until the young leave the pool (3-4 months, depending on the species and location). The terrestrial adults and juveniles can be found under cover material (logs, rocks, stumps) and in animal burrows in moist forest soils that have adequate leaf litter. Spotted turtles are seasonal users, foraging in vernal pools in the early spring. They require large wetland complexes and move between wetlands through the warm season. There are several insects, snails, fingernail clams, fairy shrimp, and other invertebrates that use vernal pools for their entire life cycle. During the dry months, these animals or their eggs remain on or under the soil surface, awaiting the return of water to the pool depression. Many other SGCN use vernal pools seasonally but do not require them.

Seeps are home to a few specialized SGCN as well as many more common species. The gray petaltail is a rare dragonfly that lays its eggs in forested seeps, where the nymphs remain and feed until reaching adulthood. Eastern Jacob's ladder is a threatened plant that is closely associated with seeps in Vermont.

Implementing the 2005 Wildlife Action Plan

VFWD contracted with the Vermont Center for Ecostudies and Arrowwood Environmental to map and inventory of vernal pools in Vermont. Approximately 5,000 vernal pools were mapped and approximately 1,200 were visited by project organizers and volunteers. The resulting mapping and database is used by the DEC Wetlands Office for regulatory purposes and has been the basis for conservation action.

VFWD has drafted conservation and management guidelines for vernal pool-breeding amphibians to provide the scientific justification for the critical nature of these pools and two "life zones" extending 100 feet and 600 feet from the pool edge. These guidelines are expected to be finalized soon and will provide the basis for site-specific vernal pool conservation and management.

Vernal pools are one of the 95 types of natural communities recognized in Vermont. Ranking specifications were developed for all natural communities, including vernal pools, in order to evaluate individual examples for their relative ecological significance and importance for amphibian breeding habitat.

Species of Greatest Conservation Need in Seeps and Vernal Pools

High Priority

Whip-poor-will (*Caprimulgus vociferus*)
 American Woodcock (*Scolopax minor*)
 Jefferson Salamander (*Ambystoma jeffersonianum*)
 Fowler's Toad (*Anaxyrus fowleri*)
 Spotted Turtle (*Clemmys guttata*)
 Wood Turtle (*Glyptemys insculpta*)
 Odonates-Bog/Fen/Swamp/Marshy Pond Group
 Freshwater Snails Group

Medium Priority

Great Blue Heron (*Ardea herodias*)
 Ruffed Grouse (*Bonasa umbellus*)
 Red-shouldered Hawk (*Buteo lineatus*)
 Prairie Warbler (*Dendroica discolor*)
 Pied-billed Grebe (*Podilymbus podiceps*)
 Blue-spotted Salamander (*Ambystoma laterale*)
 Spotted Salamander (*Ambystoma maculatum*)
 Four-toed Salamander (*Hemidactylium scutatum*)
 DeKay's Brownsnake (*Storeria dekayi*)
 Eastern Ribbonsnake (*Thamnophis sauritus*)
 Cinereus or Masked Shrew (*Sorex cinereus*)
 Smoky Shrew (*Sorex fumeus*)
 Black Bear (*Ursus americanus*)

SGCN Notes: Six plant SGCN are found in seeps and vernal pools. The SGCN invertebrate groups listed here contain numerous species. See individual species assessment

summaries in Appendix A for specific information about Species of Greatest Conservation Need listed here.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need/Category	Problem/Info Need Detail	Rank
1. Habitat Alteration	Thermal and hydrologic alterations that reduce the quality or usability of pools and seeps; modification of surrounding upland habitat needed to maintain dependent wildlife; creation of ditches and ruts that lure amphibians to unsuitable breeding habitat	High
2. Habitat Conversion	Direct loss of pool and seep habitat due to hydrologic manipulation, filling, draining, etc; loss of associated upland habitat due to development or conversion	Medium
3. Impact of roads and trails	Roads located too close to vernal pools kill amphibians as they attempt to migrate between the pools and upland habitat; loss of animals increases with traffic volume	Medium
4. Trampling or direct impacts	Destruction of and damage to amphibian eggs and invertebrate SGCN due to people and their pets entering vernal pools	medium
5. Incompatible recreation	Damage to habitat and loss of SGCN due to recreational vehicles entering vernal pools	High
6. Impacts of Roads and Trails	Trails leading to sensitive vernal pools bring recreational hikers and their pets	low
7. Pollution	Stormwater directed into pools carries sediments and contaminants that have a negative impact on this habitat and its aquatic populations	Medium
8. Disease	West Nile Virus control: Vernal pools may be annual targets of mosquito control, including the use of chemical and biological pesticides.	Medium
9. Inventory	Inventory needed for many SGCN, particularly those for which distributional and abundance information is greatly lacking	High
10. Monitoring	Monitor SGCN population trends to determine whether populations are able to persist; evaluate long-term effects of development near these habitats	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Monitor known SGCN populations and evaluate effects of development	Number of known SGCN sites monitored	ANR, EPA	SWG, EPA
Continue field investigation of vernal pools identified in statewide inventory of vernal pools and seeps important to SGCN	Number of completed field inventories	ANR, EPA	SWG, EPA
Identify areas within the state with the largest examples of seep and vernal pools for inclusion in conservation opportunity area.	Number of opportunity areas identified	ANR, VHCB, TNC	SWG
Promote conservation easements where appropriate	Number of acres of habitat protected and/or restored	ANR	LIP
Manage access at sensitive sites	Number of selected sites with managed/restricted access in place	ANR, USFWS, Green Mt. Club	

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Educate foresters, landowners, developers, and municipalities about the value of vernal pools and seeps and encourage behavior that conserves wildlife dependent on these features and the necessary surrounding habitat	Number of parties contacted	ANR, VFF, VINS, RPCs, towns	SWG, EPA
When appropriate, re-vegetate area surrounding pool or seep and restore hydrology	Number of sites restored; number of acres restored	ANR	EPA
Develop and distribute forestry guidelines for the protection and management of vernal pools and seeps	Number of forest management activities meeting vernal pool guidelines	ANR, USFWS, SAF, VWA, NRCS,	USFWS, USFS, SWG, WHIP
Technical assistance to towns and regional planning organizations to maintain and enhance vernal pools for SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns considering vernal pools and seeps in their planning.	VFWD	SWG
Develop recreational management plans for state lands where vulnerable, sensitive vernal pools and seeps occur	Number of recreational management plans adopted for state lands identified as having vulnerable vernal pools and seeps	ANR, VOGA, VASA,	
Work with VTrans and Federal Highway Administration to encourage protection of vernal pool, seep, and associated upland habitat when designing future roads; encourage the use of well-designed animal passage structures or other methods to allow safe passage for animals across existing roads	Number of cooperative projects that have avoided potential wildlife conflicts or restored safe passage	VFWD, VTrans, FHWA	

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Plan or planning entity	Goal/Scope of plan	Lead
State Outdoor Recreation Plan (SCORP)	A comprehensive recreation plan for state lands	FPR

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. *Conserving Vermont's Natural Heritage*. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Bellis, E. D. 1965. Home range and movements of the wood frog in a northern bog. *Ecology* 46: 90-98.
- Calhoun, A. J. K., and M. W. Klemens. 2002. Best development practices: conserving pool-breeding amphibians in residential and commercial developments in the northeastern United States. MCA Technical Paper No. 5, Metropolitan Conservation Alliance, Wildlife Conservation Society, Bronx, New York.
- Calhoun, A. J. K., and P. deMaynadier. 2004. Forestry habitat management guidelines for vernal pool wildlife. MCA Technical Paper No. 6, Metropolitan Conservation Alliance, Wildlife Conservation Society, Bronx, New York.

- Faccio, S. D. 2003. Postbreeding emigration and habitat use by Jefferson and spotted salamanders in Vermont. *Journal of Herpetology* 37 (3): 479-489.
- Gibbs, J. P. 1993. Importance of small wetlands for the persistence of local populations of wetland-associated animals. *Wetlands* 13:25-31.
- Petranka, J. W. 1998. *Salamanders of the United States and Canada*. Smithsonian Institution Press, Washington, DC.
- Petranka, J. W., M.P. Brannon, M. E. Hopey, and C. K. Smith. 1994. Effects of timber harvesting on low elevation populations of southern Appalachian salamanders. *Forest Ecology and Management* 67:135-147.
- Semlitsch, R. D. 1998. Biological delineation of terrestrial buffer zones for pond-breeding amphibians. *Conservation Biology* 12: 1113-1119.
- Semlitsch, R. D. and J. R. Bodie. 1998. Are small, isolated wetlands expendable? *Conservation Biology* 12: 1129-1133.
- Thompson, E. H., and E. R. Sorenson. 2005. *Wetland, woodland, wildland - A guide to the natural communities of Vermont*. University Press of New England, Hanover, NH.
- Vermont Wetlands Bioassessment Program, Department of Environmental Conservation, and Vermont Nongame and Natural Heritage Program, Department of Fish and Wildlife. 2003. An evaluation of the chemical, physical, and biological characteristics of seasonal Pools and northern white cedar swamps.
- Windmiller, B. S. 1996. The pond, the forest, and the city: spotted salamander ecology and conservation in a human-dominated landscape. Ph.D. dissertation, Tufts University, Medford, MA.
- Vermont Center for Ecostudies and Arrowwood Environmental. 2013. Vermont Vernal Pool Mapping Project 2009 – 2012, Final Report to the Vermont Department of Fish and Wildlife.
- Ferguson, M. 2006. *Conserving Pool-specialist Amphibian Habitat*. VT Fish and Wildlife Department. Draft.
- Faccio, F., M. Lew-Smith and A. Worthley, E. Sorenson and M. Ferguson. 2014. *Vernal Pool Ranking Specifications*. VT Fish and Wildlife Department.

Open Peatlands Summary

Characteristics and location

Open peatlands are wetlands that accumulate peat, a type of soil which consists of partially decomposed organic matter. These wetlands are permanently saturated with a stable water table at or near the soil surface, creating a nearly anaerobic soil environment. Seasonal flooding is generally lacking in these wetlands and mosses and liverworts are abundant. With the exceptions of Black Spruce Woodland Bogs and Pitch Pine Woodland Bogs, trees are generally absent or sparse due to the very low availability of dissolved oxygen and minerals in the soil and the saturated soil conditions. Bogs are a type of peatland with slightly raised surfaces that receive most of their water from precipitation, have acidic waters poor in minerals and nutrients, and are dominated by sphagnum mosses, heath shrubs, and in some areas black spruce. Fens, on the other hand, have slightly acidic to slightly basic mineral-rich waters from groundwater discharge and seepage, may be flat or gently sloping, and are dominated by sedges, grasses, and non-sphagnum mosses. Water in fens generally has higher oxygen concentrations than in bogs resulting in greater peat decomposition. There is a continuum, however, in the variations between bogs and fens.

Open Peatlands Natural Communities

The different natural community types in this group are all considered rare:

Dwarf Shrub Bog: These are open, acid wetlands with few trees and are dominated by heath shrubs and sphagnum moss. Size ranges from one to 600 acres in isolated kettlehole basins and as inclusions in larger wetland complexes. They occur throughout Vermont but are more common in the cooler regions.

Black Spruce Woodland Bog: Stunted black spruce trees cover 25 to 60 percent of the ground over heath shrubs and sphagnum moss. Found in cold climate areas. These bogs are generally less than 50 acres in size in Vermont and are found in the cooler regions of Vermont, including the Southern Green Mountains.

Pitch Pine Woodland Bog: Pitch pine covers 25 to 60 percent of the ground over heath shrubs and sphagnum moss. This community is known only from Maquam Bog at the mouth of the Missisquoi River. Small patches of this community are scattered across this larger wetland matrix.

Alpine Peatland: This community has characteristics of both bog and poor fen, but is distinguished by its high elevation and the presence of particular plants. It is found only on the highest peaks of the Green Mountains, particularly Mount Mansfield. By their nature, these communities are limited in size to very small patches.

Poor Fen: These are open, acid peatlands dominated by sphagnum moss, sedges, and heath shrubs. There is some mineral enrichment of surface waters. Poor fens are scattered in all biophysical regions of Vermont.

Intermediate Tall Sedge Fen: These open, slightly acid to neutral fens are dominated by tall sedges, non-sphagnum mosses, and a sparse to moderate cover of shrubs. Most examples are only several acres in size, with all known sites being less than 50 acres.

These fens are found only in areas with calcium-rich bedrock, which may occur in all regions outside of the Green Mountains.

Rich Fen: These fens are similar to Intermediate Fen but typically have shallower sedge peat and more mineral-enriched surface waters. Sedges and non-sphagnum mosses dominate, and shrubs are present. All documented examples are 6 acres or less in size. Rich Fens are restricted to areas with calcium-rich bedrock in the Piedmont, Vermont Valley, and limited areas of the Taconic Mountains.

Open Peatlands Condition

Current Condition: Open peatlands occur in a variety of situations across the Vermont landscape, from small, hydrologically-isolated basins to components of large wetland matrices. The primary problems to SGCN in open peatlands include recreation, exotic species, hydrologic alterations, climate change, and habitat conversion and degradation. Peatlands are popular destinations for outdoor recreationalists interested in experiencing unique natural areas, an activity that can prove detrimental to these fragile communities and their associated SGCNs if not properly managed. Trampling of plants is a major concern especially near urban centers and at the more accessible sites. Nutrient enrichment of runoff waters due to agriculture can lead to invasion by exotic plants as well as replacement of rare plant species by more generalist species. The integrity of bogs and fens can be limited by significant changes in adjacent land use, such as development and clear-cutting, that result in increases in runoff and changes in water quality. Activities that alter the quality and quantity of water received from the groundwater recharge zone can be devastating to fen communities. Climate change is especially a concern with the Alpine Peatlands due to rising temperatures and expansion of forest cover at high elevations. Alteration of precipitation quantity and timing patterns associated with climate change puts the peatlands at risk resulting from peat decomposition rates – a fine balance in peatlands between temperature, soil saturation, and dissolved oxygen levels, and microbial activity. Development of broadcasting facilities on mountain ridgelines also impacts this community type. Alteration of natural water level fluctuations in lakes, ponds, and streams associated with peatlands can also impact these wetlands. Prevention of natural disturbance regimes, including lightning-ignited fires, may limit the Pitch Pine Woodland Bog community.

Desired Condition (SGCN Needs): Many SGCN associated with this open peatlands, particularly some invertebrates and plants, are habitat specialists found only in these natural community types. Several SGCN dragonflies and damselflies require breeding and rearing habitat that is commonly described as bogs, fens, fen puddles, boggy ponds, boggy sloughs, and boggy streams. Many plants are found only in the wet, acid soils of bogs. Some vertebrates, such as bog lemmings (*Synaptomys* sp.) and spruce grouse are closely tied to bogs. Others, such as the blue-spotted salamander, four-toed salamander, spotted turtle, and water shrew, may rely on peatlands for habitat locally. Many of the other SGCN may utilize Open Peatlands but are not dependent on its specific characteristics (e.g., wood turtle, spruce grouse, and DeKay's Brownsnake).

Implementing the 2005 Wildlife Action Plan

Field work for a statewide inventory of Dwarf Shrub Bogs and Poor Fens and data is currently being analyzed. When analysis complete it will greatly expand our understanding of these rare natural community types and the birds associated with them.

The first statewide assessment of Vermont dragonfly and damselfly populations (collectively known as odonates) was completed in 2009. This survey (Pfeiffer, 2009) provides vital species distribution and occurrence information which has broadened our understanding of rare habitat-specialist dragonfly and damselfly SGCN. Habitat data collected as part of the study provides a comparative baseline for future population trend monitoring. Future efforts toward odonate SGCN conservation will continue to rely on the information resulting from this and future field studies.

Species of Greatest Conservation Need in Open Peatlands

High Priority

American Black Duck (*Anas rubripes*)
Spruce Grouse (*Falcipennis canadensis*)
Vesper Sparrow (*Pooecetes gramineus*)
Spotted Turtle (*Clemmys guttata*)
Wood Turtle (*Glyptemys insculpta*)
Odonates-Bog/Fen/Swamp/Marshy Pond Group
Lakes/ponds Odonata group
Moths group
Tiger Beetle group
Butterflies-Wetland Group
Hoary Bat (*Lasiurus cinereus*)
Water Shrew (*Sorex palustris*)
Northern bog lemming (*Synaptomys borealis*)
Southern Bog Lemming (*Synaptomys cooperi*)

Medium Priority

Red-shouldered Hawk (*Buteo lineatus*)
Chimney Swift (*Chaetura pelagica*)
Black-backed Woodpecker (*Picoides arcticus*)
Pied-billed Grebe (*Podilymbus podiceps*)
Lesser Yellowlegs (*Tringa flavipes*)
Blue-spotted Salamander (*Ambystoma laterale*)
Spotted Salamander (*Ambystoma maculatum*)
Four-toed Salamander (*Hemidactylium scutatum*)
Smooth Greensnake (*Opheodrys vernalis*)
DeKay's Brownsnake (*Storeria dekayi*)

SGCN Notes: Plant Species of Greatest Conservation Need not listed here: 62 species. The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem & Info Needs Category	Problem & Info Need Detail	Rank
1. Habitat Degradation	Significant land-use changes in adjacent areas can result in increases in runoff and changes in water quality (e.g. development, clear-cutting)	High
2. Habitat Conversion	Development of broadcasting facilities near alpine peatlands	Medium
3. Incompatible Recreation	Trampling of plants and soil in wetlands and on mountain tops	Medium
4. Hydrologic Alteration	Activities affecting the quantity and quality of ground water input and surface water runoff, or alter natural hydrologic regimes of associated water bodies	High

Problem & Info Needs Category	Problem & Info Need Detail	Rank
5. Impacts of Roads or Trails	Trails leading to sensitive peatlands bring recreational hikers	Medium
6. Pollution	Water quality is easily altered in peatlands and can bring about shifts in species composition (e.g., agriculture near rich fens)	High
7. Climate Change	Shifts in community composition in peatlands	Medium
8. Inadequate Disturbance Regime	Fire suppression inhibits pitch pine germination and results in shift in species composition	Medium
9. Statewide inventory of Open Peatland natural communities	Need to identify and locate best examples of these habitats that support the most SGCN	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Conduct statewide inventory of Open Peatland natural communities (Dwarf Shrub Bog and Poor Fen completed by 2015)	Number of sites inventoried	VFWD, EPA	SWG, EPA
Manage access at sensitive sites	Number of selected sites with managed/restricted access in place	ANR USFWS, Green Mt. Club	
Manage for natural disturbance regime at Maquam Bay	Work with USFWS to develop and implement a fire plan to promote this natural process	VFWD, USFWS	USFWS
Technical assistance to private landowners to maintain and enhance open peatlands for SGCN.	Number landowners incorporating SGCN into their land management.	ANR, EPA, USFWS, Landowners	LIP
Technical assistance to town and regional planning organizations to manage open peatlands for SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns considering SGCN in their planning	ANR, EPA, Regional Planning Comm.	SWG, EPA, VT Watershed Grants
Develop recreational management plans for state lands where vulnerable, sensitive open peatlands occur	Number of recreational management plans adopted for state lands identified as having vulnerable peatlands	ANR, VOGA, VASA	
Financial incentives for private landowners	Number of acres conserved	NRCS, VFWD, USFWS	NRCS, other USFWS grants
Acquisition/easement of high priority sites and their groundwater recharge areas	Number of acres acquired/enrolled	NRCS, VFWD, USFWS	NRCS, other USFWS grants
Increase enforcement of access restrictions at alpine peatlands	Number of hours of increased patrol	ANR, Green Mt. Club	

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Increase cooperation/coordination among states and provinces and develop trans-jurisdictional actions to address issues such as climate change and acid rain		State of VT, other states, CA provinces, US and CA federal governments	

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Plan or planning entity	Goal/Scope of plan	Lead
State Outdoor Recreation Plan (SCORP)	A comprehensive recreation plan for state lands	FPR

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Carle, F. C. 1994. Dragonflies and damselflies (Odonata) known to or likely to occur in Vermont. Report prepared for the Nongame & Natural Heritage Program, Vermont Fish & Wildlife Department. Waterbury, VT.
- Damman, A. and T. French. 1987. The Ecology of Peat Bogs of the Glaciated Northeastern United States: a community profile. U.S. Fish & Wildlife Service Biological Report 85(7.16).
- Glassberg, J. 1999. Butterflies through binoculars, the East; a field guide to the butterflies of eastern North America. Oxford University Press, New York, NY.
- Leonard, J. G., and R. T. Bell. 1999. Northeastern tiger beetles, a field guide to tiger beetles of New England and eastern Canada. CRC Press, Boca Raton, FL.
- Pfeiffer, Bryan (Wings Environmental). 2009 An Investigation of Odonata in Vermont Rivers and Peatlands. Submitted to the Vermont Fish & Wildlife Department 3/11/2010.
- Thompson, E., and R. Popp. 1995. Calcareous open Fens and Riverside Seeps of Vermont: Some Sites of Ecological Importance. Vermont Nongame and Natural Heritage Program, Vermont Fish and Wildlife Department.
- Thompson, E. H., and E. R. Sorenson. 2005. Wetland, woodland, wildland - A guide to the natural communities of Vermont. University Press of New England, Hanover and London.

Marshes and Sedge Meadows Summary

Marshes and Sedge Meadows provide some of the largest natural openings to be found in Vermont. These natural communities and the streams and ponds with which they are associated provide critical habitat for many species of wildlife. Often called emergent marshes, these open wetlands have less than 25 percent shrub or tree cover, and in many cases woody plants are absent. Hydrology is the single most important factor controlling these wetlands.

Marsh and Sedge Meadow Natural Communities

Six different natural community types are included in this group:

Shallow Emergent Marsh: This is a common and variable marsh type with mineral or shallow organic soils that are moist to saturated and only seasonally inundated. Several grasses, bulrushes, and Joe-pye weed may be abundant. This community is commonly associated with old beaver impoundments. This is a widespread natural community found throughout Vermont.

Sedge Meadow: These wetlands are permanently saturated and seasonally flooded. Soils are typically shallow organic muck, although mineral soils may be present in some wetlands. Tussock sedge or other sedges are dominant plants here. This common community is found throughout Vermont, most often along stream and pond margins and in beaver meadows.

Cattail Marsh: Common cattail or narrow-leaved cattail dominates these marshes. The muck or mineral soils are typically inundated with shallow standing water throughout the year, although the substrate may be exposed in dry years. Cattail Marshes range in size from less than an acre to over 500 acres along the shores of Lake Champlain. These common wetlands occur throughout the state but are most common at lower elevations.

Deep Broadleaf Marsh: Water depth in these marshes is typically over one foot deep for most of the year, although some may have only saturated soils in dry summers. Soils are organic. Common plants include pickerelweed, broad-leaved arrowhead, and giant bur-reed. This common community type is found throughout Vermont on the sheltered margins of lakes and ponds, on the slow-moving backwaters of larger rivers, and in isolated basins. The largest examples occur in lowland areas.

Wild Rice Marsh: These uncommon marshes are dominated by wild rice, with an organic soil substrate that is inundated with one to two feet of water throughout the summer. Wild Rice Marshes are found in wave-sheltered coves and on river deltas of Lake Memphremagog and Lake Champlain, and in the slow-moving backwaters of our larger rivers (Connecticut River and lower Champlain tributaries).

Deep Bulrush Marsh: These are marshes of open water along the shores of larger lakes and ponds where there is strong wave action. They are found throughout Vermont. Water depths can range from one to six feet. Soft-stem bulrush and hard-stem bulrush dominate most of these marshes, although marsh spikerush and other bulrushes may be abundant.

Marshes & Sedge Meadows Condition

Current Condition: These natural community types are not considered rare, but do provide critical habitat to many wildlife species, including SGCN. Sedge Meadows are often successional stages that would lead to forested wetlands if left undisturbed. Although they may occur in isolated basins, Marshes and Sedge Meadows are most commonly associated with water bodies (lakes, ponds, rivers) and other wetlands and, therefore, are subject to the same problems (e.g., pollution) as these associated communities. Even small examples of marshes that provide significant wildlife habitat or other functions and values are protected under Vermont Wetland Rules. Invasive exotic species are a major problem for some of these community types. Common reed and purple loosestrife can easily become established in Shallow Emergent Marshes, and water chestnut can crowd out native species in Deep Broadleaf Marshes. Alteration of the natural hydrologic regime by dam operation or creation of impoundments can significantly impact deeper water communities. Greater inventory information is needed for all these natural community types as well as further study on the identification and significance of particular problems.

Desired Condition (SGCN Needs): Marshes and Sedge Meadows support a host of wildlife species. A variety of SGCN are marsh specialists. Among others, these include many plants, dragonflies, damselflies, butterflies, and birds. Several dragonflies and damselflies require breeding and rearing habitat that is commonly described as marshy ponds, marshy edges of lakes, and marshes. Black terns, least bitterns, and soras spend the nesting season raising their young within marshes. Some other SGCN, such as spotted salamanders, and northern water snakes are commonly associated with these wetland types and may rely on them locally, but do not specifically require marshes to complete their life cycles. Pygmy shrews, smooth greensnakes, and chimney swifts are examples of more casual users that may be found foraging in marshes and sedge meadows.

Species of Greatest Conservation Need in Marshes and Sedge Meadows

High Priority

American Black Duck (*Anas rubripes*)
Black Tern (*Chlidonias niger*)
Northern Harrier (*Circus cyaneus*)
Sedge Wren (*Cistothorus platensis*)
Least Bittern (*Ixobrychus exilis*)
Vesper Sparrow (*Poocetes gramineus*)
Spiny Softshell (Turtle) (*Apalone spinifera*)
Fowler's Toad (*Anaxyrus fowleri*)
Spotted Turtle (*Clemmys guttata*)
Wood Turtle (*Glyptemys insculpta*)
Boreal Chorus Frog (*Pseudacris maculata*)
Odonates-Bog/Fen/Swamp/Marshy Pond Group
Freshwater Snails Group
Lakes/ponds Odonata group
Mayflies/Stoneflies/Caddisflies Group
Butterflies-Wetland Group
Hoary Bat (*Lasiurus cinereus*)
Pygmy Shrew (*Sorex hoyi*)
Water Shrew (*Sorex palustris*)

Medium Priority

Short-eared Owl (*Asio flammeus*)
Red-shouldered Hawk (*Buteo lineatus*)
Chimney Swift (*Chaetura pelagica*)
Bobolink (*Dolichonyx oryzivorus*)
Pied-billed Grebe (*Podilymbus podiceps*)
Sora (*Porzana carolina*)
Lesser Yellowlegs (*Tringa flavipes*)
Blue-spotted Salamander (*Ambystoma laterale*)
Spotted Salamander (*Ambystoma maculatum*)
Four-toed Salamander (*Hemidactylium scutatum*)
Smooth Greensnake (*Ophedrys vernalis*)
Northern Water Snake (*Nerodia sipedon*)
Easeterm Musk Turtle (*Sternotherus odoratus*)
DeKay's Brownsnake (*Storeria dekayi*)
Eastern Ribbonsnake (*Thamnophis sauritus*)
Long-tailed Weasel (*Mustela frenata*)
Muskrat (*Ondatra zibethicus*)
Southern Bog Lemming (*Synaptomys cooperi*)

SGCN Notes: Plant Species of Greatest Conservation Need not listed here: 24 species. The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problems/Info Need Categories	Problem & Info Need Detail	Rank
1. Habitat Conversion	Loss or fragmentation, particularly in small, unmapped (NWI) wetlands; ditching and plowing for agricultural use	High
2. Habitat Degradation	Cattle grazing	Medium
3. Hydrologic Alteration	Manipulation of the natural hydrologic regimes of associated water bodies through dam operation or impoundment can drastically impact deep water marshes in particular	High
4. Exotic Invasive Species	Crowding out of native plants and wildlife habitat by purple loosestrife, common reed, water chestnut, etc.	High
5. Pollution	Pollutants entering wetlands from runoff and tributaries can impact species and can bring about shifts in community composition	High
6. Statewide inventory of Marshes and Sedge Meadows	Inventory is needed for all natural community types, as well as further study on the identification and significance of particular problems	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Conduct statewide inventory of Marshes and Sedge Meadows	Number of sites inventoried. The number of high quality examples identified containing SGCN	VFWD, EPA	SWG, EPA
Protect wetlands not on NWI maps through alternative regulations (e.g., Act 250)	Number of acres conserved	ANR, Regional Planning Comm, ACOE	
Provide technical assistance and/or financial incentives to private landowners, towns and RPC's to maintain and enhance marsh and sedge meadows for SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number landowners incorporating SGCN into their land management, Number of towns including SGCN in their planning. Number of acres conserved	ANR, EPA, NRCS, TNC, RPC's, towns, VLCT, private landowners	NRCS, SWG, EPA, LCBP, VT Watershed Grants
Financial incentives for private landowners	Number of acres conserved	NRCS, VFWD, USFWS	NRCS, other USFWS grants
Acquisition/easement of high priority sites	Number of acres acquired/enrolled	NRCS, VFWD, USFWS	NRCS, other USFWS grants, Land trusts

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Use existing/new regulations to prevent damage of SGCN-important lake/pond-side and river-side wetlands caused by dam operation	Number of acres conserved	ANR, COE, Hydro operators, FERC	
Prevent loss of SGCN-important lake/pond-side and river-side wetlands caused by new impoundments	Number of acres conserved	ANR, COE, Hydro operators, FERC	

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Carle, F. C. 1994. Dragonflies and damselflies (Odonata) known to or likely to occur in Vermont. Report prepared for the Nongame & Natural Heritage Program, Vermont Fish & Wildlife Department. Waterbury, VT.
- Glassberg, J. 1999. Butterflies through binoculars, the East; a field guide to the butterflies of eastern North America. Oxford University Press, New York, NY.
- Laughlin, S. B., and D. P. Kibbe, eds. 1985. The Breeding Birds of Vermont. University Press of New England, Hanover, NH.
- Thompson, E. H., and E. R. Sorenson. 2005. Wetland, woodland, wildland - A guide to the natural communities of Vermont. University Press of New England, Hanover, NH.

Wet Shores Summary

Characteristics and Distribution

All of the natural communities contained within the upland shore formation occur as small patches scattered irregularly over the landscape. Four of the natural communities are widely distributed while the three rarest types are restricted to one or more biophysical regions. All the community types in this formation are non-forested and maintained in this early successional state by a combination of flooding, ice scour, and erosional processes. This makes wetland shores perhaps our most dynamic and changeable group of natural communities.

Wet Shores Natural Community Types.

This formation includes the seven following types:

Outwash plain pondshore: This is among the rarest natural communities in the state and is found only in the Southern Vermont Piedmont Biophysical Region. It occurs only on sloping shorelines that are seasonally exposed due to fluctuating water levels in the gravelly soils. The vegetation is characterized by sedge, rush, and herbaceous species, many of them annuals.

River mud shore: This is a common natural community type that occurs in all eight biophysical regions. It is restricted to slow moving rivers whose shorelines are exposed during times of low flow. This community type tends to be sparsely vegetated, primarily by annuals since the shore is often exposed late in the growing season.

River sand or gravel shore: This is a common natural community type that occurs in all eight biophysical regions. It is restricted to the swifter rivers where moving water creates sand and gravel deposits. Because of their dynamic nature they are sparsely vegetated, mostly by grasses and herbs but often with a woody component consisting of willows and cottonwood.

River cobble shore: This common natural community is widely distributed across the state along high-energy waterways. Due to their dynamic nature, this community is sparsely vegetated, mostly by grass and sedge species, but often with a woody component of willows and cottonwood.

Calcareous riverside seep: This is a rare natural community type that is known mostly from the Connecticut Valley. They are restricted to areas where calcareous groundwater seeps on to exposed bedrock on rivershores. The natural processes of flooding and ice scour serve to keep the community open while the limy seepage sustains a unique flora that includes many rare species of sedges, herbs, and bryophytes.

Rivershore grassland: This is a widely distributed natural community that occurs in more sheltered, and hence more stable, portions of our larger rivers. Since the natural river processes needed to maintain their open condition occur less frequently, this community tends to have more of a woody component of shrubs and low trees mixed in with the more abundant grasses.

Lakeshore grassland: This rare natural community type is restricted to the shores of Lake Champlain and Lake Memphremagog where it occurs on gently sloping shorelines that are kept open by waves, flooding, and ice. They tend to be very narrow in width, but may extend for considerable distances along the shore. The community is dominated by grasses, sedges, and forbs with a varying amount of woody species depending upon the frequency and intensity of the natural disturbance.

Wet Shores Condition

Current Condition: All of the natural communities within the wet shore formation are dependent upon the natural processes of flooding, wave action, and ice scour. As such, they all occur as small patches that are restricted to areas where these processes are focused. Since they are maintained in an open state, all of these natural community types provide a specialized habitat for animals and plants. Spiny softshell, spotted, and wood turtles, Fowler's toad, and tiger beetles all depend on one or more of these communities. Outwash plain pondshore and calcareous riverside seeps provide the unique habitat for plants and contain a disproportionate number of rare or Threatened species.

The primary problems to SGCN in this formation include hydrologic alteration, recreation, exotic species, and habitat conversion and degradation. Since all seven community types are dependent upon periodic disturbance by water, ice or wind, anything that prevents these natural processes from occurring would jeopardize the integrity and continued existence of the SGCN they harbor. These community types also support heavy recreational use, and trampling of vegetation is a major concern especially near urban centers and at the more accessible sites. The continual natural disturbance at these sites also provides excellent opportunity for invasive plants to become established, and recreational use adds to this potential. The river cobble shore and the two grassland types are especially subject to habitat conversion or degradation to create marinas, docks, and bathing beaches.

Desired Conditions: Although all the natural communities comprising the wet shore formation occur as small patches on the landscape, they all provide critical habitat to a number of SGCN that utilize both the aquatic and terrestrial environment or require unforested areas for basking, nesting, or foraging. A total of 22 animal and 31 plant SGCN are known to utilize the wet shore communities. To protect these sites we recommend the following activities:

Species of Greatest Conservation Need in Wet Shores

High Priority

American Black Duck (*Anas rubripes*)
Vesper Sparrow (*Pooecetes gramineus*)
Spiny Softshell (Turtle) (*Apalone spinifera*)
Fowler's Toad (*Anaxyrus fowleri*)
Spotted Turtle (*Clemmys guttata*)
Wood Turtle (*Glyptemys insculpta*)
Freshwater Snails Group
Beetles-Tiger Beetle Group
Cobblestone Tiger Beetle (*Cicindela marginipennis*)
Hoary Bat (*Lasiurus cinereus*)
Tri-colored bat (*Perimyotis subflavus*)
Water Shrew (*Sorex palustris*)

Medium Priority

Red-shouldered Hawk (*Buteo lineatus*)
Chimney Swift (*Chaetura pelagica*)
Pied-billed Grebe (*Podilymbus podiceps*)
Lesser Yellowlegs (*Tringa flavipes*)
Smooth Greensnake (*Opheodrys vernalis*)
Long-tailed Weasel (*Mustela frenata*)

SGCN Notes: Plant SGCN not listed here 31. The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Categories	Problem/Info Need Detail	Rank
Habitat Conversion	Construction of marinas, docks, bathing beaches, and other activities that remove shoreline vegetation	High
Hydrologic Alteration	Communities dependent upon wind, wave, and ice action	High
Incompatible Recreation	Intense use of shore disturbs wildlife, tramples rare plants, and introduces exotic species.	High
Invasion by Exotic Species	Non-native species can spread and degrade the habitat for wildlife and eliminate some plant species	Medium
Habitat Fragmentation	Roads and development fragment habitat along wet shores for species such as the wood turtle and Smooth Greensnake	High
Inventory	Distribution, location, and condition of this habitat are not known: A statewide inventory is needed to identify and locate the best examples of these habitats that support the most SGCN	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Conduct statewide inventory of upland shores	Number of sites inventoried. The number of high quality examples identified containing SGCN	FPR	SWG
Provide technical assistance to private landowners to prevent or mitigate hydrologic and recreational impacts to wet shores.	Number landowners incorporating SGCN into their land management	NRCS, TNC, VFWD	SWG
Manage exotic species on state owned sites and provide technical assistance to landowners regarding control of exotics	Number of sites with control activities and/or invasive monitoring. Number of acres conserved.	ANR, NRCS, TNC, EPA	LIP
Technical assistance to town & regional planning organizations to help maintain and/or enhance SGCN habitat, and to maintain natural processes and hydrologic conditions. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns/RPC's including SGCN in their planning	VFWD	VFWD
Conservation easements on higher quality sites with greatest number of SGCN or T&E listed SGCN	Number of acres conserved for SGCN	ANR, VLT, TNC	VHCB, VLT, LIP
Work with state and municipal managers to reduce and focus recreational impacts		ANR, VOGA	VFWD,

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead

New England Plant Conservation Program – various Conservation Plans	Recovery of various plant species in New England	ANR
State Outdoor Recreation Plan	A comprehensive recreation plan for state lands	FPR

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Thompson, E. H., and E. R. Sorenson. 2005. Wetland, Woodland, Wildland - A guide to the natural communities of Vermont. University Press of New England, Hanover and London

Shrub Swamps Summary

Characteristics and Distribution

All of Vermont's shrub dominated natural communities are wetlands, and most are thought to be retained in this early successional state by periodic flooding. Some of the community types, however, are likely to be more transitional and will eventually become forested. These transitional types are believed to have arisen following some type of disturbance either natural, such as a catastrophic flood or beaver activity, or artificially from past agricultural use. If beaver activity and natural flooding are allowed to continue, examples of this community should continue to replace themselves on the landscape.

Of the four natural community types included within this formation two occur as small patches while the remaining two occupy larger areas on the landscape. Only one of the communities, buttonbush swamp, is thought to be rare in the state and restricted in its distribution to four of the state's eight biophysical regions. The other three communities are distributed throughout the state.

The shrub swamp formation includes the following four natural community types:

Alluvial Shrub Swamp: This uncommon natural community type is found on alluvial soils in the floodplains of small rivers. This is a high energy, dynamic environment that receives regular flooding and ice scour. As the stream channel naturally wanders across the floodplain, the community also migrates. Senescent channels succeed to floodplain forest while alluvial shrub swamps thrive in newly established channels. While speckled alder is the dominant species here, black willow and boxelder can be very abundant under certain conditions. Ostrich fern typically dominates the ground layer although some grasses, herbs, and vines can also be common in more sheltered areas.

Alder Swamp: This is a common, widely distributed community type that occurs in a variety of settings including lakes and pond margins, backwater floodplains of rivers and streams, beaver flowerages, and poorly drained basins. Depending upon the frequency and duration of flooding, some examples may become more forested over time while others may remain shrub dominated. While speckled alder is the dominant shrub, shrubby willows, dogwoods, and young red maple may be locally abundant. Sedges and grasses along with sensitive fern and Joe pye weed typically dominate the ground layer.

Sweet Gale Shoreline Swamp: This uncommon natural community occupies shorelines of ponds and slow moving streams. This swamp typically occurs as a narrow floating mat, but the shrubs may also be rooted directly into the peaty shore. Sweet gale dominates this community, but meadow-sweet is usually also abundant. Leatherleaf may be co-dominant in more acidic, boggy conditions. Various sedge species typically dominate the ground layer.

Buttonbush Swamp: This rare natural community occurs in two different settings: on the edges of larger lakes and ponds and in poorly drained, isolated depressions – both settings in which water is retained through much of the growing season. Buttonbush is one of the few woody plants that can tolerate these seasonally flooded conditions. While in some examples buttonbush may grow so dense that nearly all other vegetation is

excluded, in other situations leatherleaf and meadow-sweet may be common. Depending upon the shrub density and degree of flooding, various herbs and grasses may become established.

Shrub Swamps Condition

Current Condition: Shrub swamps are a common wetland type and occur in a variety of situations that are either too wet or too frequently disturbed to allow trees to become established. Although some examples of Alder Swamps are becoming forested, new examples continually arise due to natural disturbance. As long as the natural processes of flooding, ice scour, and beaver activity continue unabated, shrub swamps will remain common in our landscape.

The primary problems to the communities and SGCN in this formation include habitat loss and fragmentation, suppression of the natural disturbance regime, hydrologic alteration, and invasive exotic species.

Desired Condition: Providing habitat for 30 SGCN makes shrub swamps among the more valuable community types for wildlife of concern in this state. Especially notable is the high number of amphibians included in this total. There are few plant SGCN associated with shrub swamps, however; they provide habitat for only six vascular plants and three bryophytes. Many types of shrub swamps are commonly associated with larger wetland complexes along river and streams. Maintaining the natural flooding regimes and other natural processes including beaver activity of these shrub swamps and associated forested swamps and marshes is critical to their long-term function. Maintaining upland buffers for shrub swamps are especially important for amphibian SGCN as well as for other species.

Species of Greatest Conservation Need in Shrub Swamps

High Priority

American Woodcock (*Scolopax minor*)
American Black Duck (*Anas rubripes*)
Black Tern (*Chlidonias niger*)
Vesper Sparrow (*Pooecetes gramineus*)
Spiny Softshell (Turtle) (*Apalone spinifer*)
Fowler's Toad (*Anaxyrus fowleri*)
Spotted Turtle (*Clemmys guttata*)
Wood Turtle (*Glyptemys insculpta*)
Boreal Chorus Frog (*Pseudacris maculata*)
Odonates-Bog/Fen/Swamp/Marshy Pond Group
Freshwater Snails Group
Butterflies-Wetland Group
Hoary Bat (*Lasiurus cinereus*)
Water Shrew (*Sorex palustris*)

Medium Priority

Red-shouldered Hawk (*Buteo lineatus*)
Chimney Swift (*Chaetura pelagica*)
Black-billed Cuckoo (*Coccyzus erythrophthalmus*)
Rusty Blackbird (*Euphagus carolinus*)
Pied-billed Grebe (*Podilymbus podiceps*)
Lesser Yellowlegs (*Tringa flavipes*)
Blue-spotted Salamander (*Ambystoma laterale*)
Spotted Salamander (*Ambystoma maculatum*)
Four-toed Salamander (*Hemidactylium scutatum*)
Smooth Greensnake (*Opheodrys vernalis*)
Northern Water Snake (*Nerodia sipedon*)
Eastern Musk Turtle (Stinkpot) (*Sternotherus odoratus*)
DeKay's Brownsnake (*Storeria dekayi*)
Eastern Ribbonsnake (*Thamnophis sauritus*)

SGCN Notes: Plant SGCN not listed here: 9. The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
Habitat Conversion	Development, road construction, docks, marinas	High
Habitat Fragmentation	Agriculture, roads	High
Hydrologic Alteration	Sedimentation, development in watershed, road building, dams, artificial lake fluctuations	High
Distribution of successional stages	Woodcock are negatively affected by maturing alder stands and adjacent openings.	High
Invasion by Exotic Species	Non-native species can spread and degrade the habitat for wildlife and eliminate some plant species	Medium
Inadequate Disturbance Regime	Suppression of natural processes such as eliminating beaver activity, limiting flooding, etc	High
Inventory	Distribution, location and condition of these communities are not known. A statewide inventory needs to be conducted to identify and protect the best examples	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Conduct a statewide inventory of shrub swamp natural communities	The number of high quality examples identified containing SGCN		
Provide information to State Wetlands Office & EPA	Number of sites discussed	DEC, EPA	SWG, EPA
Provide technical assistance and/or financial assistance in maintaining natural processes and hydrologic conditions to landowners, especially to municipal and private owners concerned with beaver activity.	Number landowners incorporating SGCN into their land management, Number of towns considering SGCN in their planning	NRCS, TNC, VFWD, RPC, VLCT	NRCS programs, LIP
Acquisition and conservation easements on higher quality sites with greatest number of SGCN or T&E listed SGCN	Number of acres conserved for SGCN	ANR, VLT, TNC	VHCB, VLT, LIP
Manage exotic species on state lands & provide technical assistance to other landowners regarding control of exotics	Number of sites with control activities and/or invasive monitoring	ANR, NEPCoP, TNC, NRCS	SWG

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
New England Plant Conservation Program – various Conservation Plans	Recovery of various plant species in New England	ANR
Partners in Flight Plan	Bird conservation	ANR, Audubon

References

Thompson, E. H., and E. R. Sorenson. 2005. Wetland, Woodland, Wildland—A guide to the natural communities of Vermont. University Press of New England, Hanover and London.

Upland Shores Summary

Characteristics and Distribution

All of the natural communities contained within the upland shore formation occur as small patches scattered irregularly over the landscape. Both the riparian associated natural communities occur in all eight biophysical regions of the state. In contrast, the three lakeshore natural communities are more restricted with both lake or shale cobble beach and sand dunes occurring in a single biophysical region and lake sand beach in three regions. Since all the upland shores are naturally kept open, all five natural community types provide specialized habitat for animals and plants. Riverside outcrops and sand dunes in particular provide habitat for some plants that occur nowhere else in the state. Generally SGCN have the best potential for persisting at sites with the most intact natural processes. These same sites likely provide the best and most abundant habitat for SGCN.

The upland shores formation includes the six following natural community types:

Acidic Riverside outcrop and Calcareous Riverside Outcrop: These uncommon to rare natural community types occurs throughout the state wherever bedrock is exposed along waterways, but one occurs on acidic bedrock like granite and one occurs on calcareous bedrock like limestone. They are dependent upon natural hydrologic processes that typically keep the sites open via either flooding or ice scour. These community types are sparsely vegetated, primarily by herbaceous species with only a few shrubs and vines able to withstand the regular disturbance regime – the species composition varies with the two types, reflecting the available calcium from the bedrock.

Erosional river bluff: This is a rare natural community type with a statewide distribution that is restricted to steep banks where soil is actively eroding. Both the nature of the soils and the intensity of the erosional action greatly influences the vegetative cover of these communities, but rarely are woody species frequent.

Lake or shale cobble beach: This uncommon natural community can occur on any large lake in the state, but the only significant examples occur on Lake Champlain. Due to the constant wave action and seasonal flooding and ice scour, they tend to be sparsely vegetated. Although the vegetation is mostly herbaceous, willows, cottonwood, silver maple, and ash can become established at their upper reaches.

Lake sand beach: This is a rare natural community with the most extensive examples on the shore of Lake Champlain, and only scattered examples occurring in other regions of the state. Their formation and sustenance depends upon a regular source of material this is subsequently transported and deposited by waves and/or wind. Due to the constant wind and wave action and seasonal flooding and ice scour, this community is largely kept open. Typically herbs, grasses, and low sedges dominate although willows, cottonwood, box elder, and ash often becomes established at their higher reaches.

Sand dune: This extremely rare natural community is restricted to the present and previous shoreline of Lake Champlain where dunes are situated on the leeward side of sand beaches. They are dependent upon a continual supply of depositional sand and will be adversely affected by anything that inhibits this process. Because of the shifting

nature of the substrate and the dry windy conditions, they are sparsely vegetated mostly by grasses, low sedges, and viney herbs. Cottonwoods, aspen, and gray birch eventually become established and make the dune system more stable.

Upland Shores Condition

Current Condition: All five community types within this formation are dependent upon continual disturbance by water, ice and wind and therefore occur in close proximity to lakes and rivers. They all reach their best development on the shores of Lake Champlain or other larger lakes and rivers in the state. Because they are desirable places to be, recreational use has impacted a number of our upland shores. The three lake associated shores are especially subject to habitat conversion or degradation to create marinas, docks, and bathing beaches. Trampling of plants is a major concern especially near urban centers and at the more accessible sites. The continual natural disturbance at these sites provides excellent opportunity for invasive plants to become established.

Desired Condition: Functional upland shores are primarily undeveloped sites where natural processes operate and human disturbance of SGCN is limited. Although all of the upland shores occur as small patches on the landscape, they provide a very specialized habitat that is utilized by a few SGCN and that may not be available elsewhere. Eight SGCN animals and one suite of species (tiger beetles) utilize upland shores. In addition, 33 SGCN plants are dependent upon this formation. To protect the natural communities contained within this formation we would do the following:

Species of Greatest Conservation Need in Upland Shores

High Priority

Common Tern (*Sterna hirundo*)
Spiny Softshell (Turtle) (*Apalone spinifera*)
Fowler's Toad (*Anaxyrus fowleri*)
Beetles-Tiger Beetle Group

Medium Priority

Peregrine Falcon (*Falco peregrinus*)
Chimney Swift (*Chaetura pelagica*)
Cinereus or Masked Shrew (*Sorex cinereus*)

SGCN Notes: Plant SGCN not listed here 31. The SGCN invertebrate group listed here contains numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
Habitat Conversion	Construction of marinas, docks, bathing beaches, retaining walls, rip-rap	Medium
Hydrologic Alteration	Communities dependent upon wind, wave, and ice action and supply of substrate	Medium
Incompatible Recreation	Intense use of beaches tramples rare plants, degrades dunes and introduces exotic species.	Medium
Invasion by Exotic Species	Non-native species can spread and degrade the habitat for wildlife and eliminate some plant species	Medium
Inventory	Distribution, location, and condition of this habitat are not known. A statewide inventory is needed to identify and locate the best examples of these habitats that support the most SGCN	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Conduct a statewide inventory of upland shore natural communities to identify the best sites and those with SGCN	Number of sites inventoried. Number of sites with SGCN identified	FPR	SWG
Technical assistance to private landowners to prevent or mitigate hydrologic alteration and recreational impacts and to conserve SGCN	Number landowners implementing conservation practices for SGCN	NRCS, TNC, VFWD	SWG
Technical assistance to town and regional planning organizations to prevent or mitigate hydrologic alteration and recreational impacts and to conserve SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns/organizations planning for SGCN conservation	VFWD	VFWD
Conservation easements on higher quality sites with greatest number of SGCN or T&E listed SGCN	Number of acres conserved for SGCN	ANR, VLT, TNC	VHCB, VLT, LIP
Work with state and municipal managers to reduce recreational impacts on these sites and to focus recreational impacts elsewhere.	Number of sites where recreational impacts are managed successfully.	ANR, VOGA	VFWD
Manage exotic species on state owned sites and provide technical assistance to private landowners to control exotics	Number of sites with control activities and/or invasive monitoring	ANR, NRCS	LIP

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
New England Plant Conservation Program – various Conservation Plans	Recovery of various plant species in New England	ANR
State Outdoor Recreation Plan (SCORP)	A comprehensive recreation plan for state lands	FPR

References

Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. *Conserving Vermont's Natural Heritage*. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.

Thompson, E. H., and E. R. Sorenson. 2005. *Wetland, Woodland, Wildland - A guide to the natural communities of Vermont*. University Press of New England, Hanover and London,

Outcrops & Upland Meadows Summary

Characteristics and Distribution

Outcrops and upland meadows are naturally unforested as a result of a number of factors: little or no soil, high winds, cold temperatures, and drought. Many of these factors are inter-related and work together in combination to limit tree growth. Outcrops and upland meadow are generally restricted to ridgetops and ledges where bedrock is exposed or close to the surface, and thus all the natural community types occur as small patches. They are often flat or gently sloping, but by definition, have slopes less than 60 degrees.

There are five outcrop and upland meadow natural community types:

Alpine Meadows: This very rare natural community is restricted to the highest elevations in the state where the harsh growing conditions severely restrict vegetative growth. There are only a few known examples, all restricted to the Northern Green Mountains Biophysical Region. Low herbaceous vegetation, primarily grasses and sedges, dominate although stunted fir and black spruce and various heath shrubs occur in more sheltered locations.

Boreal Outcrop: This relatively common natural community occurs at mid to high elevations and is distributed widely in the cooler areas of the state. It occurs in the Northern and Southern Green Mountains, Northern Piedmont, Northeastern Highlands, and Taconics Biophysical Regions. They are sparsely vegetated by scattered low trees, including fir, red spruce, yellow birch, red maple, heath shrubs, and grasses. In some examples, however, mosses and lichens can be abundant and even dominate.

Serpentine Outcrop: One of the rarest natural communities in the state, serpentine outcrops are restricted to the Northern and Southern Green Mountains where this rock type is exposed. Serpentine rocks and the soils derived from them are very low in most plant nutrients, instead containing high amounts of heavy metals that can reach levels that are toxic to plants. The result is a sparse flora, but also one that has adapted to these extremely harsh conditions.

Temperate Acidic Outcrop: This is a relatively common natural community that is absent from only the higher elevations and colder regions of the state. Trees, especially paper and gray birch, white and pitch pine, and red maple are frequent here although they are stunted and slow growing. Beneath them typically grow low heath shrubs, grasses, and various herbs. Mosses and lichens can also be very abundant.

Temperate Calcareous Outcrop: This is an uncommon natural community that is restricted to the warmer regions of the state; generally the Champlain and Connecticut River Valleys, the Taconics and the Vermont Valley. The community is limited to areas with calcareous bedrock and thus support a characteristic flora of lime-loving plants. Despite their exposure and resulting droughtiness, the availability of nutrients makes these outcrops more diverse than their more acidic counterparts.

Outcrops & Upland Meadows Condition

Current Condition: All of the natural communities contained within the outcrop and upland meadow formation are the result of specific conditions, and as such, they occur as

small patches and are scattered irregularly over the landscape. Only temperate acidic outcrops occur in all eight biophysical regions of the state. In contrast, alpine meadows are restricted to a single biophysical region and serpentine outcrops to two regions. Since they all are open communities within a generally forested matrix, all five natural community types provide a specialized habitat for animals and plants. They are important basking sites for reptiles, and alpine meadows, serpentine outcrops, and temperate acidic outcrops in particular provide habitat for many plants that occur nowhere else in the state.

The primary problems to SGCN in this category include recreation, exotic species, climate change, and habitat conversion and degradation. Since all five community types provide vistas, they are often a destination for hikes, skiers, and climbers. Trampling of plants is a major concern especially near urban centers and at the more accessible sites. Invasion by exotic plants, especially at the lower elevation temperate outcrops and all communities with major trail access, is increasingly a concern. Alpine meadows are affected by ski area development while both serpentine and temperate calcareous outcrops continue to be limited by mining operations. Climate change is especially a concern with the colder alpine meadows and boreal outcrops.

Desired Condition: Outcrops and upland meadows are very specialized natural communities in Vermont since they are relatively permanent openings within a forested landscape. As such they provide specific habitat requirements for a small number of SGCN, especially some species of snakes which utilize these openings as basking sites. Although they provide significant habitat for only nine SGCN and two suites of species (moths and tiger beetles), these openings are utilized by many additional wildlife species. The number of SGCN plants (95) that rely on this formation speaks to its importance in the state despite the small area that it covers. To protect these sites we would do the following:

SGCN in Outcrops & Upland Meadows

High Priority

North American Race (*Coluber constrictor*)
 Timber Rattlesnake (*Crotalus horridus*)
 Eastern Ratsnake (*Pantherophis alleghaniensis*)
 Moths group
 Beetles-Tiger Beetle Group
 Northern bog lemming (*Synaptomys borealis*)
 Southern Bog Lemming (*Synaptomys cooperi*)

Medium Priority

Chimney Swift (*Chaetura pelagica*)
 Cinereus or Masked Shrew (*Sorex cinereus*)
 Smoky Shrew (*Sorex fumeus*)

SGCN Notes: Plant SGCN not listed here: 95. The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
Habitat Conversion	Quarrying activity, development, and ski area development	Medium
Climate Change	Species generally have no higher elevations to move to	High
Incompatible Recreation	Rock climbing, hiking disturbs wildlife, tramples rare plants, and introduces exotic species.	High

Invasion by Exotic Species	Non-native species can spread and degrade the habitat for wildlife and eliminate some plant species	Medium
Habitat Fragmentation	Some species require large expanses of forestland surrounding their denning sites	High
Inventory	Distribution, location, and condition of this habitat are not known. A statewide inventory is needed to identify and locate the best examples of these habitats that support the most SGCN	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Conduct a statewide inventory of outcrop and meadow natural communities to identify the best sites and those with SGCN	The number of high quality examples identified containing SGCN	FPR	SWG
Provide technical & financial assistance to private, municipal and federal landowners to control invasive species and to minimize the impact of recreation on SGCN	Number landowners managing for SGCN. Number of acres conserved	NRCS, TNC, VFWD	SWG
Technical assistance to town and regional planning organizations to maintain and enhance outcrops and upland meadows for SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns including SGCN in their planning	VFWD	VFWD
Develop conservation easements on higher quality sites with greatest number of SGCN or T&E listed SGCN	Number of acres conserved for SGCN	ANR, VLT, TNC	VHCB, VLT, LIP
Work with hiking and rock/ice climbing groups to avoid sensitive sites. Limit hiker use and new trails on high quality state-owned sites		ANR,	VFWD,

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
New England Plant Conservation Program – various Conservation Plans	Recovery of various plant species in New England	ANR
State Outdoor Recreation Plan	A comprehensive recreation plan for state lands	FPR

References

Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. *Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#)*. Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.

Thompson, E. H., and E. R. Sorenson. 2005. *Wetland, Woodland, Wildland - A guide to the natural communities of Vermont*. University Press of New England, Hanover and London.

Cliff & Talus Summary

Characteristics and location

Cliffs are areas of exposed bedrock, with slopes greater than 60 degrees. Examples range from very small and shaded by surrounding forests to extensive sites greater than one hundred acres. Vermont's cliffs are divided on the basis of their climatic affinities and their bedrock. Climate is the factor separating boreal cliff types from temperate cliff types. The boreal types are found in the cooler regions of the state, the Northeast Highlands and the Green Mountains, though a few are found in generally warmer regions, in especially cool situations such as at high elevations or in cold valleys. The temperate types are found either at middle to low elevations or in the warmer regions of the state. Bedrock is the factor separating acidic cliff communities from calcareous cliff communities. Granites, some quartzites, and sandstones are typically acidic, whereas limestones, dolomites, calcareous schists, and some quartzites are calcareous. [Thompson and Sorenson 2000]

Talus slopes are areas of rockfall below cliffs and are characterized by an accumulation of many rocks broken off a cliff face through physical forces including freezing and thawing.

Types of Cliff and Talus Communities:

Boreal Acidic Cliff: These are high elevation cliffs, generally above 2,000 feet, found on acidic bedrock such as granite, gneiss, quartzite, or non-calcareous schist. Vegetation is usually red spruce, balsam fir, American mountain-ash, bush-honeysuckle, three-toothed cinquefoil, and hairgrass. Eastern Hemlock is absent from these cliffs. Found primarily in the cooler regions of the state, the Northeast Highlands and the Green Mountains.

Boreal Calcareous Cliff: These are high elevation cliffs, mostly above 2,000 feet, where calcareous bedrock (usually calcareous schist, but occasionally limestone or marble) combined with seepage creates conditions that favor certain calciphilic plants, some of which are quite rare statewide.

Temperate Acidic Cliff: These are lower elevation cliffs, generally below 2,000 feet, found on acidic bedrock. Characteristic vegetation includes eastern hemlock, white pine, red maple, paper birch, harebell, and heart-leaved aster. Found primarily either at middle to low elevations or in the warmer regions of the state.

Temperate Calcareous Cliff: These are low elevation cliffs in warmer areas on limestone, marble, dolomite, or calcareous quartzite. They may be moist or dry, depending on the situation, but usually do not have abundant seepage. Some characteristic species are northern white cedar, purple clematis, smooth cliff-brake, purple-stemmed cliff brake, harebell, and herb robert. Found primarily either at middle to low elevations or in the warmer regions of the state.

Open Talus: This broadly defined community type includes all areas of open rockfall. These rockfall areas usually occur below cliffs, and can be comprised of granite, quartzite, gneiss, shale, or less commonly limestone or marble.

Cliff & Talus Condition

Current Condition: Generally, cliffs and talus communities are not directly vulnerable to habitat degradation simply because they tend to be inaccessible and limited in timber or development potential. Recreational activities and intensive quarrying may be the greatest impacts to these communities where such activities occur.

Desired Condition (SGCN Needs): Cliffs and talus are often host to habitat specialists, many of which are plants directly linked to the natural community type. In general, the larger the site, the greater the likelihood that numerous SGCN plant species will exist and that they will persist. Many of the animal species associated with this community types; however, do require accessible, unfragmented habitat mosaics. Several of the animal species require the cliff and talus for nesting or hibernation, but range as far as 1000 ha from the site. Three of the species (North American Race, rock vole, and five-lined skink) specifically benefit from active management for early successional features or small openings around the sites in order to provide solar radiation. The North American Race is found on only one site in Vermont. The five cliff and talus community types provides the habitat for 100 SGCN.

Species of Greatest Conservation Need in Cliffs & Talus

High Priority

North American Race (*Coluber constrictor*)
Timber Rattlesnake (*Crotalus horridus*)
Eastern Ratsnake (*Pantherophis alleghaniensis*)
Five-lined Skink (*Plestiodon fasciatus*)
Rock Vole (*Microtus chrotorrhinus*)
Small-footed Bat (*Myotis leibii*)
Long-tailed or Rock Shrew (*Sorex dispar*)

Medium Priority

Peregrine Falcon (*Falco peregrinus*)
Eastern Ribbonsnake (*Thamnophis sauritus*)

SGCN Notes: Plant SGCN not listed here 88 species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
Habitat Conversion	Quarrying activity and poorly designed ski trails	High
Habitat Fragmentation	Wider ranging reptiles require unfragmented habitat mosaics of 1000 ha or more	High
Climate Change	Species generally have no higher elevations to move to	High
Incompatible Recreation	Rock climbing disturbs falcons and tramples rare plants	High
Distribution of successional stages	Active management for early successional openings (North American Race), young forest (rock vole), and forest openings for solar radiation (five-lined skink).	Medium
Pollution	Acid rain threatens higher elevation habitats	Medium
Research & Inventory needs	Distribution, location, and condition of this community type are not known.	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Conduct statewide inventory of cliff and talus and identify and locate the best examples of these community types that support the most SGCN	Number of sites inventoried	FPR	SWG
Provide technical assistance and/or financial assistance private landowners to maintain and enhance cliff and talus for SGCN.	Number landowners incorporating SGCN into their land management,	NRCS, TNC, FWD	SWG
Technical assistance to town and regional planning organizations for conservation practices that maintain and/or enhance habitat for SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns/RPCs considering SGCN in their planning	VFWD	VFWD
Conservation easements on higher quality sites with greatest number of SGCN or T&E listed SGCN	Number of acres conserved for SGCN	ANR, VLT, TNC	VHCB, VLT, LIP
Work with hiking and rock/ice climbing groups to avoid sensitive sites	Number of sensitive sites with programs implemented to limit encroachment	ANR, GMC, VOGA	VFWD, Access Fund

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
Peregrine falcon plan	Remove peregrine from ESA list	ANR
Draft VT Bat Conservation Plan		ANR
ANR Land Conservation Plan	ANR land acquisition	ANR
VT Recreation Plan (SCORP)	Recreation priorities throughout the state	ANR, GMC, VOGA

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. *Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#)*. Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Thompson, E. H., and E. R. Sorenson. 2005. *Wetland, woodland, wildland - A guide to the natural communities of Vermont*. University Press of New England, Hanover and London.

Grassland & Hedgerow Summary

Characteristics and location

Grasslands are landscapes dominated by grasses, sedges and forbs with little to no tree or shrub cover. Most of the larger examples of this community type are the result of current or past agricultural practices. Grassland habitats are also commonly maintained at airports, fairgrounds, landfills and industrial complexes. Smaller grasslands are found in fallow beaver flowages, seasonally flooded areas adjacent to rivers, and sandplain communities, and are covered under separate summaries.

Hedgerows are linear patches of trees or shrubs, often lining field borders or roadsides. Hedgerows enable some species to more fully utilize adjacent grassland communities (for perching, nesting, sheltering or escaping predators), while other species may occupy annual or seasonal home ranges solely within hedgerows. Hedgerows also often serve as travel or dispersal corridors connecting disjunct habitat patches.

Types of Grassland & Hedgerow Communities:

Hayfields, pastures, old fields, power line and RR right-of-ways, mowed interstate medians, airports, industrial complexes. Treed and/or brushy hedgerows lining field edges and roads.

Grassland & Hedgerow Condition

Historical Perspective: Grasslands in Vermont are primarily a result of land clearing for agriculture since European settlement of the area. It has been estimated that early successional forest (1-15 year age class) occupied from 1.1-3.0% of the regional presettlement landscape in areas of northern hardwood forest and 2.4-7.1% of the regional landscape in areas of spruce-northern hardwood forest (Lorimer and White 2003).

Current Condition: Most of Vermont's grasslands occur in the Champlain Valley and to a lesser extent the Connecticut River Valley and the area around Lake Memphremagog. There are also numerous grasslands of various types and sizes scattered across the rest of the state. Most grasslands are associated with current or past agricultural practices. There are, however, grasslands that are the result of other human activities and are maintained for specific purposes. These include grasslands associated with airports (commercial and private), landfills, fairgrounds, military reservations and industrial complexes (e.g., IBM, Husky, etc.). Most of Vermont's grasslands are in private ownership, although the state and federal governments own and manage some of these areas. The counties with the highest percentages of land in agriculture and open land are Addison (35.5%), Franklin (29.5%), Grand Isle (25%) and Orleans (22%, primarily in the area surrounding Lake Memphremagog) (U.S. Department of Agriculture. 1997).

Although agriculture practices create and maintain valuable grasslands, recent intensification of these practices has had negative impacts on their quality and availability. Small diversified farming which provided a range of suitable habitat types has given way to larger, more intensively managed farms as a result of improved agricultural techniques. Advances in equipment, fertilizers and extensive use of potent pesticides and herbicides have resulted in greater management of hayfields (early and frequent cutting which disrupts nesting activity), conversion of hayfields to row crops or legumes, and intensive grazing (LaBarr et al. 2004).

Urban and suburban development has also resulted in a loss of grasslands. This loss comes in two forms, the direct loss of grasslands as structures and lawns replace fields, and fragmentation of large grassland areas into smaller parcels rendering them insufficient for use by some breeding grassland bird (e.g., Upland Sandpiper). In Vermont, the urban and suburban growth of Chittenden county is

expanding into Franklin and Grand Isle counties to the north and Addison county to the south. As a result there is increasing pressure to develop agricultural lands important to grassland species (LaBarr et al. 2004).

Other factors contributing to loss of quality grasslands include incompatible management of grasslands in non-agricultural settings (i.e., airports). Although airport construction and management has provided suitable habitat for grassland species, mowing regimes, many of which are required by the Federal Aviation Administration (FAA) often disturb nesting activity. Also, a lack of airport expansion planning (new hangers, airplane parking, etc.) which takes into account grassland species has lead to the loss of important grassland habitat at these sites (LaBarr et al. 2004).

More is known about the effects of current conditions on grassland bird species than other SGCN taxa that use grasslands and/or hedgerows. Grassland bird species as a whole have declined steadily throughout their range. Reported results from the U.S. Fish and Wildlife Service Breeding Bird Survey show that declines of grassland birds have been consistently steeper and more widespread than any other assemblage of birds (Askins 1993, Sauer et al. 2001). In Vermont, Upland Sandpiper populations have declined precipitously (Peterson 1999) and Grasshopper Sparrows are considered rare and uncommon (Ellison 1985, Record of Vermont Birds). Both Sedge Wren and Henslow's Sparrow populations have declined to where they may no longer be breeding in the state. Other obligate grassland species, although relatively abundant (i.e., Bobolink and Eastern Meadowlark) have also show significant declines in recent years (LaBarr et al. 2004).

Desired Condition (SGCN Needs): A variety of grasslands and hedgerows are needed to conserve the suite of species dependant on these habitat types. For example, Bobolinks utilize large expanses of grassland or fallow hay fields with little or no alfalfa, high litter cover and scattered broad-leafed forbs for nest-site cover (Martin and Gavin 1995). Northern Harrier habitat includes marshy meadows, wet, lightly grazed pastures, old fields, mesic grasslands, and drained marshlands. Densest populations are typically associated with large tracts of undisturbed habitats dominated by thick vegetation (MacWhirter and Bildstein 1996). Upland Sandpipers prefer large grassland areas (20-40 ha) with a mosaic of grassland types as areas of short grass are used for feeding while areas of taller grass (10-30 cm) are used for nesting. All three of these aforementioned species benefit from grasslands that are not subjected to early (before July 15) mowing. American Kestrels nest in cavities or nest boxes in most open areas (< 30% canopy cover; Smallwood and Bird 2002). Gray Fox, New England Cottontail, Eastern Ratsnake, Smooth Greensnake and DeKay's Brownsnake all utilize hedgerows for foraging, denning or nesting, and/or as movement corridors.

Implementing the 2005 Wildlife Action Plan

Over the past decade, VFWD maintained an estimated 340 acres of permanent openings as old field shrub cover by brush mowing and burning an average of 105 acres annually to maintain this vegetation type on WMAs. Such permanent shrub openings have been shown to be extremely important to shrubland birds; Smetzer et al. (2014) estimated that "maintaining the current population size of shrubland birds under a management strategy based entirely on silviculture would require a 50–300% increase in the area of silvicultural openings, depending on the species."

Species of Greatest Conservation Need in Grasslands & Hedgerows

High Priority

Grasshopper Sparrow (*Ammodramus savannarum*)
Upland Sandpiper (*Bartamia longicauda*)
Northern Harrier (*Circus cyaneus*)
Sedge Wren (*Cistothorus platensis*)
Vesper Sparrow (*Pooecetes gramineus*)
Fowler's Toad (*Anaxyrus fowleri*)

Wood Turtle (*Glyptemys insculpta*)
North American Race (*Coluber constrictor*)
Timber Rattlesnake (*Crotalus horridus*)
Eastern Ratsnake (*Pantherophis alleghaniensis*)
Butterflies-Grassland Group
Moths Group
Eastern Red Bat (*Lasiurus borealis*)

Hoary Bat (*Lasiurus cinereus*)
 Woodland Vole (*Microtus pinetorum*)
 Pygmy Shrew (*Sorex hoyi*)
 New England Cottontail (*Sylvilagus transitionalis*)
 Southern Bog Lemming (*Synaptomys cooperi*)

Medium Priority

Peregrine Falcon (*Falco peregrinus*)
 Short-eared Owl (*Asio flammeus*)
 Chimney Swift (*Chaetura pelagica*)
 Black-billed Cuckoo (*Coccyzus erythrophthalmus*)
 Bobolink (*Dolichonyx oryzivorus*)
 American Kestrel (*Falco sparverius*)
 Purple Martin (*Progne subis*)
 Field Sparrow (*Spizella pusilla*)
 Eastern Meadowlark (*Sturnella magna*)
 Lesser Yellowlegs (*Tringa flavipes*)
 Blue-winged Warbler (*Vermivora pinus*)
 Smooth Greensnake (*Opheodrys vernalis*)
 DeKay's Brownsnake (*Storeria dekayi*)
 Eastern Ribbonsnake (*Thamnophis sauritus*)
 Long-tailed Weasel (*Mustela frenata*)
 Hairy-tailed Mole (*Parascalops breweri*)
 Common Gray Fox (*Urocyon cinereoargenteus*)

SGCN Notes: Plant SGCN not listed here: 23. The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need/Category	Problem/Info Need Detail	Rank
1. Habitat Degradation	Widespread early hay harvest (early June) and heavy grazing rotations in pastures.	High
2. Habitat Conversion	conversion of agricultural habitat to urban/suburban development	High
3. Distribution of successional stages	Abandonment and forest succession of former agricultural land.	High
4. Habitat Degradation	Removal of hedgerows to accommodate larger tractors and farm machinery.	High
5. Habitat Fragmentation	Fragmentation of habitat by roads and trails and increase use of roads and trails by motor vehicles, including ATV's, and mountain bicycles.	High
6. Inventory	Distribution and condition of this habitat are not well known. Better information is necessary regarding the timing of hay mowing in landscapes with various proportions of agriculture throughout VT.	Medium
7. Inventory	Better information is needed on the distribution of SGCN within grasslands habitats and the relative values of the various types and sizes of these habitats to the SGCN.	Medium

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Locate grassland and assess management practices on those grasslands.	Number of sites located and assessed	ANR, FSA, UVM, VA	SWG
Identify areas within the state with the largest matrix of grasslands for inclusion in grassland bird opportunity areas.	Number of opportunity areas identified	ANR, UVM	SWG
Ensure protection of opportunity areas via acquisition of conservation easements, management leases and fee title acquisition	Number of sites conserved	ANR, VHCB, TNC	VHCB, TNC
Develop education and outreach program to provide information about grassland/hedgerow dependant species and management options to enhance their populations in Vermont.	Number of maintained or enhanced sites on private land	ANR, FSA, VFB	SWG, WHIP, GRP, VDA
Promote conservation easements or incentives to landowners managing grasslands/hedgerows for SGCN.	Number of maintained or enhanced sites on private land	ANR, FSA, VFB	SWG, WHIP, GRP, VDA
Develop conservation plans at state airports where SGCN are regularly found.	Number of sites with conservation agreements	ANR, VTRANS, FAA	SWG, VTRANS
Continue to work with Vermont National Guard staff at Camp Johnson to manage grasslands to benefit grassland species.	Number of SGCN conserved at Camp Johnson	VNG, ANR	SWG
Maintain and manage grasslands and hedgerows on state and federal lands (wildlife management areas, state parks, National Wildlife Refuges, GMNF)	Number of sites reclaimed and/or managed	ANR, USFWS, USFS	SWG, PR
Manage power line ROW, road margins and related lands known or suspected to support SGCN that depend on grasslands and enhance surrounding habitat by creating and maintaining open habitat.	Number of sites reclaimed and/or managed	ANR, VETCO, GMP	SWG, VETCO, GMP
Support current efforts and develop new efforts to study distribution, productivity, and survivorship of grassland bird species in Vermont.	Number of hypothesis tested	ANR, UVM, VA, VINS	SWG, PR
Develop safe road crossings to limit road kill of snakes and turtles which use grassland habitats	Number of safe crossings developed	ANR, Towns, VTRANS,	SWG, PR, VTRANS

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
VT Grassland Bird Management Plan	Maintain and enhance grassland bird populations	VFW, WHIP
Partners in Flight	Regional Bird conservation	VFW, VA, USFWS, PIF, NABS CI
VTRANS Transportation Plans	Manage airports	VFW, VTRANS

References

- Askins, R.A. 1993. Population trends in grassland, shrubland, and forest birds of eastern North America. Pages 1-34 in D.M. Power, editor. *Current Ornithology*, Volume 11. Plenum, New York, New York, USA.
- Ellison, W.G. 1985. Grasshopper Sparrow. Pages 346-47 in S. B. Laughlin and D. P. Kibbe, eds. *The atlas of breeding birds of Vermont*. University Presses of New England, Hanover, New Hampshire, USA.
- LaBarr, M., S. Parren, J. Peterson, T. Allen, T. Donovan, A. Strong, E. Derleth, B. Crenshaw and R. Renfrew. 2004. Vermont grassland bird management and recovery plan. Vermont Fish and Wildlife Dept., Waterbury, Vermont, USA.
- Lorimer, Graig G., and Alan S. White. 2003. Scale and frequency of natural disturbances in the northeastern US: implications for early successional forest habitats and regional age distributions. *Forest Ecology and Management*, Volume 186, 41-64.
- MacWhirter, R. B., and K. L. Bildstein. 1996. Northern Harrier (*Circus cyaneus*). In *The birds of North America*, No. 210 (A. Poole and F. Gill, eds.). The birds of North America, Inc., Philadelphia, Pennsylvania, USA.
- Martin, S. G., and T. A. Gavin. 1995. Bobolink (*Dolichonyx oryzivorus*). In *The birds of North America*, No. 176 (A. Poole and F. Gill, eds.). The birds of North America, Inc., Philadelphia, Pennsylvania, USA.
- Peterson, J. 1999. The Upland Sandpiper in Vermont, 1998-1999 – a declining species. Report to the Vermont Endangered Scientific Advisory Group on Birds. Vermont Institute of Natural Science, Woodstock, Vermont, USA.
- Sauer, J. R. _____. 2001. The North American Breeding Bird Survey, Results and Analysis _____. Version _____. USGS Patuxent Wildlife Research Center, Laurel, Maryland, USA.
- Smallwood, J. A., and D. M. Bird. 2002. American Kestrel (*Falco sparverius*). In *The birds of North America*, No. 602 (A. Poole and F. Gill, eds.). The birds of North America, Inc., Philadelphia, Pennsylvania, USA.
- Smetzer, J. R., D.I. King, and S. Schlossberg. 2014. Management Regime Influences Shrubland Birds and Habitat Conditions in the Northern Appalachians, USA. *Journal of Wildlife Management* 78(2):314–324.
- U.S. Department of Agriculture. 1997 National Resources Inventory, Natural Resources Conservation Service, Washington, DC, and Statistical Laboratory, Iowa State University, Ames, Iowa.

Mines and Quarries Summary

Characteristics and location

Mines may provide many or all of the habitat qualities of natural caves and can even provide better habitat in some instances. Similarly, quarries may mimic natural cliffs, outcrops, and talus slopes. These human-created cultural habitats, due to the history of Vermont are found statewide and may augment the natural habitats available to wildlife.

Types of Mines and Quarries:

Mines in Vermont include gold, silver, iron asbestos, and talc mines.

Quarries in Vermont include marble, granite, and slate quarries

In some cases gravel pits and road cuts may provide habitat

Mines & Quarries Condition

Current Condition: Mines and quarries occur throughout the state. Some are long abandoned, some more recent, and others currently used to lesser or greater extents. The sites vary in their structural stability and some are very dangerous (large sections of the Elizabeth Mine have collapsed) Bats are known to use some mine sites as hibernacula. Peregrine falcons may nest or roost on the walls of some rock cuts. Mine vents and other vertical rock structure may provide nesting habitat for swifts. Small-footed bats might seek shelter in between and under large rock talus created by mining or quarrying operations. In some instances the sites are toxic due to leaching of mine tailings. Some sites have the entrances blocked, become dumping areas, or recreational vehicle parks.

Desired Condition (SGCN Needs): Some mines and quarry site provide conditions that certain species select. A mine that has appropriate temperatures and humidity may provide good wintering habitat for bats. Like caves, if the conditions change or if disturbances occur, the site may no longer be suitable habitat and can even cause the death of bats using the mine. Some rattlesnake reports historically have been from slate quarries in proximity to existing or historical den sites. Quarries could provide foraging and basking habitat as well as escape cover. Rock piles with abundant spaces that extend below the frost line could even provide denning sites. Sites providing necessary habitat for SGCN are important and should be conserved.

Species of Greatest Conservation Need using Mines and Quarries

High Priority

Chimney Swift (*Chaetura pelagica*)
Timber Rattlesnake (*Crotalus horridus*)
Little Brown Bat (*Myotis lucifugus*)
Northern Long-eared Bat (*Myotis septentrionalis*)
Indiana Bat (*Myotis sodalis*)
Small-footed Bat (*Myotis leibii*)
Tri-colored bat (*Perimyotis subflavus*)

Medium Priority

Peregrine Falcon (*Falco peregrinus*)
Big Brown Bat (*Eptesicus fuscus*)

SGCN Note: For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
Habitat Conversion	Closure of mine entrances and filling of quarries.	High
Habitat Alteration	Modification of mine entrances or interiors that either exclude wildlife or create unsuitable conditions	High
Habitat Conversion	External surface changes to drainage patterns or tree cover that render the mine or quarry unsuitable for wildlife use.	High
Pollution	Poisonous gasses that can infiltrate a mine or runoff that contaminate a site	High
Trampling or Direct Impacts	Direct persecution [of wildlife using mines or quarries	High
Habitat conversion	Reopening an abandoned mine or cave for extraction of mineral resources	High
Incompatible recreation	Recreational use of mines or caves used by wildlife.	High
Inventory	Distribution, location, and condition of this habitat are not fully known .A statewide inventory would add to our knowledge of sites that support the most SGCN	Med

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Conduct statewide inventory of mines and quarries important to SGCN.	Number of sites surveyed that have SGCN that are dependent on mines and quarries	VFWD, Town Conservation Commissions, AVCC	SWG, Section 6
Identify those mines or quarries important to SGCN and at risk of loss, then take actions to conserve them with priority given to structures with most vulnerable species, largest concentration of a SGCN, or the greatest number of SGCN present.	Number of protected occurrences of each SGCN using mines and quarries.	VFWD, Town Conservation Commissions, AVCC	SWG, Section 6
Raise awareness and acceptance of the need to provide cultural habitat for some SGCN that depend on mines and quarries and modify recreational and other activities.	Number of audiences reached.	Environmental Educators	
Promote conservation easements or agreements for important sites for SGCN	Number sites having conservation agreements	ANR, BCI	VHCB, VLT, LIP
Consider direct purchase of a mine or quarry if that is the most effective manner to manage for SGCN	Number of conserved SGCN that are dependent on mines and quarries	VFWD, Town Conservation Commissions, AVCC	SWG, Section 6
Provide technical assistance and economic incentives for property owners to manage mines and quarries for SGCN while protecting the health and safety of humans.	Maintained or enhanced condition of SGCN using a mine or quarry (numbers of individuals, reproductive success, survival rate)	VFWD	LIP

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Work with landowners to provide fencing and/or appropriately designed gates that exclude human intrusion and reduce liability to landowner, while maintaining SGCN using a mine or quarries	Maintained or enhanced condition of SGCN using a mine or quarry (numbers of individuals, reproductive success, survival rate)	VFWD	LIP
Educate users of mine and quarry sites and encourage avoidance of important sites when SGCN are vulnerable (e.g., bats fall through spring).	Increased understanding and acceptance of mine/quarry conservation by the public	VFWD, BCI, School programs, media	Marketing? Section 6
Encourage use of alternative sites that do not harbor SGCN	Increased understanding and acceptance of mine/quarry conservation by the public	VFWD, BCI, School programs, media	Marketing? Section 6

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
Peregrine falcon federal monitoring plan and state recovery plan	Remove peregrine from state list and prevent need to re-list	ANR/NWF
Draft VT Bat Conservation Plan	Conservation of all bats, especially those currently listed in Vermont	ANR/VFWD
Rattlesnake plan being drafted	Maintain and enhance rattlesnake populations in VT and move them toward recovery	ANR/VFWD, Jim Andrews

References

Tuttle, M.D. and D.A.R. Taylor. 1994. Bats & Mines. Bat Conservation International, Inc Resource Publ. No. 3

Subterranean Summary

Characteristics and Location

Subterranean areas are defined as below-surface natural features (mines are addressed under Cultural Habitats) that consist of both aquatic and terrestrial conditions. Because these areas are below ground, there is limited human access to locate and inventory these sites. Consequently, there is little information on their abundance, distribution, and condition.

Some of the best information on subterranean areas comes from the caving community. Members of the Vermont Cavers Association have particular interest in locating, exploring, and even surveying these areas. Some of the earlier documentation of Vermont caves is from John Scott (1959) and, more recently, Peter Quick (1994).

Most of Vermont's caves are relatively small, ranging from less than 100 feet underground to several hundred feet. Some caves contain passages that may continue far beyond what has been accessed. Most Vermont caves are solutional, meaning they have been formed through erosion from moving water.

While caves are found throughout Vermont, most of the known caves are located in southern Vermont, particularly the Taconics and Southern Green Mountains regions. These areas also are known to have the geologic features most associated with underwater springs and streams that would provide subterranean aquatic habitats.

Subterranean Condition

Current Condition: Due to the geologic nature of the habitat type, caves remain in much of their original structure. Many of the more accessible caves do exhibit signs of graffiti and evidence of the destruction or removal of cave formations such as stalagmites and stalactites. Historic accounts of some caves document the loss of beautiful formations by visitors. Currently, 3 caves are gated and locked to control human visitation.

Subterranean areas provide a very consistent environment of temperature, relative humidity, and air flow. While these variables are likely important to the overall condition, there is very limited information on these variables. Changes in structure and hydrology could greatly affect these habitat provided by subterranean areas.

There are 6 species of bats known to hibernate in Vermont caves. Bats are one of the better studied wildlife species associated with subterranean areas, and have been surveyed in caves going back into the 1930s (Trombulak et al. 2001). Trend data from hibernacula surveys does provide for some evaluation of the value of specific caves to particular bat species and populations. Recent surveys indicate that caves may hold as few as less than 10 bats to as many as over 23,000. Interest and understanding in the invertebrate community associated with caves is just beginning.

Little is known about the condition of the subterranean aquatic habitats.

The primary activities resulting in the loss or degradation of subterranean areas involve either human disturbance to either the cave structure (thereby affecting temperature, humidity, or air flow) or the species using the area and pollutants to the aquatic elements of the subterranean areas.

Desired Condition: Subterranean areas provide habitat for a small number of SGCN in the state. However, subterranean areas provide a critical habitat component for the survival of these species. Subterranean areas should remain intact, with limited human alteration or influence from above-ground pollutants. Many of the SGCN associated with subterranean areas use the sites for denning or hibernation, but also spend a disproportionate amount of the year in the surrounding area (e.g., fall swarming for bats or breeding and birthing for rattlesnakes).

A total of 8 SGCN are associated with subterranean area.

Species of Greatest Conservation Need in Subterranean Landscapes

High Priority

Little Brown Bat (*Myotis lucifugus*)
 Northern Long-eared Bat (*Myotis septentrionalis*)
 Indiana Bat (*Myotis sodalis*)
 Tri-colored bat (*Perimyotis subflavus*)
 Small-footed Bat (*Myotis leibii*)
 Timber Rattlesnake (*Crotalus horridus*)

Medium Priority

Chimney Swift (*Chaetura pelagica*)
 Big Brown Bat (*Eptesicus fuscus*)

SGCN Note: For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
Hydrologic alteration	Sedimentation, development in watershed, road building	Medium
Habitat Conversion	Roads, development, and agriculture remove SGCN habitat surrounding subterranean sites	High
Habitat Degradation	Alteration of cave structure, thereby influencing temperature, humidity, or air flow	High
Incompatible recreation	Disturbance to hibernating bats or denned rattlesnakes	Medium
Pollution	Aquatic pollutants	Medium
Inventory	Statewide inventory has been completed, but not all sites have been evaluated	Low

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Gate subterranean sites experiencing risk from unlimited human visitation	Number of sites gated	USFWS, TNC, VCA	SWG, USFWS
Conservation easements on higher quality sites with greatest number of SGCN or T&E listed SGCN	Number of acres conserved for SGCN	ANR, VLT, TNC, NCC	VHCB, VLT, LIP
Provide technical assistance and/or financial incentives to private	Number landowners incorporating SGCN into	NRCS, TNC, FWD, RPC,	NRCS programs, USFWS

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
landowners, towns and RPC's to maintain and enhance Subterranean habitat for SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	their land management, Number of towns including SGCN in their planning. Number of acres conserved	VLCT, USFWS	

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
Draft Bat Conservation and Recovery Plan	Conservation and recovery of Vermont bat species	ANR
Cave Management Plans	Management plans for specific caves in Vermont	ANR, VCA, NCC

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Scott, John. 1959. Caves in Vermont. Killbuck Independent Speleological Society. Hancock, VT.
- Quick, Peter. 1994. Vermont caves – A geologic and historical guide.
- Trombulak, Stephen, P. Higuera, and M. DesMeules. 2001. Population trends of wintering bats in Vermont. *Northeastern Naturalist*. 8(1):51-62.

Buildings & Other Structures Summary

Characteristics and location

Buildings and structures may provide habitat for wildlife, generally in the form of shelter, when they provide appropriate conditions (i.e., temperature and humidity) and are relatively secure from disturbance. Sometimes the structures provide habitat for prey species (mice) that attract the foraging SGCN (snakes). In other cases the structures may simply become an extension of the natural landscape, such basking and foraging sites for skinks. Structures used by wildlife are located throughout Vermont, but are not always known or appreciated as habitat for wildlife.

Types of Buildings and Other Structures Providing Habitat for SGCN

Barns and other outbuildings, Abandoned or little used buildings, House attics, Bridges, Dams, Power poles and other vertical structures (possibly) and Towers or tall buildings that mimic cliffs.

Condition of Buildings & Other Structures

Current Condition: Buildings and other structures may be used by wildlife under a variety of circumstances. Bats may roost in abandoned building attics, the attics of occupied dwellings, or in outbuildings or covered bridges. Peregrine falcons may nest on ledges of tall buildings, tower, or bridges although we don't have any currently nesting in such locations at present. Small-footed bats might seek shelter in between and under large rock talus used to armor dams. Osprey may nest on power poles near water and chimney swifts may build their nests inside chimneys.

Desired Condition (SGCN Needs): Some buildings and other structures provide conditions that certain species select. If the site is relatively undisturbed and secure over time, large number of some species may come to depend on the site (e.g., large bat maternity colony). Change the light regime or air circulation, and the conditions may no longer be as suitable. In some cases the surrounding area, or even the specific geographic location, may determine if a structure is used by a SGCN. Only barns located near existing skink populations will be used by that species and a power pole used by osprey for nesting has to be within flying distance of fishable waters.

SGCN Using Buildings & Other Structures

High Priority

Chimney Swift (*Chaetura pelagica*)
Eastern Ratsnake (*Pantherophis alleghaniensis*)
Five-lined Skink (*Plestiodon fasciatus*)
Little Brown Bat (*Myotis lucifugus*)
Northern Long-eared Bat (*Myotis septentrionalis*)
Indiana bat (*Myotis soldalis*)
Small-footed Bat (*Myotis leibii*)
Tri-colored bat (*Perimyotis subflavus*)

Medium Priority

Peregrine Falcon (*Falco peregrinus*)
Purple Martin (*Progne subis*)
Big Brown Bat (*Eptesicus fuscus*)

SGCN Note: For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
Inventory	Distribution, location, and condition of this habitat are not known. A statewide inventory is needed to identify and locate the best examples of these habitats that support the most SGCN	Medium
Habitat Conversion	Loss of old buildings that provide shelter for wildlife	High
Habitat Conversion	Modification of structures that exclude wildlife or create unsuitable conditions	High
Habitat Conversion	Changes to structures that may trap or kill animals (including deliberate exclusions)	High
Pollution	Use of chemicals that may poison or kill wildlife	High
Trampling or Direct Impacts	Direct persecution of wildlife using structures	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Conduct statewide inventory of buildings and structures important to SGCN.	Number of conserved sites with SGCN that are dependent on buildings and other structures	VFWD, Town Conservation Commissions, AVCC	SWG, Section 6
Identify those buildings or other structures important to SGCN and at risk of loss, then take actions to conserve or replace.	Number of protected occurrences of each SGCN using buildings and other structures.	VFWD, Town Conservation Commissions, AVCC	SWG, Section 6
Promote conservation easements or agreements for important sites for SGCN	Number sites having conservation agreements	ANR, BCI	VHCB, VLT, LIP
Consider direct purchase of a structure if that is the most effective manner to manage for SGCN (e.g., PA bat maternity colony in old church).	Number of conserved SGCN that are dependent on buildings and other structures	VFWD, Town Conservation Commissions, AVCC	SWG, Section 6
Provide appropriately designed structures in suitable locations to replace buildings and structures no longer available to SGCN. In some cases these need to be provided in conjunction with an exclusion	Number of protected occurrences of each SGCN using buildings and other structures.	VFWD, Town Conservation Commissions	SWG, Section 6
Provide technical assistance and economic incentives for property owners to manage their structures for SGCN while protecting the health and safety of humans.	Maintained or enhanced condition of SGCN using a building or structure (numbers of individuals, reproductive success, survival rate)	VFWD Wildlife Services	LIP
Provide education programs and materials that improve the public's understanding of SGCN needs and perceptions of wildlife that utilize buildings and structures.	# of audiences reached, # of people attending living with wildlife programs.	NWF	

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Encourage coexistence with SGCN using buildings and structures	Increased understanding and acceptance of building/structure conservation by the public	VFWD, BCI, School programs, media	Marketing? Section 6

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
Peregrine falcon federal monitoring plan and state recovery plan	Remove peregrine from state list and prevent need to relist	ANR/NWF
Draft VT Bat Conservation Plan	Conservation and restoration of bat population	ANR
Osprey Recovery Plan	Delist osprey and prevent need to relist	ANR

References

Tuttle, M.D. 1988/ 1994. America's neighborhood bats: understanding and learning to live in harmony with them. Univ. Texas Press.

Riparian Summary

Vermont's aquatic and shoreline landscape includes all surface waters and their adjacent streambanks, floodplains and/or lakeshores. This landscape includes lacustrine (lake) formations, fluvial (stream and river) formations, floodplain forests, and shores and marshes. This landscape also includes thousands of miles of streambank areas that are comprised of upland communities adjacent to surface waters. The aquatic and shoreline landscape is described as an interconnected system of the lacustrine, fluvial, floodplain, marsh, shore, and upland communities that comprise it for the purpose of identifying and conserving the common habitat functions these communities provide at the landscape level.

Riparian (riverbank) areas, if maintained in continuous, sufficiently wide, interconnected corridors throughout a watershed, serve as movement corridors for many of Vermont's wildlife species. Maintaining intact terrestrial communities adjacent to surface waters also serves to protect aquatic habitats. Riparian areas help protect water quality, provide organic inputs, regulate water chemistry and physical properties (such as temperature), and provide physical aquatic habitat structure (e.g., undercut banks, large woody debris). Again, because aquatic communities are often inter-connected throughout the landscape, maintaining intact riparian areas is essential to protecting aquatic communities from the headwaters to downstream receiving waters.

Habitat requirements, problems, and conservation strategies have been assessed and developed for both the landscape level, and the individual aquatic and terrestrial species' habitats that are associated with it. Many SGCN meet most of their habitat needs within the aquatic-terrestrial interface that the aquatic and shoreline landscape provides. These species, in particular, are discussed in this section.

"It is a well known fact that the best fishing is where a forest is near the shore, and best of all where the limbs overhang the water. Not only do the trees afford shelter, furnish food and prevent evaporation, but at the same time they keep the water clear and cool in the summer. In the winter the forests afford protection by lessening the severity of the winter frosts, and in all forest regions the changes of temperature are not so severe as in treeless countries and on the open plain: and the effect upon the water is even greater....But the forests not only regulate the flow of water, as above stated, but they purify the water."

- Frank H. Carleton, from the Fifteenth Biennial Report of the Commissioners of Fish and Game of the State of Vermont, 1899-1900.

Characteristics and location

Aquatic and shoreline landscapes are comprised of streams, rivers, lakes, wetlands, shorelines and floodplains that form a complex and interrelated hydrological system. This hydrological system extends up and down streams and along lakeshores from the bottom of the water table to the top of the vegetation canopy, and includes land that is directly affected by surface water (Verry 2000). Riparian areas are known for their high biological diversity. They are "characterized by frequent disturbances related to inundation, transport of sediments, and the abrasive and erosive forces of water and ice movement that, in turn, create habitat complexity and variability...resulting in ecologically diverse communities" (Verry 2000).

The landscape level includes both the terrestrial-aquatic interface and the aquatic areas found throughout Vermont, from the mountain streams to the large valley rivers and the lakes and ponds scattered throughout the landscape. The following aquatic and terrestrial areas are

associated with the aquatic and shoreline landscape (for details see the following summaries in Appendix B):

Lacustrine (“Inland” Lakes)

Lake Champlain

Lake Champlain Tributaries

Connecticut River Tributaries

Floodplain Forests

Upland Shores

Wet Shores

Swamps and Marshes

Landscape Condition

Current Condition: Nationwide an estimated 70% to 90% of natural riparian vegetation, vital to maintaining the integrity of riparian and aquatic habitats, has already been lost or is degraded due to human activities (Doppelt 1993). In Vermont, some of our rivers, streams, lakes, and wetlands still have intact riparian areas, while many others no longer have functioning riparian areas due to more than 200 years of intensive human use of the land.

In general, riparian areas in Vermont are most affected by habitat conversion, alteration, and fragmentation. Typically, steeper mountainous streams and high elevation lakes and ponds, less suited for human development, have well forested riparian areas with cold, clean water and stable stream channels and shorelines. Recreational activities and their associated development and forestry are the land uses most common in these areas that may affect riparian and aquatic species. Mid and low elevation waterbodies and their adjacent riparian areas are more likely to be impacted by human land uses, including clearing of riparian vegetation, alteration of stream channels and lakeshores, and direct inputs of toxins, excess nutrients, and sediments. These impacts are related primarily to roads, residences, commercial development, and agriculture, with agriculture being especially extensive in the lower valleys of the Champlain and Connecticut tributaries. Lacustrine areas and their associated shorelines are particularly impacted by lakeshore development, such as seasonal and permanent residences, marinas and docks, and public and private beaches. In many instances these developments have altered natural lakeshore and littoral zones resulting in the direct loss of habitats for SGCN through the addition of fill materials (sand, bottom barriers) and the removal of native aquatic vegetation.

The fragmentation of riparian habitat is extensive in Vermont, due primarily to Vermont’s roadways paralleling the stream, rivers, and lakeshores, and use of rich floodplain areas for agriculture. Historic settlement and transportation patterns and ease of construction have resulted in roads paralleling the majority of Vermont’s major waterbodies and thousands of associated bridges and culverts. This results in removal of riparian vegetation and fragmentation, both longitudinally and laterally between the waterbody and adjacent upland communities.

Desired Condition (SGCN Needs): Aquatic and shoreline areas provide several habitat functions for the species that inhabit them. Some species rely directly on both the aquatic and terrestrial components of the riparian-aquatic interface. For example otter use aquatic areas within 100 meters of water’s edge for feeding and riparian areas for denning and as travel corridors. These species move daily between terrestrial and aquatic areas to fulfill their life needs. Other species move seasonally between the aquatic and terrestrial components of the aquatic and shoreline landscape. For example, the wood turtle uses streams and rivers for overwintering, and uses adjacent riparian areas up to 300 meters from the water’s edge for foraging, breeding, nesting, and dispersal. For those species that are strictly aquatic, the

adjacent terrestrial riparian areas function to protect the aquatic areas, providing shade, organic inputs, filtering and storage of overland runoff, and bank stability.

Implementing the 2005 Wildlife Action Plan

In 2013, VFWD completed the Vermont [BioFinder](#) project, a map and database identifying Vermont's lands and waters supporting high priority ecosystems, natural communities, habitats, and species. A notable outcome of the project was a map of all aquatic features and the riparian areas/valley bottoms in which rivers and streams occur and the identification of these areas as critical conservation components for wildlife habitat, rare species, aquatic system health, and wildlife/landscape connectivity.

VFWD provided technical assistance to every Vermont Regional Planning Commission and nearly every town on a variety of wildlife and land planning related issues, including SGCN conservation, habitat blocks, and wildlife corridors. [Conserving Vermont's Natural Heritage](#) (Austin et.al. 2004) was reprinted and distribution of this planning document continues.

On a landscape level, aquatic and shoreline areas provide habitat for 41 SGCN.

Species of Greatest Conservation Need in Aquatic and Shoreline

High Priority

Bald Eagle (*Haliaeetus leucocephalus*)
Wood turtle (*Glyptemys insculpta*)
Common Mudpuppy (*Necturus maculosus*)
Silver-haired Bat (*Lasionycteris noctivagans*)
Eastern Red Bat (*Lasiurus borealis*)
Hoary Bat (*Lasiurus cinereus*)
Tri-colored bat (*Perimyotis subflavus*)
Water Shrew (*Sorex palustris*)
Odonates-Bog/Fen/Swamp/Marshy Pond Group
Freshwater Mussels Group
Freshwater Snails Group
Lakes/ponds Odonata group
Mayflies/Stoneflies/Caddisflies Group
River/stream Odonata group
Elktoe (*Alasmodonta marginata*)
American Eel (*Anguilla rostrata*)
American Brook Lamprey (*Lethenteron appendix*)
Bridle Shiner (*Notropis bifrenatus*)
Blackchin Shiner (*Notropis heterodon*)
Blacknose Shiner (*Notropis heterolepis*)
Northern Brook Lamprey (*Ichthyomyzon fossor*)
Stonecat (*Noturus flavus*)

Medium Priority

Peregrine Falcon (*Falco peregrinus*)
Pied-billed Grebe (*Podilymbus podiceps*)
Lesser Yellowlegs (*Tringa flavipes*)
Northern River Otter (*Lontra canadensis*)
Muskrat (*Ondatra zibethicus*)
Cinereus or Masked Shrew (*Sorex cinereus*)
Mottled Sculpin (*Cottus bairdi*)
Redfin Pickerel (*Esox americanus*)
Brook Trout (naturally reproducing populations)
(*Salvelinus fontinalis*)
Silver Lamprey (*Ichthyomyzon unicuspis*)
Redbreast Sunfish (*Lepomis auritus*)
Atlantic Salmon (landlocked) (*Salmo salar*)

SGCN Notes: Plant SGCN not listed here include 16 species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
Habitat Conversion	Floodplain forests, lakeshores and other riparian communities converted to agriculture, roadways, and residential/commercial development. Habitat conversion is most prevalent in low and mid elevation areas.	High
Habitat Degradation	Removal or alteration of vegetative community, ground disturbance, and manipulation of shorelines and streambanks; can lead to degradation of water quality, and loss of physical habitat structure. Habitat degradation occurs primarily in upper elevation areas, in contrast to complete habitat conversion, which is more common in mid and low elevation areas.	High
Habitat Fragmentation	Interruption of movement corridors to and from breeding, feeding, and seasonal habitats via conversion, degradation, and road mortality (herps).	High
Inadequate Disturbance Regime	Dams, drainage ditching, floodplain filling, and channel incision (floodplain abandonment) that affect flooding, erosion, and deposition processes	High
Invasion by Exotic Species	Habitat alteration from invasive plant species (e.g., Japanese knotweed, Purple loosestrife); plant inter-species competition for habitat.	High
6. Harvest or Collection, Trampling or Direct Impacts	Collection and harvest pressures; increased human activity disturbing breeding, nesting and movement.	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Develop a plan to identify and prioritize existing contiguous riparian corridors and associated wildlife habitat linkages	Increase in number of riparian habitat linkages identified and conserved	ANR, TNC, NWF, NRCS, FSA	EQIP, CRP, CREP
Technical assistance to private landowners to maintain and enhance SGCN habitat in riparian areas.	Increase in number of acres of riparian habitat restored and/or conserved by private landowners	NRCS, ANR, USFWS, FSA	WHIP, EQIP, , CREP
Financial incentives for private landowners to maintain and enhance SGCN habitat in riparian areas.	Increase in number of acres of riparian habitat restored and/or conserved by private landowners	NRCS, ANR, USFWS, FSA	WHIP, EQIP, CRP, REP, LIP
Technical assistance to town and regional planning organizations to maintain and enhance SGCN habitat in riparian areas. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Increase in number of towns incorporating riparian conservation into planning and zoning	ANR, ACCD, VLCT, AVCC, NRCS, FSA	ANR, NRCS
Technical assistance to state and federal land management agencies on riparian habitat management goals/strategies	Increase in % or number of state and federal land management plans providing for riparian conservation	ANR, VTrans, USFWS, USFS	
Work with VTrans, towns, and private landowners to identify and maintain (or restore) riparian habitat connectivity and improve aquatic organism passage	Increase in % or number of road crossings that do not impede riparian corridor movement – longitudinally and laterally	VTrans, ANR, NRCS	WHIP, VTrans, SWG
Technical assistance to landowners and conservation groups on invasive exotic management and eradication		TNC, ANR, NRCS, FSA	CRP, CREP,

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
ANR State Lands Management Plans	Management practices for ANR-owned lands	FPR, VFWD
Floodplain Forests of Vermont	Natural Community Inventory	ANR
Riparian Buffer Guidance, and Riparian Buffers and Corridors Technical Papers 1/20/2005 http://www.anr.state.vt.us/site/html/buff/anrbuffer2005.htm	Helps in the development of recommendations and designs for Act 250-regulated projects that incorporate appropriate buffer zone widths for protecting riparian functions	ANR
ANR Stream Geomorphic Assessments	Stream and riparian condition inventories	ANR

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Carleton, Frank H. 1899-1900. Why forest preservation should interest fisherman. 15th Biennial report of the Commissioners of Fisheries and Game of the State of Vermont. p. 98-110
- Doppelt, B., M. Scurlock, C. Frissell, J. Karr. 1993. Entering the Watershed: A New Approach to Save American River Ecosystems. Island Press. Washington, D.C. As farms are abandoned management strategies to maintain grassland habitat are critical.
- Verry, E. S., J. W. Hornbeck, and C. A. Dolloff (eds). 2000. Riparian management in forests of the continental Eastern United States. Lewis Publishers, Boca Raton, FL.

Streams & Rivers Summary

Characteristics and location

There are more than 7,000 miles of rivers and streams in Vermont draining 4 major watersheds: Connecticut, Lake Champlain, Hudson, and Memphremagog. The headwater streams of the western Green Mountains drain to the large rivers of the lower Champlain Valley and eventually into Lake Champlain. The eastern slopes of the Green Mountains drain primarily to the Connecticut River. Portions of the Northeastern Highlands and Northern Piedmont drain north into Lake Memphremagog. The Taconic Mountains and southern Green Mountains drain into the Batten Kill, Deerfield, Walloomsac, and Hoosic rivers. These rivers, with the exception of the Deerfield, eventually drain into the Hudson River in New York. The Deerfield drains to the Connecticut River. Despite this diversity of landscape over which Vermont's streams and rivers flow, fluvial ecosystems can be described by three general categories based on physical stream characteristics. There are various biotic communities associated with each of these physical stream types, depending on both the physical stream characteristics and the geographic location of the waterbody. For example, the large rivers of the lower Lake Champlain watershed are similar in physical characteristics to the large tributaries feeding Lake Memphremagog, but some of the species found in these two settings differ due to the repopulation patterns of aquatic species into freshwater ecosystems post-glaciation. This summary does not include discussion of the lower Connecticut River tributaries and the lower Lake Champlain tributaries below the fall-line and/or below 150 feet elevation, as these areas are covered under separate summaries.

General types of Streams & Rivers communities:

High-elevation Headwater Streams: These streams are typically located in high elevation mountainous areas. They are small in size, having small drainage areas, and are located in steep valleys (typically > 4% slope). Valleys are confined, meaning the stream channel has little or no floodplain, and upland forest communities are adjacent to the channel, typically with no distinct riparian vegetative community present. Channel bed form is usually cascade over bedrock and boulders or step-pools over boulders and cobbles. Stream flow is fast and turbulent with white water common. Stream temperatures are typically very cold. Forest canopy completely shades the stream, and the food web of the system is based on inputs of organic material from the adjacent vegetation (e.g., leaves, twigs, branches). Large trees falling into the stream channel also provide important habitat features and channel bed stability, acting as cover and causing localized scour and deposition of stream sediments. Species that typically inhabit these streams include brook trout, slimy sculpin, northern spring salamander, northern dusky salamander, two-lined salamander, and numerous aquatic insects, including stoneflies and mayflies. SGCN species uniquely associated with these ecosystems include the water shrew, some specific mayfly and Odonata species and naturally reproducing populations of brook trout.

There are some headwater streams in high elevation areas that do not meet the above description. Small, low gradient streams are often found in ridgeline saddles and bowls. These streams are typically meandering, with alternating riffles and pools and gravel and sand substrates. Adjacent wetlands are often associated with these streams. These are typically still cold water systems, due to abundant groundwater feed and cooler climatic conditions influenced by high elevation, and therefore often host many of the same species as the high gradient

headwater streams. Invertebrate communities, however, are likely to be distinct from the higher gradient systems (Burnham 2005).

Mid-elevation Streams and Rivers: These streams are typically located in mid elevation areas where the steep mountains transition to the low gradient valleys. Stream channels are small to moderate in size, and are located in moderately steep valleys (typically 2-4% slope). Valleys are semi-confined, resulting in narrow floodplains. These floodplains may have narrow bands of distinct riparian vegetation, but quickly transition into upland forest communities. Channel bed form is typically step-pool or plane bed. Step-pool channels have short vertical drops over boulders and cobbles with channel spanning pools in between, which are typically dominated by cobbles and gravels. Plane bed systems lack distinct pools, and are primarily riffles, runs, and rapids over a mix of boulders, cobbles, and gravels. Stream flow is fast and somewhat turbulent with whitewater common. Stream temperatures are typically cold to cool. Forest canopy usually shades the stream but may not form a complete canopy over the channel. The aquatic food web in these channels is based largely on inputs of organic material from the adjacent vegetation (e.g., leaves, twigs, branches), though some mosses and algae are also present, providing primary production in the waterbody. Large trees falling into the stream channel and transported from upstream provide important habitat features and channel bed stability, acting as cover and causing localized scour and aggradation of the channel bed. Species that typically inhabit these streams include brook trout, slimy sculpin, blacknose dace, white sucker, longnose dace, northern dusky salamander, two-lined salamander, and numerous aquatic insects. SGCN species uniquely associated with this habitat potentially include naturally reproducing populations of brook trout, as well as American eel, Atlantic salmon, wood turtle, river otter, water shrew, muskrat and some specific mayfly and Odonata species.

Low-elevation Large Valley Rivers: These rivers are located at low elevations in Vermont's large river valleys, such as the Winooski, Lamoille, Mississquoi, Barton, Otter, and Batten Kill. This description does not include those portions of the large Lake Champlain tributaries located below the fall-line. These river channels are moderate to large in size, and are located in low gradient valleys (typically <2% slope). Valleys are unconfined, and floodplains are broad and flat. Adjacent wetlands are common in the floodplains. These floodplains have extensive distinct riparian vegetation and often include unique natural communities, such as floodplain forest, marsh, and shoreline communities. The channel bed undulates vertically, being composed of alternating riffles and pools or dune-ripple formations. Riffle-pool systems are dominated by gravels and sands, where dune-ripple systems are usually dominated by sands and silts. Stream flow is slow and flat with whitewater rarely present. Stream temperatures are typically cool to warm. Forest canopy shades the nearbank area of the channel but does not form a complete canopy over the channel. The aquatic food web in these channels is based on inputs of organic material from the adjacent vegetation (e.g., leaves, twigs, and branches) and transported from upstream, as well as instream aquatic vegetation. Large trees falling into the stream channel and transported from upstream provide important habitat features, especially since coarser streambed substrates are typically lacking in these systems. Woody debris provides cover and substrate for aquatic biota, as well as helping to maintain channel bed stability and enhancing habitat complexity with localized scour and aggradation of the channel bed. Numerous cool and warmwater fish species inhabit these streams, including bluntnose minnow, fallfish, blacknose dace, creek chub, tessellated darter, and white sucker, as well as several mussel species. SGCN species uniquely associated with this habitat include American eel, Atlantic salmon, blackchin shiner, bridled shiner, blacknose shiner, redbfin pickerel, stonecat, giant floater, cylindrical floater,

elktoe, brook floater, wood turtle, river otter, muskrat, bald eagle, and some specific species of freshwater snails and Odonata.

Low Elevation Small Streams: These streams are small in size, but located in low gradient valleys (<2% slope) at low elevations (but above the Lake Champlain fall-line and 150 feet in elevation), and typically drain directly into a large waterbody (e.g., Lake Memphremagog, large tributaries of Lake Champlain). Valleys are unconfined, and floodplains are broad, relative to stream size, and flat. These floodplains have distinct riparian vegetation on the valley floor, and transition into upland forest communities on the valley side slopes. Adjacent wetlands are common in the floodplain. The channel bed undulates vertically, being composed of alternating riffles and pools or dune-ripple formations. Riffle-pool systems are dominated by gravels and sands, where dune-ripple systems are dominated by sands and silts. Stream flow is slow and flat. Stream temperatures are typically cool to warm. Streamside vegetation shades the channel, usually forming a closed canopy over the channel. The aquatic food web in these channels is based primarily on inputs of organic material from the adjacent vegetation (e.g., leaves, twigs, branches). Large trees falling into the stream channel provide important habitat features, especially since coarser streambed substrates are typically lacking in these systems. Woody debris provides cover and substrate for aquatic biota, as well as helping to maintain channel bed stability and enhancing habitat complexity with localized scour and aggradation of the channel bed. Typically cool and warmwater fish species inhabit these streams, such as blacknose dace and creek chub. SGCN species uniquely associated with this habitat include American eel, Atlantic salmon (landlocked), blackchin shiner, bridle shiner, redbfin pickerel, stonecat and some specific species of Odonata.

Landscape Streams & Rivers Condition

Current Condition: In general, fluvial ecosystems in Vermont are most affected by conversion, alteration, and fragmentation. Typically steeper mountainous streams at high elevations, less suited for human development, have well forested riparian areas with cold, clean water and stable stream channels. Recreational activities and their associated development, such as ski resorts, and forestry are the land uses most common in these areas that may affect stream habitats. Mid and low elevation streams and rivers are more likely to be impacted by human land uses, including clearing of riparian vegetation, alteration of stream channels, and direct inputs of toxins, excess nutrients, and sediments. These impacts are related primarily to roads, residences, commercial development, and agriculture, the latter being especially extensive in the lower valleys of the Lake Champlain and Connecticut River tributaries.

The fragmentation of fluvial ecosystems is extensive in Vermont. A recent inventory of more than 200 culverts in the White River watershed showed more than half of the culverts inventoried were barriers to the upstream movement of all fish species present in the waterbody all of the time, and the other half of the culverts inventoried were barriers to some species and/or barriers some of the time (i.e. under certain stream flows when species movement is likely to occur) (Vermont Fish and Wildlife 2004). In addition, most of Vermont's major rivers have large flood control and/or hydroelectric dams on them, with numerous smaller dams found throughout Vermont's smaller streams. Such structures influence local habitat conditions, restrict movement of aquatic species, and alter downstream flood and sediment transport processes. The Vermont Agency of Transportation is currently funding research regarding the extent of stream impediments and how to address issues such as culvert sizing and retrofits.

Some aquatic habitat degradation is due to lasting effects of historic land uses. During the last two centuries land use in Vermont has been dominated by extensive land clearing for forestry and agriculture, aggressive stream clearing of boulders and coarse woody debris for stream log driving and flood control, and by dam construction and railroad and road building. Such activities have resulted in the relocation and straightening of stream and river channels throughout Vermont, resulting in an overall decrease in available fluvial habitat. For example, a recent assessment of the upper White River watershed between Granville and Stockbridge shows that 93% (17.8 of 19.1 miles) of the length of the mainstem White River has been channelized in the past, 13 miles of which are still in channelized form (Vermont Department of Environmental Conservation 2004). In addition, the extensive removal of natural substrates, such as boulders and coarse woody debris, has reduced overall stream habitat complexity throughout the Northeast (Verry 2000). The hard armoring of channels combined with the construction of flood control dams means that many of Vermont's river channels have not regained their historic sinuosity. Furthermore, the slow regrowth of the Northeast's forests means that large woody debris contribution to stream and river channels has yet to reach historic levels (Verry 2000). Zadock Thompson, who served as Vermont's Assistant State Geologist and State Naturalist in the mid 1800's, offers first-hand insight on the impacts Vermont's intensive land use history has had on the streams and rivers of the state.

"Before the country was cleared, the whole surface of the ground was deeply covered with leaves, limbs, and logs, and the channels of all the smaller streams were much obstructed by the same. The consequence was that, when the snows dissolved in the spring, or the rains fell in the summer, the waters were retained among the leaves, or retarded by the other obstructions, so as to pass off slowly, and the streams were kept up, nearly uniform as to the size during the whole year. But since the country has become settled, and the obstructions, which retarded the water, removed by freshets, when the snow melts or the rains fall, the waters run off from the surface of the ground quickly, the streams are raised suddenly, run rapidly, and soon subside. In consequence of the water being thus carried off more rapidly, the streams would be smaller than formerly during a considerable part of the year, even though the quantity of water be the same. It is a well known fact that the freshets in Vermont are more sudden and violent than when the country was new."

Zadock Thompson, Natural History of Vermont, 1853

Desired Condition (SGCN Needs): Most of Vermont's aquatic species rely on streams and rivers that provide clean water, a diversity of in-channel habitat, and unobstructed movement upstream and downstream between habitats.

Characteristics of water quality vary in streams from clear and cold with little buffering capacity in most mountain streams to somewhat turbid and cool or warm with greater buffering capacity in the large valley rivers. Species found in the mountain headwater and mid-elevation streams are typically dependent on cold well-oxygenated waters. Some species found in the headwater streams, such as brook trout, are fairly acid tolerant. Low-elevation rivers and streams typically support species with warmer water temperature requirements and tolerance to some turbidity and nutrient enrichment.

Whether in the mountain streams or large valley rivers, most aquatic SGCN require instream cover and/or substrates for protection and colonization. Most fish species seek cover for predator avoidance and to reduce metabolic (energy) demands. Mussels need firm substrates for colonization, as do most aquatic insect species. Substrates utilized may vary from rock to sand to instream aquatic vegetation, depending on the species, but all species can suffer from excessive fine sediments in the

channel that can bury instream substrates. Loss of complexity and solid substrates for cover and colonization reduces overall habitat availability and quality. In addition, many species use instream substrates for reproduction. For example, brook trout deposit eggs in gravels on the channel bottom, whereas many shiner species utilize aquatic vegetation to spawn. Embedding of substrates, destabilization of substrates due to chronic channel instability, and direct removal of substrates all impact aquatic habitats and species. The mammal and bird species associated with streams and rivers, such as bald eagle, river otter, muskrat, and water shrew, are also impacted when aquatic species are affected, as these species rely on aquatic species as prey. In addition, muskrat, otter, and particularly water shrew, utilize undercut streambanks and other stable bank areas for denning. Chronic channel instability that results in substantial streambank erosion may reduce potential denning areas for these species.

Some of the SGCN uniquely associated with streams and rivers have extensive movement requirements, such as the Atlantic salmon and American eel, migrating from freshwater streams and rivers to the Atlantic Ocean and back again. Other species move shorter distances, but still require habitat connectivity to be able to access spawning, rearing, and seasonal habitats. There are also species, such as wood turtle and river otter, that move back and forth between the aquatic and nearby terrestrial habitats both daily and seasonally. Thus, it is important to maintain habitat connectivity both longitudinally along the river channel and adjacent riparian lands, as well as laterally between the aquatic habitat and the riparian habitat.

Ideally, Vermont's rivers and streams would provide an interconnected network of habitats in which species can move upstream and downstream as needed to fulfill seasonal and diurnal habitat needs. Instream structure would provide an abundance and diversity of habitat niches and be naturally maintained by physical stream processes over time (e.g., flooding, balanced sediment transport). Streams and rivers would be connected to the adjacent riparian habitats, which in turn function to protect and provide for fluvial habitat components, such as instream coarse woody debris and pollutant removal from surface runoff.

It is difficult to quantify the number of miles of intact fluvial and riparian habitat needed to conserve SGCN as the exact distribution of all SGCN associated with fluvial habitats is not known at this time.

Implementing the 2005 Wildlife Action Plan

In 2013, VFWD and partners completed the Vermont [BioFinder](#) project, a map and database identifying Vermont's lands and waters supporting high priority ecosystems, natural communities, habitats, and species. A notable outcome of the project was a map of all aquatic features and the riparian areas/valley bottoms in which rivers and streams occur and the identification of these areas as critical conservation components for wildlife habitat, rare species, aquatic system health, and wildlife/landscape connectivity.

The Vermont Department of Environmental Conservation's Rivers Program has completed Phase 2 mapping for most of Vermont rivers, has acquired river corridor easements, and has supported the passage of new legislation aimed at protecting river geomorphic processes.

Streams and Rivers provides habitat for 25 species and invertebrate groups of greatest conservation need.

Species of Greatest Conservation Need in Streams & Rivers Habitat

High Priority

Bald eagle (*Haliaeetus leucocephalus*)
 Fowlers toad (*Anaxyrus fowleri*)
 Wood turtle (*Glyptemys insculcata*)
 River/stream Odonata Group [dragonflies]
 Elktoe (*Alasmodonta marginata*)
 Giant floater
 Cylindrical floater
 Brook floater
 Dwarf wedge mussels Group
 Freshwater Snails Group
 Mayflies/Stoneflies/Caddisflies Group
 American eel (*Anguilla rostrata*)
 Bridle shiner (*Notropis bifrenatus*)
 Blackchin shiner (*Notropis heterodon*)
 Blacknose shiner (*Notropis heterolepis*)
 Stonecat (*Noturus flavus*)

Medium Priority

Northern river otter (*Lontra canadensis*)
 Muskrat (*Ondatra zibethicus*)
 Water shrew
 Redfin pickerel (*Esox americanus*)
 Brook trout (*Salvelinus fontinalis*)
 Atlantic salmon (landlocked) (*Salmo salar*)

SGCN Notes: Lake sturgeon is addressed in the Lake Champlain tributaries summary. Plant SGCN not listed here: 16 species. The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
Habitat Conversion	Channel straightening and maintenance of such that reduces overall stream/river miles, loss of floodplain connectivity, impoundment of river channels	High
Habitat Alteration	Floodplain and stream channel manipulation (e.g., riprap); degradation of water quality, loss of physical habitat structure, temperature alteration	High
Habitat Fragmentation	Interruption of movement to and from breeding, feeding, and seasonal habitats via alteration and conversion; roadways, and impassable dams and culverts	High
Sedimentation	Alteration of habitat (e.g., spawning areas); smothering of organisms	High
Pollution	Acid rain threatens higher elevation habitats, nutrient overloading is common in lower elevation areas, other toxins are suspected but data is unavailable to assess impacts	High
Pollution	Catastrophic spills: toxic chemicals (e.g., chlorine) and contaminants limit mid and lower elevation habitats, especially where roadways and development are in close proximity to stream channels	High
Invasion by Exotic Species	inter-species competition for habitat and food; predation on native species, loss of native riparian vegetation community from invasive competition.	High
Hydrologic Alteration	Stream flow regulation at dams, watershed development, and withdrawals alter hydrographs and instream flows	High
Inventory need	Minimal data is available on the distribution in Vermont of many fluvial-associated SGCN	Med

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Conduct inventories of known and potential SGCN sites		ANR, USFS, USFWS, TU	SWG, TU, EPA, NRCS
Provide technical assistance to anglers and other conservation groups on invasive exotic management and eradication	No new introductions of invasives exotic species that impact fluvial habitats	TNC (plants), angler groups, baitfish dealers	NRCS, LCBP
Provide technical assistance to private landowners and watershed organizations on riparian and fluvial habitat conservation	Increase in number of stream/river miles in "reference" condition, as per VTANR Stream Geomorphic Assessments	ANR, NRCS, FSA,	Clean Water Fund, LCBP, CRP, WRP, EQIP
Provide financial incentives to private landowners for conservation and protection of SGCN and their riparian and fluvial habitat	Increase in number of stream/river miles and associated riparian areas that are conserved and/or restored	ANR, NRCS, USFWS, FSA	WHIP, USFWS, CRP, CREP, WRP
Provide technical assistance to town and regional planning organizations. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Increase in number of towns incorporating riparian and aquatic habitat conservation into planning and zoning; and increase in number of stream/river miles under regulated development that are in "reference" condition, as per VTANR Stream Geomorphic Assessments	ANR, ACCD, VLCT, AVCC, TNC, watershed organizations	ACCD planning grants, LCBP, SWG
Monitor, protect and restore water quality from excessive nutrient sediment loading, other pollutants.	Miles of SGCN habitat meeting water quality standards.	ANR, USFWS, NRCS, USFS, Lake & Watershed Associations	ANR, Clean Water Fund
Support efforts to reduce the long range transport of acid rain pollutants to Vermont.	Reduction in acidity levels in monitored high elevation waterbodies	ANR, USFS, AG office, Legislature, Congress.	
Identify pollutant sources posing risks of catastrophic spills to SGCN populations and implement programs to minimize those risks		ANR, Agency of Agric., VTrans, wastewater facilities, town road managers	
Technical assistance to state and federal land management agencies to ensure consistency in program implementation and sensitivity to SGCN requirements	Increase in % or number of state and federal land management plans that provide for fluvial and riparian habitat conservation	ANR, USFS, USFWS, ACOE, VTrans	
Support efforts to manage flow regulation projects to minimize impacts on SGCN	Decrease in number of river miles with altered flow regimes	ANR, ACOE, VT Dam Task Force, USFWS, watershed orgs	LCBP, USFWS, ACOE, SWG
Provide technical assistance to VTrans, towns, and private landowners to identify and maintain (or restore) aquatic habitat connectivity	Increase in % or number of road crossings that do not impede aquatic organism movement	ANR, VTrans, Better Back Roads, USFWS, USFS, AVCC	SWG, USFWS, LCBP, VTrans

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
ANR State Lands Management Plans	Management practices for ANR-owned lands	FPR, VFWD
ANR Stream Geomorphic Assessments	Stream and riparian condition inventories	ANR
Opportunities for Action – LCBP	Aquatic resource conservation for the Lake Champlain Basin	LCBP
Strategic Plan for the Restoration of Atlantic Salmon to the Connecticut River.	“Protect, conserve, restore and enhance the Atlantic salmon population in the Connecticut River for the public benefit, including recreational fishing.”	CRASC

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Burnham, D. 2005. Vermont Department of Environmental Conservation. Personal communication.
- Thompson, Zadock. 1853. Natural History of Vermont with Numerous Engravings and an Appendix. Stacy and Jameson.
- Vermont Department of Environmental Conservation. 2004. Stream geomorphic assessment data available from the DEC River Management Program, Water Quality Division, Waterbury, VT.
- Vermont Department of Fish and Wildlife. 2004. Evaluation of Bridge and Culverts for Aquatic Organism Passage and Fluvial Function in the White River Watershed. Waterbury, VT.
- Verry, E.S., J.W. Hornbeck, and C.A. Dolloff (eds). 2000. Riparian management in forests of the continental Eastern United States. Lewis Publishers, Boca Raton, FL.

Lake Champlain Tributaries Summary

These waters include the lower-most portions of tributaries that empty into Lake Champlain. On many rivers and streams this is defined at its upstream end by the first major waterfall or cascade, called the principal fall line. On streams which do not have this abrupt elevation change, the upstream limit for is roughly 150 feet above sea level. The SGCN supported in these waters are numerous, with many found nowhere else in the state but in here and Lake Champlain. There are several factors accounting for the unique aquatic assemblages found here, including: glacial history and ancient routes of colonization from the west and south; the barrier to upstream migration presented by the principal fall line; and the generally warmer temperatures and finer substrates found here compared to those in higher elevation areas of Vermont. Unique species include many fishes and freshwater mussels, the common mudpuppy, and the spiny softshell. Key features include riffles, runs, and long pools with a variety of dominant substrate types. Small gravel, sand, and finer substrates are more dominant in the lowest reaches of these streams and rivers. Woody debris is prevalent, especially in deep holes in pool sections.

Lake Champlain Tributaries Condition

Current Condition: Some of the most heavily human-populated areas of the state occur adjacent to river sections included in this community type. While the larger volumes of water carried by the large rivers in Lake Champlain tributaries do afford a greater diluting potential than found in smaller rivers and streams, these Champlain tributaries are located in an area of the state where the intensity and frequency of insult to the aquatic habitat from human use is expected to be greater. For example, stormwater runoff reaching the lower Winooski River from developed lands is much greater than in most other Vermont fluvial communities. Stormwater runoff from developed lands increases the amount of sediments, nutrients, and contaminants that reach rivers instead of being trapped by the soil and vegetation. Floodplains function, in part, to absorb runoff and deliver it slowly to rivers through the soil. Paving of land sends water more directly to streams and rivers, in essence bypassing the floodplain. This creates a scouring effect on riverine habitat, due to the more extreme fluctuations in velocity of stormwater runoff.

Pollutants enter these rivers from various non-point sources as well. Agricultural lands located adjacent to rivers within these watersheds can contribute excessive amounts of silt, nutrients, and pesticides to the systems when adequate riparian buffers are not maintained. In such instances, excessive sediments can cover coarser river-bottom substrates needed by many SGCN, as well as covering some of these species themselves. Also, salt from roadways makes its way into rivers, degrading the water quality. These and other sources of non-point pollution are likely the greatest contributors of contaminants to these systems. These pollutants comes not only from adjacent lands, but from the entire watershed.

Accidental contaminant spills are rare, but can have immediate and devastating effects on the aquatic environment and the SGCN that live there. Chemicals, manure, industrial waste, and other potential contaminants stored in areas where they could reach these rivers or their tributaries if released are significant problems. Bridges and riverside roads and railways also present long stretches where accidental spills into rivers and streams can occur. A catastrophic contaminant spill could (and has) easily wipe out entire SGCN populations. As

with other sources of pollution, this problem comes from the watershed upstream as well as adjacent lands.

Direct loss of habitat occurs when fill material is placed on the river bottom. Examples of this include riprapping to stop toe erosion along streambanks, placement of piers or causeways to accommodate bridges, and construction of boating access facilities. Direct mortality of freshwater mussels, which live on the river bottom, is sometimes the result of these activities within lower Champlain rivers. The replacement of natural substrates with large stone provides reduced or unsuitable habitat for recolonization by bottom-dwelling animals. The construction of buildings and roads adjacent to rivers creates a hazard for the structures, increasing the potential that bank stabilization will be pursued.

Two dams on major rivers within the Lake Champlain tributaries have cut off migration for fishes and mussels, and have resulted in the loss of spawning habitat for some species. Impoundments created by these structures have altered the natural habitat from riverine to more lake-like water bodies. “De-watering” of the aquatic habitat that sometimes occurs due to atypical “hydro-peaking” dam operations leaves many benthic SGCN, particularly mussels, out of the water and exposed to the elements and predators. This can occur upstream or downstream of these structures. Existing dams located on fall lines may significantly alter the natural physicochemical regime of waters flowing downstream. The altered hydrologic regimes found below dams degrades the quality of habitat here for SGCN.

Zebra mussels that have devastated the Lake Champlain freshwater mussel community are a problem for rivers in this Lake Champlain tributaries. Adult zebra mussels have been found in the lower reaches of Otter Creek, Little Otter Creek, Lewis Creek, LaPlatte River and the Winooski River in past years. They are also present in Lake Bomoseen, whose outlet stream feeds into the Poultney River.

Desired Condition (SGCN Needs): These waters, along with Lake Champlain, support the greatest diversity of aquatic species found in the state. The larger rivers support the highest number of SGCN. Allowing these rivers to meander freely within their natural floodplains and maintaining and/or restoring natural vegetation to all or a portion of the rivers’ floodplains would significantly improve the ecological integrity of these systems, improve water quality, and significantly improve the habitat provided for many aquatic SGCN, as well as the diversity of wildlife species that rely on riparian cover movement or other habitat functions. Species include both year-round residents and those that use the rivers and streams primarily for spawning, development of young, or feeding. Minnows, freshwater mussels and snails, benthic fishes, and mammals are among those that utilize the Lake Champlain tributaries year-round, and often require a variety of habitats. Lake sturgeon, mooneye, greater redhorse, and possibly common mudpuppy are among those that depend on these rivers seasonally for reproduction. Others, such as map turtle, spiny softshell, northern watersnake, wood turtle, and bats use these waters for foraging, winter shelter, or other seasonal purposes.

Gravel/cobble substrates that are free of loose silt are required by many of the riverine species that spawn here. Eggs in contact with excessive silt are not able to adequately absorb oxygen for development. The eastern sand darter requires silt-free sand for this purpose. Substrates also need to be stable in order to support many SGCN, particular benthic organisms like freshwater mussels, darters, hibernating spiny softshells, and nesting

mudpuppies. This is often affected by stream hydrodynamics; that is, streams that are hydrodynamically imbalanced can have substrates that shift frequently and do not provide a firm footing or shelter for aquatic organisms that occur there. Small invertebrates are less abundant in silted-in or unstable stream bottoms, thus providing a reduced food source for their predators. Woody debris is an important habitat component in lower Lake Champlain tributaries, especially for aquatic insects. Historically, people removed trees and branches that fell into streams. Unfortunately, this removed the structure and habitat needed for many invertebrates and their predators, as well as basking habitat for turtles.

Two aquatic SGCN, the American eel and the Atlantic salmon, were historically able to ascend the fall line from downstream. The American eel did so to reach smaller waters upstream where the young eels would grow for several years before migrating back out to sea to spawn. Landlocked Atlantic salmon jumped the falls to reach the clean, coarse gravel substrates located upstream where they would spawn their eggs. With the construction of dams at or below the falls on all the major Lake Champlain rivers, much of the habitat needed for these two species was made unavailable to them. Reconnection of these fishes with this habitat would likely be beneficial to their long-term survival.

River otter and mink are susceptible to heavy metals and PCB's.

Implementing the 2005 Wildlife Action Plan

In 2013, the Department completed the Vermont [BioFinder](#) project, a map and database identifying Vermont's lands and waters supporting high priority ecosystems, natural communities, habitats, and species. A notable outcome of the project was a map of all aquatic features and the riparian areas/valley bottoms in which rivers and streams occur and the identification of these areas as critical conservation components for wildlife habitat, rare species, aquatic system health, and wildlife/landscape connectivity. The project mapping results for aquatic features, valley bottoms, and riparian connectivity together provide a tool for prioritizing restoration of riparian areas, including floodplain forests.

Species of Greatest Conservation Need in Lake Champlain Tributaries

High Priority

Bald Eagle (*Haliaeetus leucocephalus*)
 Lake Sturgeon (*Acipenser fulvescens*)
 Eastern Sand Darter (*Ammocrypta pellucida*)
 American Eel (*Anguilla rostrata*)
 American Brook Lamprey (*Lethenteron appendix*)
 Northern Brook Lamprey (*Ichthyomyzon fossor*)
 Silver Redhorse (*Moxostoma anisurum*)
 Greater Redhorse (*Moxostoma valenciennesi*)
 Bridle Shiner (*Notropis bifrenatus*)
 Blackchin Shiner (*Notropis heterodon*)
 Blacknose Shiner (*Notropis heterolepis*)
 Stonecat (*Noturus flavus*)
 Channel Darter (*Percina copelandi*)
 Sauger (*Sander canadense*)
 Spiny Softshell (Turtle) (*Apalone spinifera*)
 Common Mudpuppy (*Necturus maculosus*)
 River/stream Odonata group
 Wood Turtle (*Glyptemys insculpta*)
 Silver-haired Bat (*Lasionycteris noctivagans*)
 Eastern Red Bat (*Lasiurus borealis*)
 Hoary Bat (*Lasiurus cinereus*)
 Tri-colored bat (*Perimyotis subflavus*)
 Freshwater Mussels Group
 Freshwater Snails Group

Medium Priority

Peregrine Falcon (*Falco peregrinus*)
 Pied-billed Grebe (*Podilymbus podiceps*)
 Lesser Yellowlegs (*Tringa flavipes*)
 Mottled Sculpin (*Cottus bairdi*)
 Mooneye (*Hiodon tergisus*)
 Silver Lamprey (*Ichthyomyzon unicuspis*)
 Shorthead Redhorse (*Moxostoma macrolepidotum*)
 Atlantic Salmon (landlocked) (*Salmo salar*)
 Northern Water Snake (*Nerodia sipedon*)
 Eastern Musk Turtle (*Sternotherus odoratus*)
 Northern River Otter (*Lontra canadensis*)
 Muskrat (*Ondatra zibethicus*)
 Cinereus or Masked Shrew (*Sorex cinereus*)

SGCN Note: The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
Habitat Alteration	Input of sediments and nutrients from surface and stormwater runoff, and from small tributaries; caused by human land use nearby	High
Habitat Conversion	Loss of benthic habitat due to riprapping, bridge construction, boat access construction, etc. Loss of riverine environment due to impoundment.	High
Hydrologic Alteration	Changes in hydrologic and physicochemical regime due to dams and stormwater runoff. Direct loss of SGCN due to dewatering.	High
Habitat Fragmentation	Migration barriers created by dams	High
Pollution	Vulnerability to Catastrophic Spills: Bordering roadways, bridge crossings, adjacent industry, and manure pits are examples of high risk points of entry for large-scale contaminant spills	High
Invasion by exotic species	Zebra mussels are currently high risk threat to SGCN; other exotics may also be displacing native SGCN	High
Sedimentation	Alteration of habitat (e.g., spawning areas); fine sediments can embed of substrate and smother invertebrates, incubating eggs and the young of many fish species.	High
Pollution	Water quality degradation due to contaminants from agricultural fields, stormwater runoff, other point and non-point sources	High

Inventory	Inventory needed for many SGCN, particularly those for which distributional and abundance information is greatly lacking	High
Monitor	Detect SGCN population trends to help guide conservation actions and to track the effectiveness of current management	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Monitor known SGCN populations	Number of known SGCN sites monitored	USFWS, ANR, TNC, Universities, EPA	SWG, VFWD, VT Watershed Grants, EPA
Conduct inventories of rivers to detect and gather information on new SGCN populations	Number of sites/rivers with completed inventories	USFWS, ANR, TNC, Universities, EPA	SWG, VFWD, VT Watershed Grants, EPA
Protect and restore habitats on which SGCN are dependent through pollution abatement, riparian buffers, flow regulation.	Number of acres of riparian habitat protected and/or restored	LCLT, VLT, Watershed groups, USFWS, ANR, Army Corps, EPA	EPA, SWG, LCLT, VLT, NRCS, EPA, Clean Water Fund
Restore migration corridors for SGCN by removal of artificial barriers or construction of effective fish passage facilities at dams	Number of artificial SGCN migration barriers removed or provided with passageways Number of adult fish passed migrating to upstream spawning habitat (e.g., lake sturgeon, greater redhorse)	Hydro operators, FERC, ANR, Municipalities, VNRC	USFWS, NRCS
Provide for the safe and expeditious out-migration of SGCN from upstream of dams	Number of artificial SGCN migration barriers removed or provided with out-migration passageways	Hydro operators, FERC, ANR, Municipalities, VNRC	ANR, Army Corps
Prevent the introduction and spread of invasive exotic species, particularly zebra mussels	Number of sites with control activities and/or invasive monitoring. Number sites where invasive species are eliminated or controlled	LCBP, ANR Municipalities, USFWS, EPA	VT Watershed Grants, LCBP, Clean Water Fund
Provide technical outreach and financial assistance to private landowners, watershed groups and other partners to maintain or enhance habitat and tributary functions for SGCN.	Number of actions implemented to maintain or enhance tributary function for SGCN.	USDA, USFWS, EPA, NRCS, VFWD, TNC, LCBP, LCI, RPC's, Municipalities, Watershed groups	EPA, USFWS, WHIP, CRP, CREP, VT Watershed Grants, LCBP, SWG, Clean Water Fund
Provide technical outreach towns and regional planning commissions to maintain or enhance Lake Champlain tributary habitat and tributary functions for SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of actions implemented to maintain or enhance tributary function for SGCN.	USDA, USFWS, EPA, NRCS, VFWD, TNC, LCBP, RPC's, Municipalities, Watershed groups	EPA, USFWS, WHIP, VT Watershed Grants, LCBP, SWG, Clean Water Fund
Acquire conservation easements for the protection of critical SGCN habitats and maintenance or restoration of ecological functions	Number of riparian habitat acres acquired/enrolled	LCLT, VLT, ANR, TNC, NRCS	LCLT, VLT, EPA, TNC, SWG, NRCS

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Enhance coordination between government agencies and partners to ensure consistency in respective program implementation and increased sensitivity to SGCN requirements and problems to SGCN		ANR, USFWS, COE, FEMA, FHWA, NRCS, LCI, Wildlife Services, VTrans	WHIP, EQUIP, USFWS, EPA, Clean Water Fund

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
Lake Sturgeon Recovery Plan	Lake Sturgeon restoration	VFWD
Vermont's Clean Water Initiative	Water quality improvement	VDEC
Lake Champlain Basin Aquatic Nuisance Species Management Plan (2000).	Management and prevention of invasive exotic species in the basin	VTDEC, NYDEC
DEC Water Quality Division	Water quality and stream protection and restoration programs	DEC
Quebec Ministère de l'Environnement	Shared watershed for Missisquoi River	Quebec Ministère de l'Environnement
Conserving Lake Champlain's Biological Diversity 6/102005	Strategic plan focused on conserving Lake Champlain's biological diversity	TNC
Various watershed planning efforts	Watershed protection and restoration; river and lake restoration and protection	VTDEC; local/regional watershed groups

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Carle, F. C. 1994. Dragonflies and damselflies (Odonata) known to or likely to occur in Vermont. Report prepared for the Nongame & Natural Heritage Program, Vermont Fish & Wildlife Department. Waterbury, VT.
- Fichtel, C., and D. G. Smith. 1995. The freshwater mussels of Vermont. Nongame & Natural Heritage Program, Vermont Fish & Wildlife Department. Technical report 18.
- Hegman, W. et al. 1999. Estimation of Lake Champlain Basinwide Nonpoint Source Phosphorus Export. LCBP Technical Report No. 31, Grand Isle, VT: LCBP.
- Meals, D. 1990. LaPlatte River Watershed Water Quality Monitoring and Analysis Program, Comprehensive Final Report. Burlington, VT: University of Vermont, School of Natural Resources, Vermont Water Resources Research Center.
- Meals, D. 2001. Lake Champlain Basin Agricultural Watersheds Section 319 National Monitoring Program Project: Final Project Report May 1994 - November 2000. Waterbury, VT: VTDEC.
- Neddeau, E. J., M. A. McCollough, and B. I. Swartz. 2000. The freshwater mussels of Maine. Maine Department of Inland Fisheries and Wildlife.
- Stangel, P. 2004. Lake Champlain 2003 zebra mussel monitoring program. Final report. Vermont Department of Environmental Conservation. Waterbury, VT.

Lower Connecticut River Summary

Characteristics and Location

The Lower Connecticut River encompasses approximately 130 miles of the main stem from the Massachusetts state line upstream to its confluence with the Wells River and occurs almost exclusively within the Southern Vermont Piedmont biophysical region. Additionally, this summary includes the lower sections of its Vermont tributaries that are directly influenced by or have physical and/or biological similarity to the Connecticut River. The presence of a readily identifiable geological feature, such as a fall line, is not evident on all tributaries. Where the fall line is apparent, typically within a short distance from the tributary mouth (e.g., as on the Williams, Black, Ottauquechee, Waits and Wells rivers), this feature delineates the upstream extent of the Lower Connecticut River. On other tributaries (e.g., the West and White rivers), artificial structures (e.g., the lowermost dam) are used to define the upstream limit. Rivers and streams located within the Connecticut River basin but upstream of the habitat boundary are covered under the Fluvial (Stream) Summary. To a limited degree the historic distribution of several anadromous fish species native to the Connecticut River basin, namely sea-run Atlantic salmon, American shad and sea lamprey, as well as current management goals for the restoration of these fishes to the basin also define the bounds of the Lower Connecticut River.

Lower Connecticut River Condition

Current Condition: Prior to European settlement and subsequent industrial development of the Connecticut River basin, rivers and streams were free-flowing systems subject to natural flow regimes and processes. Waters ran free of pollutants, and the landscape, including riparian lands, was predominantly forested. These conditions provided habitat for both aquatic and terrestrial plant and wildlife assemblages native to the Lower Connecticut River. However, over the past 200 plus years, the river and its tributaries have been altered extensively fragmenting historic migration routes, changing natural habitats and ecological functions, as well as the current composition of the plant and wildlife communities.

Dams constructed for waterpower and flood control have greatly altered river and streams throughout the Connecticut River basin. Historic migration corridors used by Atlantic salmon, American shad, blueback herring, and American eel to gain access to critical spawning and nursery habitats have been obstructed. Long sections of the main stem and tributaries have been transformed from free-flowing waters to impoundments; and natural flow regimes are now regulated in ways that are not compatible with the habitat requirements of many aquatic species, including SGCN. Impoundments and artificial flow regimes have significantly influenced sediment transport and deposition, which in turn have altered the character, quantity and quality of various habitat types found throughout the Lower Connecticut River. Waters above and below dams are managed in ways, which result in fluctuating impoundment levels and tail water discharges. Frequently, flows released from dams are not adequate in volume or fluctuate in magnitude and duration so as to create habitat conditions unsuitable for SGCN. While water management within impoundments and free-flowing river segments may benefit habitat for a few SGCN (e.g., expose mudflats and shorelines used by feeding lesser yellowlegs during migration), fluctuating water levels can be detrimental to strictly aquatic SGCN (e.g., redbreast sunfish, common mudpuppy).

The extensive conversion of the Connecticut River from a free-flowing system to one dominated by impoundments has created habitats suitable to a variety of aquatic exotic plants and animals. Shorelines and wetlands associated with these impoundments have been invaded by phragmites, Eurasian milfoil and purple loosestrife, which have established dominant stands degrading nesting habitats needed by waterfowl, songbirds and muskrats. In recent years water chestnut was discovered in North Springfield Reservoir, which is on the Black River, a tributary of the Connecticut River. It has also been found in sections of the river in Massachusetts and Connecticut. This invasive exotic aquatic plant has been a significant environmental problem on Lake Champlain demanding large expenditure of funds and labor to keep it under control. Several fish species not indigenous to the Connecticut River, including predatory largemouth bass, northern pike, bluegill, crappie and rock bass, were introduced during the 1800s and early 1900s and have benefited from habitat formed within the impoundments. These species have altered the composition of the natural fish community of the river and have influenced ecological relationships at all trophic levels. At the present time, zebra mussels have not been found in the Connecticut River.

Prior to the federal Clean Water Act (amended in 1977) and subsequent implementation of water pollution abatement programs, a 1951 government report described the Connecticut River as the “best landscaped sewer in New England” (CRJC 2009). Over the past three decades water quality in the river and its tributaries has vastly improved habitats for aquatic SGCN. Nonetheless these waters continue to receive point and non-point source pollution (sediments, nutrients, toxic chemicals), which remain problems to aquatic habitats and the ability of the environment to support healthy, sustainable populations of SGCN, such as the eagle, fishes, freshwater mussels, and other aquatic invertebrates. Healthy aquatic systems are important to maintaining food webs not only for aquatic SGCN but also terrestrial species (eagle, bats, otter).

Development and logging along the river and tributaries has had a significant impact on riparian areas functions and benefits to SGCN. The loss of naturally vegetated (forested) riparian areas have lead to increased inputs of sediment and other pollutants to streams, increased water temperatures, channel instability, and loss of in-stream habitat structure created by the recruitment of large wood. Removal of living and dead trees (snags) from riparian lands has reduced sites for eagle nesting, roosting and perching.

Unique to the Lower Connecticut River is the existence of the Vermont Yankee Nuclear Power Station, which uses the river as a source of reactor cooling water and by state permit is allowed to discharges waste heat within specified thermal limits to the river. Excessive heat discharged to the river can potentially limit the temperature regime of the river within vicinity of the power plant to the detriment of aquatic SGCN intolerant of warm water. On a larger scale is the effect of climate change on aquatic habitats critical to certain SGCN (e.g., Atlantic salmon, American eel).

Desired Conditions (SGCN Needs): Eventual restoration and maintenance of sustainable populations of migratory native fishes to the Connecticut River basin is dependent on eliminating or mitigating artificial barriers which currently do not allow fish access to critical habitats, whether freshwater spawning and nursery areas or seawater (e.g. Atlantic salmon, American shad, sea lamprey, American eel). Dam removal would open river migration corridors, as well as restore natural flow regimes, sediment transport and other fluvial processes essential to creating and maintaining instream aquatic habitat. Where dam removal is not feasible, fish passage should be restored by retrofitting structures with fish ladders, lifts or similar devices. Existing fishways demand continued operation and maintenance to assure their effectiveness. New dam construction should be

avoided. Establishing flow regimes below dams and water level management within impoundments that mimic natural systems would benefit many of the aquatic SGCN.

The reduction of sediment inputs to the Lower Connecticut River from land development and chronic streambank erosion is important to maintaining SGCN populations, many of which depend on habitats consisting of coarse river bottom substrates (i.e., gravels and cobbles) that are not embedded by finer substrates. Riparian vegetation contributes to the reduction of these fine sediment inputs to surface waters by obstructing and slowing down overland runoff, while also reinforcing streambanks against the erosional forces of running water. Riparian areas also provide several habitat functions for species that inhabit them. Mature trees in the riparian zone provide necessary nesting sites for eagle. These trees eventually may be recruited to the river channel, creating instream habitat such as refuge cover required by common mudpuppy and redbreast sunfish.

Allowing these rivers to meander freely within their natural floodplains and maintaining and/or restoring natural vegetation to all or a portion of the rivers' floodplains would significantly improve the ecological integrity of these systems, improve water quality, and significantly improve the habitat provided for many aquatic SGCN, as well as the diversity of wildlife species that rely on riparian cover movement or other habitat functions.

The potential for new non-indigenous invasive organisms (e.g., zebra mussel, Asiatic clam, hydrilla) becoming established in the Connecticut River is a persistent problem for the native biota and habitats.

Implementing the 2005 Wildlife Action Plan

In 2013, the Department completed the Vermont [BioFinder](#) project, a map and database identifying Vermont's lands and waters supporting high priority ecosystems, natural communities, habitats, and species. A notable outcome of the project was a map of all aquatic features and the riparian areas/valley bottoms in which rivers and streams occur and the identification of these areas as critical conservation components for wildlife habitat, rare species, aquatic system health, and wildlife/landscape connectivity. The project mapping results for aquatic features, valley bottoms, and riparian connectivity together provide a tool for prioritizing restoration of riparian areas, including floodplain forests.

Species of Greatest Conservation Need in Lower Connecticut River

High Priority

Bald Eagle (*Haliaeetus leucocephalus*)
American Eel (*Anguilla rostrata*)
Anadromous Atlantic Salmon (*Salmo salar*)
Common Mudpuppy (*Necturus maculosus*)
Freshwater Mussels Group
River/stream Odonata group
Silver-haired Bat (*Lasionycteris noctivagans*)
Eastern Red Bat (*Lasiurus borealis*)
Hoary Bat (*Lasiurus cinereus*)
Tri-colored bat (*Perimyotis subflavus*)

Medium Priority

Peregrine Falcon (*Falco peregrinus*)
Pied-billed Grebe (*Podilymbus podiceps*)
Lesser Yellowlegs (*Tringa flavipes*)
Blueback Herring (*Alosa aestivalis*)
American Shad (*Alosa sapidissima*)
Redbreast Sunfish (*Lepomis auritus*)
Northern River Otter (*Lontra canadensis*)
Muskrat (*Ondatra zibethicus*)
Cinereus or Masked Shrew (*Sorex cinereus*)

SGCN Note: The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
1. Habitat conversion	Loss of riparian and in-stream habitats from land and water development projects and activities, including impoundments.	High
2. Habitat alteration	Alteration and degradation of riparian and in-stream habitats from land and water development projects, including streambank rip rapping.	High
3. Hydrologic alteration	Replacement of natural flow cycles and processes with regulated flow regimes (e.g., inadequate minimum flows, fluctuating flows) rendering riverine habitats unsuitable to certain SGCN.	High
3. Sedimentation	Habitat degradation resulting from land development and uses; dams disrupting natural sediment transport; flushing sediments from impoundments; excessive bank erosion from inadequate riparian vegetation.	High
4. Habitat fragmentation	Interruption of migration corridors to and from breeding/spawning/wintering habitats via alteration and conversion of home range; construction of dams and culverts.	High
6. Invasion by exotic species	Displacement or restructuring of native aquatic plant and animal communities by invasive organisms impacting habitat and community structure and processes.	Med
7. Pollution	Nutrient overloading and other pollutants.	High
8. Pollution	Vulnerability to catastrophic spills: Bordering roadways, bridge crossings, adjacent industry and urban centers pose high risk points of entry for large-scale contaminant spills.	High
10. Monitoring	Population and habitat monitoring: Improved data on known SGCN populations is needed to track changes in species abundance and habitat quantity and quality as may be affected by natural processes and anthropogenic factors; habitats with potential for having existing SGCN populations or SGCN restoration potential should be investigated.	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Monitor, protect and restore riparian and in-stream habitats limited or impacted by development.	Number of SGCN sites (habitats) monitored; acres/miles of undisturbed habitats protected; acres/miles of disturbed habitats restored.	ANR, USFWS, NRCS, CRJC, TNC, Power Companies	EPA
Monitor, protect and restore river and stream water quality from excessive nutrient and sediment loading and other pollutants.	Miles of SGCN habitat meeting water quality standards.	ANR, USFWS, NRCS, CRJC, TNC	EPA
Monitor, protect and restore migration and travel corridors limited or impacted by dams, culverts and roads.	Number of identified artificial migration barriers removed or mitigated; miles of critical habitat restored by removal of barriers.	ANR, CRASC, USFWS, CRJC, VTrans, Utilities	EPA, USACE
Monitor the Connecticut River and its tributaries for invasive species; prevent the introduction or spread of invasive species; implement control measures which take into account SGCN and their habitat requirements.	Number of SGCN habitats monitored for invasive species; number of SGCN habitats with plans in place designed to control invasive species and restore or enhance SGCN.	ANR, USFWS, CRJC, VY, TNC	WHIP, EQUIP, USFWS Conte Grants, EPA
Support policies and programs designed to reduce climate change.	Number of climate change policies and programs established or supported.	ANR, EPA, Other NE States	EPA
Conduct inventories to detect and gather information on new SGCN populations and their habitats.	Number of potential SGCN habitats surveyed.	ANR, USFWS, TNC, USGS, EPA	EPA

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Provide technical outreach and financial assistance to private landowners, towns, watershed and lake associations, regional planning commissions, and other partners to increase their awareness of problems to SGCN.	Number of actions implemented to maintain or enhance river function for SGCN.	ANR, USFWS, NRCS, CRJC, TNC	Farm Bill, Conte Grants, EPA
Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004) to town and Regional Planning Commissions.	Number of towns and RPC considering SGCN in their planning. Number of actions implemented to maintain or enhance river function for SGCN.	AVCC	SWG, VFWD
Acquire conservation easements for the protection of SGCN sites and maintenance or restoration of their ecological functions.	Number of SGCN habitats acquired or enrolled in land conservation easement programs.	ANR, USFWS, TNC	EPA
Enhance coordination between government agencies/partners to ensure consistency in respective program implementation and increase sensitivity to problems and requirements for SGCN.	Number of agencies and private conservation organization, which recognize and address problems to SGCN.	ANR, USFWS, USFS, NRCS, USACE, VTrans, CRJC, TNC	EPA

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
Connecticut River Corridor Plan	"That plants, migratory birds, anadromous fish, and other native birds, fish, and wildlife continue to find the Connecticut River corridor and watershed hospitable to their unique needs for clean water and connected, protected open lands and forests;"	CRJC
A Plan to Restore the Aquatic Ecosystem in the Connecticut River Watershed	"Restore aquatic ecosystem so as to recover and support migratory and native fish populations and promote natural reproduction in the Connecticut River and its tributaries."	NRCS
Strategic Plan for the Restoration of Atlantic Salmon to the Connecticut River.	"Protect, conserve, restore and enhance the Atlantic salmon population in the Connecticut River for the public benefit, including recreational fishing."	CRASC
A Management for American Shad in the Connecticut River Basin.	"Restore and maintain a spawning shad population to its historic range in the Connecticut River Basin and to..."	CRASC
Management Plan for Blueback Herring in the Connecticut River Basin.	"Restore and maintain a spawning blueback herring population within its historic range in the Connecticut River basin."	CRASC
Plan for the Restoration of Migratory Fishes to the Ashuelot River Basin, New Hampshire.	"Protect, conserve, restore, and enhance the migratory fish populations in the Ashuelot River system for both public and ecological benefits."	NHFG

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. *Conserving Vermont's Natural Heritage*. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- CRJC (Connecticut River Joint Commissions). 2009. [Connecticut River Water Resources Management Plan](#). CRJC, Charlestown, NH.

Lake Champlain Summary

Lake Champlain includes the main body of the lake and its bays and river deltas. These waters are shared with New York and Quebec. At about 120 miles in length and a maximum depth over 400 feet, this is Vermont's largest waterbody. Aquatic habitats found here are many and extensive. Among these are expansive sand-bottomed shallows, shale/cobble littoral shorelines and bays, and deep limnetic environments. Other natural communities, such as large tributaries, emergent marshes, and floodplain forests, are integral to Lake Champlain and provide a critical habitat component for many SGCN found here. This is, in general, an oligo-mesotrophic lake, with nutrient levels in different parts of the lake dependent on local soil and bedrock types, as well as the type and extent of human land use within the surrounding watershed. This lake supports the highest lacustrine diversity of any of our lakes, which is due mainly to its large size and connections (current and historical) with the Great Lakes-St. Lawrence River and the Hudson River.

Lake Champlain Condition

Current Condition: The most outstanding concerns facing this large system are water quality and habitat degradation, and invasive exotic species. The lake is within the largest watershed in Vermont and is fed by many large tributaries that drain extensive agricultural and developed lands. A significant portion of the excessive nutrients, contaminants, and fine sediments that enter streams and rivers eventually reach Lake Champlain. Water and benthic habitat quality are affected, particularly in delta areas and along the shoreline, but also within the open and deeper waters over time. SGCN that are sensitive to contaminants and those that depend on consolidated (firmly-packed) substrates may be impacted by these changes to their habitat. Development along Lake Champlain's shoreline and within smaller watersheds immediate to the lake is ever-increasing, and with it the amount of contaminants entering directly into the lake. Excessive nutrients that reach the lake from various land uses within the watershed can cause eutrophication, reducing water quality and altering food webs. Zebra mussels have had a dramatic and devastating impact on the biotic community of Lake Champlain, including populations of many SGCN. These exotic pests foul the shells of native freshwater mussels, decreasing their ability to move about and obtain food and oxygen, resulting in a slow death. Populations of native mussels have been eliminated from large areas, a scenario that has repeated itself throughout most of the lake. The only areas where native mussels have not been seriously impacted by zebra mussels are Mallets Bay, the Inland Sea, and Missisquoi Bay. Water chestnut is an invasive that has impacted aquatic communities in the lake by forming huge, dense masses that cover the water surface and crowd out species. The exotic snail *Bithynia tentaculata* now dominates much of the shale/cobble habitat in Lake Champlain, likely reducing native snail populations and altering the food web. Among other invasive exotics that loom on the horizon for Lake Champlain are round goby, alewife, quagga mussel, and rusty crayfish. Additional problems to Lake Champlain include habitat conversion and vulnerability to catastrophic contaminant spills.

Desired Condition (SGCN Needs): These waters, along with Lake Champlain tributaries, support the greatest diversity of aquatic species found in the state. SGCN supported by Lake Champlain include mid- to deep-water species like cisco and lake whitefish that require cold, well-oxygenated waters. Shallow-water species such as mooneye and sauger utilize upper portions of the lake where temperatures are often much warmer. Near-shore and benthic species like bridle shiner, pink heelsplitter, giant floater, and spiny softshell are often found in bays or in the shallows of deltas.

The lake offers a variety of habitats that provide for the many needs of aquatic species, such as refuge, food, thermal protection, and spawning substrate. The great majority of freshwater mussel species remain buried in the substrate most of their lives, where they grow, feed, produce offspring, and seek refuge from the elements. Lake sturgeon feed on lake-bottom invertebrates, only entering rivers for brief periods to spawn. Different fishes can be found occupying different strata of the lake where they find the temperatures and oxygen levels they prefer. Degradation of water quality through nutrient input, thermal shifts, or other changes can cause significant alterations in food webs and habitat availability. Similarly, excessive fine sediments entering the lake from the shoreline and tributaries blankets and degrades the benthic substrate used by many SGCN. Improvement and protection of Lake Champlain's water quality, including reduction of nutrient and fine sediment inputs, is paramount to ensure that the SGCN populations found here remain viable. Control of exotic species, including preventing new species from invading, is also of great importance to the survival of these native species.

Many SGCN utilizing Lake Champlain depend on closely associated aquatic, wetland, and terrestrial habitats to complete their life cycles. Many fish, such as lake sturgeon, greater redhorse, and mooneye are found in the lake most of the year, but spawn over rocky substrates in Champlain tributaries. Bald eagle feed in the lake but need nearby suitable nesting trees or structures to raise their young. Spiny softshells occupy the lake much of the year for basking, feeding, and over-wintering, but require adjacent beaches of sand or gravel/cobble for egg-laying. Bats feed on emerging aquatic insects over the lake, while utilizing upland roosting and nursery sites. Muskrat, river otter and mink find a rich aquatic food source within Lake Champlain and its associated wetlands, but must den above the waterline. Maintaining these connections to critical wetland, aquatic, and terrestrial habitat is key to ensuring the continuation of these SGCN in the lake.

Species of Greatest Conservation Need in Lake Champlain

High Priority

Bald Eagle (*Haliaeetus leucocephalus*)
 American Brook Lamprey (*Lethenteron appendix*)
 Lake Sturgeon (*Acipenser fulvescens*)
 American Eel (*Anguilla rostrata*)
 Silver Redhorse (*Moxostoma anisurum*)
 Greater Redhorse (*Moxostoma valenciennesi*)
 Sauger (*Sander canadense*)
 Blackchin Shiner (*Notropis heterodon*)
 Bridle Shiner (*Notropis bifrenatus*)
 Crustaceans Group
 Freshwater Mussels Group
 Freshwater Snails Group
 Spiny Softshell (*Apalone spinifera*)
 Silver-haired Bat (*Lasiorycteris noctivagans*)
 Eastern Red Bat (*Lasiurus borealis*)
 Hoary Bat (*Lasiurus cinereus*)
 Tri-colored bat (*Perimyotis subflavus*)

Medium Priority

Peregrine Falcon (*Falco peregrinus*)
 Cisco or Lake Herring (*Coregonus artedii*)
 Lake Whitefish (*Coregonus clupeaformis*)
 Mooneye (*Hiodon tergisus*)
 Silver Lamprey (*Ichthyomyzon unicuspis*)
 Shorthead Redhorse (*Moxostoma macrolepidotum*)
 Atlantic Salmon (landlocked) (*Salmo salar*)
 Lake Trout (naturally reproducing populations) (*Salvelinus namaycush*)
 Muskrat (*Ondatra zibethicus*)
 Northern River Otter (*Lontra canadensis*)

SGCN Note: SGCN plants not listed here 7 species. The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
1. Habitat Alteration	excessive sediments and nutrients from surface runoff and tributaries; caused by human land use nearby	High
2. Habitat Conversion	benthic habitat due to riprapping, bridge construction, boat access construction, etc.	High
3. Pollution	bility to Catastrophic Spills: Bordering roadways, bridge crossings, adjacent industry, and manure pits are examples of high risk points of entry for large-scale contaminant spills	High
4. Invasion by exotic species	Zebra mussels and water chestnut are currently impacting SGCN; other exotics may also be displacing native SGCN	High
5. Pollution	Water quality degradation due to contaminants from agricultural fields, stormwater runoff, other point and non-point sources	High
6. Inventory	Inventory needed for many SGCN, particularly those for which distributional and abundance information is greatly lacking	High
7. Monitor	Detect SGCN population trends to help guide conservation actions and to track the effectiveness of current management	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here.

See Chapter 6 for definitions of acronyms used in the Partners and Funding Source columns

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Monitor known SGCN populations	Number of known SGCN sites monitored	USFWS, ANR, TNC, Universities	SWG, VFWD, VT Watershed Grants
Conduct inventories to detect and gather information on new SGCN populations	Number of completed species or species-group inventories	USFWS, ANR, TNC, Universities	SWG, VFWD, VT Watershed Grants
Protect and restore habitats on which SGCN are dependent through pollution abatement, riparian buffers, etc.	Number of acres of riparian and lakeshore natural vegetation protected and/or restored Number of acres of lake habitat restored/protected	LCLT, VLT, Watershed groups, USFWS, ANR, EPA	SWG, LCLT, VLT, NRCS, EPA, Clean Water Fund
Restore migration corridors for SGCN by removal of artificial barriers to spawning habitat or construction of effective fish passage facilities at dams	Number of artificial SGCN migration barriers removed or provided with passageways Number of adult SGCN fish passed migrating to upstream spawning habitat (e.g., lake sturgeon, greater redhorse)	USFWS, Hydro operators, FERC, ANR, Municipalities, VNRC, EPA	NRCS, USFWS, Clean Water Fund, EPA
Implement an invasive species monitoring program to prevent the introduction and spread of invasive exotic species. Manage, mitigate, and/or eliminate invasive species that are detected.	Estimated percent of invasive exotic species controlled per year. Number of sites with control activities and/or invasive monitoring	LCBP, ANR, Municipalities, USFWS, EPA	USFWS, VT Watershed Grants, LCBP, EPA, Clean Water Fund
Provide technical outreach and financial assistance to private landowners, watershed groups, and other partners to maintain and enhance Lake Champlain for SGCN.	Number of actions implemented to maintain or enhance lake suitability for SGCN	EPA, VFWD, TNC, LCBP, VLCT, LCI, Watershed groups,	VT Watershed Grants, LCBP, EPA, Clean Water Fund

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Provide technical outreach to towns and regional planning commissions to maintain and enhance Lake Champlain for SGCN. Distribute <i>Conserving Vermont's Natural Heritage</i> (Austin et.al. 2004)	Number of towns considering SGCN in their planning. Number of actions implemented to maintain or enhance lake suitability for SGCN	EPA, AVCC, LCBP, RPC's Municipalities,	EPA, SWG
Acquire conservation easements for the protection of SGCN sites and maintenance or restoration of ecological functions	Number of riparian habitat acres acquired/enrolled	LCLT, VLT, ANR, TNC, NRCS, EPA	LCLT, VLT, TNC, SWG, NRCS, EPA, Clean Water Fund
Enhance coordination between government agencies and partners to ensure consistency in respective program implementation and increased sensitivity to SGCN requirements and problems to SGCN	Number of programs that incorporate SGCN conservation.	ANR, USFWS, COE, FEMA, FHWA, NRCS, LCI, Wildlife Services, VTrans	USFWS, EPA, Clean Water Fund

Coordination with other plans

See Chapter 6 for definitions of acronyms used in the lead column

Plan or planning entity	Goal/Scope of plan	Lead
Lake Sturgeon Recovery Plan	Lake Sturgeon restoration	VFWD
Vermont's Clean Water Initiative	Water quality improvement	VDEC
Vermont Lake Champlain Phosphorus TMDL Implementation Plan (Phase 1)	Reduction of phosphorous inputs to Lake Champlain	VDEC, AAFM
Lake Champlain Basin Aquatic Nuisance Species Management Plan (2002).	Management and prevention of invasive exotic species in the basin	VTDEC, NYDEC
VTDEC Water Quality Division	Lake protection and restoration programs	VTDEC
NYDEC	Lake protection and restoration programs	NYDEC
Quebec Ministère de l'Environnement	Protection of Québec's ecosystems and biodiversity; prevention, reduction or elimination of water contamination	Quebec Ministère de l'Environnement
Vermont Osprey Recovery Plan	Recovery and management of osprey within VT	VFWD
Conserving Lake Champlain's Biological Diversity 6/10/2005	Strategic plan focused on conserving Lake Champlain's biological diversity	TNC
Various watershed planning efforts	Watershed protection and restoration; river and lake restoration and protection	VTDEC; local/regional watershed groups

References

- Austin, J.M. C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson. VT League of Cities and Towns. 2004. Conserving Vermont's Natural Heritage. [A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity](#). Vermont Fish & Wildlife Department and Agency of Natural Resources. Waterbury, VT.
- Budd, L. and D. Meals. 1994. Lake Champlain Nonpoint Source Pollution Assessment. LCBP Technical Report No. 6A. U. S. Environmental Protection Agency. Boston, MA.
- Fichtel, C., and D. G. Smith. 1995. The freshwater mussels of Vermont. Nongame & Natural Heritage Program, Vermont Fish & Wildlife Department. Technical report 18.
- Good, S. 2001. Alternative Strategies for the management of non-indigenous alewives in Lake St. Catherine, Vermont. Draft report. Vermont Fish and Wildlife Department, Pittsford.

- Hegman, W. et al. 1999. Estimation of Lake Champlain Basinwide Nonpoint Source Phosphorus Export. Lake Champlain Basin Program Technical Report No. 31, Grand Isle, VT.
- Meals, D. 2001. Lake Champlain Basin Agricultural Watersheds Section 319 National Monitoring Program Project: Final Project Report May 1994 - November 2000. Vermont Department of Environmental Conservation. Waterbury.
- Neddeau, E. J., M. A. McCollough, and B. I. Swartz. 2000. The freshwater mussels of Maine. Maine Department of Inland Fisheries and Wildlife.
- New York State Department of Environmental Conservation. 1992. Agricultural Management Practices Catalogue for Nonpoint Source Pollution Prevention and Water Quality Protection in New York State. Albany, NY: New York State Nonpoint Source Management Practices Task Force.
- Lake Champlain Basin Program. 2003. Opportunities for Action: An evolving plan for the future of the Lake Champlain basin. Grand Isle, VT.
- Stangel, P. 2004. Lake Champlain 2003 zebra mussel monitoring program. Final report. Vermont Department of Environmental Conservation. Waterbury, VT.
- Vermont Department of Environmental Conservation. 2004. 2004 Vermont Water Chestnut Management Program Summary. Waterbury, VT.
- Vermont Department of Environmental Conservation and New York State Department of Environmental Conservation. 1997. Lake Champlain Diagnostic-Feasibility Study Final Report - A phosphorus budget, model, and load reduction strategy for Lake Champlain. Waterbury, VT and Albany, NY.
- Vermont Department of Environmental Conservation and New York State Department of Environmental Conservation. 2005. Lake Champlain Long-Term Water Quality and Biological Monitoring Program. Waterbury, VT and Albany, NY.

Lakes Summary (excluding Lake Champlain)

Characteristics and Location

Lakes or lacustrine areas include natural lakes and ponds throughout Vermont, which can be classified on the basis of their productivity and associated physio-chemical characteristics. Lake types discussed here include oligotrophic lakes, mesotrophic lakes, eutrophic lakes, high elevation acidic lakes, and dystrophic lakes. Lake Champlain, representing oligotrophic, mesotrophic and eutrophic habitat types, is not included in this summary due to its large size and unique species assemblages (see Lake Champlain Summary). The following descriptions of Vermont lake types are based in part on parameters provided by the Vermont Department of Environmental Conservation, Water Quality Division, Lakes and Ponds Section.

Types of Lake Communities:

Oligotrophic Lakes: These lakes are typically deep with clear, cold water; low in dissolved nutrients, such as phosphorus and nitrogen; and experience seasonal periods of temperature and oxygen stratification and de-stratification (mixing). Biochemical parameters generally characterizing this lake type are: (1) total phosphorus concentration in the summer photic zone, $<10 \mu\text{g/L}$; total nitrogen concentration, $<0.35 \text{ mg/L}$; average summer chlorophyll-a concentration, $\leq 3.5 \mu\text{g/L}$; and average summer secchi disc depth, $\geq 5.5 \text{ m}$. Another general feature of oligotrophic lakes is the lack of an extensive littoral zone. Littoral plants are scarce and plankton density is low. Several SGCN uniquely associated with this lacustrine waters are landlocked Atlantic salmon, lake trout, and round whitefish. In Vermont, lakes of this type are predominantly located in the Northeast Highlands biophysical region.

Mesotrophic Lakes: Lakes of this type are intermediary between oligotrophic (nutrient poor) and eutrophic (nutrient rich) systems. Mesotrophic lakes are shallower than oligotrophic lakes, have a well-established littoral zone supporting aquatic vegetation, and are moderately rich in dissolved nutrients. Consequently, primary productivity and plankton densities are greater than in oligotrophic systems but less than in eutrophic waters. Biochemical parameters generally characterizing this lake type are: (1) total phosphorus concentration in the summer photic zone, $10 \text{ to } 24 \mu\text{g/L}$; total nitrogen concentration, $0.35 \text{ to } <0.65 \text{ mg/L}$; average summer chlorophyll-a concentration, $>3.5 \text{ to } 7.0 \mu\text{g/L}$; and average summer secchi disc depth, $3.0 \text{ to } 5.5 \text{ m}$. Several SGCN uniquely associated with meso-eutrophic lakes are bridle shiner, blackchin shiner, redbfin pickerel, redbreast sunfish, Eastern Musk Turtle, and northern water snake. Lakes of this type are distributed throughout Vermont; however, those supporting one or more populations of SGCN tend to be represented in greater frequency in the Champlain Valley and Connecticut River biophysical regions.

Eutrophic Lakes: Lakes of this type are generally characterized as nutrient mature systems. They are richer in dissolved nutrients and generally shallower than oligotrophic and mesotrophic lakes with extensive littoral areas supporting prolific growths of aquatic vegetation. Primary productivity and plankton densities are greater than in mesotrophic lakes. Eutrophic lakes that thermally stratify are likely to experience oxygen depletion below the thermocline during summer and/or winter stratification periods. Oxygen depletion during winter can occur when ice cover prohibits atmospheric exchange of

oxygen resulting in “winter-kill” conditions. Biochemical parameters generally characterizing this lake type are: (1) total phosphorus concentration in the summer photic zone, $>24 \mu\text{g/L}$; total nitrogen concentration, $>0.65 \text{ mg/L}$; average summer chlorophyll-a concentration, $\geq 7.0 \mu\text{g/L}$; and average summer secchi disc depth, 0 to 3.0 m. SGCN associated with eutrophic lakes are similar to mesotrophic lakes with decreasing occurrence in lakes of more advanced eutrophication. Though advanced eutrophication may make unsuitable habitat for purely aquatic SGCN, the productivity of these waters may be important to terrestrial and semi-aquatic species (e.g., bald eagle, bats and northern water snake) due to the abundance of food organisms these waters are capable of producing. Lakes of this type are distributed throughout Vermont but are more likely to be at low elevations and in disturbed landscapes.

High Elevation Acidic Lakes: These are clear-water lakes generally located at elevations over 1500 feet with neutralizing capacity (ANC) less than 25 mg/L and more typically within the range of 0 to 5 mg/L. Lakes of this type are vulnerable to and in some cases are known to be adversely affected by acid deposition. These lakes are usually small and shallow, with rocky or gravelly bottoms, and little accumulated organic material. Dissolved nutrient concentrations and primary production are generally low. Relatively few SGCN are associated with high elevation acidic lakes. One possible associate is brook trout. In Vermont lakes of this type are generally distributed within the Northern and Southern Green Mountain biophysical regions.

Dystrophic Lakes: Lakes of this type are usually associated with bogs. These are characterized by brown stained water (color $>50 \text{ Pt Co}$) and are high in nutrients and humic materials. Dystrophic lakes are often acidic and may be anoxic or nearly so in the deeper waters. Relatively few SGCN are associated with dystrophic lakes with the possible exception of brook trout. Although examples of dystrophic lakes may be found statewide, generally they are more abundant in the Northern Green Mountains, Southern Green Mountains and Northeast Highlands biophysical regions.

Lake Condition

Current Condition: The lake waters represented here have notably different physio-chemical characteristics, therefore problems and changes to their water quality and chemistry may affect each lake type and species assemblages in different ways. Most oligotrophic and mesotrophic lakes in Vermont have experienced abundant lakeshore development, both historically and currently, such as seasonal and permanent residences, marinas and docks, and public and private beaches. In many instances these developments have altered natural lakeshore and littoral habitats through the addition of fill materials (e.g., sand, bottom barriers), removal of large woody debris, and removal of native aquatic vegetation for beach construction and maintenance, resulting in the direct loss of habitats for SGCN. Additionally, development has increased stormwater runoff to lakes and has elevated the input of pollutants, including sediments, nutrients, and toxic chemicals. Nutrient loading can accelerate the eutrophication process causing excessive growth of phytoplankton and other aquatic vegetation, reduced water clarity, and increased biological oxygen demand. Such water quality and habitat changes may be detrimental to certain SGCN associated with specific lakes. Many SGCN species are heavily dependent on healthy aquatic systems for food sources, such as abundant fish and/or invertebrate populations utilized by eagles, river otter, muskrat and bats.

Development of shorelands may alter habitat required by terrestrial SGCN that are associated with lacustrine areas, such as bald eagle and osprey. As an example, the reduction of mature trees by clearing within the riparian area may eliminate eagle and osprey nesting sites and reduce recruitment of woody debris into the littoral zone.

The invasion of lacustrine waters by habitat-altering exotic species and the subsequent control of these exotics may have degraded habitat for some aquatic SGCN. For example, the establishment of Eurasian milfoil in several mesotrophic lakes where blackchin shiners are known to occur has likely displaced native aquatic plant communities on which this fish species is dependent for spawning and refuge. While milfoil control activities, such as herbicide treatment, are conducted to restore lake conditions conducive to water-based recreational pursuits (boating, swimming, sport fishing), the result is loss of vegetative cover now provided by milfoil stands, increased predation on shiners by other resident fishes (e.g., bass, sunfish, pike), as well as the loss of spawning habitat. These pressures on blackchin shiner populations continue until littoral areas are adequately revegetated with native plant species, a process that may take many years.

The deliberate and accidental introduction of plant and animal species to Vermont's lakes and ponds over the past 200 years has greatly changed natural communities and their ecological functions. Many fish species, including those native to the state as well as those brought from outside, have established in waters where they did not naturally occur. For example, largemouth bass, bluegill and northern pike, all native in Vermont to Lake Champlain only, now have transplanted populations in habitats nearly statewide. Rainbow and brown trout originated from the western United States and Europe, respectively, and now are established in many lakes within the state. The distribution of these species was expanded beyond their natural range for the primary purpose of increasing sport fishing opportunities; however, in the past little consideration was given to the negative effects these species have on native ecosystems. More recently, 1997, the exotic alewife was discovered in Lake St. Catherine where previously the species did not exist. This was the first recorded occurrence of alewife in the state. The impacts this species has on native fish communities are well documented, including: (1) out-competing other planktivores for food and causing shifts in zooplankton species composition and size structure; (2) preying on the eggs and larvae of native fishes; and (3) causing significant mortality syndrome in salmon and trout fry (Good 2001). The trans-state movement and introduction of exotic species into natural habitats has become an environmental problem of national scale. Past species introductions changed the current character of many Vermont lacustrine areas, and the problem is a persistent problem for maintaining lakes and ponds in a desired condition well into the future.

Currently dystrophic and high elevation acidic lakes are somewhat less limited by direct development pressures that other lake types are experiencing. On the other hand, these lakes are particularly vulnerable to habitat alteration through the effects of acid deposition.

Desired Condition (SGCN Needs): Lacustrine areas directly and indirectly support a host of species, including aquatic invertebrates (insects, crustaceans, mollusks), fishes, reptiles, amphibians, mammals, birds and plants.

Obligate SGCN associated with oligotrophic lakes (e.g., landlocked Atlantic salmon, lake trout, round whitefish) require deep, clear, well-oxygenated water for their survival. Potential

increases in lake water temperatures due to climate change represent a problem for these oligotrophic lakes and the associated cold-water SGCN.

In contrast, species associated with mesotrophic and eutrophic lakes are more dependent on the high productivity of these lake systems to produce needed food sources and habitat complexity, such as well-established littoral communities for feeding, reproduction and refuge cover.

A number of SGCN, notably the reptiles and amphibians, have home ranges that encompass both lacustrine and terrestrial areas at particular times of the year. For example, spotted and musk turtles, which reside most of the year in lakes and ponds, leave these waters briefly for upland areas to lay eggs. Similarly, lake residing brook trout may seasonally ascend tributary streams to spawn. In contrast, Fowler's toads travel from their usual terrestrial haunts to aquatic habitats to deposit eggs along the shoreline. Forested riparian zones provide nesting and feeding perches for bald eagle and osprey. Mature trees that eventually die and are recruited into the littoral area contribute to forming refuge and basking habitats. Maintenance of water quality conditions characteristic of specific lake types is a requirement of SGCN associated and dependent on those habitats.

The desired condition for all lacustrine communities would include: 1) the existence of intact riparian conditions; 2) the existence of minimally disturbed littoral zones; 3) evolutionary (e.g trophic) processes occurring at rates not accelerated by disturbance; 4) pollutant levels (e.g sediment and toxics, including acid deposition) below concentrations that would adversely affect SGCN; 5) absence of exotic species that adversely affect SGCN; 6) unimpeded access by SGCN to habitats required for the maintenance of life cycle functions; and 7) unaltered hydrological and temperature regimes.

Implementing the 2005 Wildlife Action Plan

In 2013, the Department completed the Vermont [BioFinder](#) project, a map and database identifying Vermont's lands and waters supporting high priority ecosystems, natural communities, habitats, and species. A notable outcome of the project was a classification of lakes and ponds based on alkalinity and trophic status and identification of best examples of each type.

SGCN in Lacustrine Communities (excluding Lake Champlain)

High Priority

Bald Eagle (*Haliaeetus leucocephalus*)
American eel (*Anguilla rostrata*)
Bridle Shiner (*Notropis bifrenatus*)
Blackchin Shiner (*Notropis heterodon*)
Fowler's Toad (*Anaxyrus fowleri*)
Common Mudpuppy (*Necturus maculosus*)
Odonates-Bog/Fen/Swamp/Marshy Pond Group
Crustaceans Group
Freshwater Snails Group
Odonates-Lakes/Ponds Group
Odonates-River/Stream Group
Silver-haired Bat (*Lasionycteris noctivagans*)
Eastern Red Bat (*Lasiurus borealis*)
Hoary Bat (*Lasiurus cinereus*)
Tri-colored bat (*Perimyotis subflavus*)

Medium Priority

Peregrine Falcon (*Falco peregrinus*)
Redfin Pickerel (*Esox americanus*)
Redbreast Sunfish (*Lepomis auritus*)
Round Whitefish (*Prosopium cylindraceum*)
Atlantic salmon-landlocked (*Salmo salar*)
Brook Trout-naturally reproducing populations
(*Salvelinus fontinalis*)
Lake Trout-naturally reproducing populations
(*Salvelinus namaycush*)
Northern Water Snake (*Nerodia sipedon*)
Eastern Musk Turtle (*Sternotherus odoratus*)
Muskrat (*Ondatra zibethicus*)

SGCN Note: Plant SGCN not listed here 25 species. The SGCN invertebrate groups listed here contain numerous species. For more information about a specific Species of Greatest Conservation Need see that species' assessment summary in Appendix A.

Problems & Information Needs

See Appendix C for definitions of problem and strategy categories used here

Problem/Info Need Category	Problem/Info Need Detail	Rank
1. Habitat Conversion	Loss of riparian, shoreline and littoral habitats from land and water development projects and activities.	High
2. Habitat Alteration	Alteration and degradation of riparian, shoreline and littoral habitats from development, invasive species, and aquatic vegetation control; water level regulation; loss and inadequate recruitment of large woody debris.	High
3. Sedimentation	Alteration and degradation of habitat (e.g., spawning areas); smothering of organisms.	High
4. Habitat Fragmentation	Interruption of migration and travel corridors to and from breeding/spawning/wintering habitats via alteration and conversion home range; construction of roads, dams and culverts.	High
6. Invasion by Exotic Species	Alteration and conversion of native littoral plant communities; inter-species competition for habitat and food; predation on native species; impacts resulting from invasive species control programs and activities.	High
7. Climate change	Alteration of water and temperature regimes.	High
8. Pollution	Nutrient and sediment overloading, acid deposition and other pollutants.	High
9. Pollution	Nutrient input to lakes accelerates the eutrophication process altering normal trophic succession.	High
10. Monitoring	Population and habitat monitoring: Improved data on known SGCN populations is needed to track changes in species abundance and habitat quantity and quality as may be affected by natural processes and anthropogenic factors; habitats with potential for having existing SGCN populations or SGCN restoration potential should be investigated.	High

Priority Conservation Strategies

See Appendix C for definitions of problem and strategy categories used here

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Monitor, protect and restore riparian, shoreline and littoral habitats limited or impacted by development.	Number of SGCN sites (habitats) monitored; acres of undisturbed habitats protected; acres of disturbed habitats restored.	ANR, USFWS, NRCS, USFS, Lake and Watershed Associations	USFWS, EPA, Clean Water Fund, ANR
Monitor, protect and restore lake and pond water quality from excessive nutrient and sediment loading, other pollutants, and acid deposition.	Acres of SGCN habitat meeting water quality standards.	ANR, USFWS, NRCS, USFS, Lake and Watershed Associations	ANR, Clean Water Fund
Monitor, protect and restore migration and travel corridors limited or impacted by roads, dams, culverts, etc.	Number of identified artificial migration barriers removed or mitigated; number of migration corridors protected.	ANR, USFWS, NRCS, USFS, VTrans	USFWS, EPA
Monitor, protect and maintain known softshell turtle nesting sites; restore and protect additional nest sites.	Number of nest sites monitored, managed and protected; nest sites restored.	ANR, USDA, Wildlife Services, EPA	USFWS, EPA

Strategy	Performance Measure	Potential Partners	Potential Funding Sources
Monitor lakes and ponds for invasive species; implement programs to prevent the introduction or spread of invasive species; implement control measures which take into account SGCN and their habitat requirements.	Numbers of SGCN habitats monitored for invasive species; number of SGCN habitats with plans in place designed to control invasive species and restore or enhance SGCN; incorporation of SGCN.	ANR, Lake and Watershed Associations	
Support policies and programs designed to reduce climate change.	Number of climate change policies and programs established or supported.	ANR, EPA, Other NE States	
Conduct inventories to detect and gather information on new SGCN populations and their habitats.	Number of potential SGCN habitats surveyed.	ANR, USFWS, USFS, EPA, USGS	USFWS
Provide technical outreach and financial assistance to private landowners, towns, watershed and lake associations, regional planning commissions, and other partners to increase their awareness of problems to SGCN.	Number of actions implemented to maintain or enhance lake function for SGCN.	ANR, USFWS, NRCS, TNC	USFWS
Acquire conservation easements for the protection of SGCN sites and maintenance or restoration of their ecological functions.	Number of SGCN habitats acquired or enrolled in land conservation easement programs.	ANR, TNC, USFS	USFWS
Enhance coordination between government agencies/partners to ensure consistency in respective program implementation and increase sensitivity to SGCN requirements and problems to SGCN.	Number of agencies and private conservation organization, which recognize and address problems to SGCN.	ANR, USFWS, USFS, NRCS, VTrans, TNC, Lake and Watershed Associations	

Coordination with other plans

Plan or planning entity	Goal/Scope of plan	Lead
Vermont's Clean Water Initiative	Water quality improvement	VDEC
DEC Water Quality Division	Lake protection and restoration programs	VDEC

References

Good, S. 2001. Alternative Strategies for the management of non-indigenous alewives in lake St. Catherine, Vermont. Draft report. Vermont Fish and Wildlife Department, Pittsford.

Reid, G. K. 1961. Ecology of inland waters and estuaries. D. Van Nostrand Company, New York.

Vermont's Wildlife Action Plan

Threat & Action Definitions

Appendix C

Draft • 10/1/2015

Threat Categories for Wildlife Action Plan Revision	2
Conservation Actions for Wildlife Action Plan Revision	7

Threat Categories for Wildlife Action Plan Revision

Element number three of the eight congressionally required elements of a Wildlife Action Plan requires that states: describe the problems that may adversely affect Species of Greatest Conservation Need or their habitats and priority research and survey efforts needed to identify factors which may assist in restoration and improved conservation of these species and habitats. Problem and threats are defined as follows:

Problem: Something that is a concern and could cause a negative impact at the species, population, habitat and/or landscape levels (e.g., habitat conversion, pollution, illegal pet trade). A problem can also be the lack of information or a data gap vital to the successful management of a species.

Threat (direct): Processes or human activities “that have caused, are causing, or may cause the destruction, degradation, and/or impairment of biodiversity targets” (adapted from Salafsky et al. 2008).

Threat (indirect): The factors contributing to or enabling direct threats. Typically there is a chain of contributing factors behind any given direct threat. Synonyms include contributing factors, underlying factors, drivers, and root causes (adapted from Salafsky et al. 2008).

For the purposes of this report, problem and threat are used in a similar or related manner. For each Species of Greatest Conservation Need in the Action Plan we identified priority problems. Priority research needed to evaluate other potential problems was also identified. They are detailed in SGCN conservation summaries (Appendix A) and in habitat/ community summaries (Appendix B).

Each of the threats and problems identified in the Action Plan was assigned to one of 24 categories roughly grouped into habitat-related factors and non-habitat-related factors. These categories make it possible to search our database for similar factors impacting other species. It also makes it easier to roll-up for broad scale conservation planning. The categories were cross-walked (Appendix C) with those developed by the International Union for Conservation of Nature and Natural Resources (IUCN) (Salafsky et al. 2008) to aid in the regional roll-up of Action Plan data as recommended by the Diversity Technical Committee of the Northeast Association of Fish & Wildlife Agencies (Crisfield 2013).

The categories are not mutually exclusive and threats can often logically be placed into more than one category depending on the particular stress it causes for a species or habitat. For example, a road can fragment the habitat of grassland nesting birds, cars traveling the road can injure or kill amphibians that were crossing the road to mate in an adjacent pool, and salt spread on the road to prevent icing can wash into a stream impacting its population of Brook Trout. In this example the threats stemming from the road would be recorded in the "Habitat Fragmentation," "Impacts of Roads & Transportation Systems," and "Pollution" categories.

Threats are often species and/or habitat specific. What may negatively impact one species may benefit another. For example, if a cold water stream with a healthy Brook Trout population was dammed it might no longer support Brook Trout. That impact to the dam would be described as the "conversion of habitat" category. However, the reservoir created by the dam might make it more suitable for a warm water fish species.

Clearly life is too complex to be placed into any one box. Therefore it is important to read the full description of a factor affecting a species or habitat in the appropriate species or habitat summary. Table C.1 lists the threat categories used in this Wildlife Action Plan.

Table C.1 Threat Categories for Wildlife Action Plan Revision

#	Threat/ Problem	Description	IUCN Category
Habitat-Related Categories			
A	Climate Change	Long-term changes linked to global warming and other climate issues that can lead to major changes in habitat availability (e.g., high elevation habitats), vegetative composition and location (e.g., the movement up in elevation or north in latitude, invasion by exotic pests) as well as climate variability (e.g., change in snow depth, rainfall and/or natural disturbances).	11. Climate Change and Severe Weather
B	Habitat Alteration/ Degradation	A lessening of the quality of a habitat by human action stopping short of complete conversion (e.g., the reduction of mast production from a forest stand, the riprapping a streambank, and significant land use changes adjacent to a habitat such as replacing a forest stands on the edge of a wetland with a housing development. In the last case, the development would be a conversion of the forest stand and, if not designed properly, could also degrade the wetland). Habitat Conversion, Habitat Fragmentation, Hydrologic Alteration, Sedimentation, Pollution and Inadequate Distribution Of Successional Stages are closely related categories.	1, 7.3. Residential and Commercial Development; Other Ecosystem Modifications
C	Habitat Conversion	The complete transformation or loss of a habitat by human action (e.g., filling a wetland to create a grassy field, converting a forest into a parking lot, or damming a stream to create a reservoir). Habitat Alteration/Degradation, Habitat Fragmentation, Hydrologic Alteration, and Inadequate Distribution of Successional Stages are closely related categories.	1. Residential and Commercial Development
D	Habitat Fragmentation	The breaking up of habitats into smaller, non-contiguous patches as a result of habitat conversion (e.g., housing, commercial development, roads, utility lines). Fragmentation can: 1) render important habitats inaccessible (such as isolating a den site from a feeding site), 2) breakdown of the metapopulation structure of a species (for example grassland butterflies, spotted salamander, and tiger beetles); and, 3) degrade remaining habitat patches through edge effects that favor edge-tolerant species such as raccoons and crows, as well as invasive exotic species that can out-compete native and rare species. The result of habitat fragmentation is often increased predation, increased mortality, reduced mobility and changes in habitat micro-climates. Habitat Alteration/Degradation, Hydrologic Alteration and The Impacts off Roads and Trails are closely related categories.	1. Residential and Commercial Development
E	Hydrologic Alteration	Changes in the flow, periodicity or quality of a surface or subsurface water system (examples include a dam on a river preventing historic fluctuations in water level and mining activities causing a perennial seep to run dry). Dams can also increase water temperature. If warm water was identified as a problem then that problem would be placed in the pollution category. Hydrologic Alteration is a subset of Habitat Alteration but is a significant enough problem to warrant a separate category.	7.2 Dams and Water Management/Use

#	Threat/ Problem	Description	IUCN Category
F	Impacts of Roads & Transportation Systems	Transportation corridors that bring people, disturbance, and exotics to a habitat or directly impact a Species of Greatest Conservation Need (e.g., road kill, bird species whose mating calls are drowned out by the noise of road traffic, a road corridor that speeds the spread of an exotic invasive species. Habitat Fragmentation, habitat Alteration/Degradation and Impact of Roads are closely related categories. <i>In 2005 this category included trail impacts. For 2015 trail impact should be categories in Incompatible Recreation.</i>	4.
G	Impacts of Energy Infrastructure & Development	Threats from exploring for, developing, producing and distributing energy resources. Hydrofracturing and other natural gas extraction and distribution processes. Removal and distribution of minerals and rocks, limited to energy production. Generating and distributing power from the wind. Generating and distributing power from the sun.	
H	Inadequate Disturbance Regime	A disturbance regime is re-occurring process that disrupts a habitat, ecosystem, populations, and/or substrate causing significant change to a system (Picket and White). Many species have adapted to these disturbances and depend upon them to maintain habitats (e.g., the loss of beaver created wetlands, and a dam preventing the yearly flooding of floodplain forests that brings nutrients to the site and creates opening for early successional vegetation).	7.1. Fire and Fire Suppression (Other disturbance types need representation)
I	Invasion by Exotic Species	The introduction and spread of nuisance exotic and native species (plants and animals). These species may lead to the elimination of populations, threats to long-term stability or extirpation by out-competing a native species, displacing its food source or altering a key process or function of a habitat. Note that this category includes both exotic species and invasive native species such as cowbirds and sea lamprey. Exotic disease and parasites are addressed separately in the disease category.	8.1. Invasive Non-native/Alien Species/Diseases
J	Parcelization	Separating a large parcel into multiple smaller parcels. Parcelization is a significant driver of habitat fragmentation (and is often driven itself by tax policy). Parcelization can make it difficult to deliver management programs or present access issues which could impede actions benefiting SGCN even when fragmentation is not a problem (e.g., when a single 800-acre parcel is broken into many smaller lots some of the new landowners may choose to post their land while others may close logging roads).	
K	Sedimentation	Excessive inputs (in frequency and/or abundance) of solid material (inorganic or organic solid fragments) that are carried and deposited by wind, water, or ice to a water body. These materials have a negative impact on Species of Greatest Conservation Need through their physical presence (example: soils washing into a stream from a construction site and smothering fish eggs and other aquatic species that live in the spaces between rocks and gravel in a streambed). Sedimentation was broken out from the habitat alteration/degradation category because of its significant impact on aquatic species. Note: a problem that exerted a negative chemical impact on wildlife (e.g., road salt), would be listed in the Pollution category.	9.3.2. Ag/Forestry Soil Erosion and Sedimentation (Other sedimentation sources need representation)

#	Threat/ Problem	Description	IUCN Category
Non-Habitat-Related Threat Categories			
L	Competition	Where two or more species compete for the same limited resource (e.g., space, food) decreased survival, growth rate and/or reproduction of competing individuals is possible (e.g., fisher, bobcat and coyote have overlapping habitat needs and prey preference).	8.2. Problematic Native Species
M	Disease	Any disease causing agent such as, fungi, bacteria and viruses (e.g., rabies, West Nile disease, whirling disease, chronic wasting disease, hemlock wooly adelgid, and sudden oak death). Diseases are often transmitted by parasites, a related problem.	8.1, 8.2, 8.3?, 8.4-6. Invasive and Other Problematic Species, Genes and Diseases
N	Genetics	A reduction of survival or fecundity of a species due to inbreeding depression (the mating of close relatives) usually due to small and isolated populations, and outbreeding depression (the mating of different locally adapted populations). Examples include lake sturgeon and timber rattlesnake for whom inbreeding depression and genetic drift may be distinct possibilities due to small and isolated populations. Outbreeding depression can be a problem for native baitfish whose locally adapted genes may be swamped out by the accidental release of relatives taken from other waters in the state or elsewhere.	<i>Research needed</i>
O	Harvest or Collection	Legal or illegal taking of biological resources (e.g., hunting, trapping, collecting, fishing) for commercial, recreation, subsistence, research, or management purposes (e.g., the accidental taking of spruce grouse or American marten by a hunter or trapper, illegal collection of wood turtles for sale in the pet trade, and the harvesting of eels).	5.1, 5.2, 5.4. Hunting and Collecting Terrestrial Animals. Gathering Terrestrial Plants
P	Incompatible Recreation	Recreational activities (outside of established transport corridors) that directly impact SGCN or their habitats (e.g., include: poorly planned ORV or hiking trails that cause wildlife to abandon a den site, bird watchers getting too close to nesting loons causing loons to abandon their nest, and off-road vehicles operated outside of approved areas can run through vernal pools degrading the pool and crushing spawning amphibians and their eggs) Because of the scale of impact, the construction of a golf course or ski area would be listed in Habitat Conversion or Habitat Alteration categories respectively. Incompatible Recreation is also closely related to Impacts of Roads & Trails.	6.1. Recreational Activities
Q	Loss of Food Base or Prey Base	The disappearance of a food source important to a species' survival (e.g., lynx which feeds primarily on snowshoe hare and the whippoorwill whose primary diet of flying insects has been decreased).	<i>Research needed</i>
R	Loss of Relationship with Other Species	A species whose existence depends upon another for a process, function or product (examples include the larvae of many mussel species will attach to fish and depend on these fish for dispersal). Many insects, including butterflies and moths, have specific relationships with host plants that serve as its sole food. In some cases the host plant also conveys a chemical protection to the insect.) This category differs from the Loss of Prey Base category in that there are no alternatives (e.g., lynx and whippoorwill can take other prey, monarch butterflies won't persist without milkweed to feed on and to provide chemical protection from predators).	<i>Research needed</i>

#	Threat/ Problem	Description	IUCN Category
S	Parasitism	A relationship between two species in which one benefits (the parasite) and the other (the host) is harmed, (Smith 1980) although not directly killed (e.g., ticks, sea lamprey parasitizing lake whitefish and lake trout). Parasites may transmit diseases to the host therefore disease is a closely related problem. Examples include ticks transmitting Lyme disease and mosquitoes transmitting West Nile virus.	<i>Research needed</i>
T	Pollution	Introduction of exotic materials (other than sediments) from point and non-point sources. Includes chemicals and toxins in the air, land, and water; excess nutrients from farm and municipal sewage plants; garbage and other solid waste; radioactive materials; road salt; excessive noise; excessive heat; and light pollution that disturbs animals and disrupts migration patterns. Note: Sediments were broken out from this category because of its significant impact on some water bodies. Greenhouse gasses such as carbon dioxide and methane would be covered in the Climate Change category.	9. Pollution
U	Predation or Herbivory	Species or habitats negatively impacted by wildlife species that eat them (e.g., raccoons and skunks that eat wood turtle and spiny softshell turtle eggs and moose over-browsing vegetation and preventing the regeneration of a forest stand. This category differs from Competition in that competition is two or more species vying for the same specific resource, whereas predation is one species eating another.	<i>Research needed</i>
V	Reproductive Traits	Species whose specific reproductive strategies make it vulnerable, such as species producing very few offspring because they take a long time to reach sexual maturity and/or take a long time between reproductive events (e.g., lake sturgeon and wood turtle).	<i>Research needed</i>
W	Trampling & Direct Impacts	Non-recreational, and sometimes inadvertent, negative impacts to a species (examples include the crushing of wildlife by agricultural equipment operating in a farm field, vehicles operating off-road, the killing of rattlesnakes or bats out of antipathy for the species, increased nest abandonment by brown thrashers due to the proximity of people). Impacts to a habitat would be assigned to the Habitat Degradation/Alteration and perhaps Habitat Conversion. Incompatible Recreation and Impacts of Roads & Trails are closely related categories	5.1.3, 5.2.3, 5.4.3. Persecution/Control

Conservation Actions for Wildlife Action Plan Revision

Element number four of the eight congressionally required elements of a Wildlife Action Plan requires that states describe “conservation actions proposed to conserve the identified species and habitats and priorities for implementing such actions.”

We identified actions to address the threats and problems impacting each of Species of Greatest Conservation Need (SGCN) and their habitats. Selected actions are based on the best science available today as well as a strategic assessment of needs and priorities of all wildlife species. In the coming years, as monitoring data on SGCN and conservation actions becomes available, as priorities change, or new threats or opportunities arise, actions may need to be revisited. Not every action in this report will be eligible for State Wildlife Grant funding. Furthermore, it may not be suitable, or feasible, for the Vermont Fish & Wildlife Department to implement some of the actions in this report, however, some conservation partners may find them fitting and practical.

Actions are described in short narratives in each SGCN conservation summary (Appendix A) and in each habitat, community and landscape summary (Appendix B). Actions are intentionally broad, directional, and nonspecific so as not to constrain our selection of procedures for implementing them. For example, an action such as “provide technical assistance to landowners to maintain or improve riparian habitat for Species of Greatest Conservation Need” allows for different approaches to providing that assistance and leaves the door open to a variety of providers to implement. Where action implementation is to be funded by the State Wildlife Grant program the approach should be consistent with the Department’s mission and strategic plan, and precise procedures will be detailed in operational plans once the Action Plan is finalized.

Vermont’s Action Plan was designed for the state, not just the Fish & Wildlife Department. While the VFWD may be responsible for implementing many of the actions in this report, it could be conservation partners that are the more logical and appropriate leaders for others, due to their skills and expertise, staffing, history, location, available resources and constituencies.

Each of the actions identified in this report were assigned to one of 27 categories in six major classes. The categories were developed by the Conservation Measures Partnership (Salafsky 2005) as a means of standardizing terminology (not practices) among conservation practitioners worldwide. Many states have used these same categories to organize the strategies and actions in their Action Plan. They have also been incorporated into Wildlife TRACS (Tracking and Reporting Actions for the Conservation of Species) the US Fish & Wildlife Services’ system for tracking and reporting conservation activities. States, including Vermont, will use TRACS for all work funded through the USFWS once it is fully operational.

The action categories are used solely for the purpose of organizing and grouping strategies developed by Action Plan teams and committees. It was not our goal to create strategies for every category. A few categories were not applicable to the species or habitats in Vermont whereas others were deemed not as effective. Table C.2 lists the action categories used in this Wildlife Action Plan.

Table C.2 Conservation Actions for Wildlife Action Plan Revision

#	Strategy Type	Description
1	Land/Water Protection	
1.1	Creating Publicly-Owned Protected Areas	Setting up or expanding public parks, forests and other protected areas where wildlife conservation is a primary management objective (e.g., wildlife management areas, state forests, municipal lands)
1.2	Creating Privately-Owned Protected Areas	Setting up or expanding private reserves and other protected areas where wildlife conservation is a primary management objective (e.g., private lands managed by non-profit conservation or hunting groups)
1.3	Easements	Setting up protection of some specific aspect of a resource on public or private lands (e.g., development rights, wild & scenic river designation)
2	Land/Water/Species Management	
2.1	Protected Area Management	Generally managing protected areas where wildlife conservation is a primary management objective (e.g., site design, training park staff, managing water levels).
2.2	Compatible Resource Use	Promoting use of resource lands (where wildlife management is not a primary objective) to be compatible with conservation (e.g., promoting sustainable logging, grazing, fishing, hunting, farming, aquaculture, energy development, transportation infrastructure).
2.3	Invasive Species Control & Prevention	Dealing with invasive and/or alien plants, animals, and pathogens (e.g., developing boat wash stations, pulling noxious weeds from a habitat).
2.4	Habitat Restoration	Enhancing degraded or restoring missing habitats (e.g., clayplain forest restoration and riparian tree plantings). If a strategy specifically targets one or two species we consider it a species restoration strategy. If it specifically targets three or more species we consider it a habitat restoration strategy.
2.5	Natural Processes Restoration	Enhancing or restoring natural ecosystem functions (e.g., prescribed burns, dam removal and restoration of historic flow regimes, fish ladders).
2.6	Species Restoration	Enhancing or restoring specific plant and animal populations (e.g., translocating spruce grouse from Canada to Vermont, and erecting artificial nesting boxes/platforms for bluebirds and osprey.) If a strategy specifically targets one or two species we consider it a species restoration strategy. If it specifically targets three or more species we consider it a habitat restoration strategy.
2.7	Ex-Situ Conservation	Protecting wildlife out of its native habitats (e.g., captive breeding of bald eagles and creating regional refugia for New England cottontail rabbit).
3	Law & Policy	
3.1	Legislation	Making, changing, influencing, or providing input into formal legislation at all levels: international, national, state/provincial, local (e.g., lobbying to make the SWG program permanent, addressing greenhouse gas emissions).
3.2	Policy & Regulations	Making, changing, influencing, or providing input into policies and regulations affecting the implementation of laws at all levels: international, national, state/provincial, local (e.g., providing data to policy makers, development of wildlife harvest regulations).
3.3	Planning & Zoning	Developing, changing, influencing, or providing input into plans governing natural resource use and allocation (e.g., municipal zoning, public or private management plans for ecoregions, sites, habitats, or species, commenting on zoning plans, developing a town ordinance).

#	Strategy Type	Description
3.4	Standards	Setting, changing, influencing, or providing input into voluntary standards that govern practices (e.g., best management practices for forestry, habitat guidelines for state lands).
3.5	Compliance & Enforcement	Monitoring and enforcing compliance with laws, policies & regulations, plans, and standards (e.g., water quality standard monitoring, enforcement of ATV regulations).
4	Research, Education & Awareness	
4.1	Research	Conducting and disseminating research to improve knowledge about conservation issues (e.g., conducting inventories for SGCN, developing habitat maps, demonstration projects for sustainable forestry)
4.2	Technical Assistance, Training, Workshops	Enhancing knowledge, skills and information exchange for practitioners, landowners, stakeholders, and other relevant individuals in structured settings outside of degree programs (e.g., providing technical assistance to landowners, monitoring workshops, conferences, learning networks)
4.3	Lifelong Learning	Enhancing knowledge and skills of practitioners, stakeholders, and other relevant individuals through non-structured means (e.g., writing a how-to manual for landowners or towns and communities, stakeholder education on proper ORV use)
4.4	Awareness Raising, Communications	Raising conservation awareness and providing information through various media (e.g., websites, newsletters, presentations)
5	Economic & Other Incentives	
5.1	Linked Enterprises	Developing enterprises that directly depend on natural resources as a means of influencing behaviors, attitudes, and beliefs (e.g., ecotourism, sustainable forest product harvesting).
5.2	Substitution	Promoting products and services that substitute for environmentally damaging ones (e.g., floodplain restoration in place of dams and bank reinforcement, promoting recycling/use of recycled materials)
5.3	Financial Incentives & Market Forces	Using market mechanisms to influence behaviors, attitudes, and beliefs (e.g., forestry certification, positive incentives, negative incentives, forest banking, valuation of ecosystem services such as flood control)
5.4	Conservation Payments	Using direct or indirect payments to influence or reinforce behaviors, attitudes, and beliefs (e.g., landowner payment programs).
5.5	Non-Monetary Values	Using non-market forces to change behaviors, attitudes, and beliefs (e.g., landowner/land manager recognition awards).
6	Capacity Building	
6.1	Institutional Development	Creating or providing non-financial support, capacity building for non-profits, government agencies, and for-profits (e.g., creating new local land trusts)
6.2	Alliance Development	Forming and facilitating partnerships, alliances, and networks of organizations (e.g., Action Plan Conservation Partners, Vermont Monitoring Cooperative, Vermont Sportsmen's Federation).
6.3	Conservation Finance	Raising and providing funds for conservation work (e.g., State Wildlife Grants small grants program, private foundations, debt-for-nature swaps).

Vermont's Wildlife Action Plan

Vermont Species & Habitat Climate Vulnerability Assessment

Appendix D

Draft 10/1/2015

Introduction

During 2013 a group of Vermont Fish & Wildlife Department biologists and Chris Hilke, Climate Adaptation Program Manager for the Northeast Regional Office of National Wildlife Federation, collaborated to assess the vulnerability of fish, wildlife, plants and their habitats to climate changes. As part of this Vermont Species & Habitat Climate Vulnerability Assessment, the team investigated 18 key species, 20 upland habitats, 11 wetland habitats, and 13 freshwater habitats as shown the tables on the following pages. One Department biologist used a NatureServe tool to add in 13 more butterfly species. Species included Species of Greatest Conservation Need and important “surrogate” species that are widely considered representative of particular habitat types. Species assessments culminated in an overall vulnerability rating for climate-specific and non-climate stressors and an associated confidence score (see Rating Key at bottom of spreadsheet for details). Climate change vulnerability assessments were similarly conducted for Vermont's upland and wetland natural communities. For efficiency, the 95 natural community types were grouped into categories based on the environmental factors that drive their development and that could affect their susceptibility to climate change. Some natural community types were assessed individually and freshwater habitats were also assessed.

The most important lesson taken from this exercise is that species (and habitat) responses to climate change will not be uniform. For some, climate change may not be a significant threat, however if a species is already subjected to other stresses, climate change impacts may push that species over the edge. This is an important consideration to take into account. More information on climate change and wildlife conservation can be found in chapter 3: Climate Change & Conservation.

Vermont Species & Habitat Climate					
	H	I	J	K	
Species and Habitats	Thermal Stress	Toxic Substances- Pollution	Hanitat Fragmentation	Inventory Need	Notes
Key Species (Bold text =SGCN)					
Jefferson salamander					
Northern white cedar					Competition effects this species broadly
Fingernail clam	X	X			
Beaver			X		Keystone wetland builder - vulnerable via direct take
Bobcat					Coyote and fisher competition
Canada Lynx			X		trapping in Canada impacts Lynx in VT
Brook Trout	X	X	X		Rating is consistent w/ other states. Water temp is key, sedimentation also key issue
Wood Turtle			X		Other threats - Pet trade, agriculture and egg predation
Pearlshell mussel		X	X		
Fallfish			X		Risk by gravel scour when eggs present (May) 63F
Smelt					
Lake trout					Sea lamprey. Supplemental stocking
Bald Eagle		X			
Bicknell's Thrush					Red squirrel predation tied to cone production, limtited by patch size
Common Loon		X			direct impacts from fishing gear (hooks and lead)
Red Oak			X		A southern oak species so favored with CC, deer browse impacts seedling development
Sugar Maple					Likely climate loser over 100yr. Managed in sugar bushes which promotes over others
West Virginia White	X		X		
Bog Copper					Climate vulnerability and confidence score only performed for this butterfly species
Edward's Hairstreak					Climate vulnerability and confidence score only performed for this butterfly species
Early Hairstreak					Climate vulnerability and confidence score only performed for this butterfly species
Hackberry Emperor					Climate vulnerability and confidence score only performed for this butterfly species
Tawny Emperor					Climate vulnerability and confidence score only performed for this butterfly species
Jutta Artic					Climate vulnerability and confidence score only performed for this butterfly species
Cobweb Skipper					Climate vulnerability and confidence score only performed for this butterfly species
Mulberry Wing					Climate vulnerability and confidence score only performed for this butterfly species
Broad-winged Skipper					Climate vulnerability and confidence score only performed for this butterfly species
Black Dash					Climate vulnerability and confidence score only performed for this butterfly species
Dion Skipper					Climate vulnerability and confidence score only performed for this butterfly species
Two-spotted Skipper					Climate vulnerability and confidence score only performed for this butterfly species
Dusted Skipper					Climate vulnerability and confidence score only performed for this butterfly species

Vermont Species & Habitat Climate Vulnerability Assessment										Exposures & Key Climate												
Species and Habitats	Vulnerability			Sensitivity Factors						A	B	C	D	E	F	G	H	I	J	K	L	
	Climate Vulnerability Rating	Confidence Score	Vulnerability to non-climate stressors	Habitat specificity	Edge of range	Enviro or physiological tolerance	Interspecific or phenological dependence	Mobility	Exotic Pathogens or Invasive Species	Annual temperature - Increase	Seasonal temperature - Increase	# Hot days - More	# Cold days - Fewer	Variability - Increase	Annual precipitation - Increase	Seasonal precipitation - Variable	Heavy rainfall events - Increase	Soil moisture - Decrease	Snow - Decrease	Spring flows - Earlier	Summer low flow - Longers	
Wetland Habitats																						
Cattail Marsh	L	M	L								X				X	X						
Shallow Emergent Marsh	M	M	M								X				X	X		X				
Marsh and Sedge Meadow (Formation)	M	M	M								X				X	X					X	
Alluvial Shrub Swamp	M	M	L								X					X		X				
Basin swamps & wetlands	H	M	M								X	X					X	X			X	
Floodplains	H	M	M								X		X			X	X	X		X		
Ground water seepage & Flood swamp	L	M	L								X		X			X		X		X	X	
Open peatlands (precip-dependent)	M	M	L							X	X	X		X		X		X	X			
Open peatlands (ground-fed)	L	M	M								X	X				X		X				
Floodplain Forests	H	M	M								X		X			X	X	X		X		
Wet Shores	H	M	M								X	X					X	X			X	
Upland Habitats																						
Alpine Meadow	E	M	M							X	X	X	X		X	X		X	X			
Spruce-Fir-Northern Hardwood	M	M	M							X	X	X	X		X	X		X	X			
Northern Hardwood Forest	L	M	L								X	X	X	X		X		X				
Oak-Pine - Dry Mesic Forests & Woodlands with deeper soils	L	M	M								X	X				X	X					
Rocky Forests and Woodlands	L	M	L								X	X				X		X				
Oak-Pine Southern Rocky - Southern Dry Rocky Forests and Woodlands	M	L	L								X	X				X		X				
Outcrops and upland meadows	N	L	L								X	X				X		X				
Cliffs and Talus	L	M	L								X	X				X		X				
Upland shores	M	M	H								X	X						X		X		
Subalpine Krummholz	E	M								X	X	X		X	X			X	X			
Montane Spruce-fir	M	M								X	X	X		X	X			X	X			
Red Spruce-Heath Rocky Ridge	M	M								X	X	X		X	X			X	X			
Montane Yellow Birch-Red Spruce Forest	M	M								X	X	X		X	X			X	X			
Red Spruce-Northern Hardwood	L	M								X	X	X		X	X			X	X			
Lowland Spruce-Fir Forest	M	M								X	X	X		X	X			X	X			
Boreal Talus Woodland	M	M								X	X	X		X	X			X	X			
Cold-Air Talus Woodland	H	M								X	X	X		X	X			X	X			
Limestone Bluff Cedar-Pine Forest	L	M									X	X				X	X	X				
Transition Hardwood Talus Woodland	L	M									X	X				X	X					
Dry Oak Woodland	H	L									X	X				X	X					

Vermont Species & Habitat Climate	Change Factors														Non-Climate Stressors						
	M	N	O	P	Q	R	S	T	U	V	W				A	B	C	D	E	F	G
Species and Habitats	Ice dynamics - Changing	Fluctuating lake levels - Increase	Lake stratification	Flood events - Increase	# of short-term droughts - Increase	Storms - Increase	Fire	Growing season - Longer	Onset of spring - Earlier	Onset of fall - Later	Biological interactions	Other	List of exposures with the greatest Negative	List of exposures that might be beneficial	Acidity	Habitat Alteration- Altered Hydrology	Invasive Species	Channel Erosion- Sedimentation	Encroachment	Land Erosion	Nutrient Loading
Wetland Habitats																					
Cattail Marsh		X									X		N	T, U, V		X	X		X		
Shallow Emergent Marsh		X		X	X						X		I	T		X	X		X		
Marsh and Sedge Meadow (Formation)		X		X	X						X					X	X		X		
Alluvial Shrub Swamp					X						X	X	I, S, C	U, T		X			X		
Basin swamps & wetlands	X	X		X				X					M, P			X	X		X		
Floodplains	X	X		X		X		X	X	X	X	K					X	X	X		
Ground water seepage & Flood swamp					X	X		X			X	X		X		X	X				
Open peatlands (precip-dependent)					X	X	X				X		G, Q								
Open peatlands (ground-fed)		X			X						X		G, I, Q			X					
Floodplain Forests	X	X		X		X		X	X	X	X		K, P, N,				X	X	X		
Wet Shores	X	X		X				X	X	X	X		M, P, W			X	X	X	X		
Upland Habitats																					
Alpine Meadow					X	X		X	X		X		B, C, J			X	X				
Spruce-Fir-Northern Hardwood					X	X		X	X				D, I, J, T	F	X				X		
Northern Hardwood Forest					X	X			X		X		B, C, D, I, Q, W				X				
Oak-Pine - Dry Mesic Forests & Woodlands with deeper soils					X	X	X						Q, Z			X	X				
Rocky Forests and Woodlands					X	X	X						Q, Z		X		X				
Oak-Pine Southern Rocky - Southern Dry Rocky Forests and Woodlands					X	X	X						C, Q				X				
Outcrops and upland meadows					X							X	B, C, J			X			X		
Cliffs and Talus					X	X							C, I						X		
Upland shores	X			X	X						X		M, N, P			X	X				
Subalpine Krummholz					X						X	X	I, A	F							
Montane Spruce-fir					X				X			X	I, A	F							
Red Spruce-Heath Rocky Ridge					X	X	X		X			X	I, A, S	F							
Montane Yellow Birch-Red Spruce Forest					X	X			X	X		X	I, A	F							
Red Spruce-Northern Hardwood					X	X	X	X	X	X		X	I, A	F							
Lowland Spruce-Fir Forest				X	X	X	X		X			X	I, A	F							
Boreal Talus Woodland					X				X	X		X	I, A	F							
Cold-Air Talus Woodland					X	X		X	X	X	X	X	I, A	F							
Limestone Bluff Cedar-Pine Forest					X	X	X				X		Q, W				X				
Transition Hardwood Talus Woodland					X	X	X				X		Q, W				X				
Dry Oak Woodland					X	X	X						Q, C				X				

Vermont Species & Habitat Climate					
	H	I	J	K	
Species and Habitats	Thermal Stress	Toxic Substances- Pollution	Hanitat Fragmentation	Inventory Need	Notes
Wetland Habitats					
Cattail Marsh		X	X		invasives - Phragmites
Shallow Emergent Marsh		X	X		invasives
Marsh and Sedge Meadow (Formation)		X	X		invasives
Alluvial Shrub Swamp		X	X		
Basin swamps & wetlands		X	X		Variability of water level could impact dragonfly transformation.
Floodplains		X	X		Elm was once an important component to floodplains
Ground water seepage & Flood swamp			X		Most vulnerable species include Black ash (AB) & Hemlock (WA)
Open peatlands (precip-dependent)					Black spruce and Sphagnum - Impacts of drying
Open peatlands (ground-fed)					Sphagnum is most vulnerable species
Floodplain Forests		X	X		Boxelder is non-native and likely to expand
Wet Shores					
Upland Habitats					
Alpine Meadow					More krumholz?
Spruce-Fir-Northern Hardwood			X		Most vulnerable species include Spruce & Fir
Northern Hardwood Forest			X		Most vul is Hemlock. Oaks & hickory will benefit
Oak-Pine - Dry Mesic Forests & Woodlands with deeper soils			X		Hemlock (HWA) will decrease; oaks and hickories will increase
Rocky Forests and Woodlands			X		Possible increase in red pine; possible decrease in N. white cedar due to drought and temp increase
Oak-Pine Southern Rocky - Southern Dry Rocky Forests and Woodlands			X		increase in oaks, hickories, white pine, and pitch pine
Outcrops and upland meadows					All woody species are likely to decrease in abundance due to heat and drought stress.
Cliffs and Talus					
Upland shores					Encroachment of woody species if there is less ice-scour and flooding.
Subalpine Krummholz					Less spruce and fir - area compressed
Montane Spruce-fir					Less spruce and fir, more hardwood
Red Spruce-Heath Rocky Ridge					Less spruce and fir, more hardwood
Montane Yellow Birch-Red Spruce Forest					Less spruce and fir, more hardwood
Red Spruce-Northern Hardwood					Less spruce and fir, more hardwood
Lowland Spruce-Fir Forest					Less spruce and fir, more hardwood
Boreal Talus Woodland					Less spruce and fir, more hardwood
Cold-Air Talus Woodland					Less spruce and fir, more hardwood
Limestone Bluff Cedar-Pine Forest					
Transition Hardwood Talus Woodland					less cedar more red pine
Dry Oak Woodland					< hemlock, more pitch pine if more fires, more hickory if not

Vermont Species & Habitat Climate Vulnerability Assessment										Exposures & Key Climate											
Species and Habitats	Vulnerability			Sensitivity Factors						A	B	C	D	E	F	G	H	I	J	K	L
	Climate Vulnerability Rating	Confidence Score	Vulnerability to non-climate stressors	Habitat specificity	Edge of range	Enviro or physiological tolerance	Interspecific or phenological dependence	Mobility	Exotic Pathogens or Invasive Species	Annual temperature - Increase	Seasonal temperature - Increase	# Hot days - More	# Cold days - Fewer	Variability - Increase	Annual precipitation - Increase	Seasonal precipitation - Variable	Heavy rainfall events - Increase	Soil moisture - Decrease	Snow - Decrease	Spring flows - Earlier	Summer low flow - Longers
Freshwater Habitats																					
Medium-sized river (4-6 order)	M	M								X	X	X					X				X
Large river (7+ stream order)	L	M									X	X					X				X
High gradient, coldwater acidic, 1-2 order	H	H								X	X	X	X	X		X	X		X	X	X
High gradient, coldwater, not acidic, 1-2 order	H	H								X	X	X		X		X	X			X	X
Low gradient marsh	M	M								X	X	X	X			X	X				X
Lake Champlain valley	M	M								X	X	X		X			X			X	X
High Elevation Lake	M	L	L								X	X		X						X	
Dystrophic Lake	M	L	L								X	X		X							
Lake - Oligotrophic, Stratified	L	H	L							X	X	X	X								
Mesotrophic-Eutrophic Lake (stratified)	M	M	M							X	X	X	X			X			X		
Mesotrophic-Eutrophic Lake (unstratified)	H	M	M							X	X	X	X	X			X		X		
Unstratified lakes	M	H	H							X	X	X	X	X	X	X	X		X	X	X
Stratified Lakes	M	H	H							X	X	X	X	X	X	X	X		X	X	X

Vermont Species & Habitat Climate	Change Factors														Non-Climate Stressors						
	M	N	O	P	Q	R	S	T	U	V	W				A	B	C	D	E	F	G
Species and Habitats	Ice dynamics - Changing	Fluctuating lake levels - Increase	Lake stratification	Flood events - Increase	# of short-term droughts - Increase	Storms - Increase	Fire	Growing season - Longer	Onset of spring - Earlier	Onset of fall - Later	Biological interactions	Other	List of exposures with the greatest Negative	List of exposures that might be beneficial	Acidity	Habitat Alteration- Altered Hydrology	Invasive Species	Channel Erosion- Sedimentation	Encroachment	Land Erosion	Nutrient Loading
Freshwater Habitats																					
Medium-sized river (4-6 order)				X	X								L, C, P	F							
Large river (7+ stream order)				X	X								L, C, P	F, T							
High gradient, coldwater acidic, 1-2 order				X	X								F, H								
High gradient, coldwater, not acidic, 1-2 order				X	X	X							C, H, L	F, H							
Low gradient marsh				X	X								C, L	F							
Lake Champlain valley				X	X								C, L	F							
High Elevation Lake		X			X									B, M	X	X					
Dystrophic Lake		X		X	X							X	P, Q	B, M	X	X					
Lake - Oligotrophic, Stratified	X	X	X	X	X							X	A, B, C, D						X	X	X
Mesotrophic-Eutrophic Lake (stratified)	X	X	X	X	X	X		X			X		A, B, C, D, N				X		X	X	X
Mesotrophic-Eutrophic Lake (unstratified)	X	X		X	X	X		X			X		B, C, N	X			X		X	X	X
Unstratified lakes	X	X	X	X	X	X		X	X	X	X						X	X	X		X
Stratified Lakes	X	X	X	X		X											X	X	X		

Vermont Species & Habitat Climate					
	H	I	J	K	
Species and Habitats	Thermal Stress	Toxic Substances- Pollution	Hanitat Fragmentation	Inventory Need	Notes
Freshwater Habitats					
Medium-sized river (4-6 order)					Compositional changes may include loss of pearlshell
Large river (7+ stream order)					Compositional changes include an increase in the number of warm water species
High gradient, coldwater acidic, 1-2 order					Potential loss of coldwater stenotherms
High gradient, coldwater, not acidic, 1-2 order					Potential loss of coldwater stenotherms
Low gradient marsh					Potential loss of coldwater stenotherms
Lake Champlain valley					
High Elevation Lake					Most vulnerable species include brook trout
Dystrophic Lake					Most vulnerable species include brook trout
Lake - Oligotrophic, Stratified		X			Most vulnerable species include Lake trout & Round Whitefish
Mesotrophic-Eutrophic Lake (stratified)		X			Lake trout, smelt & pike will be most vulnerable
Mesotrophic-Eutrophic Lake (unstratified)		X			
Unstratified lakes					
Stratified Lakes					

Vulnerability Rating Key		
Code		Description
E	Extremely Vulnerable	Abundance and/or range extent in Vermont extremely likely to substantially decrease (>75% loss) or disappear by 2050
H	Highly Vulnerable	Abundance and/or range extent in Vermont likely to decrease significantly (25-75% loss) by 2050
M	Moderately Vulnerable	Abundance and/or range extent in Vermont likely to decrease (10-25% loss) by 2050
L	Slightly Vulnerable	Available evidence does not suggest that abundance and/or range extent in Vermont will change (decrease, 5 - 10% loss) by 2050
N	Not Vulnerable, No Effect	Abundance and/or range extent in Vermont likely to increase or decrease by less than 5% by 2050
B	Increase Possible or Likely	Available evidence suggests that abundance and/or range extent in Vermont is likely to increase (>15% increase) by 2050
U	Unknown/Uncertain	Available evidence not available or not conclusive at this time

Confidence Ratings		
Code		Description
L	Low	Not very confident (0-30% certainty in vulnerability score)
M	Moderate	Somewhat confident (30-60% certainty in vulnerability score)
H	High	Very confident (>60% certainty in vulnerability score)

Non-Climate Stressors	
Code	Description
A	Acidity/Pollution
B	Habitat Alteration/Altered Hydrology
C	Invasive Species
D	Channel Erosion/Sedimentation
E	Encroachments
F	Land Erosion
G	Nutrient loading
H	Thermal Stress
I	Toxic Substances/Pollution
J	Habitat Fragmentation
Enter as free text	Other

Exposures & Key Climate Change Factors				
	Code	Parameter	Trend	Projections (range = low to high emissions scenario)
Temperature	A	Annual temperature	increase	by 2050, projected increase 3.7 to 5.8°F; by 2100, 5.0 to 9.5°F
	B	Seasonal temperature	increase	by 2050, projected increase in winter (DJF) 4.3 to 6.1°F; summer (JJA) 3.8 to 6.4°F
	C	# Hot days	more	more frequent and more intense; by end of century, northern cities can expect 30-60+ days of temperatures >90°F
	D	# Cold days	fewer	reduction in days with cold (<0° F) temperatures
	E	Variability	increase	greater variability (more ups and downs)
Hydrology	F	Annual precipitation	increase	by end of century, projected total increase of 10% (about 4 inches per year)
	G	Seasonal precipitation	variable	more winter rain, less snow; by 2050, winter precipitation could increase by 11 to 16% on average; little change expected in summer, but projections are highly variable
	H	Heavy rainfall events	increase	more frequent and intense
	I	Soil moisture	decrease	reduction in soil moisture and increase in evaporation rates in the summer
	J	Snow	decrease	fewer days with snow cover (by end of century could lose 1/4 to 1/2+ of snow-covered days; increased snow density
	K	Spring flows	earlier	earlier snowmelt, earlier high spring flows; could occur 10 days to >2 weeks earlier
	L	Summer low flows	longer	extended summer low-flow periods; could increase by nearly a month under high emissions scenario
	M	Ice dynamics	changing	less ice cover, reduced ice thickness
	N	Fluctuating lake levels	increase	greater variability, greater amount of change in lake levels
	O	Lake stratification		some lakes may stratify earlier
Extreme events	P	Flood events	increase	more likely, particularly in winter and particularly under the high emissions scenario
	Q	# of short-term droughts	increase	by end of century, under high emissions scenario, short terms droughts could occur as much as once per year in some places
	R	Storms	increase	more frequent and intense (ice, wind, etc.)
	S	Fire		more likely
Phenology	T	Growing season	longer	by end of century, projected to be 4 to 6 weeks longer
	U	Onset of spring	earlier	by end of century, could be 1 to almost 3 weeks earlier
	V	Onset of fall	later	by end of century, could arrive 2 to 3 weeks later
	W	Biological interactions		could potentially be disrupted

Surveys, Monitoring, and Data Systems.....	3	Bobcat Movement and Habitat Use.....	13
Invertebrates	3	Habitat Blocks and Critical Crossings.....	13
Vermont Butterfly Atlas.....	3	Birds	14
Damselfly and Dragonfly Survey	3	Bicknell's Thrush in Northeastern Vermont	14
Bumble Bee Survey	4	Black Throated Blue Warbler Abundance and Nesting	
Freshwater Mussels	4	Productivity	14
Vermont Invertebrate Database	4	Landscape Conservation	14
Fish	5	Shoreline & Development Surveys for VT Lakes	14
Fish Surveys	5	Species Recovery	14
Fish Historical Record Scanning	5	Invertebrate Recovery	15
Amphibians and Reptiles	5	Bat Recovery	15
Town Surveys for Reptiles and Amphibians	5	Bird Recovery	15
Rare Amphibian Surveys.....	6	Other Recovery Projects.....	15
Rare Snake Surveys	6	Policy, Planning, and Land Acquisition	16
Turtle Conservation, Planning, & Management.....	6	Vermont Conservation Design	16
Vernal Pool Mapping.....	7	Getting the Lead Out.....	16
Mammals.....	7	Helping Landowners Protect Ecologically Sensitive	
Rare Carnivores	7	Lands	16
Bat Conservation and Management.....	7	Land Acquisition	16
Small Mammals of Vermont Atlas.....	8	Technical Assistance and Public Awareness	17
Birds.....	8	Community Wildlife Program	17
Breeding Bird Atlas.....	8	Helping Private Landowners Help Wildlife.....	17
Threats to Vermont's Mountain Birds.....	8	Public Opinion Survey	18
Plants and Natural Communities.....	8	Species of Greatest Conservation Need.....	18
Rare, Threatened, and Endangered Plant Monitoring.	8	Orange County Headwaters Project.....	18
Natural Community Inventories	9	Baitfish Workshop	18
Invasive Species	9	The Fishes of Vermont.....	19
Herbarium Records.....	9	Bats On Private Land.....	19
Revision of Natural Community Ranking		Natural Communities and the Plant Stewardship	
Specifications.....	9	Project	19
General Natural Heritage	10	Management.....	19
Natural Heritage Inventory	10	Aquatic Organism Passage.....	19
Structured Decision-making.....	10	Prevention of Fish Disease	20
Wildlife Management Area Inventories	10	Amphibian Crossings	20
Research	10	Beaver Baffles	20
Fishes	10	Grassland Birds At Airports	20
Lake Whitefish	10	Golden-winged Warbler	21
Stream Temperature	11	Double-crested Cormorant Control.....	21
Stream Morphology	11	Partnerships with Natural Resources Conservation	
Fish Health & Predator-Prey Dynamics	11	Service	21
Landlocked Salmon Stocking.....	11	Regional Partnerships	22
Atlantic Salmon Genetics	12	Regional Conservation Needs Grant Program	22
Lake Trout Genetics.....	12	The Staying Connected Initiative.....	22
Mammals.....	13	For the Future	22
Indiana Bat.....	13		

Taking Action: Implementing Vermont's First Wildlife Action Plan 2005-2015

A Decade of Fish and Wildlife Conservation

When the State Wildlife Grants (SWG) program began in 2001, the program came with a mandate: In order for states to continue to receive funding after 2005, they must adopt a strategic plan for prioritizing the conservation of the state's fish and wildlife with a focus on Species of Greatest Conservation Need (SGCN)—wildlife with declining populations, those that are threatened or potentially threatened, or those about which we knew so little that we had no means to assess conservation status.

The 2005 Wildlife Action Plan identified 143 vertebrates, 191 invertebrates, and 577 plants as SGCN. In order to provide guidance for the protection of these species, the plan also identified problems faced by these SGCN. While diverse, the most common of these problems included:

- Information needs and data gaps
- Loss of habitat
- Impacts of roads and trails
- Pollution and sedimentation
- Invasive species
- Climate change

The plan went on to recommend strategies for aiding SGCN to survive and thrive in the face of the above problems.

Now it is 2015, and a decade has passed since the Wildlife Action Plan was adopted. Vermont has made great strides to implement its Action Plan, and this document attempts to capture some of the success Vermont has had in accomplishing the plan's goals.

State Wildlife Grant has been a major source of funding for the projects below, though certainly not the only source. If not otherwise noted, it can be assumed that these projects were completed using SWG funds.

Surveys, Monitoring, and Data Systems

If we don't know what we have, we don't know what to conserve. While our knowledge of species has improved substantially over time, the 2005 Wildlife Action Plan identified numerous information gaps that have prevented us from creating conservation plans or implementing management strategies for many groups of wildlife. We didn't know what threats some species face or have a good grasp of their population dynamics. Over the past decade, Vermont has worked to fill many of these holes through surveys of species abundance and distribution. In many cases, trained volunteers aided in data collection efforts, amplifying the quantity of data collected and providing opportunities for public awareness. Updated data management systems also helped us to compile information, much of which can now be accessed by planners, government entities, conservation organizations and, in some cases, interested citizens.

Invertebrates

Vermont Butterfly Atlas

The Vermont Butterfly Atlas (2002-2007) surveyed the entire state and analyzed historic records and collections to document the distribution of 103 butterfly species, including 12 species new to Vermont as well as giant silkworm moths. The project was collaboration between VFWD, the Vermont Center for Ecostudies, and the Vermont Institute of Natural Sciences with more than 200 citizen scientists participating. Assessing the conservation status of each species and establishing a baseline for understanding future changes was the principle goal of the Atlas. Fifteen species were listed as Species of Greatest Conservation Need representing three ecological groups—wetlands, grasslands, and hardwood forests. We also calculated the vulnerability to climate change for 14 butterfly SGCN currently found in Vermont. Three species were found to be extremely vulnerable to climate change, five were highly vulnerable, one moderately vulnerable, three presumed stable and two could likely increase in numbers in Vermont. We hope to see the Butterfly Atlas repeated in ~2027 as a comparison to this baseline. More information, including *The Vermont Butterfly Survey, 2002 - 2007: A Final Report*, can be found at <http://www.vermontbutterflysurvey.org/>.

Damselfly and Dragonfly Survey

From 2007-2009 the first statewide field surveys, focused on peatland and large river habitat, were conducted to detect the presence of dragonflies and damselflies (collectively known as odonates) including 27 Species of Greatest Conservation Need. The survey data when compiled with existing records resulted in a total count of 142 odonate species in the state—98 dragonflies and 44 damselflies. This effort provided vital species distribution and occurrence information which has broadened our understanding of rare habitat-specialist dragonfly and damselfly SGCN. Habitat data collected as part of the study provides a comparative baseline for future population trend monitoring. Among the investigation's results were: new encounters with *Gomphus abbreviatus* (very rare to rare) on two rivers and a moderate gain in the knowledge of *Ophiogomphus* spp. (four species), particularly on the White River; discovery of at least two previously unknown populations of *Neurocordulia yamaskinensis* (uncommon), a species that had been rarely encountered in Vermont; an expanded knowledge of *Somatochlora* spp. (seven species) distribution in and around peatlands, including *S. albicincta* (very rare), *S. cingulata* (very rare to rare) and *S. franklini* (very rare to rare), revealing that Silvio Conte National Wildlife Refuge and West Mountain Wildlife Management Area have some of the highest *Somatochlora* diversity in New England; and two new peatland sites for *Williamsonia fletcheri* (very rare to rare). Future efforts toward odonate SGCN conservation will continue to rely on the information resulting from this and future field studies.

Bumble Bee Survey

When public attention was brought to bumble bee declines in the last decade, the Vermont Center for Ecostudies worked with the Vermont Fish & Wildlife Department to document distribution and determine the status of bumble bee species in the state. With the help of citizen scientists a two-year survey amassed over 10,000 current bumblebee records from more than 1,500 locations, which were then compared to historic data. Twelve of the 17 species previously reported from Vermont were collected during this period. Three species appear to be more widespread than in the past, as well as the carpenter bee, which seems to have extended its range northward. This project enabled the Vermont Fish & Wildlife Department to assess the Natural Heritage conservation status of all of Vermont's bumble bee species in 2014, resulting in three species ranked as rare, one as uncommon, eight as common, three as having only historical records, and two species lacking information adequate to assign a rank. The three rare species were added to Vermont's Threatened and Endangered Species list in 2015: the Rusty-patched Bumble Bee (*Bombus affinis*) and Ashton Cuckoo Bumble Bee (*B. ashtoni*) are now listed as endangered, and the Yellow-banded Bumble Bee (*B. terricola*) is threatened. Harmful parasites accidentally imported from Europe and neonicotinoids, a class of pesticides toxic to bees and other insect pollinators, are believed to account for the bumble bee declines

Freshwater Mussels

Freshwater mussels are now recognized as the most endangered group of aquatic species in North America. In Vermont, ten of the eighteen native species are listed under the state endangered species law, one is federally endangered, and several others are considered rare. Between 2003 and 2010, surveys sought to describe the abundance, distribution, and habits of Vermont's freshwater mussel populations. Findings included a drastic decline in Brook Floater and a dramatic increase in Eastern Elliptio species in the West River. We now know more about fish hosts for Creek Heelsplitter and Fluted-shell, and have a better understanding of the distribution of Dwarf Wedgemussel and Elktoe. VFWD is in the process of developing a multi-species freshwater mussel recovery plan based on survey findings.

Vermont Invertebrate Database

The paucity of basic information such as species presence, geographic distribution, habitat associations, and life history has limited our ability to direct conservation actions for most groups of invertebrates. Some years ago, the Invertebrate Scientific Advisory Group to the Vermont Agency of Natural Resources' Endangered Species Committee recognized that a substantial amount of information already exists in past invertebrate collecting and research in Vermont, although it is scattered among various collections, government offices, research facilities, published works, gray literature, and other sources. The group recommended a centralized repository of invertebrate species data. What followed was the Vermont Invertebrate Database, an effort that beginning in 2006 compiled over 75,000 records representing over 400 species of invertebrate gathered from museum and private collections and literature. These data have been incorporated into the Vermont Atlas of Life, a database maintained by the Vermont Center for Ecostudies to provide a baseline of information for a broad spectrum of invertebrate taxa for future invertebrate conservation and research planning efforts.

Fish

Fish Surveys

Twenty-three percent of Vermont's native fish species are currently designated as rare, and many of these have experienced population declines due to habitat disturbance, overharvesting, pollutants, and the introduction of invasive species. Between 2004 and 2011, five Vermont water bodies were sampled to document abundance and distribution of some of these rare species (Lake Champlain, Holland Pond, Black River, Poultney River, and Missisquoi River). Surveys targeted Lake Sturgeon; Muskellunge; Shorthead, Greater, and Silver Redhorse; Eastern Sand and Channel Darter; Blackchin, Blacknose, and Bridle Shiner; Round Whitefish; and Stonecat. When Stonecat was discovered in a new location in 2004, fisheries biologists wondered if there may be additional, undiscovered populations, and rigorous additional effort was placed on locating new sites, to no avail. Sturgeon spawning was detected in three of the four rivers that had been known spawning sites in the past (Sturgeon spawning was detected in the Winooski, Lamoille, and Missisquoi Rivers), but not in the fourth, Otter Creek. The Sturgeon project also included genetic sampling of the Lake Champlain population, indicating that our local sturgeon is genetically distinct from other North American populations. Significant effort was placed on Muskellunge surveys not only to assess the current population but also to determine the potential to restore the species. Muskellunge were documented in Otter Creek (population established from a stocking program in the 1980's). None were captured in the Missisquoi River. The Missisquoi River between Swanton and Highgate dams is believed to have been the last known refuge population of "native" strain musky in the state but appears to have become extirpated following a chemical spill to the river during the 1970s.

Fish Historical Record Scanning

From the late 1800s to the 1960s, records of fish management activities and observations were documented in diverse paper formats and stored in several physical locations across the state. In recent years, much time and attention has been given to rare or endangered species, species in decline, fish and wildlife species distribution, fish and wildlife community changes and fish and wildlife species genetic composition, and yet the important historic data that could inform management strategies on these topics were difficult to access. For this project, over 2,000 documents were digitally scanned with text recognition software, and over 50 years of stocking records have been compiled into a digital database and made available through the Department's internal website. The digital format, text recognition capabilities and searchable databases have enhanced our ability to access these records to better understand historic species distribution and abundance, past management activities, and cultural impacts that may have influenced current fish and wildlife populations. This information is critical to making meaningful decisions for future conservation activities.

Amphibians and Reptiles

Town Surveys for Reptiles and Amphibians

First published in 1995, the Vermont Reptile and Amphibian Atlas relies heavily on volunteer citizen scientists to submit records of reptiles and amphibians throughout the state. The Atlas has greatly added to Vermont's knowledge of herp distribution and abundance, but one challenge with this type of volunteer-driven project is that some towns remain poorly surveyed. In these locations, knowledge of amphibian and reptile presence is very limited. In 2006, the Vermont Fish & Wildlife Department applied for a State Wildlife Grant to help fill some gaps by surveying three under-represented Vermont towns. In each town, 10-15 previously undocumented species were recorded.

North American Racers, Eastern Ribbonsnakes, and Common Five-lined Skinks—all rare species in Vermont—were also surveyed statewide, resulting in many new and confirmed reports from both new and previously-reported locations.

Rare Amphibian Surveys

When the Boreal Chorus Frog was listed as a state endangered species in 1987, the cause of the population decline was—and still is—unknown. By 2007, it was unclear whether the frog still remained on the landscape. While surveys followed over the next three years, the species was not detected, leaving us to suspect that the frog may either be extirpated from the state or no longer has any viable populations here. Fowler's Toads were also subject to repeated, targeted surveys in 2008, and these also failed to produce any occurrences. This toad species was last documented in 2007, and it has now been added to the state's endangered species list. Mudpuppy was also considered for threatened status in Vermont in 2011, but the decision was made against listing the species. This decision was based upon 2008-2010 surveys in which Vermont Fish & Wildlife Department refined a methodology to more effectively survey this difficult-to-detect species and thus learned more about mudpuppy population distribution. As part of this project, genetic sequencing across the state also revealed that Lake Champlain Mudpuppies show significant differentiation from those in the Connecticut River basin, which may have been introduced from outside the state.

Rare Snake Surveys

Timber Rattlesnakes, Eastern Ratsnakes and North American Racers are all restricted to just a few locations in Vermont. The rattlesnake is listed as state endangered and the ratsnake and racer are threatened. Traffic, mowing, and intentional kill all pose threats, as does human visitation to denning areas. In 2010, SWG funded a rattlesnake telemetry study to gain insight on home range and movements, parcel utilization, genetic analyses and some population demographic information, which provided a basis for developing the state's 2015 Rattlesnake Recovery Plan. Essential to the project were staff of The Nature Conservancy, local citizen volunteers, the Herp Scientific Advisory Group, and Vermont Reptile and Amphibian Atlas staff and contributors. During the project, Snake Fungal Disease was confirmed in both of Vermont's two rattlesnake populations. In 2012, a radio telemetry study was also completed for Eastern Ratsnakes to expand the knowledge base and known Vermont range for this species. North American Racers have been surveyed numerous times since 2006, with the latest two documented sightings surfacing in 2014. A Racer Habitat Improvement Project conducted from 2007-2014 in one Wildlife Management Area (WMA) has created new racer travel cover, several experimental hibernacula, foraging cover and egg-laying substrate.

Turtle Conservation, Planning, and Management

It is uncertain what the future holds for any of Vermont's seven turtle species. The Spotted Turtle is state endangered, the Spiny Softshell is state threatened, Wood and Musk Turtles have been identified as SGCN, and even the remaining are vulnerable to boat traffic, human disturbance, egg predation, road mortality, collection, and habitat conversion. With the help of SWG, USDA Wildlife Services, many volunteers, the Friends of Northern Lake Champlain, ECHO Lake Aquarium and Science Center, Missisquoi National Wildlife Refuge, the Vermont Forests, Parks, and Recreation Department, Audubon Vermont, and Green Mountain Audubon, VFWD established a Spiny Softshell Turtle monitoring and management program to develop and implement conservation and recovery methods for softshell turtles in Vermont. Several large, communal nesting beaches are intensively managed for the Spiny Softshell, with Map, Snapping, Painted, and Musk turtles also benefiting. Initial monitoring not only enhanced knowledge of distribution but also made it clear

how to manage communal nesting areas to save eggs and hatchlings from predation, parasitic flies, and drowning. Additional turtle studies included a genetic analysis indicating that there is no female-mediated gene flow between the two Spiny Softshell subpopulations in the state and radio-tagging of Spotted Turtles to learn more about movements and habitat use. Invasive plants have been controlled at a Spotted Turtle Wetland.

Vernal Pool Mapping

Vernal pools are vital breeding habitat for a range of species, including several amphibians. A series of State Wildlife Grants helped in funding first a project that mapped potential vernal pools across the state using infrared photographs, and then to verify the mapped sites through field investigation. Work on this project is ongoing and relies on a large volunteer base as well as a partnership with Vermont Center for Ecostudies and Arrowwood Environmental. In all, some 2500 amphibian breeding sites were cataloged statewide. More information can be found at

<http://vtcostudies.org/projects/forests/vernal-pool-conservation/vermont-vernal-pool-mapping-project/>.

Mammals

Rare Carnivores

For the most part, rare carnivore studies are limited to the collection and analysis of reported citizens' sighting information related to American Marten and Canada Lynx. Between 2005 and 2015, this involved reported sightings of 44 marten (27 confirmed) and 45 lynx (9 confirmed). For American Marten and Canada Lynx—species with known populations in the state—additional survey efforts to delineate presence include winter track surveys, remote cameras, and genetic analysis. A regional lynx survey—funded by USFWS's Endangered Species Program—was also conducted in cooperation with Silvio O. Conte National Wildlife Refuge and New Hampshire Fish & Game to test a remote camera-based protocol, and so far 16 lynx have been recorded. A similar remote camera protocol was used to detect marten, with 8 successful detections. Genetic samples of rare carnivores have also confirmed one wolf, 17 marten, and four lynx. Because lynx sightings have been increasing, VFWD is also implementing a Canada Lynx Response and Handling Protocol to guide staff in their response to cases of injured, sick, or otherwise incapacitated lynx. Most of this rare carnivore work is funded through the Wildlife Restoration Program. We also received 317 citizen reports of mountain lions (none confirmed) and 6 wolves (none confirmed).

Bat Conservation and Management

There are nine species of bat in Vermont, and all have been identified as important conservation targets. In 2003, VFWD established a Bat Conservation and Management Program to learn more about distribution, abundance, and effective conservation strategies for all nine species. When White Nose Syndrome appeared in the Northeast, monitoring efforts intensified, including spring, summer, and fall mist net surveys coupled with winter cave surveys of large hibernacula. These revealed population declines in excess of 90% between 2008 and 2010 at many sites, with a lower rate of decline by 2013. An outreach effort known as “Got Bats?” was also implemented to collect citizen reports of Big Brown and Little Brown Bats in structures such as barns, attics, and bat houses, mapping over 500 maternity colonies across the state. This revealed that Little Brown Bat maternity colonies, once widespread and abundant, are now found primarily in the Champlain Valley. Radio telemetry tracking of Indiana Bats further aided in the development of a publication, *Forest Management Guidelines for Indiana Bats*, that details a series of conservation strategies developed over the course of the project. The funding for this work has been shifting from SWG to Wildlife Restoration Program funds.

Small Mammals of Vermont Atlas

When the 2005 WAP identified a need to improve our understanding of the status and needs of many of Vermont's small mammals, the Small Mammals Atlas was developed in response. In addition to compiling historic documents and museum collections, Small Mammals Atlas project staff conducted field surveys documenting 2,844 small mammal captures from 47 sites. This allowed for the construction of distribution maps for all 23 small mammal species in Vermont.

Birds

Breeding Bird Atlas

First published in 1985 using survey information from 1977-1981, a new Breeding Bird Atlas (<http://www.vermontbirds.org/>) was completed using surveys from 2003-2007, allowing for comparisons in bird distribution over time. This time, 200 bird species were documented—with accompanying range maps—including 17 species that were added since the first atlas and 14 species from the 1985 version that could not be confirmed as breeding in the state in 2007. Over 300 bird-savvy volunteers contributed expertise and nearly 30,000 hours for this Atlas. Vermont Center for Ecostudies handled project coordination with help from Audubon Vermont and its chapters, the University of Vermont, the National Wildlife Federation, and the VFWD. Additional funding from the World Climate Research Program helped make this possible. A website maintained by Patuxent Wildlife Research Center makes Vermont's breeding bird data accessible, at http://www.pwrc.usgs.gov/bba/index.cfm?fa=explore.ProjectHome&BBA_ID=vt2003.

Threats to Vermont's Mountain Birds

Bicknell's Thrush, Blackpoll Warbler, Swainson's Thrush, and several other bird species require the dense shrub thicket found only on Vermont's highest peaks as nesting habitat. This habitat type is not common, which limits the number of nests the birds can build and is often also correlated with low nesting success. Between 2007 and 2011, the Vermont Fish & Wildlife Department supported scientists at the Vermont Center for Ecostudies with the revision of the Mountain Bird Watch monitoring protocol to improve the estimating power of species abundance, distribution, and habitat preferences. This baseline information will allow the monitoring of population and habitat changes associated with climate change, habitat conversion/alteration, and pollution. A report of this study, commonly called "Birdwatch 2.0" can be found at <http://vtecostudies.org/wp-content/uploads/2014/09/VCE-MBW-USFWS-report-2013.pdf>.

Plants and Natural Communities

Rare, Threatened, and Endangered Plant Monitoring

Each year, New England Wild Flower Society and Vermont Fish & Wildlife Department train volunteers to monitor populations of regionally rare plants. In addition to locating and monitoring populations, there is currently a focus on seed collection of New England's rarest plants and identification of high priority sites for management. Vermont Fish & Wildlife Department conducts additional, regular monitoring of the endangered Northeast Bulrush and Jesup's Milk-vetch. Just this year, four new plant species were listed as endangered in Vermont: the Dwarf Birch, Tulip Tree, Whorled Milkweed, and Green Mountain Quillwort. Funding for the federally listed plant species comes, in part, from USFWS's Endangered Species Program; other funding is from Vermont's general fund.

Natural Community Inventories

Natural community inventories are a primary tool for identifying conservation targets, refining the natural community classification, and identifying important wildlife associations with natural communities. In the past decade, Vermont Fish & Wildlife Department has sought to fill data gaps by conducting inventories of Dwarf Shrub Bogs, Poor Fens, Limestone Bluff-Cedar Pine Forests, all types of softwood swamps, and Montane Spruce-Fir Forests. The Department is currently completing a statewide inventory of oak-pine-northern hardwood forest types. Results from these projects include identification of state-significant examples of natural communities that are added to the Natural Heritage Database and providing recommendations to private landowners on how best to manage the natural communities they own. State-significant examples of natural communities are considered high priority for conservation on both public and private land. Natural community inventory work remains a priority of the Department and will continue in years to come.

Invasive Species

Housed by the Vermont office of The Nature Conservancy, the iMap Invasives database tracks invasive species infestations, management activities to control them, and documentation of post-management population changes. Ongoing trainings by the Vermont Fish & Wildlife Department and The Nature Conservancy aid volunteers in correctly identifying invasive species and properly using the database. The Vermont Agency of Transportation has also partnered on the project to locate and document invasive plants along interstate highways. This work is supported via a contract with Florida State University, and the site can be found at www.vtinvasives.org. It follows up on previous work conducted in partnership with the New England Wildflower Society to train volunteers to conduct invasive plant surveys for the Invasive Plant Atlas of New England.

Herbarium Records

New England Wild Flower Society's 2003 Herbarium Recovery Project documented and annotated more than 18,600 specimens of regionally rare plants housed in 42 herbaria, and more than 90% of the data of interest to Vermont from this project has now been entered in Vermont's Natural Heritage Database. Additional specimen data from Vermont's own herbarium records has also been updated, and Vermont's Pringle Herbarium records are being scanned into the Consortium of Northeast Herbaria portal (<http://portal.neherbaria.org/portal/>) to make them more widely available. This project is paid for through Vermont's general fund.

Revision of Natural Community Ranking Specifications

Natural Community ranking specifications provide a means to consistently and objectively compare the relative importance of each occurrence of a particular natural community type across the state. New ranking specifications were developed for all 95 natural community types and a report on how to use these specifications is available for the public and consulting ecologists/biologists. The consistency in approach that is included in the natural community ranking specifications and the resulting Natural Heritage Database of state-significant natural communities has been the basis for including natural communities in recent revisions to two rules and land use provisions in Vermont: the Vermont Wetland Rules now includes significant wetland natural communities as a protected function, and Vermont's Use Value Appraisal program now allows for significant natural communities to be enrolled as Ecologically Significant Treatment Areas. Plant rarity ranks were also revised based on new records.

General Natural Heritage

Natural Heritage Inventory

Vermont's Natural Heritage Inventory documents the diversity of native plants, animals and natural communities in the state. The inventory is maintained using the "Biotics" software, which provides a common data management platform for members of the NatureServe network to achieve and maintain a unified taxonomy and consistent application of our shared data standards and methodology. The database tracks natural heritage elements including taxonomy, nomenclature, rarity ranks, habitat descriptions, threats, trends, and additional information for 1260 animal species, 1425 plant species, and 92 natural community types. In addition, there are over 10,000 populations documented, complete with GIS mapping. General information is available to the public for planning purposes through Vermont Center for Geographic Information (<http://www.vcgi.org/>) and the Agency of Natural Resources' online Atlas (<http://maps.anr.state.vt.us/ANRA/>). More detailed attribute data is made available to partners through data use agreements. NatureServe is a major partner and provides software support for Biotics and servers as a centralized repository for North American biodiversity information.

Structured Decision-making

The Vermont Fish & Wildlife Department, in partnership with the University of Vermont, developed a system for managing, archiving, sharing and analyzing wildlife data and information. This new system, known as Resources of Vermont, or ROVER, is a comprehensive database for a wealth of wildlife data in Vermont that compliments the comprehensive Natural Heritage Database that addresses rare, threatened, and endangered species and significant natural communities. While it will continue to grow and evolve it currently supports data for many species of birds and mammals. It will be used in the future to guide conservation decisions by providing more effective access to data and greater abilities to analyze and interpret information.

Wildlife Management Area Inventories

Whenever possible, Vermont uses Long Range Management Plans to guide activities in state-owned Wildlife Management Areas. Since 2004, the Vermont Fish and Wildlife Department routinely inventories bats, breeding birds, reptiles, and amphibians in state Wildlife Management Areas, to aid in the creation of well-informed management plans.

Research

The 2005 Wildlife Action Plan identified gaps not only in our knowledge of species abundance and distribution but also in potential threats species face, interactions with surrounding natural or human-altered landscapes, and the identification of conservation needs. The following are research projects seeking to answer questions that will inform the management of wildlife or their habitats.

Fishes

Lake Whitefish

In Vermont, Lake Whitefish occurs only in Lake Champlain where, prior to its closure in 1913, there was a substantial commercial seining whitefish fishery. Recent research conducted by the University of Vermont concluded the population is stable and unexploited but populations that once existed at two locations in the lake once supporting commercial harvest appear to be extirpated due to past

exploitation and/or degraded spawning habitat. The university is currently conducting research of the effects of man-made barriers (e.g. causeways) in limiting whitefish movements among bays within the lake and whether these are causing the formation of spatially separate and genetically differentiated sub-populations within the lake. This work was partially funded by a State Wildlife Grant awarded by Vermont Fish & Wildlife Department. Other possible threats to Lake Whitefish in Lake Champlain are competition with exotic species (e.g. Alewife) and Sea Lamprey parasitism.

Stream Temperature

Because of land clearing, dams, and other shifts in land use that remove shade from stream banks, some of Vermont's historic trout streams are now too warm to support healthy trout populations. With funding from the State Wildlife Grant program, the Vermont Fish and Wildlife Department conducted a series of case studies to model watershed temperature conditions and predict quality brook trout habitat. One study examined potential management scenarios on river temperatures, finding that increasing shade on just the upper 50% of the watershed has a high impact on the resulting downstream water temperatures. In a second study, metrics were developed that allow researchers to assess coldwater habitat suitability when only temperature data are available. This study produced recommendations which can guide decision making for the conservation of native salmonid populations.

Stream Morphology

Physical processes form habitat in a stream channel, such as movement of sediment or woody debris, formation of scour and depositional features, and dynamic riverbank changes. Combined with chemical constituents and biological interactions, physical habitat determines biological productivity and diversity and drives the aquatic ecosystem. By taking a detailed look at the habitat resulting from the physical processes taking place in a stream, it may be possible to understand how fluvial processes impact aquatic communities. The Vermont Agency of Natural Resources Reach Habitat Assessment has been created as an integral part of the existing Vermont Stream Geomorphic Assessment Protocols, although it is also available for stand-alone usage. It builds on previous habitat assessments by fundamentally being based on stream processes and their link to resultant channel, bank, and cover features. Indicators of key physical ecological attributes were identified for inclusion in the assessment that are necessary for aquatic organisms to carry out life cycle functions.

Fish Health & Predator-Prey Dynamics

For fisheries biologists to effectively manage wild fish, they must first understand a wide variety of ecosystem functions, including diseases, predator-prey relationships, and more. For example, whirling disease is a parasite that can be detrimental to trout and salmon populations. From 2006-2008, Brook Trout, Brown Trout, Rainbow Trout and Atlantic Salmon were tested for the disease, which was confirmed to be present in several Vermont streams. In a different study, fisheries personnel developed standard operating procedures for hydroacoustic surveys that quantify populations of forage fish in Lake Champlain. These surveys are used to monitor the prey base for larger, predatory fish so that fisheries can maintain the balance of fish species in the lake.

Landlocked Salmon Stocking

Landlocked Atlantic salmon were once abundant in Lake Champlain, but habitat degradation and over-fishing destroyed the native population by 1850. Since 1973, the Vermont Fish and Wildlife Department, along with New York and the U.S. Fish and Wildlife Service, have since committed to restoring the salmon population and fishery. This effort relies on a successful stocking program—

and there are several stocking methods that can be employed. The goal of this study was to evaluate one particular strategy to stock fry in tributaries to Lake Champlain. A rotary screw trap was employed to sample downstream migrating salmon smolts produced from fry stocking in the Huntington River, a tributary to the Winooski River and Lake Champlain. Results showed that significant numbers of smolts were produced, suggesting that fry stocking does indeed seem to be a promising restoration strategy.

Atlantic Salmon Genetics

Historically, anadromous Atlantic Salmon was native to the Connecticut River watershed but became extirpated by the early 1800s following construction of the first dam on the river in 1798 near present-day Turners Falls, Massachusetts. During the latter half of the 19th Century unsuccessful attempts to restore salmon to the river were undertaken. The most current restoration effort was initiated in 1965 and continued until 2012, when the U.S. Fish & Wildlife Service and other basin state fisheries agencies (New Hampshire, Massachusetts and Connecticut) withdrew from the program. Annual stocking of hatchery produced salmon fry continued through 2013 but since then no juvenile salmon have been released into Vermont waters, and smolt outmigration is expected to end after the spring of 2016. Even though salmon restoration has ended for the foreseeable future, it is anticipated that adult salmon will return to the Connecticut River in small numbers and, therefore, will be provided upstream passage by dams at Holyoke and Turners Falls, Massachusetts and Vernon, Vermont as part of the power utilities commitment to providing fish passage for other anadromous fishes (e.g. American Shad). Given these recent events, anadromous Atlantic Salmon will in all likelihood remain an extirpated species in Vermont. In 2011, prior to termination of the salmon program, a State Wildlife Grant funded project was undertaken to test for genetic variation in smolt production and adult returns to determine smolt and adult return production from each stocked Connecticut River tributary. Genetic testing of samples has been completed, even though with termination of the restoration program this information probably will not be used to adaptively manage stocking efforts.

Lake Trout Genetics

Although Lake Trout are indigenous to most of Vermont's larger and deeper lakes, the status of the native populations was in question due to more than a century of stocking at all these lakes using lake trout sources from outside Vermont. A decade ago the lake trout populations in several Northeast Kingdom (NEK) lakes were supported entirely or predominantly by natural reproduction despite annual stocking. This study sought to determine the degree to which a century of lake trout stocking had genetically influenced these populations. Results irrefutably show that stocking has influenced all the contemporary populations, with the demonstrable degree of influence varying from one lake to another. However, the suite of genetics tests included in the study also reveals evidence that elements of the indigenous genomes persist. Unfortunately, in the absence of comprehensive historical documentation of lake trout stocking in Vermont accompanied by genetic material from the various source populations, it is impossible to describe the *family trees* of contemporary wild lake trout at the several NEK lakes. Lake trout stocking was discontinued in 2006 at most of the lakes, and the naturally reproducing populations continue to be monitored periodically.

Mammals

Indiana Bat

In 1973 the Indiana Bat was one of the first bats recognized as endangered under the federal Endangered Species Act. Populations then continued to drop through much of the species' range, although Vermont's summer populations have remained robust. Biologists recently began to suspect the decline's primary culprit to be the loss of summer habitat—particularly the roost trees where females form maternity colonies, give birth, and raise their young—but the recipe for attractive Indiana Bat habitat remained mysterious prior to this study. University of Vermont researchers worked with Vermont Fish & Wildlife Department staff to evaluate minimum habitat and maternity site requirements for the Champlain Valley and developed a tool to predict maternal nest sites and prioritize sites for conservation.

Bobcat Movement and Habitat Use

Animals with large home ranges can be tricky to study and manage, because their territories include a network of many different habitat types. The Bobcat is no exception, and prior to this study there were a number of gaps in our knowledge of Bobcat habitat needs. To get a better sense of home range requirements, University of Vermont researchers and Vermont Fish & Wildlife Department staff captured Bobcats across a landscape that spanned urban, agricultural, and forested settings and outfitted them with GPS collars that recorded each individual's location over a 3-4 month period. This study provided critical information about how Bobcats travel through a landscape by staying under shrub cover as much as possible, allowed an estimation of minimum home range resource requirements for breeding females, and provided information to use in habitat suitability mapping and home range requirements for Bobcat conservation and landscape-scale management.

Habitat Blocks and Critical Crossings

Black Bear, Fisher, Marten, Lynx, and River Otter are among the species that rely both on large blocks of contiguous forest and secure means of crossing roads and human-impacted landscapes between these forest blocks. Although total forest cover in Vermont increased over the past century, it has again begun to decline, and it is breaking up into an increasing number of smaller blocks as a result of residential development and road construction. Because we still have a limited ability to prioritize conservation of these blocks and forest connectors, the Habitat Block project identified and mapped 4,055 habitat blocks, then evaluated them for biological and physical diversity and potential threat. Considerable work has been done to map and rank likely wildlife road crossing areas. Structural connectivity models showing potential road crossings have been developed and refined in partnership with the Agency of Transportation, and the Staying Connected Initiative is working with partners on several fronts toward a better understanding of functional connectivity. The Critical Paths project involved field-checking road crossings for signs of wildlife use, potential threat, and conservation priority, as well as several photo-monitoring projects that are capturing pictures of wildlife use of transportation infrastructure. Forest block and road crossing data has been made available to the public via the Vermont Agency of Natural Resources' online Atlas (<http://anrmaps.vermont.gov/websites/anra/>).

Birds

Bicknell's Thrush in Northeastern Vermont

Large-scale wind energy developments have recently been installed or proposed for a number of Vermont's peaks, and their arrival brought to the surface questions about the interplay between wind farms and the wildlife that inhabits Vermont's highest elevations. Because the peaks that developers believe are best suited for wind energy are largely covered by montane spruce-fir forest, this project studied possible effects from these developments on the rare Bicknell's Thrush that relies on this habitat. The wind project did not go forward, so the study was unable to ascertain the effects of building the towers. In their final report, biologists from Vermont Center for Ecostudies say that while Bicknell Thrush ecology is too complex to make a specific prediction of development impacts, both short- and long-term effects are possible, and they may differ as initial impacts give way to succession. The report includes maps showing the location and size of current and potential Bicknell's habitat in the Northeast Highlands of Vermont, to be used to assess future wind development and conservation priorities. In addition to funding from State Wildlife Grants, the Nature Conservancy, Vermont Institute of Natural Science and Vermont Center for Ecostudies each provided aid.

Black Throated Blue Warbler Abundance and Nesting Productivity

Habitat quality—a phrase used extensively in conservation and land use planning—is often gauged by wildlife population counts. However, as habitats become modified for human uses and low-quality habitats become more dominant, wildlife may be using environmental cues that no longer relate to better reproductive success, causing them to make poor habitat choices. Indeed, results of this study indicate that for Black Throated Blue Warblers, preferred habitats did not correlate to the highest fitness levels in terms of daily nest survival or annual fecundity. It seems that predation and cowbird parasitism kept “popular” habitats from being as successful as some of the less-preferred habitats.

Landscape Conservation

Shoreline and Development Surveys for Vermont's Lakes

Lakeshores in Vermont provide vital habitat for a variety of SGCN, and until recently there was no regulation to protect them. This project compared the SGCN present in undeveloped and developed lakeshore areas, finding that with the exception of aquatic plants, there were significantly fewer SGCN species present at the developed sites than the undeveloped shorelines. The study then used reserve design methodology to identify lakeshore areas that are most likely to support SGCN, producing a map that could be used to help prioritize lakeshore conservation efforts. The findings of this project also aided in efforts to get the Vermont legislature, governor, and the public to support lakeshore regulation, and in July of 2014, the Shoreland Protection Act was established.

Species Recovery

Over the past decade, 20 species (9 plants and 11 animals) were added to the state's Endangered Species list and are now receiving the additional protections that come with the listing. Because the ultimate goal of the endangered species program is recovery, some of these species have been targeted for additional action. A successful recovery restores a once endangered or threatened species to a point where protection is no longer needed, because the species can sustain itself as a natural part of its ecosystem. Creating a recovery plan is generally the first step, which might include protection or restoration of habitat, species or habitat management activities, or even translocation

of individuals from an outside population. The following are recovery projects that Vermont has undertaken in the past decade. All have been guided by the 2005 Wildlife Action Plan.

Invertebrate Recovery

The Cobblestone Tiger Beetle, the Hairy-necked Tiger Beetle, and several species of freshwater mussel are state-threatened in Vermont. After several years of surveying known populations of these invertebrates, Vermont Fish & Wildlife is in the process of completing recovery plans for the two tiger beetles and a multi-species recovery plan for freshwater mussel species, several of which are in rapid decline due to the invasion of the invasive Zebra Mussel.

Bat Recovery

Vermont Fish and Wildlife Department staff were in the process of creating a statewide bat conservation and recovery plan when two occurrences arose. First, proposals to expand wind energy facilities in Vermont brought up questions about the potential threat of wind development to bats, which needed to be better studied. In the winter of 2008, White Nose Syndrome (WNS) then appeared in Vermont, which has been detrimental to the bat population. Substantial efforts since then have gone toward research on the causes, threats, and spread of the disease. Now that the effects of both wind facilities and WNS are better studied and understood, a revised bat conservation and recovery plan can be developed, most likely within the next few years.

Bird Recovery

The past decade can claim several success stories among bird recovery efforts. Common Loon, Peregrine Falcon, and Osprey were all delisted from state-endangered status in 2005, and their recovery goals have been met. Common Tern is very close to its recovery goals and is being considered for downlisting, while the known nesting population of Bald Eagles in Vermont has grown from zero in 2007 to 18 nesting pairs with 17 fledglings in 2014. These species all continue to be monitored, with the help of Audubon Vermont, Vermont Center for Ecostudies and citizen volunteers. Other ongoing bird surveys include Black-backed Woodpecker, Whip-poor-will, Golden-winged Warbler, Grasshopper Sparrow, and a suite of wetland bird species. In addition, we are evaluating the results of Spruce Grouse Translocation efforts. Once down to just a single population of 150-300 birds, two State Wildlife Grants allowed a total of 134 birds to be captured in Maine and Quebec and then released in a small portion of the bird's former range in Victory, Vermont, twenty of which were fitted with radio transmitters to track movements. If the translocation efforts are successful, the establishment of this second sub-population will be a significant step toward downlisting the species.

Other Recovery Projects

Vermont's Rattlesnake Recovery Plan was adopted in 2015 after substantial research on the status of Vermont's two populations. The Vermont Eastern Spiny Softshell Turtle Recovery Plan was also approved and adopted in July of 2009. A draft restoration plan for Lake Sturgeon in Vermont was released in August, 2015. Federal recovery plans were developed for the Jessup's Milk-vetch and Northeastern Bulrush which are found in Vermont. The New England Wildflower Society has developed conservation plans for many high priority plant species in the region with some technical support from VFWD for species found in Vermont.

Policy, Planning, and Land Acquisition

In some cases, the 2005 Wildlife Action Plan identified that Species of Greatest Conservation Need would benefit most from statewide policies, long-range planning, or the protection of additional habitat important at a statewide regional scale.

Vermont Conservation Design

Fish, wildlife and other elements of biological diversity rely on both landscape-level features such as forest blocks, connectivity, and riparian areas, and finer-scale elements of natural communities, specific habitats, and species. In 2015, Vermont Fish & Wildlife Department, Vermont Land Trust, Vermont Forests, Parks, & Recreation, The Nature Conservancy, Northwood Stewardship Center and others produced “Vermont Conservation Design: Maintaining and Enhancing an Ecologically Functional Landscape.” This report identifies coarse-filter conservation targets for landscape scale features including forest blocks, riparian areas, surface waters, wildlife and landscape connectivity, and physical landscape diversity that are necessary to ensure the conservation of many finer scale conservation elements in the face of climate change and habitat loss. This is a significant addition to conservation planning in Vermont. Phase 2 of this project will focus on natural communities, habitats, SGCN, and rare species.

Getting the Lead Out

Lead is known to be toxic to many wildlife. The Common Loon is particularly sensitive to it, and lead toxicosis contributes to the mortality of the loon and at least 23 other species in North America. Research in the northeastern United States and Canada has documented that poisoning from lead sinkers and jigs is the leading cause of observed loon deaths and can account for 10 to 50 percent of dead adult loons found. In 2006, Vermont Fish & Wildlife Department assisted in a successful campaign to ban the sale and use of lead sinkers for fishing.

Helping Landowners Protect Ecologically Sensitive Lands

Approximately two million acres of Vermont’s forestland is enrolled in the Use Value Appraisal program, which requires active management of enrolled land. In 2009, changes to the Use Value Appraisal program allowed forest areas to be enrolled as “Ecologically Sensitive Treatment Areas,” meaning that instead of being managed exclusively for timber, they can be managed for their values as significant natural communities. At the same time, the Use Value Appraisal program was also revised to allow for enrollment and management for significant wildlife habitat. To qualify, Vermont Fish & Wildlife staff review and approve proposals based on the Department’s standards of significance for natural communities and wildlife habitat.

Land Acquisition

From 2005-2013, the Vermont Fish & Wildlife Department acquired 41 separate parcels in fee totaling 4,141 acres, either to be added to existing Wildlife Management Areas or to create new ones. The Department also acquired 2,322 acres under conservation easement during the same period. Conservation activities have focused on a few key criteria: 1) the purchase of key inholdings and lands adjacent to existing Wildlife Management Areas; 2) ecologically-significant lands such as habitat for rare, threatened or endangered species and critical wildlife habitat; and 3) public access for wildlife-dependent recreation.

Technical Assistance and Public Awareness

With over 80% of Vermont's land in private ownership and the majority of land use and development decisions made at the local or regional level, the conservation of Vermont's wildlife relies heavily on private landowners, municipal governments, regional planning groups, and coordination among state agencies and conservation organizations. In order to fulfill the goals of the Wildlife Action Plan, Vermont has therefore allocated substantial resources toward working with all scales of stakeholders in Vermont.

Community Wildlife Program

In Vermont, land use and development decisions are made at the local or regional level, through municipal or regional plans and zoning bylaws. Distributing information to municipalities and regional decision-makers is therefore a critical piece of the state's natural resources planning process. After publishing *Conserving Vermont's Natural Heritage: A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife, and Biological Diversity* in 2004, Vermont Fish & Wildlife Department created the Community Wildlife Program to provide additional support and technical assistance to interested towns, regional planning commissions, and others in incorporating the latest information and strategizing for the conservation of fish and wildlife. Since 2003, the program has provided technical assistance to all 14 of Vermont's Regional Planning Commissions and more than 150 towns, created a web guide to technical assistance opportunities available in the state, expanded Vermont Fish & Wildlife Department's work with partner organizations in several landscape-level conservation projects, and has been instrumental in providing resources and guidance for grassroots regional planning projects. From 2008-2010, this work was further supported by a partnership with the Vermont Natural Resources Council, who conducted a review of every Vermont town plan and bylaw that pertains to fish and wildlife species and habitat protections. This updates a similar assessment from 2000, enabling a review of progress over the past decade. While trends are noticeably in the direction of increased attention to wildlife conservation—with 87% of all municipalities recommending the protection of wildlife habitat in town plans—the report found a sharp disconnect between municipal plan recommendations and the implementation of those recommendations through bylaws. This report has aided Community Wildlife Program staff and other state-level planners in appropriately targeting technical assistance to municipalities. For more information on the Community Wildlife Program, see

http://www.vtfishandwildlife.com/get_involved/partner_in_conservation/community_wildlife_program.

Helping Private Landowners Help Wildlife

Because conservation in Vermont relies heavily on private landowners, VFWD has partnered with Vermont Coverts several times in the past decade to conduct landowner outreach. In addition to 23 workshops focused on protecting SGCN, a landowner orientation video and four publications were developed and distributed to 500 forest landowners. The partnership also reached out to new landowners by creating and distributing 100 “Welcome Wagon Kits”—buckets containing materials about land stewardship and wildlife conservation in Vermont. Two additional publications have targeted landowners; first, VFWD worked with the Center for Northern Woodlands Education to publish and distribute ~20,000 copies of *The Place You Call Home: A Guide to Caring for Your Land*, a magazine compiling formerly-published articles relating to land stewardship with an emphasis on technical forest management skills. Secondly, *Wildlife Habitat Management for Lands In Vermont* provides tips on recognizing wildlife habitat and then managing it to benefit wildlife in tandem with other management goals such as timber or hiking trails. This publication was the result of a partnership with the Vermont Department of Forests, Parks and Recreation and the USDA Natural

Resources Conservation Service. Using non-SWG funding, VFWD works with the same partners to aid landowners through additional means, such as by helping landowners to incorporate wildlife habitat management and conservation considerations into forest management plans submitted to the Vermont Use Value Appraisal Program.

Public Opinion Survey

While surveys show that most Vermonters support the conservation of wildlife, an average of 6500 acres of wildlife habitat are lost in the state every year through the development of land. Prior to this study, it was unclear what messages and/or actions most effectively motivated the public to support



specific conservation programs, and so data were collected to document public attitudes towards a variety of conservation strategies. In addition to creating a baseline against which we can monitor changes in public attitudes, these data aided in the development of a state communications plan and conservation message, including the “Respect. Protect. Enjoy” logo and tagline now used by the Vermont Fish & Wildlife Department.

Species of Greatest Conservation Need

The Vermont Fish & Wildlife Department provides technical support and guidance to protect Species of Greatest Conservation Need (SGCN) to those who work directly with these species and their habitats. Beginning in 2012, a State Wildlife Grant aided VFWD staff in providing technical assistance to other Vermont Agency of Natural Resources departments, private land managers and NGOs, interagency programs, and scientist researchers. Additional technical support is provided to the Vermont Endangered Species Committee and several Scientific Advisory Groups. A second State Wildlife Grant provided support at the municipal level, striving to connect citizens with the SGCN occurring in town forests and other municipally owned lands. This project worked through town partnerships to provide educational materials and technical assistance with municipal planning efforts in eight target communities.

Orange County Headwaters Project

In the summer of 2007 two University of Vermont graduate students undertook an inventory of wetlands and vernal pools in the towns of Washington and Corinth. A State Wildlife Grant, along with Connecticut River Mitigation and Enhancement Funds and Orange County Headwaters Project operating funds, provided additional support for identifying high priority areas for conservation and conducting outreach to community members. This included directing citizens to available conservation resources and researching recommendations for protecting vernal pools and wetlands while continuing active management for forest products. The program’s efforts resulted in the permanent protection of 24 parcels through conservation easements as well as enrollment of four landowners in Audubon Vermont’s Forest Bird Initiative program—a program that conducts habitat inventories and provides specific recommendations for the enhancement of forest bird habitat—and an equal number in NRCS’s Wildlife Habitat Incentives Program.

Baitfish Workshop

The harvest, transport and use of baitfish has long been identified as a high-risk vector for the spread of aquatic invasive species and fish pathogens. Wild harvested baitfish are frequently moved to and used in other waterbodies or watersheds, and anglers tend to release unused live bait when

done fishing. This workshop used the Hazard Analysis and Critical Control Point (HACCP) concept to teach risk identification and analysis to over 40 participants from Vermont's baitfish industry and state government. Presentations focused on managing the risk of inadvertently spreading invasives during day-to-day activities in their fields.

The Fishes of Vermont

Vermont supports the greatest freshwater fish diversity in New England, and the publication *The Fishes of Vermont*, a field guide to the fish species of Vermont, explains why. Researched and written by Rich Langdon (Vermont Department of Environmental Conservation) and Vermont Fish & Wildlife Department's Mark Ferguson and Ken Cox, the book describes glaciers and other forces that determined fish distribution while presenting a Vermont-specific identification key that maps and describes the life histories of all 92 fish species in the state. Fishermen, natural history buffs, conservationists and natural resource managers will all find this book useful. Its publication was funded by the Wildlife Conservation and Restoration Program—a precursor to State Wildlife Grants.

Bats On Private Land

Suitable summer roosting habitat is extremely limited for the federally endangered Indiana Bat, and much of what exists in Vermont is on private land. Vermont Fish & Wildlife staff provided technical assistance to private landowners interested in managing for Indiana Bats, which included assessing parcels for potential roost trees and desirable habitat characteristics and, when appropriate, connecting interested landowners with habitat incentives programs such as the Natural Resources Conservation Services' WHIP and EQIP.

Natural Communities and the Plant Stewardship Project

In Vermont, many significant natural communities and rare, threatened, and endangered species are located on private land. To support landowners and land managers the Vermont Fish & Wildlife Department has conducted outreach to identify these sites, map them, and then assess population levels, threats, and management needs. These visits and associated landowner communications were made possible thanks to funding from the Landowner Incentives Program over much of the past decade. Unfortunately, that federal program has been discontinued, and with it much of VFWD's financial capacity to carry on the program.

Management

In order for some species to thrive in Vermont, the 2005 Wildlife Action Plan identified a need for changes in species or habitat management. In some cases, habitats have been restored or expanded. In others, projects were implemented to mitigate conflicts between wildlife needs and human activities. Included below are some of the specific management efforts undertaken for the benefit of Species of Greatest Conservation Need.

Aquatic Organism Passage

Fish and other aquatic organisms need to move freely within stream networks to meet daily and seasonal life cycle needs. During the development of transportation systems in Vermont, numerous barriers to the movement of these populations have been created from the construction of culverts at road/stream crossings. An assessment of aquatic organism passage (AOP) at over 3000 culverts throughout the state found less than 6% were rated fully passable. Through this project we developed technical guidelines for designing AOP at road/stream crossings; conducted technical

trainings on these techniques for state, federal and municipal transportation managers, engineers, regulators and biologists; refined culvert assessment protocols and screening tools to better utilize assessment data; produced outreach materials (Stream Crossing Handbook) and worked with partners to identify, design and implement AOP enhancements at existing stream crossings. These efforts have also served to inform regulatory improvements for design and performance standards for culvert replacement and repair projects. In addition to State Wildlife Grants, this work was funded through Sportfish Restoration Programs.

Prevention of Fish Disease

Preventing the introduction of diseases to fish culture stations and to natural fish populations can be a daunting challenge. To accomplish disease prevention goals, VFWD employs a comprehensive fish health program that follows strict biosecurity practices. Specifically, these biosecurity practices seek to reduce the risk of fish pathogen introduction into fish populations and minimize the risk of pathogen spread. Examples of the tools implemented include but are not limited to: strict fish importation permit program, regulations prohibiting the live transfer of fish from one body of water to another, the use of water filtration and disinfection equipment for fish culture station water source(s), regular use of disinfectants at fish culture stations and pathological examinations of both naturally produced fish and fish produced at fish culture stations.

Amphibian Crossings

On warm, rainy nights in spring, many roads in Vermont experience an inundation of frogs and salamanders attempting to reach the vernal pools, wetlands, and ponds where they breed. When these amphibian travel routes cross roads heavily traveled by human vehicles, mortality can be extremely high. In 2011 and 2012, funding was secured from State Wildlife Grants, Vermont Agency of Transportation's Transportation Enhancement Grant, private donors, and several smaller grants to build two wildlife crossing structures at one well-used amphibian crossing site in the Town of Monkton. General monitoring of this crossing dates back to 1997 and monitoring specific to planned crossing structures began in 2011—prior to the structure's construction—to allow for future comparisons of road mortality. The crossing is expected to be built during the summer of 2015. Local partners and key supporters include the Town of Monkton Select Board, Monkton Conservation Commission, Lewis Creek Association, many volunteers, and many individuals who have financially supported the project.

Beaver Baffles

Wetlands created by Beaver are ecologically important and provide critical habitats for many Species of Greatest Conservation Need. However, Beaver wetlands also jeopardize—or are perceived to jeopardize—roads, dwellings, timber, homes, and water supplies. Beaver baffles are water control structures installed to alleviate flooding caused by beaver and satisfy the concerns of the landowner or town while maintaining as much wetland habitat as possible. From 2006-2010, 206 structures were installed and 1,785 wetland acres conserved as a result of this program, funded through State Wildlife Grants, the U.S. Fish & Wildlife Service Partners in Wildlife Program, the Duck Stamp Fund, and Vermont Agency of Transportation.

Grassland Birds At Airports

As a group, grassland birds have been on the decline, in large part because much of the area of grassland habitat has reverted from farmland to forest and woody shrub habitat. In an ongoing

project, Vermont Fish & Wildlife Department works with Vermont Agency of Transportation to develop airport plans that include the consideration of grassland birds in their mowing schedule.

Golden-winged Warbler

To find a nesting Golden-winged Warbler, you would want to look in old fields with sparse trees or shrubs with a grassy understory close to mature forest. In Vermont this type of habitat is becoming increasingly rare as forests mature, and you may have a difficult time finding one. Through funding from the NRCS's Wildlife Habitat Incentives Program (WHIP) and Conservation Innovation Grant (CIG), Vermont Audubon surveyed a transmission line corridor to prioritize suitable Golden-winged Warbler habitat. Vermont Fish & Wildlife Department then worked with Audubon and private landowners on six high-priority sites to enhance habitat for the Golden-winged Warbler and other shrubland birds. In 2014—just six months after management activities—monitoring visits found hybrid Blue-winged/Golden-winged Warblers. While no true Golden-winged Warblers have yet been found at the treatment areas, monitoring will continue on these sites for the next 3-5 years.

Double-crested Cormorant Control

Wildlife management increasingly involves mitigating unwanted wildlife impacts, and Double-crested Cormorants have recently been the target of such efforts in order to restore island habitats and protect rare bird species. New to our area, cormorants remove leaves and sticks to use as nesting material, and their acidic guano accumulates near nesting sites, eventually killing trees and shrubs—vegetation that provides important nesting habitat for other species. In 1999, VFWD began applying corn oil to cormorant eggs and, in 2004, culling adult birds to limit reproduction. These methods eliminated the nesting cormorant population on the state-owned islands by 2008. The objective of this project was to use an adaptive management approach in order to restore the former islands' habitats, and it was found that the strategy of egg-oiling and culling, in combination with habitat restoration plantings of grasses, shrubs and trees as the cormorant and gull numbers declined, was indeed the most efficient strategy. The USDA/APHIS/Wildlife Services was an integral partner in controlling cormorants on the islands, and funding was provided by VFWD license funds. Habitat restoration was further funded by the Wildlife Restoration Program.

Partnerships with Natural Resources Conservation Service

The former Wildlife Habitat Incentives Program (WHIP) and the Environmental Quality Incentives Program (EQIP) are administered and funded by the National Resources Conservation Service (NRCS) with technical aid from Vermont Fish & Wildlife Department. These programs help private landowners with the resources and expertise needed to manage their land for the benefit of fish, wildlife and overall environmental quality—be it by releasing mast or apple trees for wildlife, creating early successional habitat for nesting song birds, or controlling invasive species, these programs have helped Vermonters manage their land for wildlife. When the Wildlife Action Plan was adopted in 2005, the Vermont NRCS office quickly adopted it as a guide for its work on these programs. Over the last 10 years of this agreement, Department staff have worked with landowners on approximately 986 WHIP projects and over 220 EQIP projects throughout Vermont. This agreement is ongoing and the continued partnership is improving habitat throughout Vermont.

Regional Partnerships

Because wildlife do not recognize political boundaries, we have found a need to go beyond our own and collaborate with our neighbors. Below are a few examples of Vermont's participation in regional conservation efforts.

Regional Conservation Needs Grant Program

Since 2007, the Vermont and the other twelve Northeast states and the District of Columbia have worked together as Northeast Association of Fish & Wildlife Agencies (NEAFWA) to implement landscape-scale wildlife conservation actions identified in their collective Wildlife Action Plans through the Regional Conservation Needs Grant Program. Partner states contribute 4% of their annual State Wildlife Grants Program allocations to this fund. Through the RCN program states develop, coordinate, and implement conservation actions that are regional or sub-regional in scope; build upon the multiple regional initiatives that already exist and compliment ongoing work in individual states. Each year, NEAFWA states contribute roughly a half million dollars of State Wildlife Grant funds, leveraging another half million dollars or more from the Wildlife Management Institute and grantees. This is a substantial investment in regional scale conservation. Details about the RCS program can be found at <http://www.rcngrants.org>.

The Staying Connected Initiative

Encompassing a region of five Northeast States and three Canadian provinces, the Staying Connected Initiative is a collaborative of NGOs and local, state, and provincial agencies working to protect connectivity, with a focus on wide-ranging mammals such as moose, black bear, fisher, bobcat, lynx, and marten. The Initiative brings a landscape scale vision but tailors actions to the needs of local communities, with approaches including conservation science, land protection, technical assistance to communities, land use planning, transportation mitigation, and policy development. Since the Initiative began in 2009, region-wide accomplishments include the protection of over 300,000 acres of land, technical assistance on land use planning and policies for over 40 municipalities and several regional planning agencies, numerous GIS models for conservation planning, the identification of priority road segments, the development of best practices for mitigating road impacts to wildlife, the creation and distribution of numerous tools and reports to aid with the implementation of connectivity actions, and assistance or outreach to local groups through workshops, field trips, citizen science opportunities, and other community engagement measures. While funding for the Staying Connective Initiative includes substantial contributions from all of its partners, the Competitive State Wildlife Grants program, the Wildlife Conservation Society's Wildlife Action Opportunities Fund, and the Jessie B. Cox Charitable Fund, funding from Vermont's State Wildlife Grants program specifically targeted one wildlife linkage area between the Worcester Range and the state's Northeastern Highlands. Associated technical assistance included the hiring of a local coordinator to act as a liaison between partner organizations and the community, and the publication of an outreach guide entitled "An Enduring Place" (<http://216.92.98.160/assets/enduringplacefinal.pdf>). This guide is a celebration of place, weaving together the cultural and natural landscape in the region and calling out the regional importance of habitat connectivity. More information can be found at <http://stayingconnectedinitiative.org/>.

For the Future

In reality, the above list of projects and accomplishments represents only a fraction of the work undertaken in Vermont that has aided in the implementation of the Wildlife Action Plan. While this

list captures the efforts of Vermont Fish & Wildlife Department and a few prominent partners—with a primary focus on projects funded through State Wildlife Grants—the true efforts have been diverse, accomplished not only by multiple state agencies but by Vermont’s extensive network of conservation organizations, municipalities, regional groups, interest groups, and individuals. In the future, we would like to find a way to capture these collective efforts, in order to better represent Vermont’s progress toward conserving wildlife on our landscape.

VERMONT CONSERVATION DESIGN

MAINTAINING AND ENHANCING AN ECOLOGICALLY FUNCTIONAL LANDSCAPE



August 2015

Eric Sorenson, Robert Zaino, Jens Hilke - Vermont Fish and Wildlife Department

and

Elizabeth Thompson - Vermont Land Trust



Vermont Land Trust

CONSERVING LAND FOR THE FUTURE OF VERMONT

Acknowledgements

We thank Mark Scott (Wildlife Division Director, VT Fish and Wildlife Department), Kim Royar (Special Assistant to the Commissioner, VT Fish and Wildlife Department), Louis Porter (Commissioner, VT Fish and Wildlife Department), Michael Snyder (Commissioner, VT Department of Forests, Parks, and Recreation), Deb Markowitz (Secretary, VT Agency of Natural Resources), and Gil Livingston (President, Vermont Land Trust) for their support in undertaking and completing this project. We greatly appreciate the time and expertise contributed by each of the steering committee members and the support of their respective organizations. This project was funded by the U.S. Fish and Wildlife Service through a State Wildlife Grant as part of developing the 2015 Vermont Wildlife Action Plan.

Steering Committee

John Austin, VT Fish and Wildlife Department
Jayson Benoit, NorthWoods Stewardship Center
Jeff Briggs, VT Department of Forests, Parks & Recreation
Dan Farrell, The Nature Conservancy
Jens Hilke, VT Fish and Wildlife Department
Jon Kart, VT Fish and Wildlife Department
Jane Lazorchak, VT Fish and Wildlife Department
Paul Marangelo, The Nature Conservancy
Doug Morin, VT Fish and Wildlife Department
Steve Parren, VT Fish and Wildlife Department
Nancy Patch, VT Department of Forests, Parks & Recreation
Rose Paul, The Nature Conservancy
Kim Royar, VT Fish and Wildlife Department
Mark Scott, VT Fish and Wildlife Department
Eric Sorenson, VT Fish and Wildlife Department
Liz Thompson, Vermont Land Trust
Bob Zaino, VT Fish and Wildlife Department

Introduction

Vermont has a rich natural heritage and legacy, treasured by Vermonters and visitors alike. Strong public support for conservation is evidenced by numerous public surveys (Roman and Ericson 2015). Vermonters strongly value wildlife, nature, and the state's rural, sparsely developed landscape, including the working lands supporting forestry and agriculture.

This valued natural legacy is threatened. Development and fragmentation of natural habitats, the spread of non-native and invasive plants and animals, and the effects of a changing climate all have the potential to dramatically alter Vermont's natural landscape and the native plants and animals that rely on it. Nevertheless, thanks to the resilience of our forests and natural communities, much of the state is currently in good ecological condition. Following the severe deforestation and soil loss of the heavily agricultural 19th century, Vermont is now largely forested, with many large and intact forest blocks. Vermont's small size belies its rich diversity of species and habitats. The state has many pristine lakes and wetlands; abundant calcium-rich bedrock which supports high species diversity, rare species, and productive forests; and a range of elevations and soil types from the clay soils of Addison County to the rugged alpine summit of Mount Mansfield.

These ecological conditions and strong public support for conservation highlight the opportunity that now exists to protect biological diversity.

Maintaining and enhancing ecological function across the landscape is fundamental to conserving biological diversity. Ecological function – the ability of plants and animals to thrive, reproduce, migrate, and move as climate changes and the ability of natural ecosystems to function under natural processes – is served by high-quality terrestrial and aquatic habitat, natural connections across the landscape, a wide variety of habitat features from low elevation to high, clean water, and healthy rivers, streams, lakes, ponds, and wetlands.

We present a practical approach to protecting and enhancing ecological function into the future. This approach, a landscape-level conservation design for Vermont, is based on a rigorous scientific process using the best available data. The lands and waters identified here are the areas of the state that are of highest priority for maintaining ecological integrity. Together, these lands comprise a connected landscape of large and intact forested habitat, healthy aquatic and riparian systems, and a full range of physical features (bedrock, soils, elevation, slope, and aspect) on which plant and animal natural communities depend. When conserved or managed appropriately to retain or enhance ecological function, these lands will sustain Vermont's natural legacy into the future.

Coarse-filter Conservation Approach

It would be overwhelming to identify and manage for the individual needs of the thousands of species of plants, animals, invertebrates, and fungi in Vermont. The coarse-filter conservation approach treats larger-scale components (or “elements”) of the landscape as proxies for the species they contain (Panzer and Schwartz 1998; Molina et al. 2011; Shuey et al. 2012). If examples of all coarse-filter elements are conserved at the scale at which they naturally occur, most of the species they contain—from the largest trees and mammals to the smallest insects—will also be conserved (Hunter 1991; NCASI 2004; Schulte et al. 2006). This approach is well-documented in the scientific literature (Jenkins 1985; Noss 1987; Hunter et al. 1988; Hunter 1991; Noss and Cooperrider 1994; Haufler et al. 1996; Jenkins 1996; Poiani et al. 2000; USDA 2004).

The coarse-filter conservation approach can provide for the habitat needs of many of Vermont’s species, allowing for efficiency in conservation planning and design. This project focused on identifying landscape-level coarse filters. We have very high confidence that this conservation design identifies areas essential for the long-term functioning of Vermont’s landscape and the species it contains. However, coarse-filter conservation alone cannot adequately address all the needs of Vermont’s species, habitats, and natural communities. Very rare species, whose distribution on the landscape is too infrequent and unpredictable to be captured by most coarse filters, and species with very specific habitat needs (such as grassland nesting birds that in Vermont are only associated with very specific agricultural mowing regimes) require additional considerations. A complementary “fine-filter” conservation approach is needed for these species and habitats, and we are planning to address a conservation design for these species and habitats in a subsequent project.

Methods and Results

Our first step in applying the coarse-filter approach to this project was to list landscape scale elements that could serve as coarse filters and the finer scale elements that could effectively be captured by each. The list of finer scale elements included a broad range of ecological processes, natural communities, habitats, and species. Analyses of the effectiveness of several proposed coarse filters were compiled in a tabular format. This analysis informed our selection efforts. The table is a significant product of this project and serves as a strong conceptual foundation for identifying those landscape elements that most effectively represent the ecologically functional landscape needed to support most of the fine filter elements into the future. The table will be expanded in a later phase of the project to include many more natural communities, habitats, and species and will be the basis for identifying which of these finer scale elements are not “captured” by the landscape scale coarse filters and therefore need specific conservation and/or management attention.

Based on these first steps, we selected five landscape elements from an initially longer list as collectively being the most effective and parsimonious for maintaining an ecologically functional landscape. These are:

1. Interior Forest Blocks
2. Connectivity Blocks
3. Surface Waters and Riparian Areas
4. Riparian Areas for Connectivity
5. Physical Landscape Diversity Blocks

We also identified Wildlife Road Crossings as a key element of the conservation design. Wildlife road crossings are road segments with suitable habitat on both sides of the road. Although not actually a coarse filter, wildlife road crossings are essential to the success of the five chosen landscape elements and therefore are a critical component of maintaining and enhancing Vermont's ecologically functional landscape.

When the ecological functions of each of these landscape elements are maintained and enhanced, and when each is conserved at the appropriate scale and distribution across the landscape, the majority of Vermont's species and natural communities are very likely to be conserved even as the climate changes.

While each landscape element is important on its own, it cannot function in isolation. Maintaining or enhancing an ecologically functional landscape in Vermont depends on both the specific functions of each element, and the ability of the landscape elements to function together. Interactions between elements are what support Vermont's environment and are essential for long-term conservation of Vermont's biological diversity and natural heritage.

Each of these landscape elements is described below, and a map shows the areas identified as "priority" and "highest priority" for each. The highest priority areas are those that are critical for maintaining an ecologically functional landscape. The priority areas are also important but there is more flexibility available for conserving ecological function within these areas. The highest confidence in maintaining an ecologically functional landscape will be achieved by conservation of both priority levels for all of these landscape level elements.

The following maps of landscape level elements identify a large percentage of Vermont's lands and waters for conservation priority. We are highly confident that ecological functions must be maintained within these areas if Vermont is to have an ecologically functional landscape into the future.

Conserving Ecological Function

It is important to note that the goal for all of these areas is to maintain the ecological functions provided by that landscape element. For example, the goal for Interior Forest Blocks is to maintain the unfragmented, interior forest of these areas that provides critical habitat for many species of plants and animals. There is considerable leeway on what can happen within a forest block and still maintain interior forest function. For example, most forest management activities are compatible with maintaining the long-term interior forest functions for these blocks, providing these activities are thoughtfully planned.

Many tools can be used to achieve the overall goal of retaining ecological function. With approximately 80% of Vermont's land privately-owned, management and stewardship of private lands will be an essential path to success. Other tools include conservation easements, local planning and zoning, state regulations, and ownership by a state or federal agency or a private conservation organization. This document and these maps do not provide the detail as to which of these tools are best suited to specific places, but there are recommendations for further prioritization filters that users can apply to make these decisions.

Each section below provides guidelines on what is needed to maintain ecological functions for that element.

Landscape Element Descriptions and Maps

Interior Forest Blocks

Definition: Areas of contiguous forest and other natural communities and habitats (such as wetlands, ponds, and cliffs) that are unfragmented by roads, development, or agriculture. Forest blocks were identified, mapped, and ranked by Sorenson and Osborne (2014).

Ecological Function: Forest blocks provide many ecological and biological functions critical for protecting native species and the integrity of natural systems (Austin et al. 2004), including:

- Supporting natural ecological processes such as predator-prey interactions and natural disturbance regimes;
- Helping to maintain air and water quality and flood resilience;
- Supporting the biological requirements of many plant and animal species, especially those that require interior forest habitat or require large areas to survive;
- Supporting viable populations of wide-ranging animals by allowing access to important feeding habitat, reproduction, and genetic exchange; and
- Serving as habitat for source populations of dispersing animals for recolonization of nearby habitats that may have lost their original populations of those species.

In addition, large, topographically diverse forest blocks will allow many species of plants and animals to shift to suitable habitat within a forest block in response to climate change within the next century without having to cross developed areas to other forest blocks (Beier 2012).

Priority Areas for Maintaining an Ecologically Functional Landscape:


These are highly ranked forest blocks from all biophysical regions that provide important interior forest habitat and provide ecological support to the highest priority Forest Interior Blocks.

Highest Priority: These are the largest and/or highest ranked forest blocks from all biophysical regions that provide the foundation for interior forest habitat and associated ecological functions.

Guidelines for Maintaining Ecological Function: The primary goal is to maintain the interior forest conditions that forest blocks provide by avoiding permanent interior forest fragmentation resulting from development. Limited development on the margins of existing large forest blocks may not have significant adverse effects as long as it does not reduce connectivity between blocks and does not encroach into the forest block interior. Forest management that maintains forest structure within the block and results in a distribution of all age classes is compatible with maintaining interior forest conditions over the long term.

Further prioritization for conservation could be achieved using the following filters:

1. Areas also mapped as highest priority Connectivity Blocks.
2. Forest blocks with high “total weighted block scores” or high scores for any of the 11 individual biological or physical diversity factors (Sorenson and Osborne 2014).
3. Presence of rare or Vermont-responsibility geophysical settings, such as calcareous bedrock or clay soils.
4. High score in regional resilient sites analysis (Anderson et al. 2012).
5. High score for forest productivity (Vermont Land Trust 2007).



Map 1. Interior Forest Blocks showing Priority and Highest Priority blocks.

Connectivity Blocks

Definition: Landscape connectivity refers to the degree to which blocks of suitable habitat are connected to each other (Noss and Cooperrider 1994). Connectivity Blocks are the network of forest blocks that together provide terrestrial connectivity at the regional scale (across Vermont and to adjacent states and Québec) and connectivity between all Vermont biophysical regions. There is a high level of connectivity within individual forest blocks. The proximity of one forest block to another, the presence of riparian areas, and the characteristics of the intervening roads, agricultural lands, or development determine the effectiveness of the network of Connectivity Blocks in a particular area.

Ecological Function: A network of Connectivity Blocks allows wide-ranging animals to move across their range, allows animals to find suitable habitat for their daily and annual life needs, allows young animals to disperse, allows plant and animal species to colonize new and appropriate habitat as climate and land uses change, and contributes to ecological processes, especially genetic exchange between populations (Austin et al. 2004). Maintaining the landscape connectivity function requires both Connectivity Blocks and Riparian Areas for Connectivity, especially in highly fragmented areas of Vermont. There is general agreement among conservation biologists that landscape connectivity and wildlife corridors can mitigate some of the adverse effects of habitat fragmentation on wildlife populations and biological diversity (Beier and Noss 1998; Noss and Cooperrider 1994; Haddad et al. 2003; Damschen et al. 2006). Specifically, climate change adaptation is enhanced if the long distance movements of plants and animals is supported by a combination of short movements within large, topographically diverse forest blocks and short corridor movements between forest blocks (Beier 2012).

Priority Areas for Maintaining an Ecologically Functional Landscape:

These are the forest blocks that provide a major supporting connectivity function for the “backbone” of highest priority Connectivity Blocks. They also provide alternative pathways for connectivity, as redundancy is a critical safeguard in ensuring the long term effectiveness of the connectivity network.


Highest Priority: The terrestrial “backbone” of forest blocks is a subset of all Connectivity Blocks that provides connectivity to all biophysical regions. The “backbone” incorporates the spines of the major mountain ranges, connections outside Vermont to unfragmented habitat, and anchor blocks in fragmented biophysical regions based on abundant known occurrences of rare species and significant natural communities. Small forest blocks are included at pinch-points in the connectivity network as they are critical stepping stones.

Guidelines for Maintaining Ecological Function: Similar to Interior Forest Blocks, it is important to maintain the interior forest conditions in Connectivity Blocks by avoiding permanent interior forest fragmentation resulting from development. Connectivity within forest blocks will remain high if they remain unfragmented. For Connectivity Blocks it is also critically important to maintain or enhance the structural and functional connectivity that occurs on the margins of


these blocks where they border other blocks. This can be accomplished by maintaining forest cover along the margins and by limiting development in these areas of block-to-block connectivity.

Further prioritization for conservation could be achieved using the following filters:

1. Blocks that serve as stepping-stones in fragmented regions.
2. Anchor blocks are the largest blocks in the network and these need permanent conservation of their cores and the margins that border other blocks in order to secure the connectivity function.
3. Pinch points or bottle-necks in the connectivity network where animal movement or connectivity is narrowed due to adjacent development or fragmentation.



Map 2. Connectivity Blocks showing Priority and Highest Priority blocks.



Map 3. Connectivity Blocks and Riparian Corridors showing how the two landscape elements function together to provide connectivity in the fragmented Champlain Valley.

Surface Waters and Riparian Areas

Definition: The network of all lakes, ponds, rivers, and streams, their associated riparian zones and valley bottoms in which geophysical processes occur, and their connections to groundwater.

Ecological Function: Vermont's rivers, streams, lakes, and ponds provide vital habitat for a rich assemblage of aquatic species, including fish, amphibians, reptiles, invertebrates (e.g., insects, mussels, snails, worms, freshwater sponges), and plants. This represents an enormous contribution to Vermont's biological diversity. The ecological integrity of an aquatic system is dependent on the condition of the watershed in which it occurs, but is also critically tied to the condition of the riparian area adjacent to the stream or pond. For stability, rivers and streams must have access to their floodplains and freedom to meander within their valley bottoms or river corridors. Naturally vegetated riparian areas provide many significant ecological functions, including stabilizing shorelines against erosion, storage of flood waters, filtration and assimilation of sediments and nutrients, shading of adjacent surface waters to help moderate water temperatures, and direct contribution of organic matter to the surface water as food and habitat structure. Riparian areas are also very essential habitat for many species of wildlife that are closely associated with the terrestrial and aquatic interface, including mink, otter, beaver, kingfisher, spotted sandpiper, and wood turtle. The shorelines and riparian areas of rivers and lakes support floodplain forests, several other rare and uncommon natural communities, and many species of rare plants and animals. In addition to these ecological functions that are tied to aquatic systems, the linear network of riparian areas provides a crucial element of landscape connectivity for plant and animal movement in response to climate change (Beier 2012). Although many riparian areas and river corridors are highly altered by agriculture, roads, and urbanization, the risk of flooding serves as a natural deterrent for future development. Riparian areas also respond rapidly to restoration efforts (Beier 2012).

Priority Areas for Maintaining Ecologically Functional Landscape:

All of the aquatic network of lakes, ponds, rivers, and stream and the valley bottoms in which the rivers and streams occur; to be conserved or managed in such a way as to achieve full functioning of all natural processes.

Highest Priority: All of the aquatic network of lakes, ponds, rivers, and streams and the valley bottoms in which the rivers and streams occur, excluding developed land and including the Vermont hydrography layer and a buffer that is proportional to stream order.


Guidelines for Maintaining Ecological Function: Restoration is needed in order for Surface Waters and Riparian Areas to provide full ecological functions. Specifically, river channel equilibriums need to be maintained or restored. Natural vegetation should be maintained or restored in undeveloped riparian areas of rivers, streams, lakes, and ponds of adequate width to maintain water quality, stabilize shorelines, provide shade and biological support for aquatic systems, maintain biological diversity, and provide functional connectivity, both aquatic and terrestrial.

Further prioritization for conservation could be achieved using the following filters:

1. River Corridors as mapped by Vermont Department of Environmental Conservation (DEC).
2. Priority reaches as identified by DEC geomorphic assessments.
3. Priority reaches as identified by Flood Resilience Coarse Screen (Schiff et al. 2015).
4. Best Lakes Analysis, Lakes and Ponds Management and Protection Program, VT DEC.
5. Impaired stream reaches targeted for restoration and reference stream reaches targeted for conservation, VT DEC.



Map 4. Priority Surface Waters and Riparian Areas for Vermont



Map 5. Surface Waters and Riparian Areas for the South Burlington, Essex, and Williston area.

Riparian Areas for Connectivity (Riparian Corridors)

Definition: The connected network of riparian areas in which natural vegetation occurs, providing natural cover for wildlife movement and plant migration.

Ecological Function: In addition to supporting the integrity of the lakes, ponds, rivers, and streams that they border, naturally vegetated riparian areas are especially important for providing cover for wildlife movement and other important wildlife habitat, such as nesting habitat for birds. Many wildlife species use riparian corridors for travel to find suitable habitat to meet their life requisites, but certain species are almost entirely restricted to riparian areas, including mink, otter, beaver, and wood turtle. The linear nature of riparian areas contributes to their function as movement corridors for wildlife. Roads, development, and agricultural lands fragment the Vermont landscape. The combination of Riparian Areas for Connectivity and Connectivity Blocks provide the best available paths for connectivity across the landscape, especially in highly fragmented areas of Vermont.

Priority Areas for Maintaining an Ecologically Functional Landscape:


All of the aquatic network of lakes, ponds, rivers, and stream and the valley bottoms in which the rivers and streams occur; to be conserved or managed in such a way as to achieve full functioning of all natural processes.

Highest Priority: All of the aquatic network of lakes, ponds, rivers, and streams and the valley bottoms in which the rivers and streams occur, excluding developed land and agricultural land and including Vermont hydrography layer and a buffer that is proportional to stream order.

Guidelines for Maintaining Ecological Function: Restoration is needed to provide a fully functioning network of riparian areas that support connectivity. Restoration of natural vegetation is needed for river and stream shorelines where it does not exist now, and especially in riparian areas that provide the best available terrestrial connectivity between relatively isolated Connectivity Blocks. The width of naturally vegetated riparian areas needed to provide riparian connectivity varies from 100 feet or less on some small streams (50 feet each side) to 600 feet or more (300 feet on each side) for larger rivers or riparian areas that span long distances of otherwise unsuitable habitat.

Further prioritization for conservation could be achieved using the following filters:

1. A 300 foot buffer on rivers, streams, lakes, and ponds, which provides functional connectivity for many wildlife species.
2. Narrower buffers can provide functional connectivity for some species, such as mink.
3. A specific set of riparian areas that connect blocks in fragmented regions such as the Champlain Valley.
4. Presence of rare species or significant natural communities.



Map 6. Riparian Areas for Connectivity (Riparian Corridors) and Priority Surface Water and Riparian Areas in the vicinity of Ferrisburgh, Pantton, and Vergennes.

Physical Landscape Diversity Blocks

Definition: A set of forest blocks and other areas of natural vegetation that include physical landscape diversity features that are either rare in Vermont or under-represented in the land and water areas identified as highest priority for Interior Forest Blocks, Connectivity Blocks, and Surface Waters and Riparian Areas. The Physical Landscape Diversity Blocks complement the other block types and riparian areas in order to more fully represent the full spectrum of physical landscape diversity that is important for an ecologically functional landscape. Physical landscape diversity is represented in this conservation design by rare Land Type Associations (Ferree and Thompson 2008) and Ecological Land Units stratified by elevation, adapted from Ferree and Anderson (2008).

Ecological Function: Physical landscapes (often referred to as enduring features) are the parts of the landscape that resist change. They are the hills and valleys, the underlying bedrock, and the deposits left behind by glaciers. They remain largely unchanged when changes in land cover and wildlife occur, as plants and animals move, and even as the climate changes. However, these physical landscapes cannot continue to drive ecological processes or support plants, animals, or natural communities if they are developed or otherwise significantly altered by human activities.

If nature is likened to a dramatic play, it's possible to think of the enduring features as the stage and the individual species as the actors. The play is the natural communities, habitats and species that occur in a given place at a given time, but regardless of the action, the stage does not change. The importance of "conserving nature's stage" is that we can be much more confident in our ability to conserve biological diversity and maintain a functional landscape into the future, with the capacity to adapt and be resilient to climate change, if all elements of physical landscape diversity are represented in the conservation design (Anderson & Ferree 2010; Beier and Brost 2010; Beier et al. 2015).

Priority Areas for Maintaining an Ecologically Functional Landscape:

All are in the highest priority category.


Highest Priority: All of the identified forest blocks and other areas of natural vegetation which contain physical landscape diversity features that are either rare, under-represented in other block types or the riparian areas, or for which Vermont has a regional responsibility (especially calcium-rich bedrock).

Guidelines for Maintaining Ecological Function: Maintain or restore natural vegetation and limit development. Forest management that maintains forest structure within and results in a distribution of all age classes is very compatible with maintaining the physical landscape diversity functions.

Further prioritization for conservation could be achieved using the following filters:

1. Blocks containing the rarest landscape diversity features.
2. Blocks containing landscape features that are underrepresented on conserved lands.

3. Blocks containing limestone or other calcareous bedrock types.
4. Clay soil lands that are marginal for agricultural uses and/or adjacent to other clayplain fragments.



Map 7. Physical Landscape Diversity Blocks showing Highest Priority blocks and other areas of natural vegetation.

Wildlife Road Crossings

Definition: A section of road that crosses a wildlife corridor where the adjacent landscape quality and permeability are high, usually because the road is adjacent to a forest block, and the road is the primary impediment to animal movement. Likely wildlife road crossings are identified statewide in the habitat block project (Sorenson and Osborne 2014).

Ecological Function: Wildlife corridors (also referred to as wildlife connecting habitat) are lands and waters that connect larger patches of habitat together within a landscape and allow the movement, migration, and dispersal of animals and plants (Austin et al. 2004). Roads represent a barrier to wildlife movement and dispersal of many other species, including some plants. Sections of roads that have suitable habitat on both sides are more likely to allow wildlife movement and dispersal of other species and, therefore, these sections of roads are critical components of maintaining or enhancing an interconnected, ecologically functional landscape. Wildlife road crossings that provide connectivity over or under roads are critically important between adjacent forest blocks and along linear riparian area networks.

Priority Areas for Maintaining an Ecologically Functional Landscape:


The same as highest priority, but also including those wildlife road crossings rated moderate to high (Sorenson and Osborne 2014) that provide connections to areas adjacent to but not included within the network of Connectivity Blocks and Interior Forest Blocks.

Highest Priority: All of the wildlife road crossing segments with a moderate to high score (Sorenson and Osborne, 2014) that occur between Connectivity Blocks, Interior Forest Blocks, and within Surface Water and Riparian Areas.

Guidelines for Maintaining Ecological Function: Structural connectivity across identified wildlife road crossings is provided by the presence of forest cover, wetlands, or other natural habitat. Maintaining this natural cover is critical to maintaining structural connectivity. Maintaining or restoring natural vegetation on both sides of identified road crossing segments will maximize the effectiveness of the road crossing for connectivity. Forest management that maintains forest cover adjacent to the road is compatible with this function. Roadside development that further restricts animal movement is detrimental to connectivity and in some cases may result in a particular road crossing ceasing to provide connecting function. Road and highway structures that allow or promote animal movement, such as bridges and oversized culverts, and limiting the use of structures that impede movement, such as fences and roadside barriers, are effective in promoting wildlife passage across roads.

Further prioritization for conservation could be achieved using the following filters:

1. Areas with documented successful wildlife crossings (such as Critical Paths).
2. Known pinch points in the connected network of blocks and riparian areas.



Map 8. Wildlife Road Crossings for the Waterbury-Stowe area and including Interior Forest Blocks, Connectivity Blocks, Surface Waters and Riparian Areas.

Putting it All Together: The Ecologically Functional Landscape


Maintaining or enhancing an ecologically functional landscape in Vermont depends on conservation of all the landscape elements together: Interior Forest Blocks; Connectivity Blocks; Surface Waters and Riparian Areas; Riparian Areas for Connectivity; and Physical Landscape Diversity Blocks. It is the specific functions of each landscape level element and the complementarity of these landscape level elements functioning together that are critical for long term conservation of much of Vermont's biological diversity and natural heritage.

The following map shows the ecologically functional landscape conservation design, with all five landscape elements included.

It is our hope that this information will inform land management, local planning, and land conservation decisions throughout Vermont. Private landowners, municipalities, state agencies, and conservation organizations should find this information helpful as we all work together for a vibrant and healthy Vermont, now and into the future.

Further Information

For more detailed information on this project, its data layers, and its potential applications, please visit (BioFinder website: <http://biofinder.vermont.gov/>) or contact (Jens Hilke; jens.hilke@vermont.gov). Any of the authors can also help with interpretation of the information.



Map 9. The Highest Priority and Priority portions of the Ecologically Functional Landscape, including Interior Forest Blocks, Connectivity Blocks, Surface Waters and Riparian Areas, Riparian Areas for Connectivity, and Physical Landscape Diversity Blocks.

Literature Cited

- Anderson, M.G., M. Clark, and A. Olivero Sheldon. 2012. Resilient sites for terrestrial conservation in the Northeast and Mid-Atlantic region. The Nature Conservancy, Eastern Conservation Science. 168 pp.
- Anderson, M. G. and C. E. Ferree. 2010. Conserving the stage: climate change and the geophysical underpinnings of species diversity. *PLoS ONE* 5(7): e11554.
- Austin, J. M., C. Alexander, E. Marshall, F. Hammond, J. Shippee, E. Thompson, and Vermont League of Cities and Towns. 2004. Conserving Vermont's natural heritage: a guide to community-based conservation of Vermont's fish, wildlife, and biological diversity. Vermont Fish and Wildlife Department and Agency of Natural Resources, Waterbury.
- Beier, P. 2012. Conceptualizing and designing corridors for climate change. *Ecological Restoration* 30(4): 312-319.
- Beier, P. and B. Brost. 2010. Use of land facets to plan for climate change: conserving the arenas, not the actors. *Conservation Biology* 24:701-710.
- Beier, P. and R. F. Noss. 1998. Do habitat corridors provide connectivity? *Conservation Biology* 12:1241-1252.
- Beier, P., M. L. Hunter, and M. Anderson (editors). 2015. Special Section: Conserving Nature's Stage. *Conservation Biology* 29(3): 613-617.
- Damschen, E. I., N. M. Haddad, J. L. Orrock, J. J. Tewksbury, and D. J. Levey. 2006. Corridors increase plant species richness at large scales. *Science* 313:1284-1286.
- Ferree, C. and E. Thompson. 2008. Land Type Associations Descriptions for Vermont. Vermont Department of Forests, Parks, and Recreation.
- Ferree, C. and M.A. Anderson. 2008. Ecological Land Units. Version 11/2008. The Nature Conservancy Eastern Conservation Science Office, Boston, MA.
- Haddad, N. M., D. R. Bowne, A. Cunningham, B. J. Danielson, D. J. Levey, S. Sargent, and T. Spira. 2003. Corridor use by diverse taxa. *Ecology* 84:609-615.
- Haufler, J.B., C.A Mehl, and G.J Roloff. 1996. Using a coarse-filter approach with species assessment for ecosystem management. *Wildlife Society Bulletin* 24: 200-208.
- Hunter, M. L. 1991. Coping with ignorance: The coarse filter strategy for maintaining biodiversity. Pages 266-281 in K.A. Kohm, ed. *Balancing on the Brink of Extinction*. Island Press. Washington, D.C.
- Hunter, M.L., G.L. Jacobson, Jr., and T. Webb. 1988. Paleoecology and the coarse-filter approach to maintaining biological diversity. *Conservation Biology* 2(4): 375-385.
- Jenkins, R.E. 1985. The identification, acquisition, and preservation of land as a species conservation strategy. Pages 129-145 in R.J. Hoage ed. *Animal extinctions*. Smithsonian Institution Press. Washington, DC.
- Jenkins, R.E. 1996. Natural heritage data center network: managing information for managing biodiversity. Pages 176-192 in R.C. Szaro and D.W. Johnston eds. *Biodiversity in managed landscapes: theory and practice*. Oxford University Press. New York.
- Molina, R., Horton, T. R., Trappe, J. M., and Marcot, B. G. 2011. Addressing uncertainty: how to conserve and manage rare or little-known fungi. *Fungal Ecology* 4(2): 134-146.
- National Council for Air and Stream Improvement, Inc. (NCASI). 2004. Managing elements of biodiversity in sustainable forestry programs: Status and utility of NatureServe's information resources to forest managers. Technical Bulletin No. 885. Research Triangle Park, N.C.: National Council for Air and Stream Improvement, Inc. Accessed March 11, 2009 at: http://www.natureserve.org/library/ncasi_report.pdf
- Noss, R. F. 1987. From plant communities to landscapes in conservation inventories: a look at the Nature Conservancy (USA). *Biological conservation* 41:11-37.

- Noss, R.F. and A.Y. Cooperrider. 1994. Saving nature's legacy. Defenders of Wildlife. Island Press. Washington, D.C.
- Panzer, R., and Schwartz, M. W. 1998. Effectiveness of a Vegetation-Based Approach to Insect Conservation. *Conservation Biology* 12(3): 693-702.
- Poiani, K.A., B.D. Richter, M.G. Anderson, and H.E. Richter 2000. Biodiversity conservation at multiple scales: functional sites, landscapes, and networks. *BioScience* 50(2): 133-146.
- Roman, J. and J. Erickson. 2015. Economics of Conservation in Vermont. Gund Institute for Ecological Economics, Rubenstein School of the Environment and Natural Resources, University of Vermont, Burlington. 36 pp.
- Schiff, R., J. C. Louisos, E. Fitzgerald, J. Bartlett, and L. Thompson. 2015. The Vermont River Sensitivity Coarse Screen. Prepared by Milone & MacBroom for the Vermont Land Trust and its conservation partners, Waterbury, VT.
- Schulte, L. A., Mitchell, R. J., Hunter Jr, M. L., Franklin, J. F., Kevin McIntyre, R., & Palik, B. J. 2006. Evaluating the conceptual tools for forest biodiversity conservation and their implementation in the US. *Forest Ecology and Management* 232(1): 1-11.
- Shuey, J. A., Metzler, E. H., and Tungesvick, K. 2012. Moth communities correspond with plant communities in Midwestern (Indiana, USA) sand prairies and oak barrens and their degradation endpoints. *The American Midland Naturalist* 167(2): 273-284.
- Sorenson, E. and J. Osborne. 2014. Vermont Habitat Blocks and Habitat Connectivity: An Analysis using Geographic Information Systems. Vermont Fish and Wildlife Department, Montpelier, Vermont. 48 pp.
- United States Forest Service, USDA. 2004. Coarse filter/ fine filter planning approaches to the conservation of biological diversity. Accessed February 26, 2015 at: <http://www.fs.fed.us/emc/nfma/includes/coursefilter.pdf>
- Vermont Land Trust. 2007. Forest Project Prospecting: Developing a GIS Model to Identify Conservation Priorities. Unpublished document.



Mapping Vermont's Natural Heritage

A Guide for Municipal and Regional Planners in Vermont

Vermont Fish & Wildlife Department
2015

Acknowledgements

Thank you to all those who have worked on, edited, or added to this information packet at some point along the way. This could not have been possible without the following:

- Jamey Fidel and Kate McCarthy of the Vermont Natural Resources Council
- Jim Ryan and Steve Sinclair from Vermont Agency of Natural Resources
- The *many* town planners and conservation commissioners who have shared questions, ideas, and requests for specific information about what they would like to see in such a packet
- The partners of the Vermont Biodiversity Project, including:
 - The Nature Conservancy
 - Vermont Land Trust
 - United States Environmental Protection Agency
 - United States Fish and Wildlife Service
 - United States Department of Agriculture, Natural Resource Conservation Service
 - United States Department of the Interior, Forest Service (Green Mountain National Forest)
 - Vermont Agency of Natural Resources
 - Middlebury College
 - University of Vermont
 - The Orton Family Foundation
 - Sweet Water Trust

Funding for the development of this guide comes from the State Wildlife Grants program of the Vermont Fish & Wildlife Department.



Photo by Tom Rogers

Table of Contents

Acknowledgements	2
Background and Purpose	4
Why Maps?	5
Getting the Most Out of The Maps.....	7
Online Mapping Tools: The Vermont Agency of Natural Resources Atlas and Biofinder	7
What is the Natural Resources Atlas ?	7
What is Biofinder ?	8
How might these be used for municipal planning?	9
How do I get started?.....	9
The Maps.....	13
MAP 1: Conserved Lands and Water Resources.....	14
Mapped Layer #1: The Conserved Lands Database	16
Mapped Layer #2: Wetlands	18
Mapped Layer #3: Vernal Pools.....	21
Mapped Layer #4: Vermont Hydrography (Surface Water) Dataset.....	22
MAP 2: Land Cover	25
MAP 3: State and Regional Biodiversity Resources	29
Biofinder Components	31
Tiered Contribution to Biodiversity.....	32
MAP 4: Wildlife Resources At a Town Scale	34
Mapped Layer #1: Habitat Blocks.....	36
Mapped Layer #2: Wildlife Road Crossings.....	37
MAP 5: Fine Scale Wildlife Resources.....	41
Mapped Layer #1: Habitat Block.....	43
Mapped Layer #2: Wildlife Road Crossings	44
Mapped Layer #3: Deer Wintering Areas.....	47
Mapped Layer #4: Rare, Threatened, and Endangered Species	48
MAP 6: Physical Features.....	52
Mapped Layer #1: Rare Physical Landscapes.....	55
Mapped Layer #2: Representative Physical Landscapes.....	55
Advanced Natural Resources Inventory.....	57
Bibliography	59



Background and Purpose

Vermont is known for its beautiful forests and agricultural lands, mountain streams and scenic landscapes. This is the “Green Mountain State,” where residents often claim a love of rural places and where visitors come to ski, hunt, hike, and see the scenery. Our landscape drives our economy and is largely what makes our state so special.

Surveys support these claims. Across the nation, only Alaska outranked Vermont for participation in wildlife-related recreation—fishing, hunting, or wildlife watching—in a 2011 survey (US Dept. of Interior). In the same survey, Vermont ranked first for wildlife watching activities, with 53 percent of residents participating—over half our population—and it was estimated that more than \$704 million was spent in Vermont on fish-and-wildlife-based recreation. In 2015, a public attitude survey found that 83% of respondents agree that land use and development should be restricted to protect fish and wildlife and that 81% would like to see wildlife habitat protected even if it reduces the land use options of some landowners and developers (Duda et al). In Vermont, we like our wildlife and we want to see their continued presence on the landscape.

This strong connection Vermonters feel to wildlife has helped inform the state’s *Wildlife Action Plan*, Vermont’s all-species conservation plan. The Action Plan focuses on the conservation of the landscapes and habitats needed by Species of Greatest Conservation Need—that is, wildlife with declining or threatened populations and those about which we know very little. Among the needs and strategies identified by the plan, municipal-scale planning and conservation has emerged as a critical need.

With over 80% of the state’s land in private ownership and the majority of land use and development decisions made at the local or regional level, the protection of Vermont’s Species of Greatest Conservation Need, their habitats, and the ecological processes that support them is firmly in the hands of landowners, municipal governments, and regional planning groups. At the

same time, municipal planners must balance these wildlife needs with countless other goals, and prioritization of diverse goals and needs can be tricky.


This guide was created to support municipal planners in achieving their goals for the protection of wildlife habitats within town boundaries. In it, we share resources developed and lessons learned by many agencies and organizations throughout Vermont, combining background information about our natural landscape, planning strategies, and natural resources maps tailored to individual towns. For those wishing to dig deeper, we have provided a list of additional resources you may find helpful. Our goal is to provide planners with the knowledge and tools necessary to make wildlife-related planning decisions in their own town or region. If a community can identify and conserve the most important wildlife resources on *its own* landscape, it will also achieve goals set forth in Vermont's Wildlife Action Plan and thereby aid with the conservation of wildlife on a state and even regional scale.

Why Maps?

In 2008, the Vermont Supreme Court struck down a South Burlington zoning ordinance aimed at protecting a variety of natural resource values. The court determined that the ordinance was too vague to effectively determine what “protection” of the natural resources listed should entail, thereby making it unenforceable. Because the South Burlington ordinance was similar to those written into the bylaws of many towns throughout the state, this ruling is a call to action for all towns wishing to protect their local natural resources. If towns want their plans and bylaws to be legally defensible, they must include clear, specific, and consistent standards that define exactly what types of development are allowed and prohibited in any given area (Appeal of JAM Golf, LLC, 2008 VT 110).

A map is the first step in creating such clear, specific, and consistent standards. Before you can effectively plan, you first need to determine exactly what resources you have to work with. The better your information, the more easily you can prioritize, and the more clarity you can provide. A map is essentially an inventory of one or several components of the landscape, and this guide highlights a series of six maps created specifically to feature your town's ecological, biological, and physical resources. When combined together, these maps become even more powerful, showing how each individual map dataset relates to—and in some cases *influences*—every other.

Of course, every map also has limitations. Maps are static images, and yet they are intended to represent a continuously changing landscape. They are also intended for use at a particular scale and can become inaccurate when used at other scales. Imagine, for example, a map of the lakes present in the entire state of



Vermont. At the state level, each lake appears to be accurately placed. However, if you were to zoom in on that map, magnifying everything within the boundaries of your hometown, you may find that the “boundaries” of the lake are in fact off by 30 feet. When examining that map at the state scale, those 30 feet are unsubstantial. To a homeowner whose home is depicted as partially underwater, those 30 feet matter!

In this packet, we therefore explain in some detail the appropriate scales, data sources, and intended purpose of each map. We encourage you to read these descriptions thoroughly before including these maps or the data they contain in conservation planning efforts. When used appropriately, however, this information can open up new ways of seeing your community’s landscape and the many natural resources located there!




Photo by Dennis Curan

Getting the Most Out of The Maps

Maps and inventory may be the basis of conservation planning, but there are clearly several steps between looking at a map and having a conservation plan. In this guide, you will find many links to [*Conserving Vermont's Natural Heritage*](#), published in 2004 by the Vermont Fish and Wildlife Department and Agency of Natural Resources, and [*Community Strategies for Vermont's Forests and Wildlife*](#), a 2013 document by Vermont Natural Resources Council. Together, these books provide a background of the natural heritage features found on the maps in this guide as well as explanations of a wide range of tools a community might use to protect these resources. We encourage you to read these books alongside this guide.

Additionally, Vermont's Agency of Natural Resources has developed several online mapping tools to allow anyone with an internet connection to explore a large compilation of the state's mapping data. Using this resource, you can recreate any of the maps you see here, mix and match data, and zoom in and out to different scales on any map. Furthermore, these online tools update their mapping layers frequently, allowing you to find the most up-to-date data available. A description of how to use these tools is below.



In addition to the above resources, we also recommend agreeing on a method for capturing the goals and values of your community and including local citizens in the planning process even before you begin. Because it will ultimately be up to your community to adopt the plans and strategies you propose, it is important to be transparent about your intentions. The *Community Heart and Soul Field Guide* outlines one method for involving your community in the planning process, published in 2014 by the Orton Family Foundation and available online at <http://www.orton.org/what-we-do/what-community-heart-soul>. When combined with the scientifically-based background information outlined in this guide and by natural resources professionals, this strategy can be a powerful way to connect with citizens in your community. Of course, all strategies are not for all towns, and this is just one of many possibilities!


Online Mapping Tools: The Vermont Agency of Natural Resources Atlas and Biofinder

Found online at <http://anrmaps.vermont.gov/websites/anra5/>.

What is the [*Natural Resources Atlas*](#)?

In recent years, digital mapping resources have become increasingly sophisticated, enabling generation of abundant landscape-based information that was previously unknown. Even with these mapping developments, however, many have remained relatively inaccessible to the public, requiring expensive software or technical training. The Vermont Agency of Natural Resources developed the Natural Resources Atlas to allow easy public access to mapping information, first by compiling diverse information sources into a single, user-friendly location and then by providing a free, publicly available mapping tool.


Similar to many modern mapping programs, the Atlas uses “GIS,” or a Geographical Information System that captures, stores, analyzes, and manages a diverse array of geographical information and allows it to be viewed simultaneously. In some ways, this process works just like taking physical maps, copying them onto transparent mylar, then laying one on top of another so that a location on one lines up with the same location on another. A user can look at multiple layers at once and add—or remove—data layers as needed. In other ways, GIS is much more sophisticated than a set of transparent maps because the software not only layers the maps on top of one another but also provides tools to analyze intersections between map layers. For example, a user can see how much conserved land is covered by wetland, find locations where there is farmland within 100 feet of a river, or identify places that are mapped *both* as large, contiguous forest *and* deer wintering yard. The Atlas also allows users to generate reports, make notes, and print maps.



What is Biofinder?

Like the Natural Resources Atlas, BioFinder is an online mapping tool, but it is intended for a much more specific purpose: exploring the distribution and richness of biodiversity across Vermont’s landscape. It was designed specifically to aid those who might use natural heritage information for planning or development purposes. Also created by the Vermont Agency of Natural Resources and its partners, BioFinder maps 21 components that are known to contribute significantly to biological diversity. These components represent various scales of terrestrial and aquatic biological, ecological, and natural heritage features, and you can navigate directly to an area of interest to you to see what is present there.

When exploring BioFinder, you can choose to add each of the 21 components individually, or



there is a single layer that combines them. This “Tiered Contribution to Biodiversity” dataset is derived from the overlap of the 21 components, taking into account that some layers contribute more heavily than others to biodiversity. The result is displayed using six ranked tiers, so you can easily see locations containing the highest and lowest overlap between component layers. Users generally find the “Tiered Contribution to Biodiversity” layer a good place to begin as it points to locations important for biodiversity; they can then view individual layers to understand *why* different locations are ranked as they are.

BioFinder is also set up to create printable reports of the landscape components or the contribution to diversity of a chosen area such as the report pictured on the right.

What’s the Difference Between the Atlas and Biofinder? When would I use each?

For most mapping purposes, the Natural Resources Atlas is the tool to use. It was created as a substitute for many of the complex mapping programs used by professional cartographers, so it can be used for a wide variety of purposes.

However, Biofinder can be an extremely useful tool for those working on conservation planning efforts. Because the components mapped by BioFinder are known to contribute significantly to the conservation value of a place, planners may find this resource a useful place to begin. BioFinder gives a first-glimpse snapshot of ecologically important locations across the landscape, complete with printed reports of a chosen geographic region. The tool is also a bit simpler since the site has a focus on biodiversity specifically, so a user new to online mapping may find it easier to begin with BioFinder and then go to the Natural Resources Atlas if additional information is needed.

Functionally, the tools work in a very similar way. A user who learns to use one tool will likely understand the second tool very quickly. For this guide, all of the maps can be recreated using the Natural Resources Atlas—even Map 3, which uses data generated by Biofinder.

How might these be used for municipal planning?

As you are aware, a municipal planner must take into consideration not only features of the town’s natural landscape but also the values of citizens in the community. Because both may vary drastically from one town to the next, there is no catch-all set of maps that can display the layers that will be most useful for every town in Vermont. These mapping tools therefore allow individuals in each town to decide which information is most valuable to compile, and they can choose from a large database of available information stored in a single place.

As you look at the maps in this packet, keep in mind that you can “play” with any of them using the Atlas. We believe these maps to be a great place to start, but you may find it useful to display different sets of information together on a single map, add additional layers, zoom in to see details that may impact a single parcel or section of town, or zoom out to see how your town’s features fit into the context of the surrounding landscape.

How do I get started?

The **Natural Resources Atlas** is found online at
<http://anrmaps.vermont.gov/websites/anra5/>.

BioFinder is found at
<http://biofinder.vt.gov>.

Mapping Tip

If you're new to online mapping tools, we suggest starting with a YouTube orientation video created by the same folks who created the Natural Resources Atlas. They provide step-by-step visual instructions for using all the tools available in the Atlas:

http://www.youtube.com/watch?v=jqDr93vUC_8.

A similar YouTube video will orient you to using BioFinder:

<https://www.youtube.com/watch?v=Rpkn3laCa0o>

In fact, there are quite a few instructional videos produced by the Vermont Agency of Natural Resources' GIS mapping team. All are available on this YouTube channel:

<https://www.youtube.com/user/vtanrgis>.


When updates are available, this is also where they will be posted.

If you run into questions once you get started, try visiting the *Frequently Asked Questions* page for the Atlas: <http://anrmaps.vermont.gov/websites/anra/faq.html>.


The “help” tools within the application may also prove helpful.

Because there are instructional videos and other resources that already provide so much information (as described above), we won't use this packet as a “how to” guide. However, we do have a few tips specific to municipal planning:

Zoom to Town Tool: Because so much of what you will be doing occurs at the town level, you may find the “Zoom to Town” tool to be particularly useful. Once the Atlas is open, find the “Quick Tools” menu in the top, center of the page. Once you select the “Zoom to Town” tool from the dropdown menu, begin to enter your town's name in the box. After typing the first few letters, your town should pop up as a suggestion. Click on your town's name.



Themes: You may also find it handy to preload a theme. Themes are groups of map layers that are often used together for a particular purpose, allowing you to easily turn on a pre-chosen set of related maps. If you want to recreate the maps in this guide, for example, you can preload the “Conservation Planning Theme” into the Atlas. After loading the Atlas, click the  symbol at the bottom, left-hand corner of the page to view the “Layers” menu. Once this menu is visible, find the box that says “All Layers,” just below the “Layers” title at the top of the page. From the drop-down menu, select the “Conservation Planning.” You may find it handy to explore other pre-loaded themes on the Atlas, too. The “Tactical Basin Planning” and “New Neighborhood Planning” themes may be particularly useful for municipal planners. Other datasets are organized under each of ANR’s Departments.



Natural Resources Atlas Mapping Tip

To recreate any of the maps in this packet:

1. Open the atlas: <http://anrmaps.vermont.gov/websites/anra5/>.
2. Find your town by clicking “Quick Tools” in the top, center of the page. Use the “Zoom to Town” feature and enter your town’s name.
3. Click the  symbol at the bottom, left-hand side of the page to open the “Layers” menu.
4. Find the drop-down menu at the top of the page and select the “Conservation Planning” theme. Several new layers will appear in the sidebar.
5. One new layer is entitled “Conservation Planning.” Click on the “+” next to this layer, and boxes for Maps 1-6 appear underneath. These Maps contain the same layers and those appearing in this guide.

Navigation Aid: Throughout this document, tips for using the Natural Resource Atlas will appear in blue boxes like this one.

HTML5 and Microsoft Silverlight

A note for those wanting to maximize their online mapping capabilities

It is worth noting that when both mapping programs were created, they used the free plugin *Microsoft Silverlight*, which allowed for the programs' interactive capacity. As of 2015, the viewer is changing to *HTML5*, although the *Silverlight* version remains available for the foreseeable future. The *Silverlight* version actually has several features not offered by the new *HTML5* version—such as the ability to save a map with markup and re-open it on a different computer, which can be very handy for municipal planning—but it is being phased out because it is not supported by Chrome browsers and has limited functionality when accessed through Mozilla or Safari. The new version works on all browsers, including portable devices.

The links in this document all lead to the new *HTML5* version of the Atlas. However, because *Silverlight* allows access to features unavailable on *HTML5*, it is not being immediately discontinued and can be accessed at <http://anrmaps.vermont.gov/websites/anra>. Meanwhile, the *HTML5* version of BioFinder has not yet been released and is *only* available using Silverlight. This means that the first time you attempt to access BioFinder—or the *Silverlight* version of the Atlas—you will be prompted to download *Silverlight* and must do so before you can view the maps. To access full functionality of these programs through *Silverlight*, we recommend that you use Internet Explorer, although the program will work in all browsers except Chrome.

The Maps

There are many ways to look at a landscape, even when your focus is specific to conserving natural habitat. These six maps provide six different views. While all are zoomed to the scale of your town, the order of Maps 1-5 is designed to begin with an overview of your landscape—as though hovering in an airplane high above, looking down—then slowly add detail as you descend. Once you have gained knowledge from this up-close view, Map 6 flies you back up for a fresh look from afar.

The goal of all these maps is to provide information about your landscape that can help you to prioritize conservation goals for your town or region. We recognize that conservation planning can seem overwhelming, because there are so many possible places to start. It can also be difficult to balance wildlife needs with other community values.

The six sets of information provided on these maps are not themselves prioritized in any way. It is no more important to conserve the water resources in Map 1 than the rare or representative physical features of Map 6. Rather, these are six ways of looking at the landscape. Similarly, it is impossible to prioritize between the features mapped here at a state level, because the elements that contribute to a healthy, fully-functional landscape in one location may be completely different than in another location—not to mention that each community will contribute a different suite of values and other priorities that must be considered. However, you will likely find that after carefully examining the locations highlighted on each of the six maps, certain features begin to stand out within each town. In one town, it may be the vegetated connections along a wetland complex. In another, it may be the network of large habitat blocks linked by road crossings. As a town, see what stands out to you after you have gone through the entire guide.

Suggestion: Start by finding the map of your town at the beginning of each of the following sections. Have it handy and flip back and forth between that map and the interpretation that follows. If you're still looking for more information, try visiting the Agency of Natural Resources' Natural Resources Atlas online. In this packet, each map number contains tips on how to recreate the map in the Atlas, where you can view the map layers at different scales or add additional layers.

Once you get to know the maps, the second step in assembling inventory information is to gather local, site-specific information through on-the-ground inventories and interviews with knowledgeable residents. Some ideas for this kind of inventory are listed in the “Advanced Natural Resource Inventory” section.

MAP 1: Conserved Lands and Water Resources

This page will be replaced with MAP 1 specific to each town.




Photo by Eric Sorenson

MAP 1. Conserved Lands & Water Resources

This map has two main functions: to convey information on conserved lands, and to show where important water resources exist—streams, rivers, lakes, ponds, wetlands, and vernal pools. It is intended to provide a broad overview for your community.


The conserved lands map layer is intended to give an overview of the conserved lands in your community, both public and private. Because land is conserved by many different entities on an on-going basis, the information depicted here is not necessarily complete or up-to-date, but it can certainly help your town in making decisions about adding new conservation areas. The conserved lands data is most powerful when combined with other datasets—such as the water resources data shown here, or with data on wildlife movement or rare, threatened and endangered species. It can very interesting to see—at either a statewide or local scale—where the significant natural resources are in comparison with where the conserved lands are. Often the most important natural resources are not protected, but you can see whether this is the case in your community!

Water resources shown here include wetlands, rivers, streams, lakes, ponds, and vernal pools. These come from several distinct sources, each of which is described below. With the exception of confirmed vernal pools, information was assembled remotely¹—through the interpretation of aerial photos and other data—and is not intended to be accurate at a fine scale. However, it gives an excellent overall picture of where water appears on the landscape and can aid in planning for the protection of water resources.

¹ *The Water Resources layer was mapped by the Vermont Significant Wetlands Inventory, based on the National Wetlands Inventory.*

Natural Resources Atlas Tip:

If you try to recreate Map 1 in the [Natural Resources Atlas](#), your screen will look something like this: (*Example: Waterbury*)



Open the “Conservation Planning” Menu. Open the “+” next to “Conservation Planning” and click on “Map 1.” You can then select which layers of water resources and conserved land you would like to see on the screen.

Additional Layers:

Along with the Conserved Lands Database, you may find it useful to see land that is enrolled in the Use Value Appraisal Program. This layer is available through the Atlas, listed under the “Forests, Parks, and Recreation” heading. You may also want to see where particular natural features are in relation to conserved lands, such as Habitat Blocks or Rare, Threatened, and Endangered species (described in Maps 4 and 5).

Need help using the Natural Resources Atlas? Try the [“How Do I Get Started?”](#) section in the Introduction of this guide.

Mapped Layer #1: The Conserved Lands Database

What is the Conserved Lands Database?


The conserved lands dataset shows land that is permanently protected from development by buildings, paved roads, and most commercial infrastructure, either through conservation ownership or by virtue of a conservation easement. These data were first published by the University of Vermont Spatial Analysis Lab and developed by a partnership between many federal and state agencies and departments, the University of Vermont, and several Vermont nonprofits. Included in the map are all conservation lands owned by governmental entities at the local, state, and federal levels in Vermont, as well as land protected through conservation easements by Vermont’s nonprofit land trusts—such as The Nature Conservancy and the

Vermont Land Trust. A great attempt was made to include locally-owned town lands and the holdings of small land trusts, but users should understand that there are omissions in these areas.

The original vision for the Conserved Lands Database was to have it continually updated as the partners acquired new conservation land. However, time and money have proved limiting, so database updates have been more sporadic. Users of the database should realize that it may be a few years out of date at any given point in time.

Interpreting This Layer

While all lands in this dataset are permanently protected from development of some type, there are several classes of conserved lands that are not differentiated here. Some conserved lands are managed strictly as natural areas, with activities such as timber harvesting prohibited. Others are managed specifically *for* the production of timber and other resources but may not be developed. Others are active, working farms where normal farming activities are expected, but development is greatly restricted. In certain cases, conserved lands can be developed only for particular uses such as public recreation, as is often true with state and town parks.



The information in the Conserved Lands Database can be used at any scale larger than a few parcels, where precise boundaries are not important. Great effort was made to reconcile boundary discrepancies and mapping errors when this database was assembled. However, the quality of the maps that were digitized varied greatly. Much of the information came from sketch maps, deed descriptions, or old surveys, and these sources required a great deal of interpretation. The result is that no boundary line should be

considered precise or used to determine protection status on a fine scale. Any work of that sort would need on-the-ground presence with landowner permission. The maps also do not represent the conservation status of any parcels smaller than two acres, with the exception of Vermont Fish and Wildlife Department's fishing access areas.

Of course, permanently protected lands are not the only places that contribute to natural resources conservation. Practicing good land management or enrolling land in an established conservation incentives program can be considered conservation even without permanent protection, but these are not included on this map. As you conduct planning in your community, you may want to look further into land enrolled in Vermont's Use Value Appraisal Program ("Current Use"), the USFWS Partners for Fish and Wildlife Program, and the many incentive programs managed by the Natural Resources Conservation Service.

Please note that no inferences should be made about habitat quality or public access on conserved lands. Land may have been conserved to protect any number of different qualities: agricultural soils, views, community resources, natural areas, historic landmarks, water quality,

wildlife, or many other values. This includes large, public federal lands with publicly advertised recreational trails, and it *also* includes privately owned, 50-acre parcels with *no* public access. In using this map, please be careful to recognize and respect private property. Similarly, this map includes no information about management goals, though some public lands have management plans available.

How can a community use it?

Just like current areas of development are unlikely to change into natural areas, conserved land is unlikely to become developed. Because we know that wildlife populations are most likely to thrive if their habitats are interconnected and large, a community may want to consider the distribution of currently protected areas before planning areas of future development or conservation. From a natural resource protection perspective, it is generally best to expand upon prior investments in land conservation rather than to create a new block of conserved habitat, all else being equal. While habitat quality is not represented in this dataset, the size and interconnectedness of habitat is so important to wildlife abundance that simply having a parcel conserved elevates its general importance to resource planning.

As mentioned above, the conserved lands layer may be even more useful when combined with other datasets. For example, you might look at conserved lands alongside rare species or significant natural communities to create a snapshot of the resources already being conserved in a given area. A community may then be able to better prioritize the protection of additional natural heritage resources. Other useful map layers might include the wildlife habitats shown in Maps 4 and 5, Audubon's Important Bird Areas, The Nature Conservancy's Matrix Blocks (which show critical areas for forest interior species), or any other fine-scale resources.

Community Strategies for Vermont's Forests and Wildlife: Case Studies

There are many reasons why a family, individual, or group may want to conserve land, and every conservation decision has a unique story. [*Community Strategies for Vermont's Forests and Wildlife*](#) documents a few of these stories, found on page 28 of the book.

Mapped Layer #2: Wetlands

What Are Wetlands?

Wetlands are intermediate habitats between upland and aquatic ecosystems. Wetland communities include the vegetated, shallow-water margins of lakes and ponds, the seasonally flooded borders of rivers and streams, and an amazing diversity of topographic settings across the landscape, including basins, hillside seeps, and wet flats.

Although many definitions have been developed for the term and concept of "wetland," we consider wetlands to have three basic characteristics. First, all are inundated by or saturated with water for varying periods during the growing season. Second, they contain wet or hydric soils, which develop in saturated conditions. Finally, they are dominated by plant species that are adapted to life in saturated soils. Methodologies for identifying and delineating locations of wetland boundaries are based on these three wetland characteristics.

Wetlands are known by many common names, and in recent years these names have been applied more consistently to specific wetland types. Swamps are dominated by woody plants, either trees or shrubs. Marshes are dominated by herbaceous plants. Fens are peat-accumulating, open wetlands that receive mineral-rich groundwater. Bogs, also peat-accumulating wetlands, are isolated from mineral-rich water sources by deep peat accumulation and therefore receive most of their water and nutrients from precipitation. Each of these wetland types supports a unique group of plants and animals, many of which require these wetland habitats to survive.

Interpreting This Layer

This map layer uses the most comprehensive source of information on wetlands available: the National Wetlands Inventory (NWI). It was created to provide a broad-scale overview of where wetlands exist on the landscape. Keep in mind, however, that these maps were prepared using aerial photography. Wetlands that are hard to see on aerial photos—such as those that are forested—may not show up. Before using this map for specific planning purposes, you may want to check carefully for omissions.

How Can a Community Use This Information?

Wetlands nearly always provide beneficial wildlife habitat, although the recommended course of action may vary from one wetland to the next. Once you determine where the important wetlands are in your community, you may want to enlist a natural resources professional to conduct a field inventory and assessment. This would allow you to more fully understand the specific functions and values of the wetlands in the context of your region. Often, recommendations will focus on the extent of undeveloped buffer left around the wetland. A wetland that provides extensive habitat for breeding waterfowl or that is surrounded by steep slopes and is therefore at a high risk of erosion may benefit from a 100 to 300 foot buffer wherever possible; other wetlands may not need such an extensive riparian area to maintain function. In addition to considering the specific benefits a wetland provides to fish and wildlife, your community may also want to consider the wetland's role in flood and erosion protection, nutrient and pollution filtration, groundwater recharge, aesthetic diversity, and potential use for educational and recreational activities.

Conserving Vermont's Natural Heritage Tip

There are a variety of ways to plan for the future of the wetlands in your community. [*Conserving Vermont's Natural Heritage*](#) has a number of suggestions that might help you decide on a starting place. For example, it suggests that a town or other planning group might adopt **goals** like the following for the conservation of wetlands:

1. Protect or provide for long-term stewardship of wetlands that support significant functions and values for natural communities, rare species habitat, or wildlife habitat, and prevent additional loss of wetlands within the town.
2. Restore and/or enhance the functions and values of wetlands already affected by human disturbance.

The book goes on to provide specific recommendations of tools—both regulatory and non-regulatory—that may be useful in fulfilling these goals. See more on page 62 of the book.

It is estimated that less than 5% of Vermont is currently wetland and that nearly 50% of Vermont's historic wetland area has been lost or severely impaired due to draining, dredging, filling, or excavation activities associated with industrial, residential, and agricultural activities. Since 1995, the current rate of regulated wetland loss in Vermont is estimated at 20 acres per year, despite our state's relatively strong wetland protection laws. This does not include the loss of unregulated wetland, which is likely lost at a similar rate. The most effective way to ensure the continuation of wetlands is to provide stewardship of and/or protection for those that still exist. Fish and wildlife that depend on wetlands for their survival tend to be easily disturbed or negatively affected by human activities. Residential development close to a marsh that supports wading birds such as herons and bitterns is incompatible, for example. Domestic activities normally associated with residential development can cause disturbance, temporary displacement, or complete abandonment of the wetland by a variety of sensitive wetland-dependent wildlife.

For More Information

National Wetlands Inventory (NWI) of the U.S. Fish and Wildlife Service (USFWS): This is the agency responsible for mapping wetlands throughout the United States, including the layer displayed on Map 1. Information about the USFWS's classification system can be found on the [National Wetlands Inventory website](#)² or in the book *Classification of Wetlands and Deepwater Habitats of the United States* (Cowardin et al., 1979).

[Wetlands Section, Vermont Department of Environmental Conservation](#)³: This program regulates wetlands in Vermont, maintains VSWI maps for the state (The Vermont Significant Wetland Inventory), and provides technical assistance on wetland identification, delineation, and protection through planning and other mechanisms. It is also a source of information on the functions, values, and locations of wetlands throughout the state.

Community Strategies for Vermont's Forests and Wildlife:

Case Study

While statewide wetland protections are offered through the Vermont Wetland Rules, some towns have chosen to further protect these important natural features. The town of Warren protected wetlands and other ecologically sensitive areas by requiring a [Conservation Subdivision Design](#) in the review of major subdivisions, for example. You can read more about the town's process in *Community Strategies for Vermont's Forests and Wildlife*, on page 58.

The book also describes other tools a town could use, if this doesn't seem like the appropriate strategy for your community.

² Learn more about the National Wetlands Inventory at <http://wetlands.fws.gov>

³ The Vermont DEC Wetlands Section webpage is at <http://www.anr.state.vt.us/dec/waterq/wetlands.htm>.

Mapped Layer #3: Vernal Pools

What Are Vernal Pools?

A vernal pool is a depression in the forest floor that fills with water each spring and often dries out later in the growing season. Although trees and shrubs rarely grow in vernal pools, typical vernal pools are well shaded by the surrounding forest canopy. In the Northeast, many vernal pools start filling with fall rains, retain water, ice, and snow through the winter, and collect more water with the arrival of spring rain and snowmelt. Some are further influenced by rising groundwater in the fall and spring. The pools typically lack inlets and outlets, and while many vernal pools are dry by mid-summer, some may retain water throughout the year in wet years.



Vernal pools may take up a small amount of land area, but they are critical habitat for amphibians and very important for a number of additional species. In Vermont, vernal pool-dependent species include mole salamanders (Spotted salamander, Blue-spotted salamander, and Jefferson salamander), Eastern four-toed salamander, and wood frog. All of these species may breed in other wetlands but rely heavily on vernal pools to maintain their populations. For vernal pools to be effective breeding habitats for these amphibians, they must retain water for at

least two months during the spring and summer breeding season most years so that amphibians can complete their larval stage. The periodic drying of a vernal pool is essential, too, as this eliminates populations of predatory fish and diving beetles that prey on amphibian larvae. Other animals use pools as well, such as fairy shrimp, fingernail clams, snails, eastern newts, green frogs, American toads, spring peepers, and a diversity of aquatic insects. Fairy shrimp are thought to survive only in these temporary pools. Because the amphibians and invertebrates found in vernal pools constitute a rich source of food, various species of birds, mammals, and reptiles may be attracted to the pools. Despite their small size and temporary nature, vernal pools are highly productive ecosystems.

Interpreting This Layer

The data from this layer were collected through the Vermont Vernal Pool Project, a statewide effort to map the locations of vernal pools. These pools were initially mapped using aerial photographs; many have also been visited to confirm existence and collect data. You will notice that this information is actually divided into two similar layers: a “confirmed” vernal pools layer and a “suspected” layer. “Suspected” vernal pools have been mapped purely using aerial photographs but have not been confirmed in the field. “Confirmed” pools have been visited and enough data has been collected to say for certain that these pools exist and are used by wildlife. The locations of confirmed pools are therefore accurate to a fine scale and can be used even down to a parcel level; “suspected” vernal pools should be visited (with landowner permission) prior to taking any next steps.

How Can a Community Use This Information?

Vernal pools and the organisms that depend on them are threatened by activities that alter pool hydrology and substrate, as well as by significant alteration of the surrounding forest.

Construction of roads and other development in the upland forests around vernal pools can result in negatively affecting salamander migration and in mortality (Forman, et al., 2003).

Timber harvesting can have significant effects on vernal pools, including alteration of the vernal pool depression, changes in the amount sunlight, leaf fall, and coarse woody debris in the pool, and disruption of amphibian migration routes by the creation of deep ruts. Even when the pool is dry, alteration of the depression substrate may affect its ability to hold water and may disrupt the eggs and other drought-resistant stages of invertebrate life that form the base of the vernal pool food chain.

Community Strategies for Vermont's Forests and Wildlife:

Possible Tools

While there is no single strategy for protecting vernal pools, information about the following tools can be found in [*Community Strategies for Vermont's Forests and Wildlife*](#):

For working forests, landowners can include vernal pools as *Ecologically Significant Treatment Areas* (ESTAs) in [Forest Management or Stewardship Plans](#) (pg. 16), including those plans submitted to [Vermont's Use Value Appraisal Program](#) (pg. 19).

Because they are small in size and generally located within larger, forested parcels, municipal planners may decide to include vernal pool protections as one specific element within a larger conservation effort. For example:

- A [Conservation Zoning District](#) (pg. 41) or [Overlay District](#) (pg. 53) can include setback or buffering requirements for vernal pools.
- [Development Review Standards](#) for new developments could require a buffer around vernal pools (pg. 73).

Mapped Layer #4: Vermont Hydrography (Surface Water) Dataset

What Is the Vermont Hydrography Dataset?

The information displayed on this map is the most complete set of surface water data available in Vermont, including rivers, streams, lakes, and ponds. While wetlands were not specifically delineated in this effort, those wetlands containing open water have also been captured.

In using this layer, keep in mind that not only is the water itself important on the landscape; one of the most critical habitat elements for many wildlife is the riparian area that extends to the edge of the floodplain. Characterized by periodic disturbances—flooding, deposition of sediments, erosion, and the forces of water and ice movement, riparian habitats are highly complex and variable, and therefore extremely ecologically diverse (Verry et al., 2000). In fact, for a community faced with limited resources, prioritizing the conservation of riparian areas can be one of the best ways to help a wide diversity of wildlife species.

Throughout this section, riparian and aquatic habitats are discussed together because they are highly interdependent. When conducting planning efforts, however, please remember that only *water* is depicted on the map!

Interpreting This Layer

Because lines on this map are meant to represent the *centerline* of a stream, not the entire water body, this map may be most meaningful when used at the scale of a town or larger, to get a general sense of where water flows through the region. Also, because waterways are dynamic by nature, their exact boundaries or extent can be expected to change, both seasonally and over time. For planning purposes, your community may want to clearly define any planning or regulatory terms involving surface water through a means of measurement other than a static map the depicts one moment in time. It should also be noted that while this layer is the most complete map available of Vermont's surface waters, data were derived through the interpretation of orthophotos, aerial photographs, and topographic data and have not been checked in the field.

As mentioned above, riparian areas are not delineated on this map, and the size of these important areas varies greatly depending on topography. You may need to enlist the help of a professional ecologist to determine the functional boundaries of your area's riparian ecosystems; in general, larger bodies of water have wider riparian areas. Mountainous headwater streams are usually contained within steep, narrow valleys—with narrow riparian areas that transition into upland forests—while large streams and rivers wind through low gradient valleys, with riparian areas extending the width of the valley. Even though small streams in steep valleys have narrow bands of riparian habitat, however, the forest surrounding these streams plays an important role in protecting the riparian habitat and therefore the stream, especially where steep slopes threaten landslides, rapid storm water runoff, and hillside “gulleying.”

How Can a Community Use This Information?

Rivers, streams, lakes, and ponds are by nature diverse ecosystems, with plant and animal communities changing according to water depth, turbulence, available oxygen, and a host of other features. Shorelines contribute an additionally wide variety of plant and animal communities. These communities form an interconnected food web that ranges from tiny microorganisms to bears and humans. This web also includes reptiles and amphibians, plants, waterfowl, songbirds, bats, mink, and otter.




Photo by Eric Sorenson

One of the most effective ways to maintain high quality aquatic habitat is to protect the riparian area. A vegetated riparian area is important not only for the plants and animals that inhabit it but also for what it provides to the water below. Downed wood, leaves, and similar organic material filter into the water to become an important component of the food base and habitat structure. Mature trees in riparian areas also shade aquatic habitats, helping to control water temperatures. Riparian vegetation is crucial in filtering overland runoff, thus protecting water quality, and in stabilizing stream banks, thus preventing excessive streambank erosion and sediment buildup in aquatic habitats.

In your town, your specific conservation goals will dictate how wide an area you want to consider protecting around a stream or lake. In general, a naturally vegetated 100-foot-wide riparian buffer on each side of a stream will protect *many* of the functions associated with healthy riparian habitat, while a 330-foot buffer will protect *nearly all* the functions we value, including high-quality cover for many wildlife species.

Community Strategies for Vermont's Forests and Wildlife:

Possible Tools

Most waterways and riparian areas could be protected through strategies similar to those suggested for vernal pools, through the creation of [Forest Management or Stewardship Plans](#) (pg. 16) or adding setback or buffer requirements within a [Conservation Zoning District](#) (pg. 41), [Overlay District](#) (pg. 53), or [Development Review Standard](#) (pg. 73). In addition, you may want to consider the following, also found in [Community Strategies for Vermont's Forests and Wildlife](#):

- If landowners want to secure future protection for the wetlands, lakes, ponds, streams, or riparian areas on their land, a [Conservation Easement](#) (pg. 25) could be a valuable option.
- Because many trails parallel waterways through riparian areas, some towns choose to implement [Trail Policies](#) to protect habitat in these areas (pg. 63).

Of course, we value riparian areas for their contributions to human values, too—including water quality, recreation, education, spiritual well being, and sense of place. Riparian habitats also play a critical role in flood resilience, which has become particularly important since Vermont is experiencing increasingly more frequent flood events. Despite this, many of our rivers, streams, lakes, and wetlands no longer have functioning riparian areas due to more than 200 years of intensive human use of the land.



For additional information:

Buffers for Wetlands and Surface Waters: A Guidebook for New Hampshire Municipalities is a good source of detailed information about different buffer widths and the species and functions they protect (Chase et al. 1995).

MAP 2: Land Cover

This page will be replaced with MAP 2 specific to each town.



MAP 2. Land Cover

This map is useful at a broad scale for seeing patterns of natural land cover and land use.


This map, developed using satellite imagery, is useful at a broad scale for seeing patterns of natural land cover and land use. At a statewide scale, it is beneficial for picking out large urban areas, agricultural areas, wetland complexes, and forested areas. More locally, these data can be used to locate forested blocks, predict habitat connectivity, and see where patterns of development may hinder wildlife movement. They can also be used to distinguish hardwood forests from softwood and mixed forests, which can be helpful in predicting locations of natural communities and wildlife species.

What Is the Land Cover Layer?

The National Oceanic and Atmospheric Association's (NOAA's) Coastal Change Analysis Program (C-CAP) produces a nationally standardized database of land cover and land change information for coastal regions and adjacent uplands, including Vermont. Land cover records the natural landscape as surface components: forest, water, wetlands, urban, etc. Land use documents human uses of the landscape: residential, commercial, agricultural, etc. Land cover is documented by analyzing satellite and aerial imagery; land use is then inferred. The data used to create this map were recorded in 2011.

Natural Resources Atlas Tip:

If you try to recreate Map 2 in the [Natural Resources Atlas](#), your screen might look something like this: (Example: Waterbury)



Open the “Conservation Planning” Menu, click the “+” next to “Conservation Planning” and then “Map 2.”

Additional Layers: The [NOAA Coastal Change Analysis Program](#) created the dataset above. The program also offers data showing changes in land cover from 1996-2001 and from 2001 to 2006.

Need help using the Natural Resources Atlas? Try the [“How Do I Get Started?”](#) section in the Introduction of this guide.

Interpreting This Layer

NOAA C-CAP land cover data were developed to aid with identification of regional landscape patterns and major functional niches (habitat), and for environmental impact assessment, urban planning, and zoning applications. They can also be used to evaluate potential habitat, *if* identifying characteristics are available, such as species X needs Y acres of forested space within 500 meters of a water body. C-CAP data are not appropriate for identifying species and should be used only as a screening tool for very local or site-specific management decisions.

This map depicts twenty-two standard categories of land cover, including detailed information on wetland types:

- High Intensity Developed – Urban land cover with greater than 75 percent impervious surface
- Low Intensity Developed – Urban land cover with greater than 25 percent and less than 75 percent impervious surface

- Cultivated land – Active agriculture, orchards, and vineyards
- Grassland – Both managed and unmanaged grasslands
- Deciduous Forest – Hardwood forest with a pronounced seasonal dormancy period
- Evergreen Forest – Forest without a pronounced seasonal dormancy period
- Mixed Forest – Forest not dominated by either deciduous or evergreen species
- Scrub/Shrub – Woody vegetation less than 20 feet tall
- Palustrine Forest – Freshwater wetland forest
- Palustrine Scrub/Shrub – Freshwater wetland scrub/shrub
- Palustrine Emergent – Freshwater wetland-rooted emergent species (marsh, lilies, etc.)
- Estuarine Forest – Saltwater wetland forest greater than 20 feet (mangrove)
- Estuarine Scrub/Shrub – Saltwater wetland scrub/shrub (mangrove)
- Estuarine Emergent – Saltwater wetland emergent species (Spartina marsh, juncus grass, etc.)
- Unconsolidated Shore – Tidal flats, shoals, and intertidal areas
- Bare Land – Bare exposed rock, sand, and soil
- Water – Open water
- Palustrine Aquatic Bed – Floating vegetation and algal communities
- Estuarine Aquatic Bed – Marine algal communities
- Tundra – Permafrost and pariglacial conditions and communities
- Snow/Ice – Perennial snow and ice

Users of this map layer should keep in mind that because the above categories were mapped through the interpretation of remote images, some features are more accurately displayed than others. For example, open, similar cover types such as row crops and grasslands are not always differentiated; however, different wetland types are shown with a high rate of accuracy.

How Can a Community Use It?

This map of land cover provides a “first look” at a given landscape, identifying the abundance and distribution of general habitat types for animals and plants. Such a map also provides an initial visual inspection of the level and types of fragmentation of the landscape. Because the NOAA C-CAP land cover maps cover all of New England and New York using a standardized methodology, planners and managers can assess larger landscapes across state lines and use these maps for comparisons across geographic space as well as over time.

For More Information

[NOAA’s Coastal Services Center Northeast Landcover webpage](http://coast.noaa.gov/regions/northeast/)⁴ provides information specific to the Northeast; the Landcover Analysis webpage also includes detailed FAQs for C-CAP and other land cover mapping, and other information about C-CAP data, products and tools that may be useful in natural resource planning.

⁴ [NOAA’s Coastal Services Center Northeast Landcover webpage can be found at http://coast.noaa.gov/regions/northeast/.](http://coast.noaa.gov/regions/northeast/)

MAP 3: State and Regional Biodiversity Resources

This page will be replaced with MAP 3 specific to each town.



Map 3. State and Regional Biodiversity Resources

This map allows you to explore the distribution and richness of biodiversity of your community. It can be viewed at a town or regional scale.

This map is a bit different from other layers in that it depicts a compilation of many different ecosystem components rather than a single, concrete layer. This compilation uses [BioFinder](#), a map and database tool used to identify Vermont lands and waters that support ecologically important ecosystems, natural communities, habitats, and species.


BioFinder uses 21 datasets, each representing a different biological, ecological, or natural heritage component. A user can see each of these datasets individually, *or* you can view a layer depicting the overlap between datasets. This overlap is represented by a 6-tier classification that shows the locations on the landscape richest in biodiversity and those that have fewer known contributions to biodiversity. While only the aggregate layer is depicted on the map provided in this packet, you will likely want to see both the overlap *and* the individual components to fully understand the distribution of biodiversity across your community. We suggest beginning with the aggregate (called the “Tiered Classification to Biodiversity”) since it offers a first impression of relative geographic importance to biodiversity; then you can reference each component layer at a location of particular interest to understand why the area was assigned the tier ranking shown.

While these layers are viewable in the [Natural Resources Atlas](#), you can also visit the BioFinder website itself, at <http://biofinder.vermont.gov/>. The BioFinder website allows you to access more detailed abstracts about each dataset and print reports about the BioFinder components located in a given area.

See “[What is BioFinder?](#)” in the Introduction for more information on using BioFinder online.

Natural Resources Atlas Tip:

To recreate Map 3 in the [Natural Resources Atlas](#), your screen will look something like this:
(Example: Waterbury)



*You may notice that there is no “Map 3” on the list under “Conservation Planning.” This map is a bit more complex, so the layers have been placed elsewhere. When the “Conservation Planning” menu is selected, two layers entitled “Biofinder Components” and “Tiered Contribution to Biodiversity” appear at the **bottom** of the list of layers. “Tiered Contribution to Biodiversity” is selected here.*

Need help using the Natural Resources Atlas? Try the [“How Do I Get Started?”](#) section in the Introduction of this guide.

Biofinder Components

What are the “BioFinder Components?”

BioFinder starts with maps of 21 components that are all known to contribute to biodiversity:

Landscapes

- [Habitat Blocks](#)
- Grasslands and Shrublands
- [Rare Physical Landscapes](#)
- [Representative Physical Landscapes](#)
- Connecting Lands (<2,000 acres)
- Connecting Blocks (2,000-10,000 acres)
- Anchor Blocks (>10,000 acres)
- Riparian Connectivity

- [Wildlife Road Crossings](#)

Aquatics

- [Surface Waters & Riparian Areas](#)
- Representative Lakes
- Important Aquatic Habitats & Species Assemblages

Species & Natural Communities

- [Rare Species](#)
- Uncommon Species
- Rare Natural Communities
- Uncommon Natural Communities
- [Vernal Pools](#)
- [Vernal Pools \(Potential\)](#)
- [Wetlands](#)
- Mast Production Areas

As you may have noticed, many of these layers appear on other maps in this packet, too. If you are reading this on your screen, you can click the hyperlinks above to see explanations of what is meant by various components.



On the printed version of Map 3, we could only show the Tiered Contribution to Biodiversity aggregation. However, you may find it helpful to use the Natural Resources Atlas or BioFinder website to turn each layer on and off or mix and match to best visualize which components are important on your own landscape.

Tiered Contribution to Biodiversity

What is meant by “Tiered Contribution to Biodiversity?”

To create the Tiered Contribution to Biodiversity dataset, maps of the 21 components known to enhance an area’s biodiversity were layered together to see where there was the highest amount of overlap. Because some components contribute more than others to overall ecological importance, each component was weighted accordingly to enable any given location to be assigned a “tier” on a scale of 1 to 6. Generally speaking, higher priority tiers (Tiers 1, 2, and 3) show areas with the highest number of components present, while lower tiers show places with fewer overlapping layers, although the actual formula is slightly more complex.

Interpreting the Information

As mentioned above, the tier ranking offers a sense of the overall number of factors at play and some limited ability for comparison from one place to another. In many cases, a site’s ranking is heavily influenced by a specific component or group of components present at the site. This is because biological diversity looks so different in different places. A small handful of very rare species on the top of Mount Mansfield is considered ecologically important, as is a particularly

rich assemblage of common species in the Champlain Valley. See the [BioFinder](#) website for detailed descriptions of each of the 21 components that went into the Tiered Contribution as well as more information on how to interpret the map.

As you look at this map, it is important to keep in mind that what is displayed are locations with *identified* contributions to biodiversity. This means that locations in the highest tiers are, indeed, important for biodiversity. However, locations in lower tiers may simply not have been inventoried for particular components. While most component layers are thoroughly mapped, some layers may contain omissions; for example, many locations have never been inventoried for rare or uncommon species. Such omissions are unlikely to change the overall tier ranking substantially, but they may bump a location from one category up to the next.

How Can a Community Use This Information?

The Tiered Contribution to Biodiversity dataset points to regions of Vermont where biodiversity values are concentrated. Some of these are in remote regions and some are in urban areas, but all are locations that could serve as focal points for conservation efforts. The component layers will help in understanding what's happening in each important area.

Community Strategies for Vermont's Forests and Wildlife

Case Study

Maps provide information, but it will ultimately be up to a community to decide how the ecological values depicted on a map best fuse with other values of the community to create a viable plan. In the Mad River Valley, the Forests, Wildlife, and Communities project is an example of how one group of neighboring towns used a mapping strategy similar to the BioFinder strategy to aid in community planning efforts. You can read more about this project on page 14 of [Community Strategies for Vermont's Forests and Wildlife](#).

MAP 4: Wildlife Resources At a Town Scale

This page will be replaced with MAP 4 specific to each town.



MAP 4. Wildlife Resources at a Town Scale

This map shows the pattern of habitat across your community. Data are most appropriately used at a town or regional scale rather than for a small parcel or individual property.


The two prominent datasets in this map are Habitat Blocks (the background color) and Wildlife Road Crossings (the highlight color on certain road sections). Together these paint a picture of where the biggest blocks of contiguous habitat are located and where wildlife are likely to cross roads to move between blocks. In a sense, you can use this map to predict how some larger wildlife travel through the landscape and how easily genetic flow can occur between wildlife populations of many species.

Habitat blocks show contiguous natural cover—forests, woodlands, wetlands, old fields, and meadows—in which roads, development, and agriculture are absent. All else being equal, larger blocks of contiguous habitat generally contain greater biological diversity than smaller blocks of habitat. Therefore, finding methods to avoid fragmentation of the largest habitat blocks can be a first key step in creating a landscape conservation plan.

Over time, blocks of habitat have been getting increasingly smaller in Vermont as development pressure causes new roads to transect natural landscapes and structures to creep in from the edges. Species such as bear, bobcat, fisher, and others are increasingly using several smaller blocks rather than a single large block to get what they need for food, water, shelter and access to mates. However, this is only possible in locations containing some form of landscape connectivity—places where enough natural cover exists *between* habitat blocks for animals to feel secure enough to travel between them. The Wildlife Road Crossings layer looks at locations where wildlife are likely to cross roads in order to link together habitat blocks.

Natural Resources Atlas Tip:

If you try to recreate Map 4 in the [Natural Resources Atlas](#), your screen might look something like this: *(Example: Waterbury)*



With the “Conservation Planning” menu selected, click on the “+” next to “Conservation Planning,” then “Map 4.” This will allow you to select “Wildlife Road Crossings” and “Habitat Blocks.” The menu icons next to each layer title open a legend for that layer.

Additional Layers: Also related to wildlife connectivity is the “Stream Crossing” layer, found in the Fish & Wildlife section of the Atlas. This shows how well aquatic organisms are able to pass through culverts and other road-associated structures. You may also want to see this information alongside “Connecting Lands” and “Riparian Connectivity” layers to get a full picture of how wildlife are likely to travel through a landscape. These layers are found in the BioFinder section of the Atlas. See [Map 3](#) for more information about BioFinder layers.

Need help using the Natural Resources Atlas? Try the [“How Do I Get Started?”](#) section in the Introduction of this guide.

Mapped Layer #1: Habitat Blocks

(This information is repeated in [Map 5](#).)

What are Habitat Blocks?

Habitat blocks are areas of at least 20 acres with no roads or low densities of class III or IV roads. They contain little or no human development such as buildings, parking areas, lawns, gravel pits, active agricultural land, etc., but can be composed of any natural land cover type: various successional stages of forest, wetland, old meadow, etc. They are then categorized by size to make it easier to view them on the map.

Interpreting the Habitat Blocks Layer

While *size* is the important characteristic in this map, there isn't actually a minimum size block that is considered critical as important wildlife habitat. Blocks are best considered within the context of the landscape. A 100-acre habitat block located in Vermont's heavily-fragmented Champlain Valley may play a much more

ecologically important role than a 100-acre block in the Northeast Kingdom, where larger blocks are prevalent. The general rule of thumb is "the bigger, the better," and you can determine what "big" means in your region by viewing the habitat block layer at a regional scale using the [Natural Resources Atlas](#). The configuration of the habitat is also important. An area that is highly irregular in shape—containing a high amount of edge compared to interior forest—may be less functional for some species than habitat of the same acreage with a regular shape.

Habitat Blocks are one way to depict *contiguous habitat*. To read more about the role of contiguous habitat in our landscape, see page 39 in [Conserving Vermont's Natural Heritage](#).

To learn more about how the habitat blocks information was generated, you can find the original report from Vermont Fish & Wildlife Department and Vermont Land Trust online at <http://www.vtfishandwildlife.com/common/pages/DisplayFile.aspx?itemId=111635>. The full report is entitled *Vermont Habitat Blocks and Habitat Connectivity: An analysis using Geographic Information Systems*

Vermont's development history adds an interesting twist when we think about habitat block size. Because our areas of human settlement and development have historically—and currently—been along streams and in valleys, the largest remaining areas of contiguous habitat tend to be in upland and high elevation areas, or areas where soils are unsuitable for

agriculture or building. However, it is often those same valley bottoms where we would naturally see the greatest biological diversity. As you identify the largest areas of contiguous habitat in your town, keep in mind that they may be biased towards the uplands or other undevelopable landscape, but it is also important to include lowlands in conservation planning.

Mapped Layer #2: Wildlife Road Crossings

(This information is repeated in [Map 5](#).)

What are Wildlife Road Crossings?

Wildlife road crossings are one type of *connecting habitat*—that is, land that links larger patches of habitat within a landscape. Just as the term implies, these are areas where wildlife are most likely to cross roads. In addition to road crossing areas, connecting habitat might include riparian habitat along streams and rivers, strips of forest cover between developed areas, and even

hedgerows or fencerows through agricultural land—although only the road crossings are depicted on this map. Sometimes these areas are called "corridors," even when they are not linear, as the term implies. When

There are other types of connecting habitat present in our landscape in addition to wildlife road crossings. To read more about them, see page 48 in [Conserving Vermont's Natural Heritage](#).

several habitat blocks are connected in this way, they sometimes effectively fill the role of a larger habitat block, allowing for enhanced movement, migration, and dispersal of animals and plants.

While each species has a slightly different set of habitat elements that increases the likelihood of crossing a road, there are some general trends. Many species are most likely to cross a road when:

- Terrain is relatively flat, with no steep slopes on either side of the road;
- Wetland exists on at least one side of the road;
- Evergreen cover grows on both sides of the road;
- No houses are within 50m;
- Animals can access larger habitat blocks on each side of the road.

For some species, the presence of guardrails also significantly reduces the likelihood of crossing in a particular location, and when a bridge or large culvert is present, some species may be able to cross *under* a road rather than across the road surface.

Interpreting the Information

These data were generated by the Vermont Fish & Wildlife Department and Agency of Transportation to provide a preliminary look at where wildlife are most likely to cross Vermont roads. These departments used a computer modeling process to locate areas with a high concentration of the landscape features most closely associated with wildlife crossing areas. These features were weighted according to importance, and then road segments were given a score of 1 to 10, with 1 being the locations with the fewest associated features and 10 being the locations with the most.

While the crossings depicted on this map are sometimes very small in size and can be viewed at a fine scale, please keep in mind that these are locations of *probable* wildlife road crossings—that is, places that contain landscape features associated with wildlife crossing areas. Additional field investigation is needed to confirm the frequency at which wildlife *actually* use these potential crossings.

In 2006, the Town of Salisbury set out to determine which of the town's roads were most heavily used by wildlife to cross from one patch of habitat to another. Wildlife road tracking surveys were conducted by Conservation Commission members and local volunteers over the next several years, data were analyzed and mapped, and eventually, the town changed their town plan to reflect their newfound information.

Learn more at

<http://www.townofsalisbury.org/town-departments/conservation-commission/news-articles>.

While the layer is entitled “Wildlife Road Crossings,” it should also be noted that the map's focus is on terrestrial species with larger home ranges. There is certainly a lot of wildlife movement that *isn't* captured by this computer model, such as amphibian crossing areas or aquatic organism passage. This layer also doesn't show the connecting habitat between the road and the bigger blocks, only the road itself. Regardless, this offers a first step in addressing a likely pattern of wildlife movement across your town.

As mentioned earlier, road crossings are only one form of habitat connectivity, as roads are only one of the features that can fragment a landscape. Your community may want to identify other potential connecting habitats through the interpretation of aerial or orthophotos and/or by enlisting a natural resources professional. Once you have identified wildlife road crossings and other potential connecting features in your community, the next step may be to conduct field assessments to confirm which areas are most heavily used by wildlife.

As you examine this layer, you may also want to keep in mind that this map makes no representation of the level of safety present at different road crossing areas. An unsuccessful wildlife crossing attempt can be disastrous for both the human and the animal involved in a collision. Some crossing areas are therefore considerably better choices than others in terms of road crossing success, such as road sections with lower speed limits or straighter roads that provide heightened visibility to drivers. Even more ideal are places where wide culverts or bridges provide enough space for animals to cross under roads rather than across them. As you consider strategies that allow for wildlife connectivity in your community, you may simultaneously be able to address these safety concerns.

How Can a Community Use These Layers?

The inclusion of both large habitat blocks and wildlife road crossings in a conservation plan will benefit a wide range of native plant and animal species. For example, many larger mammals like moose, bobcat, black bear, fisher, and otter require large home ranges to find the food, shelter, and access to mates they require. Not only do these animals need to travel great distances to fulfill their high-calorie dietary needs on a daily basis, but their primary food sources tend to change seasonally, and they may need to move from one habitat type to another throughout the year in order to survive. Of equal importance is the prevention of inbreeding by allowing for genetic crossover between populations. For large animals, this is often only possible in places with large habitat blocks and links between the blocks.

Bird species such as barred owl, hermit thrush, ovenbird, and many others also require forest interior to ensure nesting success. In the absence of such habitat, these birds are subject to increased nest predation from raccoons, skunks, squirrels, and chipmunks as well as nest parasitism from brown-headed cowbirds. Many native migratory songbird populations are now in decline due in part to the loss of contiguous forest habitat. Even smaller animals such as spotted salamanders will benefit from the protection of large habitat blocks, because they, too, use connecting habitat in the spring to move from their hibernation sites to breeding pools.

Community Strategies for Vermont's Forests and Wildlife

Once the locations where wildlife are most likely to travel have been identified, your community may want to think about whether regulatory or non-regulatory strategies are most appropriate for you. Possible regulatory tools include [Overlay Districts](#) (pg. 50) or [Subdivision Regulations](#) (pg. 54), about which you can read more in *Community Strategies for Vermont's Forests and Wildlife*. You may also find it beneficial to provide clear definitions in your town plan for phrases such as "Habitat Block," "Contiguous Forest," or "Wildlife Travel Corridor." Sample definitions for these and other phrases are given on [page 71](#).

In addition to the species mentioned above, landscape-level conservation efforts benefit *most* species, because a large habitat block comprises many diverse smaller-scale habitats, each of which is thereby conserved. The ability of many plant and animal species to shift geographically or adapt to a changing climate is also enhanced through the protection of both large blocks and connecting features.

Keep in mind that while conserving corridors has great merit, conserving threads of vegetative cover within a developing landscape will not necessarily maintain an area's ecological values and biological diversity. Corridors alone are unlikely to meet species' habitat needs; they support ecosystem functions only in conjunction with large areas of undeveloped land containing diverse habitat. Similarly, one area of large, contiguous habitat provides a much greater ecological benefit than two smaller blocks of half the area, joined by a corridor.

As you examine the habitat available to wildlife in your community, you may find the following useful in evaluating and prioritizing different areas:

- *Size:* As a general rule of thumb, larger habitat blocks are likely to have more ecological value. They often also provide greater benefits to the civic community through opportunities to access forest resources, hunting, or recreational use.
- *Condition:* Areas that contain diverse natural habitat types normally have a greater variety of plant and animal species. While condition can be difficult to ascertain, the



- BioFinder "Tiered Contribution to Biodiversity" scores depicted on [Map 3](#) may help you in identifying places with a high potential for biodiversity.
- *Landscape Context:* Habitat areas that are near other patches of natural vegetation and are well buffered from substantial fragmenting features like roads, development, or agricultural land function better as wildlife habitat for many species.
- *Connectivity:* Connecting features can link blocks together to effectively function as somewhat larger blocks. While larger blocks generally remain better for wildlife than a series of linked smaller blocks, these features can allow a broader diversity of wildlife to inhabit human-populated areas.

MAP 5: Fine Scale Wildlife Resources

This page will be replaced with MAP 5 specific to each town.



Photo by John Hall

MAP 5. Fine Scale Wildlife Resources

This map shows several datasets developed to be accurate at a very local scale—even down to individual parcels.

This map shows four data layers: Habitat Blocks, Wildlife Road Crossings, Deer Wintering Areas, and Rare, Threatened, and Endangered Species. Other than the Habitat Blocks, all layers can be analyzed on a fine scale even down to the level of an individual parcel. Habitat Blocks are shown here only as a background layer, in order to provide an element of statewide perspective.


Data at the local scale can help round out the patterns seen in [Map 3](#) and [Map 4](#). We recommend starting with those two maps to gain an understanding of the general patterns of habitat and biodiversity across your town or region. This map can then provide additional details about specific areas of interest.

As noted in Map 4, the largest areas of contiguous habitat in Vermont are found primarily in uplands, even though lower elevation wetlands, riparian areas, and other habitat types are ecologically just as important. This map provides a way to identify areas of biological interest that may or may not correspond to larger habitat blocks.

Please note that while the primary layers depicted on this map may be appropriate at a local scale, they aren't comprehensive. For example, a mark depicting a Rare, Threatened or Endangered (RTE) species is spatially accurate, but the absence of an RTE marker is *not* a definite sign that there are no rare species present. This is because the entire state has never been inventoried for all RTE species, so there are inevitably points missing from the database.

Natural Resources Atlas Tip:

If you try to recreate Map 5 in the [Natural Resources Atlas](#), your screen might look something like this: (Example: Waterbury)



With the “Conservation Planning” menu selected, click on the “+” next to “Conservation Planning,” then “Map 5.” This will allow you to select “Wildlife Road Crossings” and “Habitat Blocks.” Please note that for this map, the data layers included on the Natural Resources Atlas are slightly different than the layers printed on the map provided.

Need help using the Natural Resources Atlas? Try the [“How Do I Get Started?”](#) section in the Introduction of this guide.

Additional Resources: The [Fluvial Erosion Hazards](#) Website from the Department of Environmental Conservation also has information appropriate for this local scale. While this information is not yet available as a layer that you can upload to the Natural Resources Atlas, it may certainly still be helpful.

Mapped Layer #1: Habitat Block

(This information is repeated in [Map 4](#).)

What are Habitat Blocks?

Habitat blocks are areas of at least 20 acres with no roads or low densities of class III or IV roads. They contain little or no human development such as buildings, parking areas, lawns, gravel pits, active agricultural land, etc., but can be composed of any natural land cover type: various successional stages of forest, wetland, old meadow, etc. They are then categorized by size to make it easier to view them on the map.

Interpreting the Habitat Blocks Layer

While *size* is the important characteristic in this map, there isn't actually a minimum size block that is considered critical as important wildlife habitat. Blocks are best considered within the context of the landscape. A 100-acre habitat block located in Vermont's heavily-fragmented Champlain Valley may play a much more ecologically important role than a 100-acre block in the Northeast Kingdom, where larger blocks are prevalent. The general rule of thumb is "the bigger, the better," and you can determine what "big" means in your region by viewing the habitat block layer at a regional scale using the [Natural Resources Atlas](#). The configuration of the habitat is also important. An area that is highly irregular in shape—containing a high amount of edge compared to interior forest—may be less functional for some species than habitat of the same acreage with a regular shape.

Habitat Blocks are one way to depict *contiguous habitat*. To read more about the role of contiguous habitat in our landscape, see page 39 in [Conserving Vermont's Natural Heritage](#).

Vermont's development history adds an interesting twist when we think about habitat block size. Because our areas of human settlement and development have historically—and currently—been along streams and in valleys, the largest remaining areas of contiguous habitat tend to be in upland and high elevation areas, or areas where soils are unsuitable for agriculture or building.

To learn more about how the habitat blocks information was generated, you can find the original report from Vermont Fish & Wildlife Department and Vermont Land Trust online at <http://www.vtfishandwildlife.com/common/pages/DisplayFile.aspx?itemId=111635>. The full report is entitled *Vermont Habitat Blocks and Habitat Connectivity: An analysis using Geographic Information Systems*

However, it is often those same valley bottoms where we would naturally see the greatest biological diversity. As you identify the largest areas of contiguous habitat in your town, keep in mind that they may be biased towards the uplands or other undevelopable landscape, but it is also important to include lowlands in conservation planning.

Mapped Layer #2: [Wildlife Road Crossings](#)

(This information is repeated in [Map 4](#).)

What are Wildlife Road Crossings?

Wildlife road crossings are one type of *connecting habitat*—that is, land that links larger patches of habitat within a landscape. Just as the term implies, these are areas where wildlife are most likely to cross roads. In addition to road crossing areas, connecting habitat might include riparian habitat along streams and rivers, strips of forest cover between developed areas, and even hedgerows or fencerows through agricultural land—although only the road crossings are depicted on this map. Sometimes these areas are called "corridors," even when they are not linear, as the term implies. When several

There are other types of connecting habitat present in our landscape in addition to wildlife road crossings. To read more about them, see page 48 in [Conserving Vermont's Natural Heritage](#).

habitat blocks are connected in this way, they sometimes effectively fill the role of a larger habitat block, allowing for enhanced movement, migration, and dispersal of animals and plants.

While each species has a slightly different set of habitat elements that increases the likelihood of crossing a road, there are some general trends. Many species are most likely to cross a road when:

- Terrain is relatively flat, with no steep slopes on either side of the road;
- Wetland exists on at least one side of the road;
- Evergreen cover grows on both sides of the road;
- No houses are within 50m;
- Animals can access larger habitat blocks on each side of the road.

For some species, the presence of guardrails also significantly reduces the likelihood of crossing in a particular location, and when a bridge or large culvert is present, some species may be able to cross *under* a road rather than across the road surface.

Interpreting the Information

These data were generated by the Vermont Fish & Wildlife Department and Agency of Transportation to provide a preliminary look at where wildlife are most likely to cross Vermont roads. These departments used a computer modeling process to locate areas with a high concentration of the landscape features most closely associated with wildlife crossing areas. These features were weighted according to importance, and then road segments were given a score of 1 to 10, with 1 being the locations with the fewest associated features and 10 being the locations with the most.

While the crossings depicted on this map are sometimes very small in size and can be viewed at a fine scale, please keep in mind that these are locations of *probable* wildlife road crossings—that is, places that contain landscape features associated with wildlife crossing areas. Additional field investigation is needed to confirm the frequency at which wildlife *actually* use these potential crossings.

In 2006, the Town of Salisbury set out to determine which of the town's roads were most heavily used by wildlife to cross from one patch of habitat to another. Wildlife road tracking surveys were conducted by Conservation Commission members and local volunteers over the next several years, data were analyzed and mapped, and eventually, the town changed their town plan to reflect their newfound information.

Learn more at

<http://www.townofsalisbury.org/town-departments/conservation-commission/news-articles>.

While the layer is entitled “Wildlife Road Crossings,” it should also be noted that the map's focus is on terrestrial species with larger home ranges. There is certainly a lot of wildlife movement that *isn't* captured by this computer model, such as amphibian crossing areas or aquatic organism passage. This layer also doesn't show the connecting habitat between the road and the bigger

blocks, only the road itself. Regardless, this offers a first step in addressing a likely pattern of wildlife movement across your town.

As mentioned earlier, road crossings are only one form of habitat connectivity, as roads are only one of the features that can fragment a landscape. Your community may want to identify other potential connecting habitats through the interpretation of aerial or orthophotos and/or by enlisting a natural resources professional. Once you have identified wildlife road crossings and other potential connecting features in your community, the next step may be to conduct field assessments to confirm which areas are most heavily used by wildlife.

As you examine this layer, you may also want to keep in mind that this map makes no representation of the level of safety present at different road crossing areas. An unsuccessful wildlife crossing attempt can be disastrous for both the human and the animal involved in a collision. Some crossing areas are therefore considerably better choices than others in terms of road crossing success, such as road sections with lower speed limits or straighter roads that provide heightened visibility to drivers. Even more ideal are places where wide culverts or bridges provide enough space for animals to cross under roads rather than across them. As you consider strategies that allow for wildlife connectivity in your community, you may simultaneously be able to address these safety concerns.

How Can a Community Use These Layers?

The inclusion of both large habitat blocks and wildlife road crossings in a conservation plan will benefit a wide range of native plant and animal species. For example, many larger mammals like moose, bobcat, black bear, fisher, and otter require large home ranges to find the food, shelter, and access to mates they require. Not only do these animals need to travel great distances to fulfill their high-calorie dietary needs on a daily basis, but their primary food sources tend to change seasonally, and they may need to move from one habitat type to another throughout the year in order to survive. Of equal importance is the prevention of inbreeding by allowing for genetic crossover between populations. For large animals, this is often only possible in places with large habitat blocks and links between the blocks.

Bird species such as barred owl, hermit thrush, ovenbird, and many others also require forest interior to ensure nesting success. In the absence of such habitat, these birds are subject to increased nest predation from raccoons, skunks, squirrels, and chipmunks as well as nest parasitism from brown-headed cowbirds. Many native migratory songbird populations are now in decline due in part to the loss of contiguous forest habitat. Even smaller animals such as spotted salamanders will

Community Strategies for Vermont's Forests and Wildlife

Once the locations where wildlife are most likely to travel have been identified, your community may want to think about whether regulatory or non-regulatory strategies are most appropriate for you. Possible regulatory tools include [Overlay Districts](#) (pg. 50) or [Subdivision Regulations](#) (pg. 54), about which you can read more in *Community Strategies for Vermont's Forests and Wildlife*. You may also find it beneficial to provide clear definitions in your town plan for phrases such as "Habitat Block," "Contiguous Forest," or "Wildlife Travel Corridor." Sample definitions for these and other phrases are given on [page 71](#).

benefit from the protection of large habitat blocks, because they, too, use connecting habitat in the spring to move from their hibernation sites to breeding pools.

In addition to the species mentioned above, landscape-level conservation efforts benefit *most* species, because a large habitat block comprises many diverse smaller-scale habitats, each of which is thereby conserved. The ability of many plant and animal species to shift geographically or adapt to a changing climate is also enhanced through the protection of both large blocks and connecting features.

Keep in mind that while conserving corridors has great merit, conserving threads of vegetative cover within a developing landscape will not necessarily maintain an area's ecological values and biological diversity. Corridors alone are unlikely to meet species' habitat needs; they support ecosystem functions only in conjunction with large areas of undeveloped land containing diverse habitat. Similarly, one area of large, contiguous habitat provides a much greater ecological benefit than two smaller blocks of half the area, joined by a corridor.

As you examine the habitat available to wildlife in your community, you may find the following useful in evaluating and prioritizing different areas:

- *Size:* As a general rule of thumb, larger habitat blocks are likely to have more ecological value. They often also provide greater benefits to the civic community through opportunities to access forest resources, hunting, or recreational use.
- *Condition:* Areas that contain diverse natural habitat types normally have a greater variety of plant and animal species. While condition can be difficult to ascertain, the BioFinder "Tiered Contribution to Biodiversity" scores depicted on [Map 3](#) may help you in identifying places with a high potential for biodiversity.
- *Landscape Context:* Habitat areas that are near other patches of natural vegetation and are well buffered from substantial fragmenting features like roads, development, or agricultural land function better as wildlife habitat for many species.
- *Connectivity:* Connecting features can link blocks together to effectively function as somewhat larger blocks. While larger blocks generally remain better for wildlife than a series of linked smaller blocks, these features can allow a broader diversity of wildlife to inhabit human-populated areas.

Mapped Layer #3: Deer Wintering Areas

What are Deer Wintering Areas?

Deer wintering areas vary in size from a few acres to over a hundred acres and provide essential relief to deer from winter conditions. Covered by a dense cover of mature or maturing softwood trees, they provide protection from deep snow, cold temperatures, and wind. These areas are characterized by a favorable aspect (south-



facing, or perhaps southeast or southwest-facing but rarely north), they generally occur at moderate elevations, and they are found in places with low levels of human disturbance in winter. Tree cover is most often from hemlock and white pine in the southern part of the state, and white cedar, spruce, and fir in the north. Deer inhabiting these areas can minimize how much energy they lose in the winter, thus enhancing survival.

From one year to the next, wintering areas do not change significantly, so these areas can be used by generations of deer over many decades if habitat conditions are maintained. Deer annually migrate—often several miles—from fall habitats to wintering areas, and a single large deer wintering area can occasionally attract deer from a radius of several towns.

Interpreting the Information

Deer wintering areas were identified in 2010 using aerial observations, infrared aerial photos, and ground confirmation. Additional areas are added to the database as they are discovered. It is important to keep in mind that not all deer wintering areas have been mapped, and that changes in forest cover and land use change an area's use as a deer yard. If you suspect an area serves as deer winter habitat that is not mapped, we recommend that you contact us.

How Can a Community Use This Information?

In addition to benefits for deer, dense softwood stands provide critical winter shelter and food supplies for a variety of other wildlife species including porcupines, snowshoe hare, fox, fisher, coyotes, bobcats, crows, ravens, red and white-winged crossbills, and many others. Logging can be either detrimental or beneficial to the habitat depending on the harvest method employed and the overall sensitivity shown by the logger and landowner to maintaining these areas of dense softwood cover.

The conservation of deer wintering areas is essential to maintaining and managing white-tailed deer in Vermont. Deer wintering areas make up a relatively small percentage of the land base of most towns; only 8% of the forested landscape of Vermont has been mapped as deer winter habitat. However, residential, commercial, or industrial development

within or adjacent to these areas decreases the amount of winter habitat available to deer and can eventually reduce the number of deer within the area. Without adequate winter habitat, northern populations of deer would be subject to extreme fluctuations due to heightened levels of winter mortality during moderate and severe winters.

Additional information on deer winter habitat requirements and management recommendations can be found in the publication [*Wildlife Habitat Management for Vermont Woodlands, a Landowner's Guide*](#), which is available from the Vermont Fish and Wildlife Department.

Mapped Layer #4: Rare, Threatened, and Endangered Species

What Are Rare, Threatened, and Endangered Species?

A rare species is one that has only a few populations in the state and faces threats to its continued existence in Vermont. Threats might include loss of habitat, harassment, collection, or

suppression of natural processes, such as fire. The Vermont Fish and Wildlife Department uses a ranking scheme that describes species rarity within the state, ranging S1 (very rare) to S5 (common and widespread). Species are assigned a rank based on the number of known examples, the population size, and the degree to which the populations are threatened.

If you're interested in seeing a complete list of Vermont's rare species and their respective ranks, go to

<http://www.vtfishandwildlife.com/common/pages/DisplayFile.aspx?itemId=229831>.

Endangered and threatened species are defined by both state and federal law. Vermont's [*Endangered and Threatened Species Rule*](#) (10 V.S.A. Chap. 123 section 5401) maintains a record of listed species, and the federal Endangered Species Act maintains a similar [list](#).

According to the Vermont Rule, endangered species are those “whose continued existence as a viable component of the state’s wild fauna or flora is in jeopardy.” Threatened species are “species whose numbers are significantly declining because of loss of habitat or human disturbance, and unless protected will become an endangered species.”

As of July 2015, there were 314 species of plants and animals in Vermont that are protected by the Vermont Endangered Species Law and assigned a status of either threatened or endangered. Some of these species that occur in Vermont also have a federal status of threatened or endangered, and are protected by the Federal Endangered Species Act (P.L. 93-205).

Interpreting the Information

The data shown on this layer were compiled through the Vermont Fish & Wildlife Department’s Natural Heritage Inventory (NHI), which is the state’s contribution to a greater, regional database of conservation information maintained by the organization NatureServe. Unlike many of the layers in this packet, all of the information depicted about rare, threatened, and endangered species represents field-confirmed, geographically accurate data points.

Planners should be aware that when a point is used to represent a rare species observation, the size of the population should not be assumed to be small. A mapped point may represent only a few square yards, but it could also indicate a large wetland, a river stretch over a mile long, or an extensive ridge top. Usually, more specific data is recorded in the NHI database, and this may even be included in the mapping attributes, but you may need to do research beyond seeing the initial point on the map to get this information. As you look for information, keep in mind that those conducting the field inventories sometimes record additional notes, such as threats or management needs like an invasive species that is affecting a rare species. Contact VTFWD for more information.

While rare, threatened, and endangered species *locations* are provided on this map, you will notice that plant and animal species information is missing. Nationwide, this information is omitted from mapping efforts so as not to jeopardize the survival of sensitive species. Some rare organisms are sought after for collection, others are targeted as unwanted (such as the timber rattlesnake), and others draw attention from those attracted to the uniqueness of the species, which can sometimes disturb the species’ natural habitat or behavior (as can happen with the peregrine falcon or bald eagle). In addition to the potential damage to the species, these

behaviors can also be disruptive to land owners and managers. As a result, each RTE location mapped here is labeled generally as a plant or animal. Landowners, managers, as well as town planning and conservation commissions can contact the Vermont Fish & Wildlife Department for additional information, but the information is not provided to the general public. Natural communities are similarly mapped, but attribute information *does* include the name of the natural community type.

How Can a Community Use This Information?

Rare native species in Vermont, such as Indiana bat, common loon, spiny softshell turtle, goldenseal, and sweet coltsfoot are an important part of Vermont's natural heritage. Rare species can play crucial roles in ecosystems, with other species relying on them for their survival. Many of these species are admired and appreciated by people for their beauty, sounds, or mere presence on the landscape.

Most of these species in Vermont are rare because they are on the edge of their range or they are separated from the main population by a large distance.

Conserving Vermont's Natural Heritage details many strategies that might be helpful in keeping rare, threatened, and endangered species on our landscape. To read more, see [chapter 6, page 106](#).

For example, the spiny softshell turtle is found in Vermont in parts of Lake Champlain, and the next nearest population is in the St. Lawrence River. The majority of the population is found west of New York. A number of our rare species occur in unique habitat types or rare natural communities, such as Lake Sand Beach or Calcareous Riverside Seep. More wide-ranging animal species, like osprey, are considered rare when their overall populations consist of small numbers of breeding pairs.

Because the planet in general—and possibly Vermont specifically—is experiencing the loss of species at a rate never before experienced, those species most at risk of extinction, extirpation, and endangerment serve as barometers of the state of the environment (Defenders of Wildlife, 1998). Protecting and restoring rare, threatened, and endangered species represents one of the most difficult present-day conservation challenges in Vermont. If we are to see the continued presence of these species in our state, we may need to address their needs at all levels of planning—local, regional, and statewide.

Each town harbors its own set of rare and uncommon species that contributes to the overall diversity of the state. Even though Vermont is a small state, it has varied terrain, aquatic systems, elevations, wetlands, geology, and natural communities. It may be that the rare species found in your community are located in habitats that are ecologically unique at the state level—even when they don't seem particularly rare in a local context. In this way, viewing this fine-scale layer can actually point to locations critical within a larger context.

Community Strategies for Vermont's Forests and Wildlife:

Possible Tools

Fine-scale resources such as deer wintering areas or habitat for rare, threatened, and endangered species could be protected using the following tools, found in [*Community Strategies for Vermont's Forests and Wildlife*](#):

Several [Federal and State Assistance Programs](#) (pg. 32) will help with technical and financial assistance for the implementation of some conservation measures for rare, threatened, and endangered species.


For working forests, landowners can include critical habitat—such as that required by rare, threatened, and endangered species—as *Ecologically Significant Treatment Areas* (ESTAs) in [Forest Management or Stewardship Plans](#) (pg. 16), including those plans submitted to [Vermont's Use Value Appraisal Program](#) (pg. 19).

You can also include fine-scale resource protections within larger municipal goals:

- In a [municipal plan](#) (pg. 42), specific goals or policies can be stated for deer wintering areas and critical habitat for rare, threatened, and endangered species.
- Planning goals can be backed up with bylaws including a [Conservation Zoning District](#) (pg. 41) or [Overlay District](#) (pg. 53) that specifies setback or buffering requirements for deer wintering areas or other sensitive wildlife habitat.
- [Development Review Standards](#) for new developments can include specific requirements for fine-scale habitats, such as building envelopes located to exclude critical wildlife habitat (pg. 36).

MAP 6: Physical Features

This page will be replaced with MAP 6 specific to each town.



MAP 6. Physical Features

This map gives a big-picture view of how physical features are geographically distributed.


Physical landscapes (often referred to as “enduring features”) are the parts of the landscape that resist change. They are the hills and valleys, the underlying bedrock, and the deposits left behind by glaciers. They remain largely static when natural or human-induced changes in land cover and wildlife occur, as plants and animals expand or contract their ranges, and even as the climate changes. If nature were likened to a dramatic play, the physical landscape is the stage, and the plants, animals, and other species are the actors.

But *unlike* the stage used in a playhouse, physical landscapes are also driving forces behind ecological processes. They heavily influence which plants, animals, and natural communities appear and thrive—and they can therefore help us in understanding biological diversity and in predicting habitat conditions and species presence. Physically diverse landscapes support diverse natural communities and species (Anderson & Ferree, 2010), and so one way to ensure that biological diversity persists on our landscape is to conserve a variety of physical landscapes.

The purpose of this map is to provide some context for your town’s physical landscape. While the map is zoomed in to a town scale, the physical landscapes depicted are important at a regional level. There are two types of physical landscape shown: “representative” physical landscapes, and “rare” physical landscapes. Both are places likely to support a diverse set of natural communities and species.

Natural Resources Atlas Tip:

If you try to recreate Map 6 in the [Natural Resources Atlas](#), your screen might look something like this: (Example: Waterbury)



With the “Conservation Planning” menu selected, click on the “+” next to “Conservation Planning,” then “Map 6.” This will allow you to select the “Rare Physical Landscapes” and “Representative Physical Landscapes” layers.

Need help using the Natural Resources Atlas? Try the [“How Do I Get Started?”](#) section in the Introduction of this guide.

What Is Meant By a “Physical Landscape?”

The physical landscape includes:

- Bedrock: the rock that underlies everything we see on the surface
- Surficial materials: the gravel, sand, silt, clay, or peat that sits on top of the bedrock
- Elevation
- Topography or Landforms: cliffs, coves, summits, flats, etc.

Individually, each physical feature influences the ecological landscape in a unique way. When they are mixed and matched, the resulting patterns can be quite complex. In order to describe the numerous combinations of physical features found across Vermont’s landscape, we use “Land Type Associations,” or LTAs, the results of a computer model developed by The Nature Conservancy and Vermont Land Trust, originally for use by the Green Mountain National Forest (Ferree & Thompson 2008).

This model has combined the physical landscape components listed above with additional factors such as soil types and potential natural vegetation to create a visual representation of what Vermont’s “stage” looks like in different areas. For example, the LTAs of the Green Mountains illustrate variations in the steep, high-elevation terrain with acidic bedrock and rocky glacial

deposits, while the Champlain Valley LTAs generally feature combinations of flatter, calcium-rich clay plains.

Mapped Layer #1: Rare Physical Landscapes

“Rare” physical landscapes are the Land Type Associations that are least commonly found across Vermont’s landscape, each covering less than 4.5% of the state’s land area. Rare physical landscape types often correspond to the presence of rare species or natural communities, so locations where these conditions are found often add to the state’s overall biodiversity. Rare associations include the following:

Rare Land Type Association	Percent of Vermont’s Land Area
Calcareous Metamorphic High Hills/Low Mountains	0.3
Connecticut River Valley - Lake Hitchcock Sediments	0.8
Enriched Slopes	0.6
Granitic Basin	0.4
Granitic High Hills/Low Mountains	0.5
Granitic Mid-Elevation Hills	1.1
Marine-Lacustrine-Glaciofluvial Coarse Sediments	0.9
Precambrian Plateau	2.2
Upper Mountain Slopes/Mountaintops	2.7
Valley Floor Glacial Lake/Marine Plains	4.4
Vermont Escarpment	0.8
Water-deposited glacial sediments along major riverways	2

Mapped Layer #2: Representative Physical Landscapes

For each Land Type Association not included on the “rare” list, high-quality examples were selected throughout the state based on condition and patch size. These examples are considered “representative” physical landscapes. Representativeness is an effective tool in conservation planning by allowing us to identify at a *local* level areas that contribute to biodiversity at a *state* level. Common species and natural communities are every bit as important as the rare species we often focus on, but without datasets like this, it can be difficult to include their importance on a map.

Interpreting the Information

We haven’t inventoried (and *can’t* inventory) every species in every location in the state—and even if we could, natural systems are in a constant state of fluctuation. This leaves us unable to accurately quantify biodiversity. Including physical features in conservation planning is a way to capture biodiversity in the absence of such an inventory, as they portray the “ecological potential” of the landscape regardless of present-day land use or land cover resulting from human alterations to the natural vegetation.



Land Type Associations are very useful at a statewide or multi-state scale where a high degree of accuracy is unnecessary. At a town scale, they should be interpreted with caution. At this scale, the boundaries between two LTA types cannot be considered highly geographically accurate. However, the physical landscapes map can be used as an initial bird's eye view to help you think about the local landscape in a new way as you decide where you may wish to encourage or limit growth or other planning strategies.

How Can A Community Use This Information?

As mentioned above, physically diverse landscapes correspond to diversity in species. Conserving wildlife habitat within rare and representative landscapes can therefore encourage a diversity of species to flourish. This is particularly true in the face of global climate change. As changes occur over time, plant and animal species adjust their ranges to more climatically suitable conditions. Areas of diversity in the physical landscape will allow for these adjustments to be made more easily, and these areas are likely to continue as the stage for biological diversity even as species composition changes.

As you look at this map, you may also notice that the areas highlighted may be quite different from the other maps in this guide. This is because the other layers tend to reflect current landscape condition, where habitat exists *now*. One interesting way to look at the Physical Features map is to think about how biodiversity may have been historically distributed prior to European or even Native American settlement, keeping in mind that physical diversity often corresponds to biological diversity. Similarly, you could use this map to understand how biodiversity *could* be distributed in the future in undeveloped areas. Essentially, this map provides a lens for erasing current land use patterns to allow you to think about the ecological potential of the land. If your community is interested in restoration or land conservation efforts, you may find this map particularly enticing.



Advanced Natural Resources Inventory

While we hope that your community has found useful information in the six maps provided, we're aware that every town in Vermont has a different set of values *and* a different landscape. The information that will help you most may have been omitted here. Luckily, there are other opportunities for getting an in-depth view of your local landscape. Be aware that some of these opportunities may come with a price tag, since they involve on-the-ground fieldwork and more advanced computer analysis. While some field data can be gathered by volunteers with minimal professional guidance, other information is best collected by (or under the close supervision of) a natural resources professional. Towns may therefore want to start by assigning a priority ranking to each new information need. Even if it takes several years to get the new information, it is a worthwhile effort; the reward is better planning for the future.

The following is a list of ideas (by no means exhaustive) of projects a town might undertake to get a better sense of what resources are present. A detailed inventory could include such things as:

- Natural community mapping
- Water resources evaluation (e.g. water quality studies)
- Wetland evaluation/prioritization studies
- Significant wildlife habitat assessment
- Agricultural lands assessment
- Managed forest lands inventory
- Undeveloped shorelines inventory
- Cultural features inventory (e.g. archaeological and historic sites, recreation areas, scenic areas, designated scenic roads)
- Unique geological resources mapping

In addition to the above, keep in mind that some of the information included on the six inventory maps of this packet may also benefit from on-the-ground field checking. Do the streams in your town have functional riparian areas? Are the boundaries of the land cover data accurate at a local level? Are there current threats facing wildlife road crossing areas? These questions can be most effectively answered by a field inventory.

As you undertake your inventory, remember that while some landscape elements are not static, a map can depict only a snapshot in time. Development of a new building site may change the size of a habitat block, and land use and land cover of a given location change routinely due not only to human alterations but also because of natural succession as forests grow up and mature over time. In addition to seeking updates on the Natural Resources Atlas, inventories that you undertake in your community may also benefit from routine repetition. In some cases, the success of a particular planning goal even *necessitates* a map update! For example, if you implement strategies to restore or enhance riparian habitat, success may require that you update your riparian habitat map on a routine basis. Each map can then become a tool that helps you track progress.

Ultimately, it will be up to your town to decide which information is most important to *you*. It is by combining your knowledge of the local natural resources with your town's own goals and interests that you will be able to create a natural resources plan that you are able to successfully implement within your community.



Bibliography

Anderson M.G. and C.E. Ferree. 2010. *Conserving the Stage: Climate Change and the Geophysical Underpinnings of Species Diversity*. Available at https://oak.ucc.nau.edu/pb1/Anderson_and_Ferree.2010.Climate_Change.pdf. PLoS ONE 5(7): e11554.

ANR Maps & Mapping. Vermont Agency of Natural Resources. Available at <http://www.anr.state.vt.us/site/html/maps.htm>, 2004. Web. 2 August 2015.

Austin, J. M., et al. 2004. *Conserving Vermont's Natural Heritage: A Guide to Community-Based Planning for the Conservation of Vermont's Fish, Wildlife and Biological Diversity*, Vermont Fish and Wildlife Department and Agency of Natural Resources, Waterbury, VT.

Chase, V., L. Demming, and F. Latawiec. 1995. *Buffers for Wetlands and Surface Waters: A Guidebook for New Hampshire Municipalities*. Concord, NH: Audubon Society of New Hampshire.

Doppelt, R., et al. 1993. *Entering the Watershed: A New Approach to Save America's River Ecosystems*. Washington, DC: Island Press.

Endangered Species. Available at <http://www.fws.gov/endangered/>. Department of the Interior, U.S. Fish and Wildlife Service.

Fidel, J., et al. 2013. *Community Strategies for Vermont's Forests and Wildlife*. Available at <http://vnrc.org/programs/forests-wildlife/guide/>. Vermont Natural Resources Council, Montpelier, VT.

Fish and Wildlife Considerations in Local Planning: A Vermont Review. September 2002. Waterbury, VT: Agency of Natural Resources, Department of Fish and Wildlife.

Fluvial Erosion Hazards Areas. 2010. Available at http://www.watershedmanagement.vt.gov/rivers/docs/rv_vtfqhqa.pdf. Waterbury, VT: Vermont Agency of Natural Resources, Department of Environmental Conservation, Vermont River Management Program.

Forman, R.T.T., et al. 2003. *Road Ecology – Science and Solutions*. Washington, DC: Island Press.

George, S., et. al. 1998. *State Endangered Species Acts: Past, Present, and Future*. Washington, DC: Defenders of Wildlife.

Local Planning and Zoning Options for Water Quality Protection. September 2001. Waterbury, VT: Vermont Agency of Natural Resources, Department of Environmental Conservation, Water Quality Division.

Native Vegetation for Lakeshores, Streamsides, and Wetland Buffers: What You Need to Know to Re-establish or Enhance a Buffer Strip Along Water and

Wetlands in Vermont. 1996, 2nd printing. Waterbury, VT: Agency of Natural Resources, Department of Environmental Conservation.

“Procedure on ANR Floodway Determination in Act 250 Proceedings.” Available at <http://www.vtwaterquality.org/riversmtg.htm>. Waterbury, VT: Vermont Agency of Natural Resources, Department of Environmental Conservation, Water Quality Division.

Rare and Uncommon Animals of Vermont: Vermont Natural Heritage Inventory. 2015. Available at <http://www.vtfishandwildlife.com/common/pages/DisplayFile.aspx?itemId=229831>. Montpelier, VT: Vermont Agency of Natural Resources, Fish and Wildlife Department.

Reay, R.S., et al. 1990. Management Guide for Deer Wintering Areas in Vermont. Waterbury, VT: Vermont Department of Fish and Wildlife, Forests, Parks & Recreation.

Riparian and Lakeshore Buffer Procedure Technical Papers. 2002. Waterbury, VT: Vermont Agency of Natural Resources, Department of Environmental Conservation, Water Quality Division.

Sorenson, E. and J. Osborne. 2014. Vermont Habitat Blocks and Habitat Connectivity: An analysis using Geographic Information Systems. Available online at <http://www.vtfishandwildlife.com/common/pages/DisplayFile.aspx?itemId=111635>. Montpelier, VT: Vermont Agency of Natural Resources, Fish and Wildlife Department and Vermont Land Trust.

“Technique Guidance for Determining Floodway Limits Pursuant to Act 250 Criterion 1(D).” Available at <http://www.vtwaterquality.org/riversmtg.htm>. Waterbury, VT: Vermont Agency of Natural Resources, Department of Environmental Conservation, Water Quality Division.

Thompson, E.H., and E.R. Sorenson. 2000. Wetland, Woodland, Wildland: A Guide to the Natural Communities of Vermont. Montpelier, VT: The Nature Conservancy and Vermont Department of Fish and Wildlife. Distributed by University Press of New England, Hanover, NH.

Thompson, E.H. 2002. Vermont’s Natural Heritage: Conserving Biological Diversity in the Green Mountain State: A Report from the Vermont Biodiversity Project. Montpelier, VT: The Nature Conservancy.

Tiner, R.W. 1987. Preliminary National Wetlands Inventory Report on Vermont’s Wetland Acreage. Washington, DC: Department of the Interior, U.S. Fish and Wildlife Service.

U.S. Department of the Interior et al. 2011 National Survey of Hunting, Fishing, and Wildlife-Associated Recreation. Available at <https://www.census.gov/prod/2012pubs/fhw11-nat.pdf>.

Vermont Endangered and Threatened Species Rule. 2015. Available at http://www.vtfishandwildlife.com/UserFiles/Servers/Server_73079/File/Learn

[More/Living with Wildlife/TE_Rule_2015.pdf](#). Montpelier, VT: Vermont Agency of Natural Resources.

Vermont Stream Geomorphic Assessment: Protocol Handbooks. Available at <http://www.vtwaterquality.org/riversmtg.htm>. Waterbury, VT: Vermont Agency of Natural Resources, Department of Environmental Conservation, Water Quality Division.

Verry, E.S., et al. 2000. Riparian Management in Forests of the Continental Eastern United States. Washington, DC: Lewis Publishers.

Wildlife Habitat Management for Vermont Woodlands. 1995. Waterbury, VT: Vermont Agency of Natural Resources, Departments of Fish and Wildlife and Forests, Parks and Recreation.



BIG GAME MANAGEMENT PLAN 2010 - 2020

CREATING A ROAD MAP FOR THE FUTURE

 **VERMONT**
FISH & WILDLIFE DEPARTMENT
(802) 241-3700 / www.vtfishandwildlife.com





Most of the programs described in this report are funded through the *Federal Aid in Wildlife Restoration Program*. This program was initiated in 1937 as the Federal Aid In Wildlife Act and created a system whereby taxes are paid on firearms, ammunition and archery equipment by the public who hunts. Today this excise tax generates more than one hundred million dollars each year that are dedicated to state wildlife restoration and management projects across the United States. The state of Vermont uses these monies for acquiring land, and for restoring and managing wildlife. These excise tax dollars, coupled with state hunting license fees have been the predominate source of money funding the successful restoration and management of Vermont's wildlife resources.

Vermont Fish & Wildlife Department

Agency of Natural Resources

103 South Main Street, 10 South


Waterbury, Vermont 05671-0501

(802) 241-3700 / www.vtfishandwildlife.com



The MISSION of the Vermont Fish & Wildlife Department is the conservation of fish, wildlife, plants and their habitats for the people of Vermont.

Many Vermont Fish & Wildlife Department programs receive federal aid in fish and/or wildlife restoration. Under Title 6 of the 1964 Civil Rights Act and Section 504 of the Rehabilitation Act of 1973, the U.S. Department of the Interior prohibits discrimination on the basis of race, color, national origin, or handicap. If you believe that you have been discriminated against in any program, activity, or facility described herein, or if you desire further information, please write to the office of Equal Opportunity, U.S. Department of the Interior, Washington, D.C. 20240.




Vermont's Big Game Management Plan

Creating a road map for the future.

I am very pleased to announce the completion of the 2010 – 2020 Vermont Big Game Management Plan. This plan will guide the Vermont Fish & Wildlife Department in its conservation and management of the state's deer, moose, black bear, and wild turkey populations during the next ten years. These four big game species provide tremendous hunting opportunities in Vermont as well as countless hours of wildlife viewing for all Vermonters who love and enjoy wildlife.

This big game management plan is the culmination of a long and deliberative process that melded the science of wildlife management with the interests of Vermont residents. This plan could not have been successfully completed without the benefit of citizens that responded to our surveys, attended our public meetings, and provided their comments, concerns, and ambitions for the future of our big game species. I also would like to acknowledge all Department staff who worked so very hard putting it together and listened to the views of Vermonters. They also had to deal with the tedious tasks involved in preparing the numerous drafts that led to this final document.

In the end, I am confident the implementation of this Vermont Big Game Management Plan will assure that deer, moose, black bear, and wild turkey management will improve and ensure that these species will be enjoyed in this state for generations to come.



Wayne A. Laroche, *Commissioner*
Vermont Fish & Wildlife Department
December 2009



Table of Contents

Executive Summary	vi
Chapter 1: Introduction	1
Historical Perspective	3
Management Issues of General Concern	6
Timeline of Important Dates in Vermont	
Wildlife Management	9
Chapter 2: White-tailed Deer	
Management History	10
1997-2006 Plan Accomplishments	12
2010-2020 White-tailed Deer Management	
Issues, Goals, and Strategies	15
Habitat Loss and Assessment	15
Population Goals	16
Hunter Participation, Satisfaction	
and Antler Point Restrictions	27
Bag Limits	30
Muzzleloader and Archery Season Modifications	30
Captive Deer Hunting/Deer Farming	32
Disease Surveillance and Management	33
Locally Overabundant Deer Populations	34
Two-year Regulation Cycle	35
References	36
Chapter 3: Moose	
Management History	40
1998-2007 Plan Accomplishments	42
2010-2020 Moose Management Issues,	
Goals, and Strategies	46
Regional Population Goals	46
Moose/Human Conflicts	48
Moose Hunting Opportunities	48
Moose Viewing	50
Moose Habitat	50
Deer-Moose Competition and Forest Impacts	51
References	52

Chapter 4: Black Bear

Management History	53
1997-2006 Plan Accomplishments	54
2010-2020 Black Bear Management Issues,	
Goals, and Strategies	55
Bear Population Size and Distribution	55
Bear Habitat Conservation	57
Human/Bear Conflicts	59
Bear Management Strategies and Season Structure	60
References	64

Chapter 5: Wild Turkey

Management History	65
1999-2008 Plan Accomplishments	67
2010-2020 Wild Turkey Management Issues,	
Goals, and Strategies	67
Turkey Population Assessment	67
Public Satisfaction With Current Population Levels	68
Fall Turkey Hunting	68
Wild Turkey/Human Conflicts	70
Turkey Habitat Management and Conservation	71
Perception Regarding the Interaction Between Deer and Wild Turkeys, Ruffed Grouse and Wild Turkeys, and Various Predators and Wild Turkeys	72
Developing and Maintaining an Informed Public is Crucial to the Management Success of the Wild Turkey Project	74
References	74



Wild animals, or wildlife, by Vermont law, belong to the people of Vermont. Conserving and managing Vermont's wildlife resources on behalf of the public are obligations of the Vermont Fish & Wildlife Department. The Department has a long history of managing Vermont's big game species. This long-range management plan will help identify goals, and management objectives to insure that conservation needs of the species and the interests of the public are effectively addressed. Below is an overview of the management issues, goals and strategies for each big game species.

2010-2020 White-tailed Deer Management Issues, Goals, and Strategies

ISSUE 1. Habitat Loss and Assessment

GOAL: To monitor changes in habitat quality and quantity and perform public outreach regarding habitat management techniques, so concerned citizens may help to secure their deer herd's future.

Management Strategies

- 1.1 Update inventory of deer wintering areas for local, regional, and state habitat planning and protection efforts.
- 1.2 Stress the importance of habitat conservation with outreach efforts to various segments of the public such as farmers, educators, hunters, forest managers, and land planners.
- 1.3 Work closely with foresters and entomologists to prevent, manage, and eliminate the threat of the hemlock woolly adelgid.

ISSUE 2. Population Goals

GOALS: 1) Maintain deer densities using regional population objectives.
2) Monitor biological characteristics of habitat and deer that can change in response to deer herd size through time.
3) Adjust antlerless deer harvests to alter population levels as necessary to achieve population objectives.

Management Strategies

- 2.1 Maintain and evaluate regional population goals, established during this planning period, that are based on deer densities that recognize a lower limit that is unsatisfactory to the public and an upper limit that is ecologically unsustainable.
- 2.2 Monitor deer herd health by collecting body condition data from hunter-harvested and road-killed deer.
- 2.3 Consider establishing habitat suitability criteria to define areas of suitable deer habitat within WMUs so that consistent and reliable density estimates can be made while allowing for habitat area estimate updates as new land-cover maps become available.
- 2.4 Evaluate bowhunter surveys to better estimate regional buck:doe and fawn:doe ratios; compare fawn production estimates to autumn fawn:doe ratios to estimate summer fawn survival, and use buck:doe ratios to estimate adult doe population through reference to the unbiased buck population estimate.
- 2.5 Continue remapping and surveying deer wintering areas so that available habitat is quantified and localized winter deer density is better documented.
- 2.6 Work with foresters to develop data-driven methods for assessing localized deer overabundance problems that might lead to development of localized deer management methods. Data must provide measures of forest condition.
- 2.7 Provide outreach to landowners regarding methods that may minimize damage and encourage reduction in locally overabundant deer populations. Investigate feasibility of a formal program to connect hunters with landowners to address locally overabundant deer populations.
- 2.8 Develop strategies to maintain enough big game registration stations to make big game reporting convenient for hunters.
- 2.9 Seek statutory changes to realign boundaries of select WMUs.

ISSUE 3. Hunter Satisfaction and Antler Point Restrictions

GOAL: Employ biologically responsible, socially responsive, and adaptive management of the deer herd.

Management Strategies

- 3.1 Collect adequate yearling buck data (weights, antler beam diameter, and number of points) from the youth hunt to detect and track any changes in the buck population resulting from the current antler-point restriction (two points-on-one-antler minimum), and evaluate biologically acceptable alternatives if needed.
- 3.2 Evaluate a model assessment using genetic data to examine the likelihood of altering the genetic diversity of the buck population via the current antler restriction.
- 3.3 Inform the hunting public about deer management issues and results of antler-point restrictions and gather input concerning deer management and hunter satisfaction.

ISSUE 4. Bag Limits

GOAL: Provide suitable utilization of deer as food and provide opportunity to hunt deer in a way that maximizes potential for effective deer population management but does not overstress the heavily harvested buck population.

Management Strategies

- 4.1 Provide the public with ample opportunity to harvest white-tailed deer for food and other utilitarian purposes.
- 4.2 Advocate for an appropriate deer bag limit that allows maximum hunter opportunity while achieving deer population management strategies.

ISSUE 5. Muzzleloader and Archery Season Modifications

GOAL: Provide suitable opportunity to hunt deer in a way that maximizes the potential for effective deer population management but does not interfere with hunters during youth weekend or rifle and other fall hunting seasons.

Management Strategies

- 5.1 Evaluate feasible options to expand antlerless deer-only hunting opportunities prior to the regular rifle season. These options will include, but are not limited to, an early muzzleloader season, expanded archery season, and increases in archery bag limits.
- 5.2 During the fall and winter of 2009-2010, survey public opinion on the various management options to achieve antlerless harvest objectives prior to the rifle season and develop a proposal of recommended hunting season changes for the Vermont Fish and Wildlife Board in 2010.

ISSUE 6. Captive Deer Hunting/ Deer Farming/ Cervid Importation

GOAL: Implement new captive hunting regulations and work with other state agencies to minimize the chance of introducing and transmitting diseases via captive deer.

Management Strategies

- 6.1 Evaluate the effectiveness of the captive hunting facility regulation.
- 6.2 Work with the Agency of Agriculture, Foods, and Markets and the deer farming industry to promote and enforce disease free importation and husbandry practices.

ISSUE 7. Disease Surveillance and Management

GOAL: Monitor disease issues and respond when necessary to protect the health of wildlife and/or humans.

Management Strategies

- 7.1 Work with associated branches of government (for example, Agency of Agriculture, Department of Health) to monitor and control disease agents and deer populations where and when it is appropriate.
- 7.2 Contribute to the national Chronic Wasting Disease (CWD) surveillance effort.
- 7.3 Monitor the progress of Hemorrhagic Disease as it moves toward the Vermont border.
- 7.4 Work closely with the Agency of Agriculture to ensure dairy farms and domestic deer farms maintain their tuberculosis-free status.
- 7.5 Investigate a prohibition on the use of deer-urine-based scent lures and, if appropriate, implement a public informational effort on the justification.
- 7.6 Inform Vermonters as to the gravity of CWD and repercussions if introduced into our environment through the dissemination of Vermont's CWD Response Plan.

ISSUE 8. Locally Overabundant Deer Populations

GOAL: Promote awareness that hunting is the only practical option to reduce localized overabundant deer populations.

Management Strategies

- 8.1 Demonstrate the effectiveness of archery hunting to reduce locally overabundant deer in Vermont's suburban environments.
- 8.2 Provide communities with up-to-date and comprehensive information on deer overabundance and consider community views when deciding how to best manage deer problems in suburban, agricultural, and forested areas.

- 8.3 Encourage communication and cooperation between antlerless deer hunters and landowners that seek relief from locally overabundant deer.

ISSUE 9. Two-year Regulation Cycle

GOAL: Consider a more efficient two-year regulatory cycle that allows for annual adjustments when environmental factors deem it appropriate.

Management Strategies

- 9.1 Provide outreach to legislators, board members, and hunters to develop an understanding of the rationale behind deer management and proposed actions to improve management.
- 9.2 Evaluate the benefits and deficiencies of implementing a two-year regulation cycle for deer season recommendations.

2010-2020 Moose Management Issues, Goals, and Strategies

ISSUE 1. Regional Population Goals

GOAL: To maintain regional populations of healthy moose at or below cultural carrying capacity.

Management Strategies

- 1.1 Maintain a statewide fall post-hunt population of between 3,000 and 5,000 moose.
- 1.2 Maintain a sex ratio of between 40 to 50 bulls per 100 adults (moose of at least age-class one).
- 1.3 Maintain an adult age-class distribution of at least 25% of at least age-class four.
- 1.4 Maintain an average ovulation rate of more than 1.15 for cows age class of at least three.
- 1.5 Assess relative moose habitat condition of individual WMUs or regions of the state using forest inventory data and a GIS-based Habitat Suitability Index Model.
- 1.6 Reduce and maintain WMU E moose densities to 1.75 moose per square mile (approximately 1,000 moose post-hunt).
- 1.7 Reduce and maintain WMU D2 moose densities to 1.0 moose per square mile (approximately 600 moose post-hunt).

- 1.8 Allow slow population growth in WMUs I, L, P and Q while not exceeding one moose per square mile.
- 1.9 Stabilize moose population in other WMUs at current levels.

ISSUE 2. Moose / Human Conflicts

GOAL: To minimize motor vehicle/moose collisions and other forms of damage caused by moose.

Management Strategies

- 2.1 Develop and implement a policy for Department response to “nuisance” moose.
- 2.2 Continue to cooperate with the Vermont Agency of Transportation (VTRANS) to erect warning signs at traditional moose highway crossings.
- 2.3 Cooperate with VTRANS in implementing roadside brush-clearing projects to improve visibility at the most dangerous moose crossings, when feasible.
- 2.4 Cooperate with VTRANS to investigate the use of new technology that may help reduce moose/vehicle collisions.
- 2.5 Continue with annual press releases to remind motorists of moose hazards during seasons of increased moose movements.

ISSUE 3. Moose Hunting Opportunities

GOAL: To maximize quality moose hunting opportunity.

Management Strategies

- 3.1 Provide quality moose hunting opportunity in all WMUs where feasible.
- 3.2 Coordinate with large property owners to enhance moose hunter access.
- 3.3 Provide information to hunters on how they can share moose meat with needy households throughout Vermont.
- 3.4 Conduct outreach efforts prior to any significant reduction in total permit numbers made in response to moose population changes.
- 3.5 Provide public opportunity to harvest moose for food and other utilitarian purposes.

- 3.6 Maintain and improve hunter satisfaction by managing a preference point lottery system.
- 3.7 Propose to implement a limited special archery-only moose hunting opportunity.

ISSUE 4. Moose Viewing

GOAL: Provide safe and quality moose viewing opportunity.

Management Strategies

- 4.1 Construct at least one moose observation tower with a parking area near a state highway in the Northeast Kingdom region and investigate other locations in other regions.
- 4.2 Include moose in a guide to wildlife viewing sites on the Department’s website.

ISSUE 5. Moose Habitat

GOAL: Maintain necessary habitat to support 3,000 to 5,000 moose on a sustained basis.

Management Strategies

- 5.1 Implement field studies to investigate, measure, and monitor the degree of moose and deer browsing within selected WMUs.
- 5.2 Provide natural resource professionals and landowners with moose habitat management guidelines.

ISSUE 6. Deer-Moose Competition and Forest Impacts

GOAL: Balance the nutritional needs of regional moose and deer populations with the need for adequate forest regeneration.

Management Strategies

- 6.1 Develop a study to assess the carrying capacity for moose and deer on Vermont’s forestland.
- 6.2 Develop a decision making process that assists managers in determining the appropriate mix of moose and deer densities for a given WMU based on cultural and ecological factors.

2010-2020 Black Bear Management Issues, Goals, and Strategies

ISSUE 1. Bear Population Size and Distribution

GOAL: Identify an appropriate bear population objective that ensures the viability of a wild, free-ranging bear population, provides for hunting opportunities, and satisfies human social expectations and tolerances for nuisance bear occurrences.

Management Strategies

- 1.1 Update and re-evaluate Vermont's black bear population model to reflect the most current harvest and biological parameter data available.
- 1.2 Evaluate and develop hunting season structures that align population estimates with biological data, habitat limitations, and public satisfaction data to sustain a bear population between 4,500 and 6,000 animals.

ISSUE 2. Bear Habitat Conservation

GOAL: Maintain a no net loss of function and value of existing bear habitat.

Management Strategies

- 2.1 Maintain and enhance habitat protection efforts through Act 250, wood-to-energy harvest review, work with town and regional planning commissions, land acquisition, and other conservation methods.
- 2.2 Provide technical assistance in managing for critical bear habitat in the Use Value program.
- 2.3 Revise and update "A Landowner's Guide, Wildlife Habitat Management for Vermont Woodlands" to include habitat management recommendations for black bears.

ISSUE 3. Human/Bear Conflicts

GOAL: Minimize the overall number of negative interactions occurring between bears and humans to achieve acceptable levels of human safety and social acceptance.

Management Strategies

- 3.1 Update statewide policy for handling black bear/human conflicts.
- 3.2 Improve and disseminate outreach/education materials and messages for minimizing human/bear conflicts.
- 3.3 Monitor bear/human conflicts and explore new strategies for reducing the number of complaints from the public.
- 3.4 Use permitted houndsmen with trained bear hounds to haze bears and keep them wary of humans.

ISSUE 4. Bear Management Strategies and Season Structure

GOAL: Optimize public hunting opportunity for the utilization of bears for food and other appropriate purposes and ensure hunter satisfaction within biologically sustainable regulations.

Management Strategies

- 4.1 Hunting season management strategies and season structure will be evaluated and adjusted to maintain the population goal of 4,500 to 6,000 bears. Changes in hunting season structure will consider, when necessary, the use of season length, regionalization, or incremental changes to season bag limits to achieve population goals.
- 4.2 Work with partner organizations on issues related to bear management as they are raised throughout the management plan period and develop specific strategies to address them. Such strategies may range from legislative changes to educational efforts.

2010-2020 Turkey Management Issues, Goals, and Strategies

ISSUE 1. Turkey Population

GOAL: To adequately assess Vermont's wild turkey populations and trends.

Management Strategies

- 1.1 Annually collect and assess turkey harvest data to determine trends as well as summer/fall turkey

sighting survey data in order to direct future management decisions.

- 1.2 Conduct the public annual Internet turkey brood survey along with the Department staff summer turkey survey.
- 1.3 Continue the turkey program's investigation into the genetic variability and structure of the statewide population.
- 1.4 Evaluate new wild turkey population estimation methods and models for use in Vermont.
- 1.5 Evaluate the use of a public Internet survey to assess winter flock sightings.

ISSUE 2. Public Satisfaction with Current Population Levels

GOAL: Assess public and hunter satisfaction with current turkey population levels and management program.

Management Strategies

- 2.1 Provide statewide spring bearded-bird-only seasons (including the Youth and regular May season) and limited fall either-sex hunting seasons in WMUs that can sustain a fall harvest so as to provide for population stability.
- 2.2 Prioritize high quality spring hunting over additional fall harvest opportunity.
- 2.3 Manage fall turkey harvests through changes in fall hunting season length within WMUs depending upon stability or growth of three-year average spring harvest densities, except in WMU A Champlain Islands where inadequate forest cover exists to sustain a fall firearm harvest.

ISSUE 3. Fall Turkey Hunting

GOAL: To provide appropriate opportunity for sustainable fall hunting while maintaining current levels of high quality spring turkey hunting.

Management Strategies

- 3.1 Provide public opportunity to harvest wild turkey for food and other utilitarian purposes.
- 3.2 Facilitate healthy, abundant spring turkey populations that are stable using modest, fall hunting seasons/bag limits to control the

population. When the three-year spring average harvest density reaches the specific threshold value, liberalization of fall hunting in a WMU may be called for (initiate shotgun seasons, extend gun seasons).

- 3.3 Consider reducing the current guideline for the threshold as to when fall gun hunting opportunities could be initiated in a new WMU, from the three-year average spring harvest density of one bird per square mile, to an average harvest density of .75 bird per square mile.
- 3.4 Lengthen the current fall seven-day shotgun season to a nine-day season.
- 3.5 Expand the fall shotgun season to include WMUs H1, D1, and B with a nine-day shotgun season.
- 3.6 Expand the fall archery turkey season, coinciding with the opening of the deer archery season, to allow archery hunting statewide.
- 3.7 Investigate establishing a new separate "Fall Gun Season Only" tag.

ISSUE 4. Wild Turkey/Human Conflicts

GOAL: To minimize and manage agricultural damage and nuisance turkey incidents.

Management Strategies

- 4.1 Provide property owners with access to coordinated services of personnel trained to deal with nuisance turkey issues including wildlife biologists, game wardens, and USDA Wildlife Services staff to assist with nuisance complaints via technical guidance/assistance on techniques to minimize/discourage damage.
- 4.2 Conduct follow-up site visits to nuisance complaint sites when necessary and provide hazing equipment to help ameliorate persistent nuisance situations.
- 4.3 Solicit assistance from local volunteers through the Vermont Chapter of the National Wildlife Turkey Federation (NWTf) to help provide on-the-ground assistance to landowners via hazing and behavior modification efforts.
- 4.4 Assist USDA Wildlife Services staff with development of educational materials to inform and educate farmers about techniques for minimizing conflicts.

- 4.5 Compile and evaluate wild turkey damage complaint reports from farmers, state game wardens, biologists and wildlife service personnel to document problems, management approaches and results.
- 4.6 Develop/modify a standard set of protocols/guidelines/solutions to perceived and actual conflicts caused by wild turkeys (nuisance animals, agricultural damage).

ISSUE 5. Turkey Habitat Management and Conservation

GOAL: To encourage conservation and appropriate habitat management practices to support and sustain Vermont's wild turkey population.

Management Strategies

- 5.1 Continue efforts on wildlife management areas and other public lands to provide habitat demonstration areas to promote appropriate commercial and noncommercial vegetation management practices beneficial to turkeys and other wildlife. This includes the use of prescribed fire and other management practices to establish and maintain long-term mast production areas.
- 5.2 Provide technical information and assistance regarding turkey habitat management to private landowners and other land managers, town planning commissions via staff biologists, habitat demonstration projects, LIP and WHIP program lands, etc.
- 5.3 Update the "A Landowner's Guide, Wildlife Habitat Management for Vermont Woodlands" and make this document available on the Internet and in published copy as well.
- 5.4 Work with the NWTF regional biologists and chapter volunteers on development of the North American Wild Turkey Management Plan.
- 5.5 Work with partnering organizations on high priority projects and issues.

ISSUE 6. Perception Regarding the Interaction Between Deer and Wild Turkeys, Ruffed Grouse and Wild Turkeys, and Various Predators and Wild Turkeys

GOAL: To improve the public's knowledge, awareness, and understanding of the role of the wild turkey and its interactions within the ecosystem.

Management Strategies

- 6.1 Promote sound scientific principles regarding inter-species competition and predator-prey relationships through a variety of outreach methods including public speaking events, web-based information and links, and print and broadcast media.

ISSUE 7. Developing and Maintaining an Informed Public is Crucial to the Management Success of the Wild Turkey Project.

GOAL: To ensure continued information exchange and program acceptance by keeping the general public, state and federal agencies informed on the status of the wild turkey resource in Vermont.

Management Strategies

- 7.1 Disseminate wild turkey project information to the public/media professionals via biological reporting stations, teacher workshops, private and public landowner visits/conferences, slide/video presentations, mail correspondence, popular and technical reports, etc.
- 7.2 Use the Department's library to fill all public requests for its video production "The Wild Turkey in Vermont" as well as its wildlife study guide "The Wild Turkey Education Kit."
- 7.3 Continue involvement with standing professional committees, regulatory bodies and cooperative agreements with nongovernmental organizations to assist the Department with meeting the goals and objectives of this plan.

BIG GAME PLAN - INTRODUCTION

Managing Wildlife — A Public Trust

Under federal and state law the management of wildlife falls under the concept of public trust, which means that it is considered a resource that must be preserved and protected for public use. Unlike Europe's feudal system during the Middle Ages, wildlife does not belong to a royal family or a government. Nor can individuals possess wild animals as a commodity as pets or farm animals. The Public Trust Doctrine, based on English Common Law and upheld by the United States Supreme Court, is the principle upon which natural resources, such as wildlife, are conserved in the public interests and for reasonable use by current and future generations. The Vermont Fish & Wildlife Department (the Department) is obligated to conserve and manage Vermont's wildlife resources on behalf of the public. Vermont law entrusts the stewardship and management of wildlife resources to the Department in accordance with the Public Trust Doctrine to ensure this principle is carried out.

The principle of wildlife as a resource that is managed in public trust by state and federal governments is the foundation of what is known as the North American Wildlife Conservation Model. The Model holds that by placing wildlife in the public trust the value that is derived is not merely personal profit. The motive for harvesting wildlife is not one of simple profit as it was in the nineteenth century when market hunting was rampant, but instead, one of broad public benefit and sound and sustainable wildlife and habitat management. This Model has served wildlife and the public well for more than 100 years. As a result, game species such as the four big game species featured in this plan have flourished. Under this Model, the public is involved in the decision-making process, and for this reason, it has been embraced across North America. In keeping with these basic principles of wildlife management and conservation in North America, the mission of the Department is "...the conservation of fish, wildlife, plants, and their habitats for the people of Vermont."

About This Plan

To carry out this mission the Department's long-range management plan identifies issues, goals, and strategies that insure that a balance between the conservation needs of the species and the interests of

the public is effectively addressed. The plan has three major objectives:

- Conserve, enhance and restore Vermont's natural communities, habitats, and plant and wildlife species along with the ecological processes that sustain them.
- Provide a diversity of safe and ethical fish- and wildlife-based activities and opportunities that allow hunting, fishing, trapping, viewing, and the utilization of fish, plants, and wildlife resources consistent with the North American Wildlife Conservation Model.
- Maintain safe fish- and wildlife-based activities while limiting harmful human encounters with fish and wildlife species, and provide general public safety service incidental to our primary fish and wildlife duties.

Management of Vermont's four big game species has been combined into a single, comprehensive big game plan. This will provide the public with easy access to all information related to big game management. It will also help ensure that a more comprehensive assessment of the overlapping and divergent management needs of each big game species are holistically considered and coordinated to improve overall management. In addition, the process of developing a single, comprehensive big game management plan is more cost effective and efficient than four separate planning efforts.

The Process for Developing The "Ten-Year Big Game Management Plan"

This plan is based on currently available and relevant biological and ecological data associated with each of the four big game species and their habitats. A survey of 1,000 randomly selected Vermont residents was also conducted to gather public opinion related to deer, moose, bear and wild turkey management. Respondents were asked their views on many topics such as habitat protection, game species population size preferences, and property damage from wildlife (results of the survey can be found on the Department's website: www.vtfishandwildlife.com/library/)

A series of open house style public meetings were held in five locations around the state during the

summer of 2008 and a web page was developed to allow people to discuss issues and offer opinions to Department staff. This was followed by two public meetings and a month-long public comment period to allow Vermonters to submit opinions regarding draft management plans. Approximately 200 people attended the meetings, wrote letters and e-mails, used the on-line comment option, or made phone calls to express their views. The majority of comments pertained to deer management, with comments varying widely across the topics of season lengths, bag limits, and appropriate antler point restrictions. All of the comments were reviewed and considered by the Department and, as much as possible, assimilated into commonly voiced themes. While biologically responsible wildlife management must come before public opinions, there will always be aspects of wildlife management that can be decided by public sentiment. From the beginning, public feedback steered many aspects of the Department's ten-year planning efforts, and public feedback will continue to help shape our goals and objectives. Provided below are summaries of the issues raised by the public in response to the draft plans for each of the big game species.

WHITE-TAILED DEER

The Department received input on several potential strategies to address the need to harvest more female deer in select parts of the state. First, where female deer are locally overabundant, it may be desirable to encourage bow hunters to fill a second, or even third, archery tag by taking an antlerless deer. This could be achieved by liberalizing the archery bag limit to three deer, with one of these possibly being a buck. All hunters would still be subjected to the annual bag limit that is currently three deer. The option to "tag-out" with three deer during archery season and forfeiting further deer hunting in Vermont in that calendar year would be the choice of the hunter if he or she were successful during this season. It is also noteworthy that additional archery tags may help manage localized deer populations where firearm ordinances restrict the ability to harvest antlerless deer during the Youth Weekend and muzzleloader seasons.

Second, there was considerable interest voiced for an early antlerless-only muzzleloader season that would occur sometime before the regular rifle season. Such a season would only be open to those individuals holding an antlerless-deer permit for muzzleloader hunting. Although the Department is sensitive to the various concerns expressed by hunters, landowners,

and other nature enthusiasts regarding this policy, the Department needs to explore ways to harvest more antlerless deer in some areas when and where consecutive mild winters allow the deer herd to grow beyond our ability to control it with existing antlerless deer hunting seasons. A brief antlerless-only early muzzleloader season is a method to consider with other potential benefits from removing more antlerless deer earlier in the season.

Because of increased interest in an early muzzleloader season and antler-point restrictions, a survey on these topics was circulated at the July public hearings. As recommended by some of the attendees, the survey was also posted on the Department's website. The survey response was substantial, numbering nearly 600 submissions. Additional public surveys that solicit opinions on the use of early muzzleloader season and/or the archery season to achieve female deer harvest objectives will guide the Department's management approaches in the future.

The public provided mixed reviews of a special crossbow season or allowing crossbows during the archery season. Given the consideration of other, more popular antlerless harvest enhancements, such as the early muzzleloader season and a lengthened archery season, the use of crossbows as an additional hunting implement does not appear to be supported by the public at this time.

In preparing the final plan, there were two areas in particular that appeared to require further scientific documentation. Participants requested more information about chronic wasting disease (CWD), how it is transmitted, and what it means for CWD-free and CWD-infected deer populations. Of particular interest was how this might apply to deer-urine-based scent lures. Although widespread live-testing for CWD still remains unfeasible with high probabilities of false-negative results (indications of a disease-free animal when it is actually infected), methods to detect the infectious protein (prion) in animal fluids is advancing. As a result of these advancements, recent studies have found the CWD-causing prion in urine and other excretions and body parts of infected deer. In this final plan we provide additional and current references to pertinent scientific literature and results on this subject.

The other topic needing more supporting documentation was the issue of antler-point restrictions. Again, we provide additional sources of information cited in the text. Many of the studies

cited are available as complete documents on the Internet. Copyrighted studies on the Internet appear as abstracts that may be purchased through the journal in which they are published.

MOOSE

Among the persons commenting on the Department's website, seven made comments related to the draft moose plan. The only issue that drew much attention was the proposed special archery season for moose; six people expressed support for this season, and none were opposed.

The other major issue where public input was specifically solicited in the plan was the proposed management for slow growth of the moose population in the central and southern "mountain" wildlife management units (WMU) of I, L, P, and Q. Only one web comment addressed this issue, and it was in favor of the proposed direction.

Six written moose comment forms were collected from open houses held in July, 2009. Five of these were in support of the special bow season and one was opposed. Similar levels of support (seven in favor, one opposed) were voiced at the May, 2008 open houses. None of the comments addressed population desires for the southern mountains, except for one respondent who desired fewer moose in WMUs H1, H2, D1, D2.

Six respondents commented on the moose lottery. One liked the present system; two thought it was unfair because some families have won multiple times while others have never won. One individual thought bonus points should be earned during the three-year waiting period, one wanted a two-year wait instead of three-years, and one person felt applicants should possess a Vermont hunting license before they could enter the lottery.

BLACK BEAR

Two bear management issues received the most comment from 15-20 respondents. The first was opposition to a regulation requiring minimum registration standards for bear hunting guides or hunting guides in general as a means to address concerns for fee-for-bear hunting. Some felt it would diminish a person's opportunity to earn money or offer a potential mentoring experience for an inexperienced hunter. Others felt a guide registration system did not address the fee-for-bear hunt guiding concerns.

The second most frequent comment concerned nuisance bear situations, especially those involving birdfeeders. Most felt the Department had a good message regarding the removal of feeders but needed to be more aggressive with its advertisement and insistence with compliance. Beehive owners expressed some concern about higher bear populations in the Champlain Valley where apiaries are numerous. The consideration of regional management zones for bear seasons may be an appropriate tool for addressing this concern.

Several respondents wrote to say the bear population was "about the right size," or that it appeared to be growing, and the population goal was appropriate. Others felt the population was too high in parts of the state and suggested managing bear populations by regions to address these differences, while still others felt the bear population was too low.

Comments were received regarding opposition to bear hunting, especially with the aid of dogs. The Department believes it cannot achieve and maintain the proposed bear population objectives without the use of regulated hunting of these animals.

WILD TURKEY

A number of substantive comments were received during the public comment period for the draft wild turkey management plan. These comments ranged from suggestions for a variety of spring and fall hunting season expansion proposals to "maintaining the status quo" to comments on the availability of check stations for reporting harvest. The general focus of comments pertained to the opportunity to expand fall hunting opportunity. One comment of interest suggested a separate fall bird tag to enable hunters to harvest a fall bird who might not otherwise participate in the spring season. A second comment of interest suggested opening the fall season concurrently with archery deer season to permit greater opportunity to hunt turkeys.

Historical Perspective

The following is a brief overview of the historical influences on wildlife in Vermont. It traces some of the most important elements of early land use activity and cultural trends that have affected the state's wildlife and its habitat.

Prior to European colonization in North America in the early 1600s, human activity affected the landscape very little. Native Americans did not

have the technology, other than fire, with which to create landscape-level changes in their environment. Thus, sporadic, naturally occurring events such as hurricanes, earthquakes, and wild fires were the primary forces affecting large geographical areas. Even these major events merely served to set back forest communities to earlier stages of ecological succession. With seed stock and soil still in place the forest communities were always capable of replacing themselves.

Indigenous tribes hunted, fished, trapped, and tended corn fields and small gardens on the banks of major rivers throughout Vermont and New England. Some species of fish and wildlife were very abundant. The passenger pigeon, for example, was so plentiful that it was reported the birds “blacken the sky” with their numbers and broke tree branches when they set down to roost. This single species accounted for 25 to 40% of all birds living in the United States. According to some records, there were 3 billion to 5 billion passenger pigeons at the time Europeans began arriving on the continent. Ducks, geese, deer, moose, and many other species were also plentiful. For native communities wildlife was a primary source of sustenance and socially and culturally important. But just as today, wildlife populations fluctuated through the years and varied with the seasons, and there were times of wildlife scarcity.

The balance between wildlife and human activity, however, changed dramatically in the 1800s with the influx of European settlers. Following establishment of the colonies, the human population increased steadily in Vermont. Just prior to Vermont becoming the fourteenth state, its population was estimated at 85,425. Over the course of the following ten years, the population doubled to 154,465. Just 50 years later, it doubled again (Table 1.1). European settlers changed the wildlife equation in several important ways.

Unregulated market hunting and hunting wildlife for profit rather than for subsistence contributed to a rapid decrease in many species. Another factor was the settlers’ demand for lumber and firewood, as well as land to convert to agricultural use. Throughout Vermont’s early history the landscape has shifted with changes in farming – from sheep to dairy farming, from grass crops to a corn crop. But on a larger scale, farming transformed the land from forests to open pastures. At

one point in our history, the land went from 95% forested to 63% nonforested, eliminating most or nearly all suitable habitat for some species. This, along with the unregulated harvest of wildlife, took a significant toll on many wildlife populations that depended on forestland habitat. By the mid-1800s, many of the species that had been very abundant began to decline or disappear from the landscape. The passenger pigeon, mountain lion, wild turkey, moose, and wolf became extinct, while deer and bear populations were limited to forested remnants of the state.

As early as 1847, famed conservationist and resident of Woodstock, Vermont, George Perkins Marsh remarked on the speed with which this transition to a nonforested landscape occurred. The ecological damage sustained by farming and logging, noted Marsh, was “too striking to have escaped the attention of any observing person.” Governor Urban M. Woodbury angrily proclaimed before the State Legislature in 1894, “Owners of timber lands in our state are pursuing a ruinous policy in the method used in harvesting timber.” The Governor recognized that the deterioration of forestland in Vermont also meant an insecure future for the state’s major industry: lumber and wood products. “There is no more valuable crop produced from the land than timber,” Woodbury commented in the same speech. “Every decade will see timber more valuable and it is of great importance to the state as a whole... that some measure should be adopted to lessen the wanton destruction of our forests.” Although Marsh and Woodbury were early observers of the fact that Vermont’s economy was tied to the resources and aesthetic qualities of its forests, public awareness and concern regarding the effect of certain land practices on the natural environment did not fully emerge until the turn of the twentieth century.

As concern for the loss of species took root among citizens in Vermont and across North America, actions began to be taken to restore the wildlife species that had been lost. Deer were one of the first species to be protected by state laws. In 1865, the hunting of deer in Vermont became illegal and remained so for the next 32 years. During this period,

Table 1.1 Vermont population from US Census Bureau statistics 1790 - 2005.

Year	1790	1800	1850	1900	1950	2005
Population	85,425	154,465	314,120	343,641	377,747	623,050

seventeen white-tailed deer were transplanted from New York into the state, which provided breeding stock that rebuilt the deer herd. The most important change, however, that led ultimately to successful restoration of white-tailed deer and other species was the abandonment of farms that allowed the land to revert back to forests. The combination of improving habitat conditions, legal protection, and lack of significant mortality factors other than winter conditions resulted in a rapid recovery of the deer population. The rapid success of this restoration effort led to the opening of a limited, regulated deer hunting season in October of 1897.

As Vermont entered the era of active wildlife management, the deer population continued to grow as habitat expanded and improved. Throughout the first half of the twentieth century, deer numbers increased and their range expanded. During this time of restoration, a bucks-only harvest regulation was used to maximize the growth rate of the deer population. Because only bucks were harvested for a period of more than fifty years, the buck-only harvest restriction moved from being a population management tool to becoming a Vermont deer hunting tradition.

The white-tailed deer population responded well to the bucks-only regulation and expanded so quickly that in less than 50 years the buck harvest grew from 103 deer in 1897 to more than 4,000 deer in 1940. So rapid was the population growth that by 1946 wildlife biologists had already begun to observe negative impacts on habitat quality caused by large numbers of deer. In this same year, the Department released the publication "The Time is Now" as an attempt to inform the people of the new situation and the problems that the future would hold if growth of the deer population was not limited. Biologists recognized that the harvest of female deer was the only way to control total deer numbers. Unfortunately, Vermont's bucks-only harvest tradition had become well established by this time and the hunting public would not accept harvests of female deer as the solution.

This difference of opinion engendered an infamous period of deer management in Vermont that became known as the "deer wars." Over the next nearly 50 years, public outcries occurred on and off as biologists attempted to implement deer management changes. Although most deer hunters today recognize the importance of harvesting female deer to limit growth of the deer herd and protect deer habitat, deer

management remains an area of great public concern and continued contention.

The other three big game species did not recover as quickly as the deer. Bear populations recovered slowly for several reasons. Livestock owners considered them a nuisance. Not only was bear hunting unregulated, but Vermont state law offered bounties for animals that were killed from 1831 until 1941. The first laws limiting the harvest of bears were not implemented until about 1950. Rapid recovery of forest habitat along with limited harvest of bears proved to be a boon to black bear recovery. Black bears are now distributed throughout most of Vermont.

Moose also may have completely vanished from Vermont at one point. When a young bull was shot in March 1899, at Wenlock (now Ferdinand) in Essex County, the local newspaper reported it as "a strange animal" and "the last moose in Vermont." The shooting was actually illegal because the 1896 Legislature had established a closed hunting season on moose. Moose recovery lagged behind deer and bear due to a lack of suitable forest and wetland habitat. But as the forestlands recovered and wetland habitat expanded with the return of beaver populations, moose habitat also expanded. The Department estimated that in the early 1960s about 25 moose existed in Essex County. The moose population grew steadily over the next 30 years. By the 1990s, moose were abundant enough to support a limited, controlled regulated hunt.

In 1993, the Department issued 30 moose permits in Essex County and conducted the first regulated moose hunt in the state's history. Today, the moose population has fully recovered and has reached a level where regulated hunting is a tool needed to keep the population in balance with its habitat and to protect private property and public safety. The Department's current management aims are to keep the moose population in balance with available habitat and to provide abundant hunter harvest and citizen viewing opportunities.

By the mid-1800s, wild turkey was another species that had disappeared from the Vermont landscape. Thirty-one wild turkeys from New York were stocked into Vermont in 1969 and 1970. From this point on, the turkey population grew so fast that the first modern turkey hunt was established only three years later in 1973. Less than 40 years later, the turkey population has expanded throughout the state and continues to grow in numbers with record

harvests occurring annually. Today these birds can be found in nearly every town of the state with a total population thought to number approximately 50,000 to 55,000 birds. This was an unexpected outcome. Early biologists believed that Vermont's long winters and deep snows would limit the distribution of wild turkeys to the Champlain and Connecticut River Valleys where winters are less severe, acorns are plentiful, and agriculture provides a source of winter food. Wild turkeys proved to be more adaptable than anticipated, however, and today they are found even along the Canadian border in Essex County. In fact, wild turkeys have expanded their range across the border into Quebec, Canada.

The twenty-first century begins with approximately 75% of Vermont's landscape being forestland. A half-century of science-based regulation has restored many wildlife species, including game species. Conservation and management issues, however, still confront our deer, moose, bear, and wild turkey populations. Although these issues more often relate to overabundance than to scarcity and recovery, they are no less daunting. The issues surrounding our wildlife in this century are now focused on maintaining wild and robust populations in balance with their habitats while providing abundant opportunity for the public to use and enjoy. Today the issues we face involve an ever expanding human population and the activities that accompany it (Table 1.1). Bears in backyards, moose in urban areas, turkeys damaging agricultural feed crops, and deer eating the next generation of forests have now replaced the old issues of wildlife scarcity. The loss and fragmentation of habitat associated with development presents new challenges to the conservation and management of deer and other species of Vermont's wildlife. If land ownership in Vermont continues to be divided into ever smaller parcels, available space to hunt and opportunity to access game will become an increasing challenge.

The Benefits of Fish and Wildlife Based Outdoor Opportunities

Hunting, fishing, and trapping are important outdoor activities culturally, socially, economically, and ecologically. These activities conducted under regulated seasons provide for sustainable utilization of fish and wildlife resources statewide. Currently 30% of Vermonters fish or hunt (over 80,000 hunters and 121,000 anglers), a higher participation rate than skiing (19%). Recent surveys report that Vermont is third nationally (behind

Alaska and Maine) in per capita participation by the public in hunting, fishing, trapping, feeding and observing wildlife. Over 600,000 pounds of white-tailed deer, 192,000 pounds of moose, and 15,000 pounds of black bear meat are harvested annually from the forests and wetlands of Vermont. Wildlife related outdoor activities accounted for 5% of Vermont's gross state product in 2001, with nearly \$300 million spent on fishing and hunting alone. These expenditures particularly benefit rural areas of the state and occur when tourism is typically low in Vermont. Within the context of this ten-year plan, the Department examines four of Vermont's big game species with the goal of managing these as assets to perpetuate into the future for the various cultural, social, economic, and ecological values they bring to the state of Vermont.

Management Issues of General Concern

1. Habitat Loss. Loss of critical habitat, such as deer yards and bear feeding areas, can occur as a consequence of development that fragments habitat as well as results in mortality from increased animal movement and motor vehicle collisions. Maintaining an adequate supply of quality, inter-connected habitats in a variety of forms (for example, young forests or wetlands) that sustains viable wildlife populations is one of the most significant conservation challenges given today's issues of sprawl and parcelization of land. For example, it is estimated that a black bear in Vermont requires 10,000 acres of land to successfully meet its annual life needs. Therefore, it is essential that sufficient habitat be maintained, managed and connected through travel corridors in order to sustain a healthy, productive population of black bear.

2. Hunter Demographics. During the last 100 years, regulated hunting has served to effectively provide people with food in terms of a sustainable, renewable wildlife resource and a continuous opportunity to be afield pursuing game. It has also served as a highly effective tool to regulate population size to levels that are compatible with habitat limitations and human expectations. Nationwide, hunters have declined over the past decade while the general population has grown (U.S. Department of Interior 2006). While the national average for annual hunting participation declined to only 5% (U.S. Department of Interior 2006), it was

14% in Vermont (Duda et al., 2007). About 41% of Vermonters have hunted at some time (Duda et al., 2007), indicating that hunting remains an important tradition here. Concern remains that reduced numbers of hunters may make it difficult to harvest enough deer to control the population in the future.

Since 1997, various youth hunting seasons for big game have been established to promote opportunities for youth to participate in hunting under the mentorship of an adult hunter. Youth Weekend seasons now exist for deer and wild turkey. Interest and support among adult hunters for these programs remains high.

3. *Public Access to Land.* Private lands remain very important to most Vermont hunters. One study estimated 30% of Vermonters still travel less than five miles one-way to hunt deer (Duda et al., 2007). Public lands open to hunting are under various ownerships and are distributed widely across Vermont with a total of more than 800,000 acres under state or federal management. The Vermont Agency of Natural Resources manages more than 333,000 acres of this total as wildlife management areas, state forests, and state parks. The Agency also holds easements on over 123,000 acres of conserved commercial forestlands that guarantee public access. The Green Mountain National Forest and Silvio Conte National Wildlife Refuge comprise most of the federally owned public lands in the state.

The value of private lands for hunting and other public access is recognized by laws ranging from Vermont's strict landowner liability laws to statutes granting landowners who own at least 25 acres a preferred status for receiving antlerless deer muzzleloader permits. Because of the latter, the Department is opening up more private land for hunting by offering these landowners first choice in the kind of permits that are issued for hunting on their land. The Department encourages hunters to ask permission and be respectful of private lands even when lands are not posted to ensure that Vermont's heritage of free access to private lands for hunting may continue indefinitely.

4. *Privatization of Wildlife.* Privatization of wildlife resources threatens fair chase hunting

wherever it occurs. When private landowners erect high fencing and charge a fee for the opportunity to hunt, the privatization of a public wildlife resource has occurred. Access for pay or lease hunting systems that restrict land access to those having the money to pay for it is a similar but less direct form of privatization. As demonstrated in much of Texas, lease hunting systems result in reduced hunting pressure and an inability for state wildlife agencies to manage overabundant deer populations (Haskell 2007). In accordance with the founding principles of this nation and the state of Vermont, it is the Department's responsibility to prevent privatization of Vermont's public wildlife resources and ensure the public's right to hunt.

5. *Human-Wildlife Conflicts.* The Department faces increasing conflicts between humans and wildlife. The four big game species present unique cases involving nuisance and other human conflicts. The Department addresses these issues in a consistent fashion for big game species in accordance with the following principles: Protection of human health and safety is first. Second, we must handle the animal involved responsibly when it must be confronted, displaced/removed from the scene, or euthanized. When these two guidelines are met, public acceptance is usually achieved.

6. *Loss of Big Game Check Stations.* These facilities perform a vital data collection service to the Department and provide a convenient means for hunters to legally register their game. The number of check stations has steadily decreased during the last ten years to a point where some hunters now have to drive 30 or more miles to legally report their game. There are a number of reasons for the decline including the time required to record a harvested animal, the small fee received for the effort, and change in ownership of stations. While hunters and others visit the check stations during hunting seasons and make purchases of materials, goods, and products, in some instances the agents believe this ancillary business is insufficient to cover the costs of participating as a reporting station. Big game registration and sale of licenses are a tremendous benefit to the Department and to the hunting public. The Department is examining a variety of strategies to correct this situation.

7. Access to Game. As Vermont's population approaches 650,000, land continues to be developed and subdivided into smaller parcels, resulting in less available habitat for wildlife and fewer opportunities for hunters to access private land. Houses and people now occupy areas that were once open to hunting, posing a safety risk that limits the area where hunters may use their firearms. Posting no trespassing signs on private property also limits the amount of lands available for hunting. To stem the loss of access to game, the Department remains committed to public land acquisition programs (for example, Forest Legacy) that contribute to the acreage available for public hunting. The Department also recognizes the increasing negative impacts of a third-party fee for hunting on private lands. In these cases, individuals or groups of individuals lease hunting privileges to a sought-after hunting location from a landowner and charge clients for the exclusive use of the land. Or the landowner charges a select few directly for use of the land. In either case, access to game is just as restricted as if the land were developed or posted. This reduces hunting opportunity for the hunter without the financial means to buy into the hunting privilege. Examples of this have been readily seen with waterfowl hunting and more recently with bear and deer hunting. As the willingness to pay to hunt increases more and more opportunity will be lost to the general hunting public. Furthermore, the redistribution of hunting pressure due to fee hunting will likely become inconsistent with game species management goals.

Enacting rules against fee for hunting may appear to be a simple solution, but private property rights require that this type of response be carefully weighed before moving in the direction of regulation. It is also important to distinguish the difference between a fee for hunting versus a fee for a guided hunt. The former involves restricting access to hunting land while the latter, as in the case of moose hunting guides, offers a service but does not prevent non-paying hunters access to hunting space. Efforts need to be increased during the next ten years to address the fee-for-exclusive-hunting tide. These efforts must include outreach towards landowners by the Department, organized sporting groups and individual hunters.

8. Wildlife Management Unit (WMU)

Realignment. Wildlife management unit boundaries were established in 1979 to regulate deer harvest on a geographical basis where deer densities mirrored the effects of habitat quality and winter severity. Since that time, WMUs have been applied to the management of moose and wild turkey populations on a regional basis. Bears range across such large areas of land that individual WMU boundaries have lesser value as a management tool. Groups of WMUs, however, can be established that may provide feasible opportunities to manage this species on a regional basis.

Unit boundaries, however, do not in every case align with natural boundaries of population abundance of big game species, particularly deer. To more effectively manage deer populations, it is necessary to periodically reassess and realign unit boundaries. A detailed description of the proposed changes in WMU boundaries is provided in Chapter 2, "Deer Management Plan," Issue 2 Population Goals. It is important to note any realignment of WMU boundaries will apply to all big game species.

Habitat loss and an aging hunter population are significant barriers to meeting the goals of this plan and to wildlife conservation in general. The economic and social forces affecting these changes are diverse and will be part of the Department's focus in addressing these new conservation challenges. The Department will consider a variety of opportunities to address these issues including but not limited to the following:

- Increase hunter recruitment and participation through a variety of strategies, such as introducing families to safe shooting through workshops sponsored by 4-H.
- Develop outreach materials for private landowners to introduce them to the wildlife management services offered by the Department, the rules and regulations concerning hunting on private property, and acquaint them with the traditions of Vermont's rural culture associated with hunting.
- Improve hunter access to land through a variety of strategies including creating

incentives to reduce parcelization of private property.

- Ensure that Project WILD, Project WET, and Project Learning Tree materials are in the hands of all elementary school teachers.
- Improve and expand hunter education opportunities through a variety of strategies including creation of a mentored hunting program.
- Help adjacent landowners form a community-based land access program to expand hunter access to land.
- Expand habitat management and conservation programs on public and private land to benefit big game and other wildlife species.
- Continue to raise public awareness of and appreciation for the benefits of sustainable harvest of wildlife for food as a renewable natural resource that is good for people and the environment.

Time line of important dates in Vermont Wildlife Management	
1609	Samuel de Champlain is the first European to see Lake Champlain.
1640-1760	French Canadians slowly begin early European colonization in Champlain Valley.
1761-1791	English colonization of Vermont rapidly expands.
1791	Vermont becomes the 14th state, there are 85,425 people living in the state (1790 Census), landscape is 80% forested. State Constitution gives inhabitants the right to hunt, fish and trap.
1800	US Census reports there are 154,465 people living in Vermont.
1831	First bounties for bear are enacted by State Legislature.
1865	US Civil War ends, over 300,000 people live in Vermont, state is 37% forested, it is illegal to hunt deer.
1887	The last known native catamount is killed in Barnard.
1897	The first modern deer season is held, 103 deer are harvested in a 30-day season.
1904	The forebear of the Vermont Fish & Wildlife Department is created.
1941	Bear bounties repealed.
1946	"The Time is Now" written by Department biologists in response to rapidly growing deer herd.
1950	In 100 years Vermont population has grown very little, 377,747 people reside in the state.
1951	First regulation of bear harvest.
1953	First archery season is held.
1962	First sections of Interstate Highway System completed (Montpelier to Burlington).
1963	Limited antlerless permits issued.
1969	First wild turkeys (17) are reintroduced in Pawlet.
1970	Statewide land-use and development law (Act 250) passed.
1973	First wild turkey hunt held in parts of Addison and Rutland Counties. 579 permitted hunters harvest 23 birds.
1979	Modern deer management era begins, antlerless permits are issued by Wildlife Management Units (WMUs).
1986	First muzzleloader deer season is held in December.
1990	First Deer Management Plan is written.
1993	First regulated moose hunt in Vermont history held in October, 30 permits are issued.
2003	First Youth Day deer season is held the Saturday before regular deer season.
2005	Legal buck definition changes after 108 years from 1, 3-inch antler to 2 points on one side.
2008	Vermont is 75% forested, 625,000 people live in the state, ten-year planning cycle begins for deer, moose, bear, and wild turkey populations.

WHITE-TAILED DEER

I. Management History

Catastrophic conditions in both the deer population and habitat had already developed by the time Vermont's modern-day management program had begun in 1963. Buck-only deer hunting, which had been the tradition since 1897, allowed the deer population to grow rapidly and reach the biological carrying capacity (Seamans 1946, Garland 1978, Miller and Wentworth 2000). In Windham, Windsor and Rutland counties, the deer herd reached an overabundant and unhealthy state during the 1940s. The sporadic and small antlerless harvests between 1963 and 1970 removed less than 5% of the total deer herd (estimated at 250,000 deer). This proved to be insufficient to curtail growth and prevent the herd's impending collapse. As had occurred in other deer populations in other parts of the country earlier in the century, the consecutive harsh winters of 1969 and 1970 severely affected the health and abundance of Vermont's deer herd, which was already compromised by years of chronic overpopulation. In poor physical condition and without a sufficient nutrition base, Vermont's deer population would continue to fluctuate in response to winter conditions throughout the 1970s. Although reduced to only half of its former size through the 1970s, the deer population of about 120,000 animals lacked the vigor and supporting habitats to rebound. Allowing the habitat to recover by holding the deer population at a low or moderate level was the only realistic solution



WAYNE LAROCHE

to the chronic infirmity within the population created by the long-term over use of deer habitats.

In 1979, the Department began an ambitious deer population recovery effort. This effort occurred in three phases. During the first phase, the deer population was intentionally reduced to a level even below what remained after the winter mortality of the late 1970s. The second phase through the mid-1980s maintained the population at a relatively stable, low-density level to allow habitats to recover their ability to support a larger deer population. The third phase allowed for a gradual increase in the population to sustain annual deer harvests of 15,000 to 20,000 animals, while monitoring measures of herd health. By and large, this plan was successful. The habitats recovered and measurements of deer health such as antler beam diameter, weight, and reproductive rate improved (Table 2.1, Fig. 2.1).

Table 2.1 Reproductive rates of incidentally-killed adult (at least one year-old) female deer examined during winters in Vermont.

Year	# Doe	# Pregnant	Percent Pregnant	# Live Fetuses	# Fetuses per Doe
1963	99	82	83%	121	1.22
1966	115	97	84%	122	1.06
1972	139	121	87%	188	1.35
2001	121	115	95%	199	1.64
2004	78	72	92%	110	1.41
2008	119	108	91%	172	1.45
1963-72*	353	300	85%	431	1.22
2001-08	318	295	93%	481	1.51

*From Garland (1978)

Improvements did not come without a cost, however. The decade of the 1980s saw some of the smallest buck harvests since 1946. Legislation was passed in 1990 that prevented antlerless deer seasons from occurring during the November rifle season. Given this new constraint, the Department set out to make adjustments as to how deer management would be conducted in the years to come. Because the adult females drive the reproductive potential in a deer herd, effective population management means managing the number of does.

Responding to this challenge, the Department moved to involve the public more deeply in deer management decisions than ever before. Based on buck harvest objectives derived from averages of the 1970s and the results of a general public survey, a draft management plan was presented to interested citizens. The plan contained information about historic buck harvests, deer health statistics, and population trends through time on a WMU basis. The first deer management plan concluded with a selection of harvest objectives (within parameters set by the Department) that considered views of the public. With these objectives in hand, the Department proceeded to make annual antlerless harvest recommendations based on regional harvests. With antlerless deer numbers now being controlled using archery, muzzleloading, and youth hunters, the question remains: will the existing season structure and harvest limits be sufficient to take enough does in the future to prevent excessive population growth during mild winters? The answer may be most of the time, as long as multiple consecutive mild winters do not occur. Some of the time, however, it may be necessary to take additional measures to ensure that the deer herd does not become overly abundant.




Figure 2.1 Statewide mean-average yearling buck weights (dressed) 1948–2008. In 2008, biological check stations were held during youth weekend, so biologists could again get weight data that were representative of the population, because spike-antlered deer can still be taken during youth weekend.

During the 1990–1995 planning period, buck harvests increased significantly. Light antlerless harvests and mild winters during three of the five years of this planning period were largely responsible for this rapid response. Buck harvests met, or consistently exceeded, the harvest objectives in 15 of the 24 WMUs during this time. On a statewide basis, the statewide harvest objective was exceeded twice, and twice was within 1% of the objective. The overall size of the deer population increased as indicated by the 45% increase in buck harvest. Although the 1996 deer population estimate was between 120,000 and 140,000 animals, the health indices of antler beam diameter and body weight did not decline. Yet, continued growth at the pace experienced during the 1990s would have put the future of the deer herd and its habitats at risk. Indeed, a modest decline in yearling buck weights in the late 1990s (Fig. 2.1) preceded another herd decline resulting from severe winters in 2001 and 2003. The difference this time was that habitat had improved through the 1980s, and the herd was in better overall physical condition to rebound rapidly during this current decade (see data on following page).

1997-2006 Plan Accomplishments

In addition to seeking a balance between human demand for deer and the environmental consequences of too many deer, the Deer Management Plan for the State of Vermont 1997-2006 had five specific tasks to address.

► Task 1. Protect Deer Yards

❖ **Action:** Given the importance of deer wintering areas (DWAs) to the state's deer herd, the Department has vigorously defended against the loss of wintering habitat to human development. This is done through Vermont's land-use and development law known as Act 250, which requires an evaluation of a project's impacts on wildlife habitat. As a result, the Department worked with developers to modify development plans to lessen the impact to wintering deer. During this plan period, Department biologists reviewed 971 impacting deer wintering area projects, totaling 25,542 acres, and of these 91% (23,338 acres) were protected as part of the Act 250 regulatory process (Table 2.2).

Because only a small percentage of land development in the state actually requires an Act 250 permit, the majority of development is regulated at the local level or not at all. When

consulted, the Department worked closely with town and regional planning bodies to assure DWA protections were incorporated at the local level.

► Task 2. Population/Buck Harvest Objectives/ Adequate Antlerless Harvests

❖ **Action:** Balancing the demands of the people for more deer with the demands of the forest for fewer deer is the continual dilemma every northeastern fish and wildlife management agency faces. During the 1997-2006 planning period, the Department established an annual total buck harvest objective of 11,650. It was estimated that a buck harvest of this size would be generated by a deer population density of 18–20 deer per square mile. Assuming the buck harvest to be directly related to the overall deer population, it would serve as a good indicator of when the deer population increased or decreased. With this goal each year the Department would recommend an antlerless permit allocation distribution by WMU to adjust for population growth or loss resulting primarily from the previous year's harvest and winter severity index (WSI). Table 2.3 illustrates the relationship between the change in buck harvest (and assumed change in deer population) and

the corresponding change in antlerless permit numbers.

► Task 3. Antlerless Permit Application Process

❖ **Action:** A prominent concern expressed by hunters prior to the 1997-2006 plan was the ability of an individual to make multiple applications for an antlerless permit thereby increasing his or her odds of being drawn. Recognizing that this issue of fairness was very important to a majority of hunters, the Department recommended to the Vermont Fish and Wildlife Board a regulation change that limited an individual to one antlerless permit application. This change became effective for the hunting seasons of 1997.

► Task 4. Promoting Hunting Culture

❖ **Action:** Although a free youth hunting license had been available since 1993, Vermont's downward trend in sales continued to follow the national decline. Growing concern for the decrease in the number of hunter served as an impetus to advance a youth hunting opportunity (Fig. 2.2). With the support of the deer hunting community, the Vermont Legislature passed a measure designating the Saturday before the regular deer season as Youth Hunting Day. The first Youth Hunting Day occurred in 1997. Seeking to expand the opportunity for youth, especially considering all of the alternative activities available to them on a Saturday, the Legislature expanded the Youth Season to include Sunday as well. The first youth weekend was held in 2003. Early enthusiasm for youth hunting reached its peak in the year 2000. It was followed by a period of decline mirrored by adult participation. This suggests that factors beyond

Table 2.2 Summary of Act 250 DWA acres with Department involvement (1997-2006).

Year	# Projects involving DWA	Total Wintering Area Acres	Acres Impacted	Acres Conserved or Protected	Pct. Acres Protected per Year
1997	89	3,087	266	2,821	91%
1998	115	3,132	348	2,784	88%
1999	114	3,281	281	3,000	91%
2000	107	2,154	198	1,956	91%
2001	78	1825	205	1620	89
2002	116	3,484	180	3,304	95%
2003	132	2,888	222	2,666	92%
2004	94	2,169	270	1,899	88%
2005	92	2,125	265	1,860	88%
2006	112	3,222	174	3,048	94%

1997-2006 Plan Accomplishments (continued)

Table 2.3 Buck harvest, antlerless harvest, and WSI relationship for the period 1997-2006.

	1997	1998	1999	2000	2001	2002	2003	2004	2005*	2006*
Buck Harvest	12,596	12,641	11,907	12,610	9,409	11,023	9,194	7,648	4,956	7,805
Antlerless Harvest	7,240	7,427	7,876	7,888	5,602	5,609	5,334	4,277	3,590	4,877
Winter Severity	37.3	29.9	35.6	34.0	73.3	23.6	83.9	62.2	44.7	15.2

*New antler point and bag limit regulations in effect

a special hunting season, perhaps the same influences from suburbanization that adult hunters are faced with, are affecting young hunter recruitment and retention (Fig. 2.3). In 2009, the Vermont General Assembly removed the Vermont residency requirement. This now allows any eligible youth to participate in the Youth Hunting Weekend as long as he or she has obtained landowner permission and is accompanied by an adult with a valid Vermont hunting license.

The Department also initiated several other successful programs and activities that encourage hunting, outdoor activities, and appreciation of our forests and wildlife. The "Outdoors Woman" and "Outdoor Family" programs were aimed at educating and exposing women and families to outdoor sports and skills. The Department partnered with the Vermont Outdoor Guides Association to sponsor a yearly "Doe Camp" to introduce women to outdoor hunting skills. A two day retreat, "The Future of Hunting in Vermont" at Castleton State College in 2006, brought together over 80 people from many youth organizations, sporting groups, and academic and government institutions to discuss challenges and solutions associated with barriers to hunting and recruiting new hunters. And finally, the Department created a pilot project called "Working for Wildlife" that establishes partnerships with sporting clubs to work on projects associated with habitat improvement, landowner relations, and conservation education.

► Task 5. Quality Deer Management (QDM)

❖ **Action:** Vermont deer hunters with an interest in hunting older-aged bucks (3.5 years or more) with well-developed antlers approached the Department during the development of the 1997-2006 Deer Management Plan. According to the definition used in the management plan, Quality Deer Management (QDM) is described as a management




Figure 2.2 Resident hunting license sales, 1987–1996.




Figure 2.3 Resident hunting license sales during the 1997-2006 deer management plan period.

technique used to shift the age structure of the buck population from one dominated by young (1.5 years old) males to a population with a higher proportion of older-aged (at least 3.5 years old) bucks.

To further assess these components and develop an objective approach for designing a QDM program that was intended to balance deer population and habitat and increase the numbers of older bucks, the Department assembled a nine-member panel of deer hunters in January of 1998. Following their

1997-2006 Plan Accomplishments (continued)

seven months of research and deliberations, the QDM advisory panel identified 14 elements to be included in the QDM program (Table 2.4) and four possible alternatives (Table 2.5).

The panel recognized the implementation challenges and the significant amount of effort required to make institutional changes to a long-standing traditional deer season. To avoid making premature decisions about deer season changes, the panel recommended that strong, broad-based support of hunters and landowners be present before implementing any changes in season format. The panel also recommended that QDM be implemented at the WMU level and not vary in design from one unit to another.

Upon being presented with these recommendations, the Department decided first to assess hunter satisfaction. Results of the 1998 survey indicated most respondents (63%) indicated they were "Very Satisfied" or "Satisfied" with their deer hunting experience in the last five years (Table 2.6). Hunters preferred to retain the then current deer hunting season format (one 3-inch antler minimum) over any QDM restrictions that might be implemented by a 57% to 41% margin, with 2% reported as "Undecided."

Following a review of what the QDM panel produced and the hunter opinion survey, it was decided to table further consideration of any changes to the season format. However, following poor hunting seasons in 2001, 2003, and 2004 related to the severe winters of 2001 and 2003, hunter satisfaction decreased significantly. Another hunter satisfaction survey was

completed in 2003 to assess interest in "QDM," or what was then being labeled as "Comprehensive Deer Management" (CDM).

Results of the 2003 survey indicated that, in general, since 1998 more hunters were still satisfied with their deer hunting in Vermont than those who were not (42% "Satisfied" vs. 31% "Dissatisfied"). However, when compared to the 1998 survey where 63% were "Satisfied" vs. 20% "Dissatisfied", there clearly had been a shift towards greater dissatisfaction. When asked of their support for greater antler restrictions to protect more young bucks, 66% supported and 24% opposed this idea with 10% reporting "neutral" (VFWD 2004).

With the results of the survey showing hunters' support for increasing the proportion of bucks afield, the Department renewed its effort to meet this goal.

A series of public hearings were held, and the Fish and Wildlife Board was given authority by the Legislature to set deer hunting regulations, with the exception of the November rifle season, as they do for all other fish and wildlife species.

An antler-point restriction regulation to promote CDM was put into effect by the Board beginning with the 2005 hunting seasons. The new hunting regulation also reduced the

Table 2.4 Elements of QDM , by relative importance, identified by the QDM advisory panel.

1. Hunter participation
2. Ecological integrity
3. Recruitment of young hunters
4. Ease of implementation
5. Endorsement of residents in WMU
6. Quality of hunting experience
7. Balanced Buck:Doe ratio
8. Balanced age structure
9. Maximum sustainable yield
10. Acceptance by the nonhunting population
11. Increased body weight of individual deer
12. Increased antler size on bucks
13. Equal hunting opportunity for all hunters
14. Genetic improvement of the deer herd

Table 2.5 Methods for QDM implementation recommended by the QDM Advisory Panel.

1. Retain the current season structure. The present hunting seasons include the elements of QDM and can be defined by the individual hunter.
2. Restrict the buck harvest by changing the definition of a 'legal buck' from a deer with at least one, 3-inch antler, to a deer with at least 3 antler points.
3. Restrict the annual bag limit from 3 deer of either sex (with appropriate permits) to 3 deer with no more than 1 buck per year.
4. Restrict the annual bag to one deer per year and include antlerless deer, by permit, as part of the 1 deer bag.

Table 2.6 1998 and 2003 survey results for the distribution frequency (%) of hunting satisfaction.

	1998	2003
Very Satisfied	17.3	5.8
Satisfied	46.1	35.7
Neutral	16.9	26.8
Dissatisfied	13.8	23.8
Very Dissatisfied	5.9	7.9

annual bag limit from three deer to two and redefined a legal buck to a deer having at least two points on one side. The points were defined as the terminal point and one other point measuring at least one inch from the main beam.

II. 2010-2020 White-tailed Deer Management Issues, Goals, and Strategies

Many Vermonters would like to have more deer than is advisable under the new deer density objectives, and many others would like to have fewer. The rationale for the deer density objectives are provided in the supporting text that follows. Ultimately, the proper balance maintains ample harvests of deer as well as forest products over the long term. It is apparent that localized deer management issues are mounting in Vermont and methods are needed to support more localized deer management to relieve foresters, gardeners, and farmers from locally overabundant deer populations. The overall goal of deer management in Vermont is to manage Vermont's deer herd to sustain viable populations consistent with biological, social, and economic considerations.

ISSUE 1. Habitat Loss and Assessment

GOAL: To monitor changes in habitat quality and quantity and perform public outreach regarding habitat management techniques, so concerned citizens may help to secure their deer herd's future.

White-tailed deer populations vary widely through time and space in response to varying habitat and landscape conditions as well as weather, hunting intensity, predators, and disease. Changes in any of these factors complicate the ability to track deer populations, but the factors most important in determining population size are habitat conditions and winter severity. Hunting, as a form of predation, can be used as a tool to control the deer population in Vermont as long as enough does can be taken.

Optimum deer habitat has been described as a mosaic of fields and forests (Halls 1984). In areas with high quality habitats, deer can live in an area as small as one square mile. Within this area, the diversity and arrangement of plant species provide a setting for deer to feed, bear young, and find shelter and concealment. The greatest concentrations of deer in Vermont are found in agricultural areas of the state (having the highest carrying capacity for deer) with a mix of field and forest. Reduced numbers of deer occur in remote aging forestlands, especially in large

blocks of forests at high elevations where diversity and quality of food plants are reduced and extreme snow depths frequently occur. For these reasons, Vermont's lower elevation areas tend to have higher densities of deer. The differences in both the habitat quality and the density of deer in different areas of the state are the reason and basis for the state being divided into wildlife management units.

Deer wintering areas, or "DWAs," are habitats that provide shelter for deer in periods of extreme cold and deep snows. These areas are usually comprised of stands of softwood tree species, such as hemlock, spruce, fir, cedar, and pine, and they range in size from less than 100 to more than several thousand acres. Within these critical areas, combinations of vegetative and topographic factors create micro-climates that favor survival of deer through the harshest season of the year. These areas are essential to the survival of our deer during severe winters. Wintering areas do not usually change significantly from year to year and may be used by many generations of deer over many decades if appropriate habitat conditions are maintained. Deer exhibit a great deal of fidelity to individual wintering areas. When cover is removed, deer don't always move to another area and are more likely to succumb to harsh weather.

Department wildlife biologists first identified and mapped Vermont's deer wintering areas during the 1960s and updated the maps in the mid-1980s. Since that time, Vermont has lost some of this important habitat to residential development and even more has been affected by winter recreational trails and logging. The Fish & Wildlife Department biologists endeavor to protect and enhance deer wintering areas through negotiations with land developers during the Act 250 land use regulatory process by working with municipal and regional planners to recognize these areas as being sensitive habitats and by coordinating with landowners, foresters, and loggers to maintain and improve conditions within these essential wintering habitats. The Department uses strict guidelines for logging and maintaining DWAs on state-owned Wildlife Management Areas and has recently updated the "A Landowner's Guide, Wildlife Habitat Management for Vermont Woodlands" (VFWD 2009) designed to provide guidance for interested landowners.

In addition to being concerned with the habitat losses caused by people, the Department is also closely monitoring the spread of invasive plant and insect

species that could affect deer habitats. One species that has potential to alter large amounts of deer wintering habitat is the hemlock woolly adelgid. This insect kills eastern hemlock. If this insect becomes established in Vermont, it could have far reaching effects on the state's hemlock-dominated forests and DWAs. Hemlock trees provide superior cover for wintering deer. Department biologists are closely monitoring the occurrence of this harmful insect with help from state foresters. There is some concern that warmer winters and extended growing seasons may allow the movement and colonization of this tree pest northwards up the Connecticut River valley.

Management Strategies

- 1.1 Update inventory of deer wintering areas for local, regional, and state habitat planning and protection efforts.
- 1.2 Stress the importance of habitat conservation with outreach efforts to various segments of the public such as farmers, educators, hunters, forest managers, and land planners.
- 1.3 Work closely with foresters and entomologists to prevent, manage, and eliminate the threat of the hemlock woolly adelgid.

ISSUE 2. Population Goals

GOALS: 1) Maintain deer densities using regional population objectives.
2) Monitor biological characteristics of habitat and deer that can change in response to deer herd size through time.
3) Adjust antlerless deer harvests to alter population levels as necessary to achieve population objectives.

DEER DENSITY

Vermont statutory law states that “an abundant, healthy deer herd is a primary goal of fish and wildlife management” (Title 10 V.S.A. §4081(c)). This is the foremost charge of deer population management in Vermont. The deer herd is kept healthy by preventing overabundance with carefully planned antlerless deer harvests.

The population density of a deer herd affects the general health of the animals, the sustainability of its habitat, and the probability of human and animal

conflict. The following discusses the factors that the Department considers when setting management objectives: the sex ratio between bucks and does and biological and cultural carrying capacities. It also discusses how the Department gathers data that is used to determine deer harvests.

Sex Ratio

Adult white-tailed deer females typically produce twin fawns if summer and autumn nutrition are adequate (Ozoga and Verme 1982, DelGiudice, et al. 2007). If successful, the Department's management strategy should maximize the reproductive potential of does. Sex ratios that are highly skewed in favor of does can result in does remaining barren through the first estrous thus delaying pregnancy for the entire year because there are too few bucks to tend all does (Mysterud et al. 2002). The gregarious nature of female deer and courting nature of breeding bucks typically allow a sex ratio of one buck to three does to be sufficient to breed all does in a population (Table 2.1; Demarais et al. 2000). Populations that are heavily hunted require more does than bucks in order

Table 2.7 Number of road-killed adult (at least 1 year-old) male and female deer registered by game wardens in Vermont

Year	# Males	# Females	# Females per Male	# Males per 100 Females
1971	274	1,057	3.86	25.9
1972	414	1,394	3.37	29.7
1973	419	1,252	2.99	33.5
1974	381	1,095	2.87	34.8
1975	361	1,208	3.35	29.9
1976	318	1,091	3.43	29.1
2000	434	1,244	2.87	34.9
2001	325	1,225	3.77	26.5
2002	257	974	3.79	26.4
2003	299	1,010	3.38	29.6
2004	255	889	3.49	28.7
2005	299	953	3.19	31.4
2006	357	1,012	2.83	35.3
2007	459	1,149	2.50	39.9
2008	471	1,239	2.63	38.01
1971-76*	2,167	7,097	3.28	30.5
2000-05	1,869	6,295	3.37	29.7
2006-08	1,287	3,400	2.64	37.9

*From Garland (1978) describing a period of buck-only hunting.

to produce the excess of offspring needed to sustain harvests.

Many hunters in Vermont believe that there are too few bucks to completely breed all does. Statewide data from deer road-kills has consistently demonstrated that a sex ratio of a little over three does per buck exists in Vermont (Table 2.7). Sex ratios can also be estimated from survival estimates determined from age data. White-tailed does commonly live many productive years in Vermont (Fig. 2.4) while bucks typically live only a few years (Fig. 2.5). In general, does have about 75–85% annual survival while annual survival of bucks is about 25–40%. Model results confirm the road-kill estimates that before the antler restriction (AR) in 2005, statewide prehunt sex ratios were about 3.25 does per buck.

Increased yearling survival following the AR has changed the buck to doe ratio. Prehunt sex ratios are now estimated to be about 2.75 does per buck state-wide. With more than 50% of legal bucks harvested annually, it is expected that the number of does per buck increases post-harvest. Localized differences are expected to always exist around the state.

Biological and Cultural Carrying Capacities

In determining the optimal size of the deer herd, biologists consider the concept of carrying capacity — biological and cultural. The term biological carrying capacity (BCC) refers to the maximum number of animals that an environment can support without detrimental effects. The quality and quantity of available habitat determines the BCC. The cultural carrying capacity (CCC) is more subjective. It is determined by assessing the values people place on wildlife versus the liabilities created by overabundant wildlife populations. While BCC has only an upper limit, CCC has both an upper and lower limit because most people desire that there not be too few, but not too many deer to cause damage. Hunters and the general public want enough deer to satisfy their

hunting and viewing desire while too many cannot be ecologically sustained and are considered to be a nuisance to humans.

Biological Carrying Capacity and Maximum Sustainable Yield

When deer herds approach or exceed an area's biological carrying capacity, the animals' health is affected. Wildlife managers have determined that deer herds managed at densities below BCC are healthier and in balance with their habitat. This concept of maximum sustainable yield (MSY) is the point




Figure 2.4 Laboratory-determined ages of 427 female white-tailed deer from the 2003–2006 annual Vermont harvests.




Figure 2.5 Ages of 10,300 male white-tailed deer from 1995–2004 as determined by Vermont biologists at check stations during the opening weekend of rifle season.

within the biological carrying capacity curve when the density of a herd is in balance with its habitat and when fawn recruitment is at its maximum level. A population at BCC recruits as many fawns as it loses adults, so it has no harvestable surplus. At MSY there are fewer deer overall in a population. Does produce more fawns (Table 2.1, pg 10), and fawns have a much better neonatal and overwinter survival.

Deer and their habitats are unhealthy when at a BCC level, but healthy and productive near MSY. This principle is particularly important in northern environments where body condition of deer going into winter can be critical to over-winter survival and where the existence of too many deer can do extensive damage to wintering habitats. Managing near MSY, rather than BCC, helps minimize the boom and bust cycle of the deer herd in Vermont and can be expected to sustain greater deer harvests in the long term (Fig. 2.6). A healthy deer herd with healthy habitats can recover from bad winters or over harvesting much faster than a deer herd and habitat in poor condition.

Cultural Carrying Capacity

Owing to their beauty and athleticism, deer populations are often too low to meet the general public's desire to view these animals. From a cultural perspective, when deer populations become too large, conflicts such as damage to landscape flowers and shrubs, agricultural and forestry losses, deer-vehicle collisions, and transmission of human pathogens,

such as Lyme disease, can occur. In these cases a deer population may be below its biological carrying capacity (BCC), posing little threat to the long-term sustainability of their habitats, but at the same time above its cultural carrying capacity (CCC) if property losses or disease prevalence are deemed too high. Deer populations can also be below CCC when hunters and other outdoor enthusiasts feel that they see too few deer.




Figure 2.6 Deer population size and growth rate at biological carrying capacity (BCC) versus maximum sustainable yield (MSY)

To find the proper balance between the highs and lows of CCC, the Department conducted a public opinion survey in 2007 to assess the people's deer abundance preferences (Fig. 2.7). The assessment was analyzed at many different levels including, where one lived, one's gender, and whether or not one hunted. The results of the survey suggest that nearly half of all Vermonters are generally satisfied with the number

of deer in their county. Thirty-two percent of the respondents felt the deer population should be increased with only 5% of respondents feeling the deer population should be decreased. Fifteen percent either "did not know" or had "no opinion."

When the response to deer population change is analyzed by subgroups, similar interests were found. Of those who said they had hunted in the past five years, 66% felt the deer population should be increased and 27% felt it should remain the same. Of those who did not hunt during the last five years, 22% felt the deer population should be increased and 54% felt it should remain the same. Greater Chittenden County residents were more




Figure 2.7 Vermont public's opinion regarding deer population change over the next ten years

likely (58%) to want deer populations to remain the same than their more rural counterparts. This suggests that the deer population may be approaching CCC in Vermont's most populous county. On the other hand, more people (48%) in the Northeast Kingdom (Orleans, Caledonia, and Essex counties) want more deer. This suggests that deer numbers are not near the CCC in that region of the state.

When asked about property damage from wildlife, 14% of the respondents indicated they had suffered a loss to their automobile and 21% had incurred loss of landscape, ornamental or vegetable garden. But when these respondents were compared with those who had not incurred any damage of any kind, responses were remarkably similar for both groups when asked about their opinions of deer population size. Forty-six percent of the respondents incurring damage felt the deer population should remain the same while 48% of respondents that had not incurred any damage felt the same way. These data suggest that, in general, the upper CCC limit, with localized exceptions, is not currently an important issue to the Vermont public.

Responsible deer management dictates that a deer herd's relation to BCC be considered before CCC is considered. In this circumstance, biological measures (for example, birth rates, antler development) inform the Department about the deer population goals. In most of Vermont, deer population goals, as measured by deer per square mile, can currently be achieved through traditional regulated hunting seasons. In cases where deer numbers are below BCC, but CCC demands fewer deer, traditional hunting seasons may not always be effective in satisfying CCC. In some cases, population goals may need to be described in terms other than deer per square mile, for example: motor vehicle collisions, Lyme disease rates, or number of crop damage complaints. These measures may need to be used to set population goals in some local areas if Vermont's deer and human populations continue to grow. Special methods to reduce deer numbers, such as those described in the "Locally Overabundant Deer Populations" section, could be required in the future.

Cultural carrying capacities will likely become increasingly important in the future as a consideration in setting deer density objectives in parts of Vermont where the human population density is growing fastest. In Connecticut where high deer densities (greater than 50 per square mile) are associated with high incidence rates of Lyme disease, CCC may require long-term deer density

objectives to be set as low as 10 deer per square mile (Kilpatrick and LaBonte 2007). This is a situation that may be preventable in Vermont if we are able to maintain densities at or below 20 deer per square mile in regions such as Bennington County that are prone to Lyme disease (see Vermont Health Department statistics for Lyme disease cases in Vermont). Reduction in deer densities may reduce the abundance of Lyme disease-carrying ticks (*Ixodes scapularis*). Very few ticks were found in Maine where deer densities were lower than 18 per square mile (Wilson et al. 1990, Rand et al. 2003, 2004). On the other hand, total elimination of deer can lead ticks to feed more intensively on rodents and result in higher densities of disease-positive nymph-stage ticks (Perkins et al. 2006). Once again, finding the proper balance between too many and too few deer seems to be the best way to ensure that a healthy ecosystem exists with a minimum of human conflicts.

Body Condition and Deer Densities

The number of deer per square mile that Vermont's landscapes can support is a value that shifts across the landscape and through time as habitat quantity and quality change. Often deer themselves are a main cause of this change as they degrade habitat when they become too numerous. Thus, biologists usually rely on biological measures of the deer themselves, such as reproductive rates, weights, and yearling antler beam diameters, to gauge the relationship between the deer herd and their habitat.

Population objectives going forward should be based not only on deer harvest numbers but also on the body condition of deer. Many states and deer management systems monitor deer herd characteristics, such as reproductive rates, yearling antler beam diameter, and fawn weights to track population health (Miller and Wentworth 2000, Williamson 2003) (Fig. 2.8). These data can be used to measure the impacts of and changes in deer populations that follow severe winters (Fig. 2.9). Although tracking changes in the body condition of deer provides a way of recognizing times when there is a need to harvest more deer, it is often after damage to habitat has already occurred. Changes in body condition of deer do not provide a means to determine how many deer should actually be harvested (Fig. 2.10).

In the long-run, if deer harvests are tailored to ensure that deer body condition remains good, deer will weigh more and winterkill will not be as great during severe winters. Deer in good condition will

also produce at an optimal recruitment rate that is just above intermediate levels of abundance relative to BCC (Miller and Wentworth 2000; Fig. 2.6). This management strategy will dampen the boom and bust cycle of deer in Vermont.

It appears that a sustainable harvest of deer having good body condition may be approximately the harvest level that occurred in the mid-1990s, and again in 2008. This is a total deer harvest of approximately 17,000 deer per year (Fig. 2.11). Vermont has never sustained annual harvests of 20,000 deer for very long. Harvest of 20,000 deer per year in Vermont, given current hunting pressure and deer reproductive potential, is probably indicative of an overabundant deer herd. Buck harvests frequently exceeded deer management objectives in the late 1990s following a series of mild winters. Lessons from the 1990s and scientific studies suggest that perhaps 20% of does may need to be harvested during times of mild winter in order to stabilize herd growth when winters fail to do so (Dusek et al. 1989, Giles and Findlay 2004). Historically, less than 10% of adult does have been harvested annually in Vermont. Regional estimates are made using the same method. Adding up these regional estimates results in a total deer population estimate that is very similar to the estimate calculated above for the whole state (Table 2.9 pg 26, Fig. 2.12).

Habitat and Deer Densities

White-tailed deer play a significant role in the ecology of Vermont's forests. As herbivores (plant eaters), they disperse seeds and as prey, they allow other species to survive. The influence of deer in our forests is considered so significant that researchers and wildlife managers regard them as a "keystone" species in the Northeast. Deer browsing has profound implications for the structure and function of forested ecosystems. If deer were removed from the system, a wide variety of changes would ripple through the forest. However, overabundant deer populations can also be a negative force within the forest ecosystem.




Figure 2.8 Annual average fawn weights (with 95% confidence limits) as reported by hunters to check stations from 1997–2008. All years exclude fawns reported over 99 pounds. The decadal trend-line minimizes the distance between the annual points and the line itself. With bio-check stations now during youth weekend, the Department will investigate the use of fawn weights as a more sensitive indicator of herd health, similar to the use of yearling buck weights.




Figure 2.9 Statewide winter severity indices (WSI) in Vermont from 1980–2008. The horizontal dashed line equals a long-term average of about WSI=50. From 1 December through 15 April, one point per day is given when snow depth is at least 18 inches, and a point is given when temperatures drop below 0°F. The Department maintains 38 volunteer weather stations statewide.

Deer densities vary throughout North America as well as within Vermont and are largely in response to habitat and weather conditions that affect reproductive and survival rates and food availability (Halls 1984, Crête 1999). Young forests provide better habitat for white-tailed deer than old-growth forests. A mix of field and forest is more favorable




Figure 2.10 Pre-hunt total deer population estimates (+/- 15%) for Vermont from 2000–2007. Rapid population growth from 2005–2007 demonstrates tremendous growth potential of Vermont’s healthy deer herd given a mild winter as in 2006 and restricted antlerless deer harvests.




Figure 2.11 Annual total deer and antlered buck harvests in Vermont from 1995–2008. Boneless meat is represented as 100s of pounds, so the ten-thousand-line equals one-million pounds of meat. Pounds of boneless meat assume that hanging weight (skin, head, and feet removed) is 75% of field-dressed weight and edible meat is 75% of hanging weight.




Figure 2.12 Regional pre-hunt deer population density estimates for Vermont in 2007 (see Table 2.9) in relation to statewide upper and lower population density objectives (i.e., horizontal dashed lines near 13 and 18 deer per square mile).

than large unbroken forest tracts. For instance, deer at similar density will have less impact on forest vegetation and habitat condition in areas having some agricultural croplands compared to areas that are entirely forested (Horsley et al. 2003). Areas having greater and more prolonged snow loads during winter can be expected to have greater winter-kill than in areas having less snow. For these and other reasons, sustainable deer densities vary throughout North America and within Vermont. Thus, it is sensible to manage deer to achieve various deer density objectives regionally throughout Vermont in accordance with climate and habitat conditions that are influenced by soil type, topography, weather, and human land-use practices.

Optimal deer density varies across the landscape and through time. Studies from northeastern North America have found that general patterns associated with deer density, however, do exist. Since the mid-1900s, deer density in much of the eastern United States, including southern Vermont, has been high enough to negatively impact forest vegetation. Long-term deer densities exceeding 20 per square mile are capable of altering forest plant communities, threatening endangered plant species, reducing ground-level hiding cover and forage for other wildlife species, and reducing abundance of nesting birds (McShea and Rappole 2000, McGraw and Furedi 2005, Côté et al. 2006). At densities greater than 20.5 deer per square mile, managed forest habitats in northwestern Pennsylvania were altered enough to exclude many songbird species (DeCalesta 1994).

Forest conditions, including deer forage availability, at any point in time are related to past as well as current land and forest management practices. Forest management practices affect the capacity of the forest to accommodate deer. Certain forestry practices may be used to encourage forest regeneration in locations where deer browsing is of concern. For example, one study recommended increasing the size of clear-cuts to larger than two acres as a way to provide for sufficient forest regeneration by producing more than the deer could eat (Akins and Michael 1995). Indeed, more research is needed on forest management practices that are effective in the presence of deer. Researchers have found that northern Pennsylvania hardwood forests were able to successfully regenerate with no shift in tree species composition at deer densities of 13–21 per square mile as long as suitable “deer forage” was at least moderately abundant

(Marquis et al. 1992). On the other hand, when deer food availability was high, successful forest regeneration occurred at deer densities as high as 21–31 deer per square mile (Marquis et al. 1992). Agricultural lands interspersed with forest lands enhance the availability of suitable forage for deer and can increase the density of deer that can be sustained without impacting forest regeneration. A deer density of 18 deer per square mile was suggested to ensure regeneration of desired tree species in the absence of agricultural influences (Tilghman 1989).

Following the end of Vermont’s state-wide deer reduction campaign of the 1980s, deer numbers increased through the 1990s and once again reached high densities in many parts of the state even with increased antlerless harvests. In some parts of the state, deer populations grew to levels that again began to impact forest regeneration. In southeastern Vermont, deer have consumed much of the palatable and merchantable hardwood regeneration of oak, maple, and ash. In addition, the region has experienced a proliferation of invasive species that are not palatable to deer such as buckthorn and barberry. As a result, both the invasive species and deer browsing on the more limited food supply have compounded the impacts on the native forest species. Similar effects, although not as dramatic, may be observed in other parts of the state (Fig. 2.12). For these reasons, the densities of deer that the habitat can support in southeastern Vermont may be more limited than in other parts of the state.

Deer density in any given area typically changes with the seasons. In northern climates, the onset of snow and colder temperatures force deer to vacate their larger summer and fall ranges and concentrate in higher densities in deer wintering areas. Quantity and quality of both winter and nonwinter deer habitat, as well as severity of weather conditions, determine the density of deer that any region can sustain through time. Good summer feeding conditions result in bigger and fatter deer that survive winter better. Good winter habitat minimizes thermal and other stressors that burn energy and result in mortality.

Because optimum deer density varies depending upon regional conditions, any determination of optimal deer density objectives for Vermont should be based upon data that considers both summer and winter habitat while accounting for regional differences in winter severity, winter habitat condition and availability, and the land use considerations of

landowners. Applying all of these factors in managing for a pre-determined prehunt summer and autumn deer density objective is a method that will best provide for optimal body condition as deer go into Vermont's unpredictable winters. This is the best way to minimize boom and bust deer density cycles.

WILDLIFE MANAGEMENT UNIT (WMU) REALIGNMENT

Antlerless deer harvests began being regulated by WMU in 1979 under a permitting system allocating permits to hunt in the 17 newly formed WMUs, which are defined in state statute. Seven of the WMUs were separated by the Legislature into two sub-units in 1983. Changes in deer populations and a reassessment of existing habitat conditions warrant refining the boundaries of select WMUs in order to facilitate more effective management of the deer population in the WMU. Revisions being considered are described below and illustrated in the map (Fig. 2.13).

- a. Adjust the boundaries of the WMUs in southeastern Vermont to more accurately reflect the difference between the Connecticut River Valley habitat and the habitat associated with the physiographic region. This would merge WMUs M1 and O1 to form the Eastern Foothill unit (new WMU M) and WMUs M2 and O2 to form the Connecticut River unit (new WMU O). WMU Q would have I-91 as an easterly boundary in the town of Guilford. East of I-91 would become part of WMU O.
- b. Extend the boundary of J2 northward to US Route 2 to remove an agricultural area from WMU E because habitat in agricultural areas is generally more productive than that found elsewhere in Essex County. Combine the remaining mountainous portion of H2 with H1 to form a new WMU H.
- c. Merge WMUs K1 and K2. The area of K1 is too small to yield harvest numbers large enough to be effectively used in scientific data analyses. These two WMUs closely resemble the habitat types of their respective neighboring WMUs and can be included into a new WMU K.
- d. Move a portion of the boundary between WMU D1 and D2 to the east to put more of the Lake Memphremagog agricultural lands into WMU D1, which is most similar in land use and habitat condition.

DATA GATHERING

In order to allocate permit numbers and direct other management actions at the WMU level, data sources such as hunter sighting rates, antlerless tag fill rates, and local observations are used to fine-tune management actions. For example, some property owners would like the Department to manage overabundant deer at the level of individual properties (for example, extra doe permits for landowners). This could be an option for dealing with localized problem areas having high deer densities. However, any system would need to be scientifically credible, practical, effective, and consistent with the overall deer management strategy. Should such a system be devised it should be based on data measured from vegetation, not by sightings of deer (Mitchell et al. 1997, Augustine and DeCalesta 2003). The Department has found that localized problems of deer overabundance can often be dealt with by getting landowners to provide access to their land and work with hunters to take antlerless deer during archery, muzzleloader, and youth seasons.

Vermont has recently begun using a "mark-recapture" method for deer population estimation. Coupled with new buck:doe ratio data collected from road-killed deer and fawn:doe ratios determined through bow hunter observations, deer biologists are improving their ability to estimate annual deer population composition and density at the state-wide and regional level. Because smaller amounts of data have less predictive power than larger amounts of data, it has now been determined that current data at the WMU level is not sufficient for these techniques to be used to make accurate population estimates at the WMU level. For this reason, WMUs having the most similar deer densities are being grouped into regional units for regional population estimation purposes (for example, Northeastern Highlands, Lake Champlain Valley).

The following provides an example of how the deer population numbers for the state of Vermont can be estimated. Analysis of deer age data (Figs. 2.4 and 2.5) determines that Vermont has a statewide prehunt buck:doe ratio of 1:2.75. Age data reveals that yearling bucks make up about 52% of the antlered buck harvest. Approximately 50% of all yearling bucks have spike antlers as determined by data collected by biologists at check stations prior to 2005 (26% of total buck population has spikes). Thus, a prehunt legal buck population in 2007 of 19,286 indicates a total buck population of about 26,062




Fig. 2.13 Current and proposed WMU boundaries

if spike-antlered yearlings are included (Table 2.8). Given an estimate of 2.75 does per buck, the adult doe population is estimated to be 71,670 does. In 2007, 4,484 adult does were harvested amounting to 6% of the adult doe population. Assuming 1.5 fawns are produced per does of at least 1 year-old (Table 2.1) and assuming a 55% fawn survival rate through early autumn (Ballard et al. 1999, Haskell et al. 2007), there would have been about 59,130 fawns in the deer herd prior to harvest in 2007. The summer fawn survival estimate is the most uncertain of the estimates used in this model. However, by combining these estimates, it is possible to estimate the total prehunt deer population for 2007 which adds up to about 157,000 ($\pm 20,000$ 90% CI; Fig. 2.10), or 20.5 deer per square mile of deer habitats.

The Department currently also uses the mark-recapture technique to estimate prerifle hunt legal buck population size (results in Tables 2.8 and 2.9). The Department's technique is essentially a removal model where probability of "recapture" is set to zero. The deer are in a sense "marked" when they are registered at the check station during the 16-day rifle season when the hunter reports the WMU and day that the deer was harvested. By combining this data with daily hunter effort estimates gathered from hunter surveys, the mark-recapture model can be used to estimate the daily probability that a deer will be harvested, and ultimately, the number of deer that remained after the annual harvest. Adding the number of deer harvested to number of deer

estimated as not harvested yields a prerifle-hunt population estimate of legal bucks. This application of the mark-recapture method may be uniquely applicable to Vermont for three reasons: 1) mandatory registration of all legally harvested deer ensures that a very complete accounting of actual harvest exists; 2) an adequate return rate of hunter effort surveys exists (demonstrating Vermont hunters' dedication to sound deer management); and 3) the harvest rate of bucks during the rifle season often exceeds 50% of the total buck population (Table 2.8). All three of these conditions must be met for this technique to produce valid results. At this time, Vermont may be the only state that meets all of these conditions.

It is the Department's goal to make deer management in Vermont as scientific and data-driven as possible, but this effort will at times be limited by staff and other resources. Professional judgment provided by Vermont's wildlife biologists will always be necessary to augment the hard science of wildlife management.

DEER DENSITY OBJECTIVES

Based upon the information gathered on the issues presented above, the Department intends to set prehunt deer density objectives for each of the regions in Vermont. These will serve as a baseline from which to work in the future (Table 2.10). In setting these density objectives, it is recognized that they must vary even within a region of the state. For example, the northeastern part of the state may sustain a total deer density of 13 deer per square mile. However,

Table 2.8 Population estimates ("N-hat") of legal bucks in Vermont before the rifle harvest and then corrected for bucks taken in earlier seasons to get pre-hunt estimates. Total harvest rates respond to population size and license sales, and post-hunt buck populations may be important to consider for wintering deer. In all years, post-hunt numbers presented assume no sources of mortality during hunting seasons other than registered harvests.

Year	Pre-rifle Legal Buck N-hat	Rifle harvest	Rifle harvest rate	Early youth and archery bucks	Pre-hunt Legal Buck N-hat	Total buck harvest	Total harvest rate	Post-hunt N-hat	Post-hunt N-hat with spikes
2000	22,428	10,256	0.46	1,816	24,244	13,120	0.54	11,124	11,124
2001	16,102	7,588	0.47	1,123	17,225	9,522	0.55	7,703	7,703
2002	11,619	8,720	0.75	1,428	13,047	10,956	0.84	2,091	2,091
2003	9,575	6,868	0.72	1,623	11,198	9,196	0.82	2,002	2,002
2004	12,283	5,594	0.46	1,420	13,703	7,654	0.56	6,049	6,049
2005*	8,263	3,957	0.48	728	8,991	5,002	0.56	3,989	7,833
2006	11,395	5,964	0.52	1,319	12,714	7,807	0.61	4,907	9,733
2007	17,979	6,839	0.38	1,307	19,286	8,955	0.46	10,331	16,873

*Antler restriction changes definition of a legal buck for 2005–2007, excluding spike-horns from the initial buck population estimate (N-hat) and other estimates until the final post-hunt column.

Table 2.9 Prerifle season legal buck population estimates (N-hat) by region in 2007 and then corrected for bucks taken before the rifle season for pre-hunt estimates. Note variable harvest rates and pre-hunt density estimates among regions. Final pre-hunt estimated population density includes all bucks, does, and fawns as described in the text. Total state “Buck N-hat” estimate (and following population estimates) is the sum of models run for each region separately; it does not exactly match the model for the state as a whole (Table 2.8), but it is close and well within the 95% confidence intervals.

Region	WMUs	Buck N-hat	Rifle harvest	Rifle harvest rate	Mi2	Rifle harvest/ Mi2	Early youth and archery bucks	Pre-hunt bucks	Pre-hunt density (bucks/ mi2)	Total buck harvest	Total buck harvest rate	Estimated population pre-hunt density
Lake Plains	A,B,F1,F2	2,102	1,251	0.595	1,001	1.25	317	2,419	2.42	1,731	0.716	19.6
Mountains	C,G,I, L,P	3,675	1,062	0.289	1,930	0.55	158	3,833	1.99	1,376	0.359	16.1
Northeast	D1,D2,E	2,625	1,037	0.395	1,539	0.67	172	2,797	1.82	1,333	0.477	14.8
East-central	H1,H2, J1,J2	5,668	1,645	0.290	1,542	1.07	316	5,984	3.88	2,131	0.356	31.5
Western Foothills	K1,K2,N	2,005	1,008	0.503	685	1.47	200	2,205	3.22	1,302	0.590	26.2
Eastern Foothills	M1,M2, O1,O2,Q	2,461	828	0.336	1,178	0.70	144	2,605	2.21	1,082	0.415	18.0
State	All	18,536	6,831	0.369	7,874	0.87	1,307	19,843	2.52	8,955	0.451	20.5

WMUs D1, D2, and E may be able to sustain deer densities of 18, 13, and 8 deer per square mile, respectively. Densities will even vary locally within WMUs. The Department recognizes that it cannot manage deer densities directly

at any local small scale level. One of the working assumptions underlying small scale deer management in Vermont and other states is that many hunters, especially archers, will congregate in localized areas having higher deer densities within WMUs. This assumption is substantiated by harvest data from Vermont towns.

From previous experiences, the Department can set population goals that include regional deer densities. Statistical advancement in wildlife science made in recent years now allows for accurate estimates of deer density without incurring the high costs. Fine-tuning regional population estimates to small scale WMU-level estimates will be possible using data such as antlerless tag fill rates and hunter sighting rates of

Table 2.10 Deer population density objectives by Vermont regions for the planning period of 2010-2020.

Region	WMUs	Deer Habitats (mi ²)	Density Goal Range (deer/mi ²)		Population Goal Range (deer/mi ²)	
Lake Plains	A,B,F1,F2	1,001	16	21	16,000	21,000
Mountains	C,G,I,L,P	1,930	13	18	25,100	34,750
Northeast	D1,D2,E	1,539	10	15	15,400	23,100
East-central	H1,H2,J1,J2	1,542	15	20	23,100	30,800
W. Foothills	K1,K2,N	685	15	20	10,300	13,700
E. Foothills	M1,M2,O1,O2,Q	1,178	10	15	11,800	17,700
State	All	7,874	13	18	101,700	141,100

deer. The Department will be attempting to track deer densities at the state, regional, and WMU levels using a variety of methods that include the following:

- 1) Population estimation models using harvest and hunter effort data
- 2) Catch-per-unit-effort prehunt population estimation
- 3) Road-kill data for adult sex ratios, reproductive rates, and fawn recruitment through winter to provide necessary data for various analyses
- 4) Bow hunter surveys to determine autumn buck:doe and fawn:doe ratios and sighting rates
- 5) Rifle hunter surveys to gather deer sighting rate data

- 6) Age data to assist in determination of survival estimates and sex ratios
- 7) Change-in-ratio methods using road-kill data

A well established tenet of deer population biology is that altering survival rates of adult females is the most effective way of altering the trajectory of a deer population (Gaillard et al. 2000, Haskell and Ballard 2007). Only by regulating the antlerless deer harvest, 80% of which is typically made up of adult does, will it be possible to meet Vermont's deer population density objectives.

Maintaining Vermont's deer population density at ecologically sustainable levels is the only way to ensure the health and vigor of Vermont's deer herd, native forest, and necessary deer habitats (for example, deer yards). A deer herd in balance with its habitat will have few negative impacts on other wildlife species, the forest and agricultural industries, and will minimize conflicts with people. It will, it is hoped, also prevent periodic boom and bust cycles of deer abundance that have characterized the history of deer in Vermont.

This overall message is not new and cannot be over-emphasized. It has been widely promoted by the Department since at least the mid-1900s (Seamans 1946). Because prehunt population density estimation can only occur after data from the autumn deer seasons and because the impact of the oncoming winter is unpredictable, the task of determining appropriate antlerless harvest objectives for the next fall is a necessarily reactive process. While winter may always be an unpredictable factor, the development of predictive population models is expected to improve through time with additional data and experience. It is hoped the future will provide the tools to make deer management more proactive than reactive.

Management Strategies

- 2.1 Maintain and evaluate regional population goals, established during this planning period, that are based on deer densities that recognize a lower limit that is unsatisfactory to the public and an upper limit that is ecologically unsustainable.
- 2.2 Monitor deer herd health by collecting body condition data from hunter-harvested and road-killed deer.
- 2.3 Consider establishing habitat suitability criteria to define areas of suitable deer habitat within WMUs so that consistent and reliable density

estimates can be made while allowing for habitat area estimate updates as new land-cover maps become available.

- 2.4 Evaluate bowhunter surveys to better estimate regional buck:doe and fawn:doe ratios; compare fawn production estimates to autumn fawn:doe ratios to estimate summer fawn survival, and use buck:doe ratios to estimate adult doe population through reference to the unbiased buck population estimate.
- 2.5 Continue remapping and surveying deer wintering areas so that available habitat is quantified and localized winter deer density is better documented.
- 2.6 Work with foresters to develop data-driven methods for assessing localized deer overabundance problems that might lead to development of localized deer management methods. Data must provide measures of forest condition.
- 2.7 Provide outreach to landowners regarding methods that may minimize damage and encourage reduction in locally overabundant deer populations. Investigate feasibility of a formal program to connect hunters with landowners to address locally overabundant deer populations.
- 2.8 Develop strategies to maintain enough big game registration stations to make big game reporting convenient for hunters.
- 2.9 Seek statutory changes to realign boundaries of select WMUs as proposed above.

ISSUE 3. Hunter Satisfaction and Antler Point Restrictions

GOAL: Employ biologically responsible, socially responsive, and adaptive management of the deer herd.

The Department continually monitors deer hunter opinions. Although opinions will vary widely among hunters, collecting their observations and views is a useful "tool" in managing the deer herd. The Department gains insight into the "will of the people" via five annual public meetings held in the spring as well as through many public outings at reporting stations, sporting shows, game clubs, and various other venues. Daily contacts between state game wardens and the public also provide rapid

feedback from the public to the Department. Since 1999, the Department's annual hunter effort surveys and periodic opinion polls have provided both general and specific feedback that may be focused on some pressing, current issue. In recognizing the value, and absolute necessity of listening to the people, the Department has made it a goal to continue to improve methods for public input.

HUNTER SATISFACTION

Generally, the effects of winter severity on the deer herd correlate with changes in deer population density. Data since 1970 demonstrate that fluctuations in rifle season buck harvests have fairly predictably paralleled changes in winter severity (Fig. 2.14). This suggests that winter severity has continually influenced deer density in Vermont.

Anecdotal feedback from hunters, as well as increased license sales in 2007 and 2008, suggest hunter satisfaction has improved greatly since 2006. As the deer population rebounded, hunters have seen more deer and harvests have increased (Fig. 2.15). While biologists understand that perhaps the single greatest influence on hunter satisfaction is how many and how often deer are seen, there is a growing interest in the qualitative characteristics of Vermont's deer population.

ANTLER POINT RESTRICTIONS

In 2005, Vermont established a new antler restriction (AR) designed to "spare" a larger portion of yearling bucks and allow them to mature to an older age. Although this regulation was intended to change the age structure of the buck population by increasing the proportions of bucks in older age classes, it also slightly increased the total number of bucks and ratio of bucks to doe.

Prior to establishment of Vermont's antler restriction regulation, about 50% of each year's crop of yearling bucks in Vermont were spike-horns. This regulation protected these yearlings and resulted in a surge of two-year-old bucks and smaller increases in other age classes (Fig. 2.16).




Figure 2.14 Relative annual change in rifle season harvest from one year to the next predicted by winter severity in Vermont from 1970–2004.




Figure 2.15 Number of white-tailed deer seen per 10 hours of hunting time as reported by Vermont hunters from 2000–2008.




Figure 2.16 Age structure of bucks sampled by Department biologists during opening weekend of rifle seasons 2000–2007 as determined by tooth wear and replacement. Buck sample (n=248) for 2008 taken from buck heads submitted by cooperating meat-cutters during rifle season for disease testing.

Not only have older buck populations increased under the new AR but the weights of harvested bucks have increased. Before the antler restriction, the average field-dressed weight of bucks checked by biologists was 125 pounds. By 2007, the average weight increased to 138 pounds. In 2007 9,000 bucks were harvested yielding 117,000 pounds more of field-dressed deer and 50,000 pounds more of edible deer meat than the same number of bucks harvested in 2003.

For the first time, the quality (that is, the antler and/or body size) of deer has begun to compete with the quantity of deer as a driver of satisfaction among Vermont hunters. The Department continually monitors social acceptance and biological integrity of the statewide antler restriction experiment. Already, new concerns related to the “quality of deer” have surfaced as some hunters and scientific publications have expressed concern that protecting the smaller yearlings from harvest could have an effect on the gene pool of the deer herd (Harmel et al. 2001, Strickland et al. 2001, Coltman et al. 2003, Demarais et al. 2005, Festa-Bianchet 2007, Coltman 2008). There are, however, several reasons why, at least in the short term, adverse effects on the gene pool are not likely:

1. Does contribute 50% to genetic recombination.
2. Twin fawns have different sires about 20% of the time, and in general, it is normal for small bucks to breed does (Sorin 2004).
3. Mature and heavy does tend to breed early, which may occur before the rifle season when most bucks are harvested (Haskell et al. 2008).
4. Dominant male deer are polygamous, they breed many does, which may mean they breed early and sire disproportionately more male than female offspring (Gomendio et al. 2006, Roed et al. 2007).

Also, many confounding environmental factors, such as food availability and winter severity, can affect antler size and shape, particularly deer population density as it relates to nutrition (Harmel et al. 2001, Williamson 2003, Keyser et al. 2005, Gomez et al. 2006, Strickland and Demarais 2008).

The Department has not yet conducted thorough research into the issue of deer population genetics to be able to determine whether this issue needs to be addressed. Prudence dictates that we monitor the results of this statewide experiment closely for signs of

change. Future research and knowledge may suggest the need to modify the antler restriction to better manage for the future.

The youth deer hunt has become particularly important as a source of unbiased data on bucks. Because youths can take any yearling buck, data from the youth hunt provides a sample of the entire yearling buck population and provides data that is comparable to data collected during seasons before the antler restriction. By comparing data from pre- and post-AR harvests, it will be possible to detect any changes that may result from the antler restriction that might have some potential future effect on the deer herd. Based on assessment of pre-AR data, the current AR of two points on one side protects about 50% of yearling bucks while an AR of three points on one side would protect about 90% of yearlings. In the future, a three-point on one side AR could be considered if genetic issues were found to be of concern or if hunter preference for older aged bucks was to increase.

The antler restriction has worked to slightly increase the age structure of bucks because it has increased yearling survival rate during the hunting season, a time when yearling bucks are most vulnerable to mortality. The antler restriction is not expected to increase the number of four-year old or older bucks because the harvest rate of two-year old and older bucks remains high. In the future, some modification of the current restriction to three-points on one side, some slot limit, or other regulation to achieve desired harvest and population objectives may be appropriate.

There are also ways other than antler restrictions that can be used to increase survival rates of bucks. Alternatives include several ways to restrict hunting opportunity of bucks, such as reduced seasons, restrictive weapons, and reduced bag limits (see Issue 4: Bag Limits). The main cause of mortality of Vermont bucks, 76% of the total buck harvest, is during the rifle buck season when, in fact, only one buck can be taken. Even if there was a need or hunter support to change this proportion, it would require a legislative change. The rifle deer season is set by statute and cannot be changed by the Fish and Wildlife Board. The Department will remain open to the use of all effective methods understanding that implementation is dependent on public acceptance.

Management Strategies

- 3.1 Collect adequate yearling buck data (weights, antler beam diameter, and number of points)

from the youth hunt to detect and track any changes in the buck population resulting from the current antler-point restriction (two points-on-one-antler minimum), and evaluate biologically acceptable alternatives if needed.

- 3.2 Evaluate a model assessment using genetic data to examine the likelihood of altering the genetic diversity of the buck population via the current antler restriction.
- 3.3 Inform the hunting public about deer management issues and results of antler-point restrictions and gather input concerning deer management and hunter satisfaction.

ISSUE 4. Bag Limits

GOAL: Provide suitable utilization of deer as food and provide opportunity to hunt deer in a way that maximizes potential for effective deer population management but does not overstress the heavily harvested buck population.

One of the Department's objectives is to provide as much opportunity as is sustainably possible to hunt, fish, trap, and view wildlife in Vermont. In particular, restoring and increasing hunting opportunities and participation is one of our foremost goals during this planning period that follows a period in which hunter participation has declined.

Vermont's bag limit of three deer per calendar year has been a topic of some controversy among hunters since the poor deer season of 2001. Despite data consistently demonstrating the three-deer bag limit has very little effect on the overall harvest (Table 2.11), hunters were able to persuade the Fish and Wildlife Board to reduce the bag limit to two deer for the purpose of increasing the size of the deer population. The real impact of this action was a reduction in hunting opportunity and a reduction in the amount of time hunters spent afield. An unintended consequence of the change was a reduction in the number of female deer harvested because hunters did not wish to sacrifice an opportunity to hunt bucks during the rifle season by taking antlerless deer.

As history demonstrates, the third deer provided additional opportunity and an incentive for hunters to go deer hunting while very few deer, especially

Table 2.11 Percent of successful hunters harvesting 1, 2, or 3 deer for the period 2000 – 2008.

Year	1 deer	2 deer	3 deer	Deer Harvest
2000	83%	14%	3%	20,498
2001	83%	15%	1%	15,065
2002	85%	13%	2%	16,261
2003	88%	10%	2%	14,528
2004	90%	8%	2%	11,925
2005	93%	7%	X	8,546
2006	92%	9%	X	12,682
2007	89%	11%	X	14,516
2008	84%	15%	2%	17,046

bucks, were actually ever bagged as a third deer. Returning to a three deer limit in 2008, once again, afforded Vermont hunters more days afield and improved the harvest of does.

Management Strategies

- 4.1 Provide the public with ample opportunity to harvest white-tailed deer for food and other utilitarian purposes.
- 4.2 Advocate for an appropriate deer bag limit that allows maximum hunter opportunity while achieving deer population management strategies.

ISSUE 5. Muzzleloader and Archery Season Modifications

GOAL: Provide suitable opportunity to hunt deer in a way that maximizes the potential for effective deer population management but does not interfere with hunters during youth weekend or rifle and other fall hunting seasons.

While hunter participation in the rifle season has remained consistently high at 88% over the past decade, participation in alternative seasons has increased. Hunter participation in the muzzleloader season increased from 32% in 1996 to 43% in 2007 while participation in archery also increased from 27% to 33% (Duda et al. 2007). One survey found that more Vermont deer hunters (48%) preferred the muzzleloader season occurring after the rifle season than those who preferred a season occurring before the rifle season (30%).

The timing and length of the archery season or any proposal for an early muzzleloader season should be

carefully considered given the need for a special youth weekend before the rifle season and the interests of landowners. Since there is already a heavy harvest of bucks in Vermont, any early muzzleloader season should be tailored to the task of controlling doe numbers. Archery hunters tend to hunt from tree-stands more than muzzleloader hunters whose weapons have greater range. Many muzzleloader hunters prefer the late season because it provides greater likelihood that snow will be on the ground to improve tracking and visibility of deer.

The Department plans to enhance efforts to gather and use archery deer hunter observation data under the assumption that archers in tree stands observe deer at closer range and will be able to provide reliable observations, such as fawns per doe and buck to doe ratios. If these data prove useful, it will benefit all deer hunters.

Many Vermonters have expressed the opinion that more antlerless deer should be harvested before the November rut and December muzzleloader seasons suggesting that an early season could reduce the amount of browse consumed by 1,500 or more antlerless deer that would otherwise be harvested five or six weeks later. Most antlerless deer are currently being taken during early archery and youth seasons prior to the existing muzzleloader season. Taking more antlerless deer early in the season may be desirable.

One way to do this is to open a weekend or a few days to antlerless-only muzzleloader hunting prior to the regular rifle season, which could increase the number of antlerless deer taken before the regular rifle season. It is possible that this might also increase muzzleloader participation and the fill rate of antlerless deer tags as well as improve the Department's ability to manage Vermont's deer herd in areas where deer densities are high. The challenge is to create an early muzzleloader season without disturbing hunters participating in the other seasons — youth weekend, archery, turkey, small game, and rifle. This is a task that would require careful research and considerable input from the various user groups. The same arguments could be made for expanding the archery season. Many of the same challenges would also need to be addressed.

Because the Department relies on archery and muzzleloader hunters to harvest antlerless deer, it is prudent to regain their participation and ensure an



ROBERT CAMERON

ability to manage deer densities in Vermont. Archery and muzzleloader license sales declined from 74,193 in 2000 to 36,322 in 2005 as deer populations and hunting opportunity declined. Numbers rebounded to 43,585 in 2007 as deer numbers and opportunity again increased. In addition, longer archery seasons in neighboring states of New Hampshire, Massachusetts, Maine, and New York may have contributed to the decline in archery hunters in Vermont. It seems wise to investigate potential conflicts between seasons in neighboring states and then to assess how the situation in these states encourages or discourages nonresident participation in Vermont's early archery season.

Other means of increasing archery participation are through expanding archery season length or increasing archery season bag limits. Both strategies enhance the ability to harvest antlerless deer where needed, including areas with locally overabundant deer populations or where firearm ordinances restrict opportunities to harvest antlerless deer during the youth weekend or muzzleloader season.

Petitions to the Fish and Wildlife Board and the Department have asked for consideration to make crossbows legal for general use in Vermont and to expand archery seasons. Currently, only individuals who can show evidence of a physical disability that restricts the ability to draw a compound bow are permitted to use crossbows in Vermont. While legalizing the use of crossbows during the archery

season could increase the Department's ability to harvest does, mixed public response to the concept, however, suggests that this harvest management tool should be deferred until it is determined that other, more popular harvest strategies will not achieve population objectives.

In 2005 baiting and feeding deer was made illegal after a lengthy regulatory process involving a great deal of public involvement. When deer are baited or fed, there are serious concerns of disease threat and improper feeding methods that are actually detrimental to deer. This practice also alters the natural digestive system and movement patterns of deer. Although there are still hunters who want to bait deer, the Department believes baiting poses a threat to the health of Vermont's deer herd and does not want to reopen this issue.

Management Strategies

- 5.1 Evaluate feasible options to expand antlerless deer-only hunting opportunities prior to the regular rifle season. These options will include, but are not limited to, an early muzzleloader season, expanded archery season, and increases in archery bag limits.
- 5.2 During the fall and winter of 2009-2010, survey public opinion on the various management options to achieve antlerless harvest objectives prior to the rifle season and develop a proposal of recommended hunting season changes for the Vermont Fish and Wildlife Board in 2010.

ISSUE 6. Captive Deer Hunting/ Deer Farming/ Cervid Importation

GOAL: Implement new captive hunting regulations and work with other state agencies to minimize the chance of introducing and transmitting diseases via captive deer.

In 1986, Vermont passed legislation authorizing the inclusion of certain deer species in agriculture as



MIKE CARPENTER

part of a modern, diversification effort. Fallow deer and red deer were identified as domestic deer species and were legalized to import, possess, and propagate in Vermont the same as any domestic farm animal. Since then, fallow deer, red deer, and elk have been legally imported for agricultural purposes and have been propagated at captive hunt facilities.

The concern with introducing other deer species centers on the potential for spreading disease. Since 1986, Chronic Wasting Disease (CWD) has emerged as a new disease on the national front that threatens Vermont's deer herd. CWD is a disease of the central nervous system similar in nature to "Mad Cow Disease." There is no known vaccine or cure and always results in the death of animals that contract it. This disease cannot be detected in live animals until the disease symptoms have appeared. One of the more troubling characteristics of CWD is that it can lie dormant in an individual animal for years before symptoms appear. Thus, the presence of the disease can go undetected until years after an animal has been transported to a new farm or location.

Animals infected with CWD can be brought into the state by deer farmers, captive hunt facility owners, and even an unsuspecting hunter who has legally harvested a deer or elk from outside of Vermont. The state has established laws and regulations governing the transportation and importation of live deer as well as deer carcasses and other cervids from states where CWD is known to occur. The Agency of Agriculture, Food, and Markets regulates animals

used for agricultural purposes and the Department of Fish & Wildlife enforces the regulations that govern any animal imported or possessed for the purposes of hunting. These regulations prescribe veterinary inspections, health certificates, and other measures that mediate the threat of CWD.

Before 2000, CWD was thought to be mostly concentrated in parts of Colorado and Wyoming, but more extensive surveillance has resulted in discovery of CWD in 12 additional states and 2 Canadian provinces. Long-distance movement of the disease has most likely been due to the transport of captive deer and elk (Williams et al. 2002, Sigurdson and Aguzzi 2007, Miller 2008). Recent scientific research strongly suggests that CWD can be transmitted through ingesting feces from infected animals. Scientists also believe that it is transmitted through animal-to-animal contact and through contact with an environment that has been contaminated with the infectious prion (a mutant protein). Scientists believe the spread of the prion occurs via lymph tissues, blood, saliva, feces, and urine and can persist in soils for years. For this reason scientists are concerned that if a captive deer has the disease and escapes from a facility, the disease can spread to free-ranging deer populations with devastating results. (Miller and Williams 2003; Miller et al. 2004; Seeger et al. 2005; Mathiason et al. 2006, 2009; Johnson et al. 2007; Andrievskaia et al. 2008; Gonzalez-Romero et al. 2008; Safar et al. 2008; Sigurdson 2008; Angers et al. 2009; Haley et al. 2009; Maddison et al. 2009; Race et al. 2009).

Given the history of CWD-prevalence among captive deer herds, it seems prudent to address the spread of captive deer urine across the landscape. The risk of establishing any new disease into Vermont's native deer and moose population is of great concern to the Department. The eradication of any disease from free-ranging wildlife is nearly impossible and extremely costly. The potential loss of these animals and a way of life enjoyed by many Vermonters is incalculable. The Department believes that prevention is the only suitable option for dealing with CWD.

Management Strategies

- 6.1 Evaluate the effectiveness of the captive hunting facility regulation.
- 6.2 Work with the Agency of Agriculture, Foods, and Markets and the deer farming industry to promote and enforce disease free importation and husbandry practices.

ISSUE 7. Disease Surveillance and Management

GOAL: Monitor disease issues and respond when necessary to protect the health of wildlife and/or humans.

According to state statute, "...the protection, propagation control, management and conservation of fish, wildlife, and fur-bearing animals in this state is in the interest of the public welfare, and that safeguarding of this valuable resource for the people of the state requires a constant and continual vigilance" (Title 10 V.S.A. §4081(a)). As human and deer populations expand or are transported with relative ease, the risk of disease transmission increases and with it the Department's ability to fulfill its statutory charge. Some diseases do not present a serious consequence to wildlife or humans. However, some diseases associated with deer such as chronic wasting disease (CWD), Lyme disease, hemorrhagic disease (HD), tuberculosis (Tb), and babsiosis, present risks to humans, as well as deer.

CWD, as discussed in Issue 6, is a fatal disease of the nervous system that afflicts white-tailed and mule deer, elk, and moose. It has no known cure or vaccine and can have a long incubation period. Hemorrhagic disease is a deer disease that is common in the Southeast and the Midwest. Twenty years ago the disease was only known to exist south of Pennsylvania and New Jersey (Davidson and Nettles 1997). In 2007, confirmed cases of HD were reported in Albany County, New York, in the Hudson River drainage basin that extends into southwestern Vermont. Although HD is well understood, it is not a disease that can be readily managed. It is a viral disease that is transmitted by a small biting midge fly, often called "no-see-ums." The disease occurs in warm months. As the first frosts of autumn occur, the disease abates as the flies die off for the season. Deer often survive HD, but it can cause localized, periodic, and sometimes heavy mortality. This is a disease that will bear watching in the future as global temperatures change and result in the northward spread of the vector of this disease.

One of the Department's goals is to "limit harmful or fatal human encounters with fish and wildlife species, and provide general public safety service incidental to our primary fish and wildlife duties." Lyme disease, babsiosis, and Tb are capable of crossing from wildlife

over to other species including humans. Although cattle are more closely associated with Tb distribution in North America, deer are capable of sustaining this bacterium in the wild and acting as a reservoir, having the potential to infect and re-infect cattle and human populations. Michigan has spent millions of dollars attempting to eradicate Tb from cattle and wild deer populations. This case is a clear example of how once a disease enters wild animal populations, it is nearly impossible to eradicate.

The incidence and distribution of Lyme disease in Vermont has steadily increased in the last decade and shows no signs of abating. This disease is caused by a mycobacterium transported by a complicated relationship between black-legged ticks, white-footed mice, and deer. Populations of all three of these species have grown as the landscape has become more suburbanized, creating favorable habitats for these species in close proximity to concentrated human populations. Lyme disease infection begins with a tick bite that transmits the bacteria. The site of the bite often erupts into a “bulls-eye” rash that sometimes is accompanied by fever. As the rash soon disappears, the individual may believe that he/she has no disease. The disease, however, has merely moved to the next stage, which can lead to debilitating joint disease in humans and dogs if left untreated.

The use of urine from captive deer as a scent lure is legal in Vermont. Given the possible presence of CWD in captive deer that appear healthy and excretion of infectious prions in urine (see Issue 6 and References for citations of supporting scientific literature), it may be prudent to address the spread of captive deer urine across the landscape where disease-free native deer could contact the infectious agent. With recent advances in prion-detection methods, it is now unquestionable that scent lures originating from captive deer urine and used by hunters pose a risk of introducing CWD into CWD-free areas such as Vermont. Artificial, or synthetic, scents pose no such risk and have been commercially available since at least 2004.

Vermonters may be unaware of the seriousness of this particular disease issue and how it is transmitted from captive deer to wild populations. Dissemination of the Department’s CWD Response Plan may help educate the public. The plan includes identification of a CWD-positive free-ranging deer (deer or moose) and calls for total extermination of free-ranging deer within a five-mile radius for several years – that area

is equal to 79 square miles or about two Vermont towns. If infected deer continue to be found in the area, the control-area radius is then extended to ten miles – an area equal to 314 square miles. This is standard protocol among CWD-free states and provinces in North America. This disease has the potential to greatly impact populations of deer, deer hunters, and deer watchers alike — it is not to be taken lightly.

Management Strategies

- 7.1 Work with associated branches of government (for example, Agency of Agriculture, Department of Health) to monitor and control disease agents and deer populations where and when it is appropriate.
- 7.2 Contribute to the national CWD surveillance effort.
- 7.3 Monitor the progress of Hemorrhagic Disease as it moves toward the Vermont border.
- 7.4 Work closely with the Agency of Agriculture to ensure dairy farms and domestic deer farms maintain their tuberculosis-free status.
- 7.5 Investigate a prohibition on the use of deer-urine-based scent lures and, if appropriate, implement a public informational effort on the justification.
- 7.6 Inform Vermonters as to the gravity of CWD and repercussions if introduced into our environment through the dissemination of Vermont’s CWD Response Plan.

ISSUE 8. Locally Overabundant Deer Populations

GOAL: Promote awareness that hunting is the only practical option to reduce localized overabundant deer populations.

Ordinances in urban and suburban communities may restrict normal hunting activities, which prompt landowners to also post land against hunting. Deer, however, can live and propagate successfully in many of these environments. Without natural or human predation, deer populations grow quickly. This overabundance often results in increased foraging on agricultural or residential plantings, deer-vehicle collisions, and incidences of Lyme disease (McShea et al. 1997, Schwabe and Schuhmann 2002). As Vermont’s human population continues to grow, the



DAVE ADAMS

expanding suburban setting will cause deer-human conflicts to become more and more common.

There are a variety of nonlethal and lethal options for mitigating conflicts with human residents and managing overabundant white-tailed deer in suburban environments (DeNicola et al. 2000). Nonlethal measures include trap and transfer, fencing, sulphur-based plant sprays, and other aversive measures such as noise makers and flashing lights. Trap and transfer methods incur many risks ranging from injury to captured animals to impacts upon the social stability of receiving deer populations. All of these nonlethal methods are impractical for alleviating localized deer overabundance problems (Buck et al. 2009).

Lethal measures include a myriad of controlled hunting strategies that limit the hunter's location, time of day, and implement (for example, bow-and-arrow, crossbow, muzzleloader, or shotgun). Implements that have a limited discharge range, for example, bows, are perceived by the public as being more acceptable for use in close proximity to buildings and people. Alternative hunting strategies can also effectively and safely reduce deer numbers. Experience from urban areas in other states has demonstrated that most residents who opposed alternative hunts before implementation actually came to support the hunts once they were applied successfully (Deblinger et al. 1995, Frost et al. 1997, Mitchell et al. 1997, McDonald et al. 1998, Kilpatrick and Labonte 2003).

Archery hunters have proven to be an effective general management tool for deer in Vermont and in other states as a way to control suburban deer populations (Kilpatrick and Walter 1999, Kilpatrick and Labonte 2003). Suburban residents may be more supportive of alternative hunts when they are allowed to restrict hunting activity on their own property and when archery hunters involved in the hunt have completed

a state-certified hunter safety course including a test for shooting proficiency (Kilpatrick et al. 2007). In 2006, there were 19,173 archery permits sold in Vermont resulting in a harvest of 2,553 deer for an overall success rate of 13%, which is similar to that for rifle hunting. Of the 2,553 deer harvested during the 2006 archery season, 59% were adult does. As previously discussed, increasing the harvest of adult does is the most effective way to reduce a deer population when this becomes the desired management objective.

Management Strategies

- 8.1 Demonstrate the effectiveness of archery hunting to reduce locally overabundant deer in Vermont's suburban environments.
- 8.2 Provide communities with up-to-date and comprehensive information on deer overabundance and consider community views when deciding how to best manage deer problems in suburban, agricultural, and forested areas.
- 8.3 Encourage communication and cooperation between antlerless deer hunters and landowners that seek relief from locally overabundant deer.

ISSUE 9. Two-year Regulation Cycle

GOAL: Consider a more efficient two-year regulatory cycle that allows for annual adjustments when environmental factors deem it appropriate.

As a means to reduce costs of deer management, increase management continuity, and make regulations more consistent from year to year for hunters, the Department will investigate the feasibility of a two-year regulatory cycle instead of the one-year cycle it now operates. This could save time and money developing and printing deer hunting regulation changes every year. This approach is used in other states, New Hampshire, for example.

Management Strategies

- 9.1 Provide outreach to legislators, board members, and hunters to develop an understanding of the rationale behind deer management and proposed actions to improve management.
- 9.2 Evaluate the benefits and deficiencies of implementing a two-year regulation cycle for deer season recommendations.

REFERENCES - WHITE-TAILED DEER

- Akins, J. W., and E. D. Michael. 1995. Impact of clear-cut size on white-tailed deer use and tree regeneration. *Proceedings of the Eastern Wildlife Damage Control Conference* 6:185–195.
- Andrievskaia, O., J. Algire, A. Balachandran, and K. Nielson. 2008. Prion protein in sheep urine. *Journal of Veterinarian Diagnostic Investigations* 20:141–146.
- Angers, R. C., T. S. Seward, D. Napier, M. Green, E. Hoover, T. Spraker, K. O'Rourke, A. Balachandran, and G. C. Telling. 2009. Chronic wasting disease prions in elk antler velvet. *Emerging Infectious Diseases* 15:696–703.
- Augustine, D. J., and D. DeCalesta. 2003. Defining deer overabundance and threats to forest communities: from individual plants to landscape structure. *Ecoscience* 10:472–486.
- Ballard, W.B., H.A. Whitlaw, S.J. Young, R.A. Jenkins, and G.J. Forbes. 1999. Predation and survival of white-tailed deer fawns in north-central New Brunswick. *Journal of Wildlife Management* 63:574–579.
- Buck, J. M., C. Kandoth, and S. Predl, Editors. 2009. *An Evaluation of Deer Management Options*. Northeast Deer Technical Committee.
- Coltman, D.W. 2008. Molecular ecological approaches to studying the evolutionary impact of selective harvesting wildlife. *Molecular Ecology* 17:221–235.
- Coltman, D.W., P. O'Donoghue, J. T. Jorgenson, J. T. Hogg, C. Strobeck, and M. Festa-Bianchet. 2003. Undesirable evolutionary consequences of trophy hunting. *Nature* 426:655–658.
- Côté, S. D., T. P. Rooney, J.-P. Tremblay, C. Dussault, and D. M. Waller. 2004. Ecological impacts of deer overabundance. *Annual Review of Ecology Evolution and Systematics* 35:113–147.
- Crête, M. 1999. The distribution of deer biomass in North America supports the hypothesis of exploitation ecosystems. *Ecology Letters* 2:223–227.
- Davidson, W. R., and V. F. Nettles. 1997. *Field manual of wildlife diseases in the southeastern United States*. Second edition. Southeastern Cooperative Wildlife Disease Study, University of Georgia, Athens.
- Deblinger, R. D., D. W. Rimmer, J. J. Vaske, and G. M. Vecellio. 1995. Efficiency of controlled, limited hunting at the Crane Reservation in Ipswich, Massachusetts. Pages 75–79 in J. N. McAnich, ed. *Urban deer: a manageable resource?* 1993 Symposium of the North Central Section of the Wildlife Society, St. Louis, MO.
- DeCalesta, D. S. 1994. Effect of white-tailed deer on songbirds within managed forests in Pennsylvania. *Journal of Wildlife Management* 58:711–718.
- DelGiudice, G. D., M. S. Lenarz, and M. Carstensen Powell. 2007. Age-specific fertility and fecundity in northern free-ranging white-tailed deer: evidence for reproductive senescence? *Journal of Mammalogy* 88:427–435.
- Demarais, S., K. V. Miller, and H. A. Jacobson. 2000. White-tailed deer. Pages 601–628 in S. Demarais and P. R. Krausman, editors. *Ecology and management of large mammals in North America*. Prentice-Hall, Upper Saddle River, New Jersey.
- Demarais, S., B. K. Strickland, L. E. Castle. 2005. Antler regulation effects on white-tailed deer on Mississippi Public Hunting Areas. *Proceedings of the Southeastern Association of Fish and Wildlife Agencies* 59:1–9.
- DeNicola, A. J., K. C. Vercauteren, P. D. Curtis, and Scott E. Hyngstrom. 2000. *Managing white-tailed deer in suburban environments: a technical guide*. A publication of Cornell Cooperative Extension, the Wildlife Society, and the Northeast Wildlife Damage Research and Outreach Cooperative.
- Donovan, T.M., and M. Alldredge. 2007. Exercises in estimating and monitoring abundance. <http://www.uvm.edu/envnr/vtcfwru/spreadsheets/abundance/abundance.htm>
- Duda, M. D., M. Jones, A. Criscione, C. Craun, T. Beppler, T. Winegord, A. Lanier, S. J. Bissell, and J. B. Herrick. 2007. *Public opinion on wildlife species management in Vermont*. Report by Responsive Management for the Vermont Fish & Wildlife Department, Waterbury, VT.
- Dusek, G.L., R. J. Mackie, and J.D. Herriges. 1989. Population ecology of white-tailed deer along the lower Yellowstone River. *Wildlife Monographs* No. 104.
- Festa-Bianchet, M. 2007. Ecology, evolution, economics, and ungulate management. Pages 183–202 in T. E. Fulbright and D. G. Hewitt, eds. *Wildlife science: linking ecological theory and management applications*. CRC Press.

- Frost, H. C., G. L. Storm, M. J. Batcheller, and M. J. Lovallo. 1997. White-tailed deer management in Gettysburg National Military Park and Eisenhower National Historic Site. *Wildlife Society Bulletin* 25:462–469.
- Gaillard, J.-M., M. Festa-Bianchet, N. G. Yoccoz, A. Loison, and C. Toïgo. 2000. Temporal variation in fitness components and population dynamics of large herbivores. *Annual Review of Ecology and Systematics* 31:367–393.
- Garland, L. E. 1978. 1978 Deer season prospects, and Vermont's experience with a buck-only season since 1897. Vermont Fish and Game Department, Montpelier.
- Giles, B.G., and S.C. Findlay. 2004. Effectiveness of a selective harvest system in regulating deer populations in Ontario. *Journal of Wildlife Management* 68:266–277.
- Gomendio, M., A. F. Malo, A. J. Soler, M. R. Fernandez-Santos, M. C. Estes, A. J. Garcia, E. R. S. Roldan, and J. Garde. 2006. Male fertility and sex ratio at birth in red deer. *Science* 314:1445–1447.
- Gomez, J. A., T. Landete-Castillejos, A. J. Garcia, and L. Gallego. 2006. Importance of growth during lactation on body size and antler development in the Iberian red deer (*Cervus elaphus hispanicus*). *Livestock Science* 105:27–34.
- Gonzalez-Romero, D., M. A. Barria, P. Leon, R. Morales, and C. Soto. 2008. Detection of infectious prions in urine. *FEBS Letters* 582:3161–3166.
- Haley, N. J., D. M. Seelig, M. D. Zabel, G. C. Telling, and E. A. Hoover. 2009. Detection of CWD prions in urine and saliva of deer by transgenic mouse bioassay. *PLoS ONE* 4(3): e4848. doi:10.1371/journal.pone.0004848.
- Halls, L. K., editor. 1984. White-tailed deer: ecology and management. Stackpole Books, Harrisburg, Pennsylvania.
- Harmel, D. E., J. D. Williams, and W. E. Armstrong. 2001. Effects of genetics and nutrition on antler development and body size of white-tailed deer. Texas Parks and Wildlife Department, Kerr Wildlife Management Area, Hunt, TX.
- Haskell, S. P. 2007. Ecology of sympatric deer species in west-central Texas: methodology, reproductive biology, and mortality and anti-predator strategies of adult females and fawns. PhD dissertation, Texas Tech University, Lubbock, TX.
- Haskell, S. P., and W. B. Ballard. 2007. Modeling the growth phase: potential effects of wolves and radio collars. *Journal of Wildlife Management* 71:619–627.
- Haskell, S. P., W. B. Ballard, D. A. Butler, N. M. Tatman, M. C. Wallace, C. O. Kochanny, and O. J. Alcumbrac. 2007. Observations on capturing and aging deer fawns. *Journal of Mammalogy* 88:1482–1487.
- Haskell, S. P., W. B. Ballard, D. A. Butler, M. C. Wallace, T. R. Stephenson, O. J. Alcumbrac, and M. H. Humphrey. 2008. Factors affecting birth dates of sympatric deer in west-central Texas. *Journal of Mammalogy* 89:448–458.
- Haskell, S. P., D. A. Butler, W. B. Ballard, M. J. Butler, M. C. Wallace, and M. H. Humphrey. 2009. Deer density estimation in west-central Texas: old versus new ground techniques with mark-resight as a comparative baseline. *Proceedings of the Western States and Provinces Deer and Elk Workshop 7*: in press.
- Keyser, P. D., D. C. Gyun, and H. S. Hill. 2005. Population density-physical condition relationships in white-tailed deer. *Journal of Wildlife Management* 69:356–365.
- Horsley, S. B., S. L. Stout, and D. S. DeCalesta. 2003. White-tailed deer impact on vegetation dynamics of a northeastern hardwood forest. *Ecological Applications* 13:98–118.
- Johnson, C.J., J.A. Pedersen, R.J. Chappell, D. McKenzie, and J.M. Aiken. 2007. Oral transmissibility of prion disease is enhanced by binding to soil particles. *PLOS Pathogens* 3:874–881.
- Kilpatrick, H. J., and A. M. Labonte. 2003. Deer hunting in a residential community: the community's perspective. *Wildlife Society Bulletin* 31:340–348.
- Kilpatrick, H. J., and A. M. Labonte. 2007. Managing urban deer in Connecticut: a guide for residents and communities. Second edition. Bureau of Natural Resources, Wildlife Division, Hartford, CT, USA.
- Kilpatrick, H. J., A. M. Labonte, and J. S. Barclay. 2007. Factors affecting bowhunter access in suburban areas. *Journal of Wildlife Management* 71:2102–2105.
- Kilpatrick, H. J., and W. D. Walter. 1999. A controlled archery deer hunt in a residential community: cost, effectiveness, and deer recovery rates. *Wildlife Society Bulletin* 27:115–123.
- Lancia, R. A., J. D. Nichols, and K. H. Pollock. 1996. Estimating the number of animals in wildlife populations. Pages 215–253 in T. A. Bookhout, ed.

- Research and management techniques for wildlife and habitats. Fifth edition. The Wildlife Society, Bethesda, Maryland.
- Maddison, B. C., C. A. Baker, H. C. Rees, L. A. Terry, L. Thorne, S. J. Bellworthy, G. C. Whitlam, and K. C. Gough. 2009. Prions are excreted in milk from clinically normal scrapie-exposed sheep. *Journal of Virology* doi:10.1128/JVI.00051-09 (online ahead of print)
- Marquis, D. A., R. L. Ernst, and S. L. Stout. 1992. Prescribing silvicultural treatments in hardwood stands of the Alleghenies. United States Forest Service, Northeast Forest Experiment Station, Radnor, Pennsylvania, USA. General Technical Report NE-96. 108 pp.
- Mathiason, C. K., S. A. Hays, J. Powers, J. Hayes-Klug, J. Langenberg, S. J. Dahmes, D. A. Osborn, K. V. Miller, R. J. Warren, G. L. Mason, and E. A. Hoover. 2009. Infectious prions in pre-clinical deer and transmission of chronic wasting disease solely by environmental exposure. *PLoS ONE* 4(6): e5916. doi:10.1371/journal.pone.0005916.
- Mathiason C.K., J.G. Powers, S.J. Dahmes, et al. 2006. Infectious prions in the saliva and blood of deer with chronic wasting disease. *Science* 314: 133-136.
- McDonald, J. E., Jr., M. R. Ellingwood, and G. M. Vecellio. 1998. Case studies in controlled deer hunting. New Hampshire Fish and Game Department, Durham, NH.
- McGraw, J. B., and M. A. Furedi. 2005. Deer browsing and population viability of a forest understory plant. *Science* 307: 920-922.
- McShea, W. J., and J. H. Rappole. 2000. Managing the abundance and diversity of breeding bird populations through manipulation of deer populations. *Conservation Biology* 14:1161-1170.
- McShea, W. J., H. B. Underwood, and J. H. Rappole, editors. 1997. *The science of overabundance: deer ecology and population management*. Smithsonian Institution Press, Washington D.C., USA.
- Miller, K.V., and J.M. Wentworth. 2000. Carrying capacity. Pages 140-155 in S. Demarais and P. R. Krausman, editors. *Ecology and management of large mammals in North America*. Prentice-Hall, Upper Saddle River, New Jersey.
- Miller, M.W. 2008. Prions in the wild: epidemiology and ecology of chronic wasting disease. *Proceedings of the Western Deer and Elk Workshop* 7(in press).
- Miller, M.W., and E.S. Williams. 2003. Horizontal prion transmission in mule deer. *Nature* 425:35-36.
- Miller, M.W., E.S. Williams, N.T. Hobbs, and L.L. Wolfe. 2004. Environmental sources of prion transmission in mule deer. *Emerging Infectious Diseases* 6:1003-1006.
- Mitchell, J. M., G. J. Pagac, and G. R. Parker. 1997. Informed consent: gaining support for removal of overabundant white-tailed deer on an Indiana state park. *Wildlife Society Bulletin* 25:447-450.
- Mysterud, A., T. Coulson, and N. C. Stenseth. 2002. The role of males in the dynamics of ungulate populations. *Journal of Animal Ecology* 71:907-915.
- Ozoga, J. J., and L. J. Verme. 1982. Physical and reproductive characteristics of a supplementally-fed white-tailed deer herd. *Journal of Wildlife Management* 46:281-292.
- Perkins, S. E., I. M. Cattadori, V. Tagliapietra, A. P. Rizzoli, and P. J. Hudson. 2006. Localized deer absence leads to tick amplification. *Ecology* 87:1981-1986.
- Potvin, F., and L. Breton. 2005. From the field: testing two aerial survey techniques on deer in fenced enclosures: visual double-counts and thermal infrared sensing. *Wildlife Society Bulletin* 33:317-325.
- Potvin, F., L. Breton, and L. P. Rivest. 2004. Aerial surveys for white-tailed deer with the double-count technique in Quebec: two 5-year plans completed. *Wildlife Society Bulletin* 32:1099-1107.
- Rand, P. W., C. Lubelczyk, M. S. Holman, E. H. Lacombe, and R. P. Smith. 2004. Abundance of *Ixodes scapularis* (Acari: Ixodidae) after the complete removal of deer from an isolated offshore island, endemic for Lyme disease. *Journal of Medical Entomology* 41:779-784.
- Rand, P. W., C. Lubelczyk, G. R. Lavigne, S. Elias, M. S. Holman, E. H. Lacombe, and R. P. Smith. 2003. Deer density and the abundance of *Ixodes scapularis* (Acari: Ixodidae). *Journal of Medical Entomology* 40:179-184.
- Race, B., K. Meade-White, R. Race, and B. Chesebro. 2009. Prion infectivity in fat of deer with chronic wasting disease. *Journal of Virology* doi:10.1128/JVI.01127-09 (online ahead of print)
- Roed, K. H., O. Holand, A. Mysterud, A. Tverdal, J. Kumpula, and M. Nieminen. 2007. Male phenotypic quality influences offspring sex ratio in a polygynous ungulate. *Proceedings of the Royal Society B: Biological Sciences* 274:727-733.
- Safar, J. G., P. Lessard, G. Tamguney, Y. Freyman, C. Deering, F. Letessier, S. J. DeArmond, and S. B. Prusiner. 2008. Transmission and detection of prions in feces. *Journal of Infectious Diseases* 198:81-89.

Schwabe, K. A., and P. W. Schuhmann. 2002. Deer-vehicle collisions and deer values: an analysis of competing literatures. *Wildlife Society Bulletin* 30:609–615.

Seamans, R. A. 1946. The time is now: a pictorial story of Vermont's deer herd. Federal Aid in Wildlife Restoration Project No. 1-R, Vermont Fish and Game Service, Montpelier, VT, USA.

Seeger, H., M. Heikenwalder, N. Zeller, J. Kranich, P. Schwarz, A. Gaspert, B. Seifert, G. Miele, and A. Aguzzi. 2005. Coincident scrapie infection and nephritis lead to urinary prion excretion. *Science* 310:324–326.

Sigurdson, C. J. 2008. A prion disease of cervids: chronic wasting disease. *Veterinary Research* 39:41 doi:10.1051/vetres:2008018.

Sigurdson, C.J., and A. Aguzzi. 2007. Chronic wasting disease. *Molecular Basis of Disease* 1772:610-618.

Sorin, A. B. 2004. Paternity assignment for white-tailed deer (*Odocoileus virginianus*): mating across age classes and multiple paternity. *Journal of Mammalogy* 85:356–362.

Strickland, B. K., and S. Demarais. 2008. Influence of land composition and structure on antler size of white-tailed deer. *Journal of Wildlife Management* 72:1101–1108.

Strickland, B. , S. Demarais, L.E. Castle, J.W. Lipe, W.H. Lunceford, H.A. Jacobson, D. Frels, K.V. Miller. 2001. Effects of selective-harvest strategies on white-tailed deer antler size. *Wildlife Society Bulletin* 29:509–520.

Tilghman, N. G. 1989. Impacts of white-tailed deer on forest regeneration in northwestern Pennsylvania. *Journal of Wildlife Management* 53:524–532.

U.S. Department of the Interior, Fish and Wildlife Service and U.S. Department of Commerce, U.S. Census Bureau. 2006. National survey of fishing, hunting, and wildlife-associated recreation.

VFWD. 2004. Comprehensive deer management survey results and analysis. Vermont Fish & Wildlife Department, Waterbury.

VFWD. 2009. Wildlife habitat management for Vermont woodlands – a landowners' guide. Fourth edition. Vermont Fish and Wildlife Department, Waterbury. Available on the Department web site, www.vtfishandwildlife.com, by summer 2010.

Williams, E.S., M.W. Miller, T.J. Kreeger, R.H. Kahn, and E.T. Thorne. 2002. Chronic wasting disease of deer and elk: a review with recommendations for management. *Journal of Wildlife Management* 66:551-563.

Williamson, S. 2003. White-tailed deer harvest management and goal setting in the Northeast. The Wildlife Management Institute. Washington, D.C.

Wilson, M. L., A. M. Ducey, T. S. Litwin, T. A. Gavin, and S. Spielman. 1990. Microgeographic distribution of immature *Ixodes dammini* ticks correlated with deer. *Medical and Veterinary Entomology* 4:151–159.



MOOSE

I. Management History

In the 1700s when New England was beginning to be settled, the Vermont landscape was 95% forest. As forest dwellers, abundant moose populations roamed freely. Early town records and explorers' accounts indicate that the animal was widely distributed throughout Vermont. The French Canadians and Abenaki Indians who raided Deerfield, Massachusetts, in 1704 cached meat from 20 moose at a site on the Connecticut River near Brattleboro to provide food for their return march home to Canada (Williams 1707). An Abenaki hunter who lived near Crystal Lake in Barton also told of killing 27 moose and many beaver in that vicinity in the winter of 1783-1784 (Collins 1903).

Native Americans and European colonists killed moose opportunistically throughout the year for food. As Vermont's population grew the unregulated hunting of moose played a part in their disappearance from the state by the nineteenth century. Probably of far greater importance, however, was the loss of moose habitat when the native forests were converted to agricultural lands. This land conversion (forest into fields) began in about 1800 and peaked by 1880 after which only 37% of Vermont remained forested. By the late nineteenth century Vermont's remaining woodlands were concentrated along the higher elevations of the Green Mountains and in Essex County. Moose had become so rare that when a young bull was shot in March 1899, at Wenlock (now Ferdinand) in Essex County, newspaper reports called it "a strange animal" and "the last moose in Vermont."

During the twentieth century, hill farms went out of business on a vast scale. Forests gradually covered hard-won fields, and moose began to reappear in Vermont. By the 1960s, 25 moose were thought to exist in Essex County. By 1980, forests covered 80% of the land area of the state, and moose numbers had increased to a point where they were regularly seen in Essex County. Moose were also observed in neighboring counties. The absence of predation on moose by mountain lions and wolves, as well as by humans allowed rapid population growth. By 1990,



WAYNE LAROCHE

moose were abundant enough to support a limited, regulated hunt. The size and age structure of the moose population approximated populations in areas of North America where regulated hunting was routine.

Modern moose management began in Vermont in 1992 with the adoption of the state's first plan that used biological data derived from studies conducted in the state and the results from studies conducted on moose in nearby states and provinces of Canada. Public opinion was solicited via a series of public meetings held throughout the state during 1991 and 1992.

Vermont's first modern moose season was a three-day hunt held in 1993 in wildlife management unit (WMU) E in which 30 permits were issued and 25 moose were taken. In 1995 the season was expanded to include a second area, D2, and the season was lengthened to four days including a weekend. WMU E was subdivided into two parts prior to the 1996 season in order to distribute the moose harvest more uniformly across this area. In the new units E1 and E2, some of the hunters were issued antlerless-only licenses in order to achieve an equal adult sex ratio in the harvest and to take cows to stabilize the size of the herd by reducing the number of young moose entering the population in those WMUs. Antlerless-only permits have been issued in these units every year since 1996. Four additional units were opened to moose hunting in 1997.

No changes were made in the 1998 moose season because the Department was in the midst of drafting a new ten-year Moose Management Plan. Public comment concerning the new plan was obtained via mail and telephone surveys, open houses, public meetings, and written comments. To expand public benefits, the final plan called for further expansion of the area open for moose hunting whenever appropriate. Continued growth of the moose herd has resulted in expansion of moose hunting into a total of 17 WMUs, with 78% of the state open to regulated moose hunting.

By the early 2000s, the moose population in WMU E was causing significant damage to forest regeneration. Estimated moose densities were nearly double the target levels set in 1996, yielding population densities of about 1.75 moose per square mile.

Moose densities well over 3 per square mile in WMU E were overbrowsing forest regeneration, not only to their own detriment, but also to the detriment of other wildlife species that utilize low growing trees and shrubs for food and cover. Landowners, especially large industrial forestland owners whose livelihood and investment depends on a healthy and growing forest, were especially anxious to see moose densities reduced.

Large increases in permit numbers issued in units E and D2 were prescribed for the 2004 season (Table 3.1) in an attempt to move toward the goal of returning the moose density in these areas to their 1996 and 1999 levels, respectively. By this time, moose had approached the biological carrying capacity of the habitat.

Today, moose hunting in Vermont is regulated by a special license that limits the permit holder to a specific WMU. A moose harvest objective is determined each year for each WMU, and a specific

Table 3.1 Vermont Moose Season Results 1993 - 2008

YEAR	PERMITS ISSUED	MOOSE HARVESTED	% HUNTER SUCCESS	UNITS OPEN
1993 ¹	30	25	83	E
1994	40	28	70	E
1995 ²	75	61	81	D2, E
1996 ³	100	78	78	D2, E1, E2
1997	165	100	61	Above plus C, D1, H1 & H2
1998	165	97	59	
1999	200	120	60	Above plus G, I & J1
2000	215	137	64	
2001	229	155	68	
2002	365	221	61	Above plus B, J2, L, M1 & P
2003 ⁴	440	298	68	Above plus O1
2004	833	539	65	Above plus Q
2005 ⁵	1,046	640	61	
2006	1,115	648	58	
2007 ⁶	1,251	592	47	Above plus M2
2008	1,251	605	48	
Totals	7,520	4,344	58	

¹ 3-day, mid-week season.

² Season lengthened to 4 days and opening day moved to Saturday.

³ Antlerless-only permits issued for the first time. WMU E split into subunits E1 and E2.

⁴ Season lengthened to 6 days.

⁵ Season split into two 6-day periods; antlerless permit holders in D2, E1 & E2 hunt 2nd week.

⁶ Second season lengthened to 9 days.

number of licenses are issued to achieve target harvests. The license allows a party of up to two hunters, and an optional guide, to take a single moose during a season held in mid- to late-October. Hunters are selected by random draw from a large pool of applicants who apply prior to the license drawing. Licenses are either-sex or limited to cows and calves as necessary to achieve area-specific population goals (Table 3.1)



GINNY GARRISON

1998-2007 Plan Accomplishments

Vermont's second moose management plan was adopted in 1998 and incorporated biological data on the herd gathered between 1980 and 1997 along with public input. The results of the public input revealed that Vermonters generally wanted to see more moose statewide while wanting to stabilize populations in the Essex County area. Vermonters desired the benefits of a healthy moose population, but they also expressed serious concerns regarding moose/vehicle collisions and the upward trend in human conflicts with moose.

The ten-year moose management plan adopted in 1998 was designed to address the interests and concerns of the public and strive for a healthy, expanding moose population in balance with its habitat. The objectives and accomplishments of that plan are summarized below.

► **Objective 1. To maintain a healthy, viable moose population in Vermont.**

Vermont's statewide moose population was estimated at 2,100 animals in 1997. This objective included six strategies:

Strategy 1.1 *Maintain a minimum fall population of at least 500 moose.*

❖ **Action:** The Department estimated that the state-wide moose population following the 2007 moose hunt was about 4,000 animals. This number more than met the minimum objective of 500, but in the Northeast Kingdom region of the state the moose population grew at a rate that was unsustainable ecologically.

Strategy 1.2 *Maintain an adult sex ratio of 40 – 60 bulls per 100 adults.*

❖ **Action:** Harvest and mortality reports provide the information on the sex ratio

and age structure of Vermont's moose population. This data suggested that the adult male to female sex ratio was very close to a normal, 50:50.

Strategy 1.3 *Maintain an adult age-class distribution of at least 25% greater than age four.*

❖ **Action:** The Department kept track of nonhunting, or "incidental," mortalities occurring within the moose's "biological year" to determine an age-class distribution. The biological year (BY) for moose begins June 1, at the time of the annual birth pulse of calves, and ends May 31 of the following year. Figure 3.1 illustrates the age structure of Vermont's moose population over a five year period. Although the graph has a normal shape or curve, the percentage of younger moose, age classes one through three, declined from 71% in the early 1990s to 58%. This is likely due to the decrease in reproductive rate noted earlier resulting from increased moose density. Forty-two percent of these moose were more than four years of age.

❖ **Strategy 1.4** *Continue to monitor various biological indices, such as carcass weight, beam diameter, ovulation rate, and occurrence of parasites.*

❖ **Action:** The Department monitored the health of the moose herd throughout the state and found the physical condition of the animals was deteriorating. Biologists found that over the previous nine years Vermont's cow moose ovulation rates had dropped dramatically (Fig. 3.2). Other indicators of the moose herd's health were the decline in the dressed carcass weight of yearling bulls and the smaller beam diameter for yearlings (Fig 3.3 and 3.4). These trends strongly indicated that the moose herd was exceeding its BCC in some parts of the state, most notably in WMUs E1 and E2. Biologists also watched for diseases and health related issues caused from two common parasites, the winter tick and the roundworm. There were, however, no apparent health effects from either of these parasites during this period.




Figure 3.1 VT moose ages for legal and non-hunting mortalities for calendar years 2003 –2007.

1998-2007 Plan Accomplishments (continued)




Figure 3.2 Comparison of ovulation rates for legally harvested cows from three time periods




Figure 3.3 Yearling male carcass weight from Vermont moose harvests.




Figure 3.4 Yearling beam diameter from Vermont moose harvests.

Strategy 1.5 Develop a model to assess relative moose habitat suitability at the WMU or regional level throughout the state.

❖ **Action:** The Department assisted a graduate student at the University of Vermont (UVM) who modified a moose habitat suitability index (HSI) model that was used to evaluate moose habitat in WMUs E and I (Koitzsch 2000). The HSI values (1.0 equals ideal habitat) were estimated to be 0.64 and 0.34 for WMUs E and I, respectively. The Department expanded use of this model and sought funding and partnerships with research institutions to perform this work.

Strategy 1.6 Consider implementing field studies to investigate and monitor moose browsing in selected WMUs.

❖ **Action:** With the assistance of staff biologists, a UVM graduate student investigated the incidence of moose bark stripping on mountain ash throughout the state (Scharf and Hirth 2000). This study found that one third of mountain ash trees in northern regions were wounded by moose bark stripping. Also, forest inventories conducted on 85,000 acres of private timber lands in Essex County indicated 25% of the plots were browsed, 68% of which were heavily browsed.

➤ **Objective 2.** To provide for the controlled growth of Vermont's statewide moose population in all WMUs except for the Northeast Kingdom region where population stabilization is desired.

Strategy 2.1 Continue to utilize annual, mid-October, regulated moose hunts to stabilize the moose population in WMUs E1, E2, and D2 at 1996 levels.

❖ **Action:** The 1998 moose plan called for stabilization of the moose populations in the

1998-2007 Plan Accomplishments (continued)

Northeast Kingdom region. Density goals were about 1,000 moose in WMU E and 400 in WMU D2. Improved moose estimation methods revealed that the number of permits was still too few to achieve the objectives. Data gathered by the Department produced estimates of moose densities above the goals and continued evidence that moose were overbrowsing their range in the Northeast Kingdom.

Permit numbers for this region continued to increase during the plan period, partly in response to declining hunter success rate which made it necessary to issue more permits in order to meet harvest objectives, and partly in response to the need to expedite population reductions to protect forest habitat.

Strategy 2.2 *Continue to utilize regulated moose hunts to slow rate of growth of the moose population in WMUs C, D1, H1 and H2.*

❖ **Action:** Moose hunting first occurred in WMUs C, D1, H1, and H2 in 1997. The combined population estimate for these units had remained relatively stable since 2001 at about 700 moose.

Strategy 2.3 *Utilize regulated moose hunts to slow rate-of-growth of the moose population in WMUs G, J1, and I beginning in 1999.*

❖ **Action:** Moose hunting was initiated in WMUs G, J1, and I in 1999. The estimated moose population for these units increased from 290 (2001) to 370 (2004) and has since been successfully reduced to an estimated current population of 300 moose.

► **Objective 3. To maximize benefits from Vermont's moose population within acceptable social and biological limits.**

The Department continued to work toward balancing an abundant moose population and sustainable habitat with protection of the forest and prevention of conflicts with humans. During this planning period, the Department employed several strategies. These strategies involved regulating hunting, working with landowners to open access to hunting, and promoting habitat management through public outreach, education, and activities.

Strategy 3.1 *Continue with annual moose hunts in WMUs C, D1, D2, E1, E2, H1, and H2*

Strategy 3.2 *Open WMUs G, J1, and I to limited hunting beginning in 1999.*

Strategy 3.3 *Annually evaluate the potential for regulated moose hunting opportunities in other WMUs.*

❖ **Action:** All three of these strategies from the 1998 moose plan were implemented through regulation.

Strategy 3.4 *Coordinate with large property owners to find ways to enhance moose hunter access.*

❖ **Action:** The Department worked with large industrial forest landowners in the Northeast Kingdom to facilitate the opening of gates during the moose season and with the Vermont Horse Council and the Vermont Department of Forests, Parks & Recreation to facilitate the use of draft horses to haul moose carcasses out of roadless areas.

Strategy 3.5 *Promote the "Hunters Sharing the Harvest" program to moose hunters as a*

way of providing moose meat to needy households throughout Vermont.

❖ **Action:** The Department annually provided a 50-page guidebook to each moose hunting permit holder that included a description of the "Hunters Sharing the Harvest Program" and listed some examples of local food shelves that could store and distribute moose venison to their patrons. (There is currently no organized program or system to track donations.)

Strategy 3.6 *Cooperate with natural resource professionals and landowner organizations in dissemination of moose habitat management guidelines.*

❖ **Action:** In 1995, the Department, in cooperation with the Vermont Department of Forests, Parks & Recreation, published a booklet entitled: "A Landowner's Guide - Wildlife Habitat Management for Vermont Woodlands" (Regan and Anderson 1995). This publication, which includes a chapter on moose habitat, was made available to state biologists, private consulting foresters, and landowners through forest management workshops.

Strategy 3.7 *Develop and implement educational displays explaining Vermont's moose management for use at fairs, outdoor shows, and moose check stations.*

❖ **Action:** In 2001 the Department produced five sets of a seven-panel poster-board display covering many aspects of moose life history and management. These sets have since been used annually at moose weighing stations and in other outreach venues.

Strategy 3.8 *Construct at least two moose observation towers*

1998-2007 Plan Accomplishments (continued)

with parking areas near state highways in the Northeast Kingdom region, contingent on funding partnerships with the private business sector, regional chambers of commerce, and/or governmental tourism agencies.

❖ **Action:** The Department began work in 2006 on siting and designing a moose viewing tower off State Highway 105 in the Essex County town of Ferdinand.

Strategy 3.9 *Cooperate with a private interest in the publication of a "Vermont Moose Watcher's Guide."*

❖ **Action:** A professional wildlife photographer and author from Maine published the "Moose Watchers Handbook" in 2001, which included directions to popular moose viewing sites in Maine, New Hampshire, and Vermont (Silliker Jr. 2001).

► **Objective 4. To minimize negative interactions between humans and moose.**

Strategy 4.1 *Utilize annual limited-entry moose hunts to either stabilize or slow the growth rate of regional moose populations as noted above under Objective 2.*

❖ **Action:** The number of nonhunting moose mortalities steadily increased through the early part of the past ten years. Nonhunting moose mortalities during the last several years have seemed to decrease in the face of increased numbers of permits. Many of these mortalities (41%) occurred in the Northeast Kingdom units of D2, E1, and E2.

Strategy 4.2 *Develop and implement a policy for Department response to "nuisance" moose by 2000.*

❖ **Action:** To address damage caused by moose to livestock fencing, maple sap tubing, and

Christmas tree plantations, a Commissioner's rule was enacted in 1996 that under certain conditions allows a landowner suffering property damage to shoot the moose. To try to avoid this situation, the Department assisted the United States Department of Agriculture, Wildlife Services office in Berlin, Vermont, in developing an informational brochure describing possible ways to curb moose damage.

The Department also developed a protocol for sick or diseased moose that posed a potential hazard to public safety. These situations can arise when sick moose wander into urban areas, farmyards, or busy highways. A Department protocol for dealing with all "nuisance" moose still needs to be completed in the next planning period.

Strategy 4.3 *Continue to cooperate with the Vermont Agency of Transportation (VTrans) to erect warning signs at traditional moose highway crossings.*

Strategy 4.4 *Cooperate with the VTrans in implementing at least three roadside brush-clearing projects to improve visibility at the most dangerous moose crossings, where feasible.*

❖ **Action:** The Department worked with VTrans to evaluate several methods of reducing moose/vehicle collisions. VTrans considered the advice of the Department for the placement of moose crossing signs and the clearing of roadside brush adjacent to frequently used road-side salt licks in order to enhance the ability of approaching motorists to detect moose

Strategy 4.5 *Continue with annual press releases to remind motorists of moose hazards and explore potential for including a warning message with helpful driving tips concerning deer and moose collisions in the Department of Motor Vehicle's Driver's Manual and in all new vehicle registrations or renewals.*

❖ **Action:** The Department issued biannual press releases to newspapers and broadcast media each year to advise motorists during times of the year when movement of moose poses the greatest hazard to motorists. The Department also partnered with the Vermont Frost Heaves PBA basketball team to raise driver awareness concerning the hazard of moose on highways.



TOM MERRIFIELD

II. 2010-2020 Moose Management Issues, Goals, and Strategies

The overall goal of moose management in Vermont is to manage Vermont's moose to sustain healthy, viable populations consistent with biological, social, and economic considerations, and provide maximum hunting opportunities.

ISSUE 1. Regional Population Goals

GOAL: To maintain regional populations of healthy moose at or below cultural carrying capacity.

The Department uses several approaches in estimating moose populations — surveys, mortality data, and aerial censuses. Two annual hunter surveys (one for deer hunters and one for moose hunters), annual moose hunter success rates, and nonhunting mortality records provide the basis for the Department's moose permit allocation recommendations. Observations and knowledge provided by state game wardens, foresters, biologists, and landowners are also considered when making decisions and recommendations.

Since 1999, Vermont has conducted deer and moose hunter surveys that provide a measure of relative moose density trends by WMU across the entire state. The deer hunter survey asks hunters to identify and record the number of bulls, cows, calves, or moose of unknown sex or age that are observed. The moose hunter survey requests hunters to report any preseason scouting activities. Hunters are asked to record the number of scouting trips they took; the number of hours they spent scouting; and the number of moose they saw during these trips. The numbers are standardized to determine the average number of moose sighted per hundred hours scouting.

Moose hunter success rate is calculated as the percentage of all permit holders that harvested and registered a moose. Success rates are calculated annually for each WMU that is open to hunting. The current year hunter success rate is compared to the previous year to assess changes at the WMU level, considering number and type of permits issued. Hunter success can be affected by individual hunters' effort (time spent afield), weather conditions during the hunt, moose behavior, population levels, and the accessibility of moose to hunters (for example, the distribution of roads and trails in moose habitat).

The moose sighting rate from deer hunter surveys in WMU E has declined during the past four years, thus, the estimated moose density has also declined. The population density estimate for November 2008 was 2.59 moose per square mile, an estimated 1,526 moose. With the current permit quota, the target density for WMU E should be achieved following the 2010 hunting season. Moose sighting rates for D2, after remaining fairly stable for several years, finally decreased in November 2008. Using a rolling three-year average for deer hunter survey moose sighting rate data, the moose density in D2 is currently estimated to be 1.16 moose per square mile. It is possible that by maintaining the current permit quota of 340 for one more year that the D2 population may closely approach the target density of one moose per square mile. Permit numbers have been steadily increased in these units from 30 in 1998 to 110 in 2009. The combined population estimate for these units has remained relatively stable since 2001 at about 700 moose.

The Department maintains a statewide database of all reported nonhunting moose deaths. Nonhunting mortality data is collected and reported on a biological year basis that begins on June 1, after most of the moose calves have been born, and ends on May 31. Summaries of nonhunting moose mortalities are prepared each year and assessed prior to development of season recommendations. This information also helps us assess changes in moose numbers through time.

New Hampshire conducts aerial censuses using forward-looking infrared (FLIR) cameras that enhance the ability to observe moose on the ground and has developed a model that provides more accurate estimates of moose populations. Although potential differences in topography, road access, hunter behavior, and other factors could influence moose sighting rates between northern New Hampshire and northeastern Vermont which could affect the applicability of this model in Vermont, the Department has found the model to be useful in estimating moose densities in the state. The Department is seeking to conduct its own aerial FLIR count of moose in Vermont to verify that the New Hampshire model provides accurate estimates in Vermont. Flights are scheduled for December 2009.

Moose hunting has expanded into several additional WMUs since 1999 as populations have grown large enough to sustain hunting (see Table 3.1, page 41). As moose have become more abundant, public

attitudes toward the moose herd have changed over the nine-year period (1996-2007) as well. Results from the 2007 statewide telephone survey indicated, with some regional variations, that overall Vermonters (54%) want to see the moose population remain the same, 19% want to see it increased, and 10% want to see it decreased. Analyses of the data, with the 17% “don’t know/no opinion” responses removed, is shown in Table 3.2.

Table 3.2 Public opinion on desired regional moose population size by region of residence, in percent (sample size in parenthesis).

Region*		Decrease	Remain the Same	Increase
Northeast Kingdom	(99)	31 (31)	54 (53)	15 (15)
Greater Chittenden	(268)	9 (24)	69 (184)	22 (60)
Central Vermont	(243)	11 (26)	64 (156)	25 (61)
Southern Vermont	(246)	10 (25)	66 (161)	24 (60)

*Northeast Kingdom: Caledonia, Essex and Orleans Counties

Greater Chittenden: Franklin, Chittenden and Grand Isle Counties

Central Vermont: Addison, Lamoille, Washington and Orange Counties

Southern Vermont: Rutland, Bennington, Windsor and Windham Counties

With this public feedback in mind, the Department proposes to maintain regional moose numbers at their current levels in most areas of the state, with the exception of the Northeast Kingdom region, where moose numbers need to be reduced to a level below biological carrying capacity, and in a few WMUs where an increase in moose populations may be acceptable (WMUs I, L, P, and Q, and perhaps others). The Department will solicit more public input on this issue prior to setting final objectives on moose herd numbers for WMUs. Web-based questionnaires will be used early in this management plan cycle to solicit public input.

Based on November 2008 population estimates for each WMU (Fig. 3.5.), the Department will make adjustments in two units. The Department proposes continuing with a population target of 1,000 moose in WMU E (1.75 moose per square mile), but to readjust the target for WMU D2 from 400 to 600 moose. This new objective for WMU D2 equates to one moose per square mile, which should be well




Fig. 3.5 Estimated moose population by WMU from sighting rates of 2006 -2008 November deer seasons.

below biological carrying capacity. This change is proposed because areas of current overbrowsing in D2 are limited, whereas historically higher moose densities (1.4 moose per square mile, 800 total) created overbrowsing. With the growing importance of moose hunting in this region, 600 moose may be an acceptable population level to area residents.

Management Strategies

- 1.1 Maintain a statewide fall post-hunt population of between 3,000 and 5,000 moose.
- 1.2 Maintain a sex ratio of between 40 to 50 bulls per 100 adults (moose of at least age-class one).
- 1.3 Maintain an adult age-class distribution of at least 25% of at least age-class four.
- 1.4 Maintain an average ovulation rate of more than 1.15 for cows age class of at least three.
- 1.5 Assess relative moose habitat condition of individual WMUs or regions of the state using forest inventory data and a GIS-based Habitat Suitability Index Model.
- 1.6 Reduce and maintain WMU E moose densities to 1.75 moose per square mile (approximately 1,000 moose post-hunt).
- 1.7 Reduce and maintain WMU D2 moose densities to 1.0 moose per square mile (approximately 600 moose post-hunt).
- 1.8 Allow slow population growth in WMUs I, L, P and Q while not exceeding one moose per square mile.
- 1.9 Stabilize moose population in other WMUs at current levels.

ISSUE 2. Moose / Human Conflicts

GOAL: To minimize motor vehicle/moose collisions and other forms of damage caused by moose.

As the moose population has expanded, so have the negative interactions with humans. Damage to fences and maple sugaring equipment are common problems. More and more moose are finding their way into developed neighborhoods or becoming habituated to humans. Both situations are rarely resolved without significant public disturbance and usually result with the moose's demise.

Vehicle collisions are the most serious human/moose encounters. Although deer collisions are far more common and often result in costly damage to vehicles, they rarely result in serious human injury. Moose collisions, on the other hand, often result in serious human injury or even death. The Department is continually looking for ways to reduce the number of motor vehicle collisions with moose. Currently, there are approximately 77 signed crossing areas statewide. Many of these signs carry a 40 mph speed advisory per the Department's recommendation.

The number of nonhunting moose mortalities steadily increased through the early part of this decade. Nonhunting moose mortalities during the last several years have decreased slightly (Fig. 3.6) with increased numbers of permits. Many of these mortalities (41%) occurred in the Northeast Kingdom units of D2, E1, and E2.

The Department began drafting a protocol for dealing with moose that are not sick but pose a

threat to public safety. When finalized, this protocol will conform to the Department's umbrella policy governing how it handles "nuisance" or "hazardous" wildlife in general (Regan 1998). Under the umbrella policy, humane treatment of animals is an important consideration. Euthanasia is recognized as sometimes being the only cost-effective and practical response.

Management Strategies

- 2.1 Develop and implement a policy for Department response to "nuisance" moose.
- 2.2 Continue to cooperate with the Vermont Agency of Transportation (VTRANS) to erect warning signs at traditional moose highway crossings.
- 2.3 Cooperate with VTRANS in implementing roadside brush-clearing projects to improve visibility at the most dangerous moose crossings, when feasible.
- 2.4 Cooperate with VTRANS to investigate the use of new technology that may help reduce moose/vehicle collisions.
- 2.5 Continue with annual press releases to remind motorists of moose hazards during seasons of increased moose movements.

ISSUE 3. Moose Hunting Opportunities

GOAL: To maximize quality moose hunting opportunity.

HUNTING SATISFACTION

Feedback the Department receives from various sources indicates a favorable satisfaction rate from moose hunters on the present structure and timing of hunting seasons. No major changes are being proposed in the current new plan.

Management Strategies

- 3.1 Provide quality moose hunting opportunity in all WMUs where feasible.
- 3.2 Coordinate with large property owners to enhance moose hunter access.
- 3.3 Provide information to hunters on how they can share moose meat with needy households throughout Vermont.
- 3.4 Conduct outreach efforts prior to any significant reduction in total permit numbers made in response to moose population changes.




Fig. 3.6 VT non-hunting moose mortalities for biological years 1980 – 2007.

3.5 Provide public opportunity to harvest moose for food and other utilitarian purposes.

MOOSE PERMIT LOTTERY

An average of 10,448 Vermonters have applied for a moose permit each year for the last five years. Beginning with the 2007 permit lottery, hunters who had applied the previous year but had not won a permit were awarded an extra chance, or “bonus point.” Unsuccessful applicants now accumulate a bonus point for each year they apply and fail to draw a permit. Each “point” adds another occurrence of their name into the lottery pool of applicant names increasing their odds of winning a permit. This recent change has helped satisfy those who have applied for many years without success and seems to have been favorably received by moose hunters. No further changes to the permitting process are currently being contemplated.

Vermont has issued a relatively high number of permits in recent years to reduce the moose population in the Northeast Kingdom. For the 2009 season, 1,230 permits were proposed statewide with 940 allocated to WMUs D2, E1, and E2 alone. Once population goals are reached in the Northeast Kingdom, the number of permits issued may be reduced.

Management Strategies

3.6 Maintain and improve hunter satisfaction by managing a preference point lottery system.

SPECIAL ARCHERY SEASON

Bow-hunting enthusiasts have encouraged the Department and the Fish and Wildlife Board to consider a special archery-only season for moose. Although bows can be used in the current moose season, some archers feel they might have more success in calling moose into close range if they were able to hunt during the peak of the rut and without competition from more mobile firearm hunters. Because of this interest, the Department included the following question in the 2007 telephone survey: “Currently, moose may be harvested during the season with rifles, handguns, muzzleloaders, bows, or shotguns. Do you support or oppose establishing an archery-only season for moose in Vermont in addition to the regular moose hunting season?”

This question was asked only of survey participants who were hunters. Of 252 respondents, 50% were opposed (39% strongly opposed and 11%



moderately opposed) and 39% were supportive (23% strongly and 16% moderately). Four percent neither supported nor opposed the idea while 7% answered “Don’t know.” The 39% of responding hunters corresponds closely with the proportion of Vermont hunters who bow hunt for deer, so it seems likely that most opponents are not archery hunters. Most of the respondents opposing an archery moose season were probably concerned that their chances of winning a moose permit in the regular season lottery would diminish. A similar opposition was expressed prior to the initial deer archery season in Vermont. Subsequently, many rifle hunters took up archery hunting, and the deer archery season became widely accepted. Archery deer season has subsequently added a significant recreational opportunity for Vermont’s deer hunters.

In reality, a limited archery season would have minimal impact on chances for a regular moose season hunter to win a lottery permit because permit numbers are based upon harvest objectives and the success rate of hunters. Archers are expected to have a lower success rate and would be expected to take a small portion of the target moose harvest.

The Fish and Wildlife Board received several petitions in the spring of 2008 for the establishment of a special archery season for moose. Consequently, the Department will propose a Board regulation to establish a short moose archery season, potentially beginning the first Saturday in October. The season might run for nine days with perhaps 50 permits issued via a lottery. Success rates will likely be less than 30%, so the archery moose harvest would be expected to take less than 20 moose statewide. This small harvest would have minimal biological impact on the moose population even if it was in addition to the regular permits set by harvest objective.

Management Strategies

3.7 Propose to implement a limited special archery-only moose hunting opportunity.

ISSUE 4. Moose Viewing

GOAL: Provide safe and quality moose viewing opportunity.

A public opinion survey found that nearly 57% of Vermont residents participated in viewing or photographing wildlife (Duda and Young 1996). White-tailed deer are the most viewed and photographed (89% of respondents). Due to their large size, interesting features, and historical scarcity, viewing moose remains a special thrill for most Vermonters. Moose can often be easily observed and photographed from vehicles while feeding along roadside salt licks or shallow wetlands. People frequently make special trips to the Northeast Kingdom and other areas to observe moose thereby contributing to the economy in rural areas of the state.

The Department answers many inquiries each year concerning when and where to observe moose. Efforts are underway to place a moose viewing tower at a favorite viewing spot east of Island Pond. This project should continue to move forward with completion expected by 2010.

Management Strategies

- 4.1 Construct at least one moose observation tower with a parking area near a state highway in the Northeast Kingdom region and investigate other locations in other regions.
- 4.2 Include moose in a guide to wildlife viewing sites on the Department's website.

ISSUE 5. Moose Habitat

GOAL: Maintain necessary habitat to support 3,000 to 5,000 moose on a sustained basis.

The moose is a northern forest species and uses different habitats during various seasons of the year. In general, moose prefer thick, brushy habitat for concealment and as sources of abundant food. Lowland softwood forests, beaver ponds, and other shallow bodies of water are favorite spring and summer habitats for moose. During the hot summer months, moose can suffer from overheating and must have access to dense shade or water for cooling. Moose also use ponds to escape biting insects and

predators. Moose frequent upland hardwood or mixed forests during the fall and winter. Younger age classes of these forest types provide abundant browse, especially in recently cutover areas. Managing habitats specifically for moose is difficult because this species has a large home range (4 to 10 square miles).

Moose are not as social as deer. Although it is not uncommon to encounter several moose together during the post-rut period, by late winter moose are usually either solitary or found in groups of two or three animals. These small individual groups of moose may each seek out middle-aged to mature softwood stands where they can escape deep snows and severe winter weather.

Moose habitat management is typically a by-product of areas where commercial logging has occurred and produced abundant browse. Forested landscapes that are actively managed therefore contribute to productive moose range. Clearcutting more than 50% of a moose home range within a few years, however, can result in an unfavorable balance of forest age classes which may cause moose populations to decline (Girard and Joyal 1984).

While clearcuts may provide plenty of food, moose prefer to remain close to cover. Thus, there is relatively less browsing within the interior of larger clearcuts, particularly during the winter, than within areas closer to forest shelter. The browse within clearcuts of ten acres or less in size maximizes browse availability to moose. Special habitats that may be critical to moose survival or productivity include late-winter concentration areas, aquatic feeding areas, and salt licks.

Logging practices in Vermont over the past few decades have generally had a favorable impact on moose, especially in the Northeast Kingdom. Timber harvesting in this region increased significantly during the 1980s. Hardwood browse became abundant even in many of the former softwood stands (Moulton et al. 1984).

Many large private forestlands throughout the state are currently enrolled in Vermont's Use Value Appraisal program and/or are under working forest easements. These legal instruments mandate sustained timber harvesting, which benefit moose. Most of the larger state forests and wildlife management areas also have active timber harvesting and habitat management plans designed to sustain a diversity of habitat conditions. Thus, the quality of forested moose habitats in Vermont should remain good

for many years. Exceptions may occur on the “wilderness” designated areas of federal lands, such as the Green Mountain National Forest, which tend to minimize the early successional forests favored as forage for moose. When possible, the Department will advocate for active management to provide for all seral stages of forest vegetation and adequate amounts of early successional habitat to provide for moose and other wildlife species that favor younger forests. In isolated cases, loss of small areas of older softwood trees might be detrimental to wintering moose. In the past, the Department has been able to obtain cooperation from industrial forestland owners in reserving some of these important winter moose habitats from timber harvest.

Vermont also has a wetlands protection law that often affords protection of these important habitats. Thus, natural and roadside salt licks are not likely to disappear in the foreseeable future. Increasing human development, however, is likely to continue to slowly erode moose habitat in Vermont. More important than actual loss of acres of moose habitat will be increases in human/moose conflicts expected as residential development and road systems extend into moose habitat.

Private landowners who wish to consider moose habitat in their land management plans can receive habitat management recommendations from the Department of Fish and Wildlife. A booklet entitled “A Landowner’s Guide, Wildlife Habitat Management for Vermont Woodlands” is scheduled to be updated and reprinted in 2010.

Management Strategies

- 5.1 Implement field studies to investigate, measure, and monitor the degree of moose and deer browsing within selected WMUs.
- 5.2 Provide natural resource professionals and landowners with moose habitat management guidelines.



ISSUE 6. Deer-Moose Competition and Forest Impacts

GOAL: Balance the nutritional needs of regional moose and deer populations with the need for adequate forest regeneration.

White-tailed deer and moose play a significant role in the ecology of Vermont’s forests. As herbivores (plant eaters), seed dispersers, and prey, they can have a large impact on other plants and animals in forest systems. The presence of these animals has profound implications for the structure and function of forested ecosystems. If deer and moose were to disappear from the forest system, a wide variety of changes would ripple through the forest.

Deer and moose feeding habits are a significant influence on the ecology of the forest. Deer have been estimated to eat between four and ten pounds of plant matter each day while moose may eat more than 40 pounds per day. In winter, both species prefer the twigs of many hardwood and softwood trees. In summer, deer focus their feeding on a variety of green herbaceous plants while in the fall, fruits, nuts, and seeds make up an important part of their diet. In summer, moose continue feeding on hardwood and softwood trees but also eat succulent, sodium rich, aquatic vegetation in or near swamps, bogs, and wet forest edges. Browsing by deer and moose is a natural and desirable aspect of Vermont’s forest ecology, but too many deer and/or moose in a given area can cause problems for forests and people.

As the moose population has increased, the question of how to determine carrying capacity for both species separately and in combination has become a challenge. The Department needs to develop new ways to assess forest habitat and its capacity to support both moose and deer while maintaining a healthy native forest. There is also a need to monitor changes in the forest at various scales across the state and through time.

Management Strategies

- 6.1 Develop a study to assess the carrying capacity for moose and deer on Vermont’s forestland.
- 6.2 Develop a decision making process that assists managers in determining the appropriate mix of moose and deer densities for a given WMU based on cultural and ecological factors.

REFERENCES - MOOSE

- Collins, E.D. 1903. A history of Vermont. Ginn & Company, Publishers. The Athenaeum Press. Boston.
- Duda, M.D. and K.C. Young. 1996. Vermont resident's opinions and attitudes towards species management. Responsive Management, Inc. Harrisonburg, VA.
- Girard, F. and R. Joyal. 1984. L'effet des coupes a blanc mecanisees sur l'original dans le Nord-ouest due Quebec. *Alces* 20:3-25.
- Koitzsch, K. B. 2000. Application of a moose habitat suitability index model to wildlife management units in Vermont. M.S. Thesis, University of Vermont, Burlington.
- Moulton, W., J. Horton and J. St. Arnould. 1984. Northeast Kingdom regeneration cut survey. Vermont Department of Forests, Parks and Recreation. St. Johnsbury, VT.
- Regan, R. J. and G. Anderson, eds. 1995. A landowner's guide – wildlife habitat management for Vermont woodlands. Vermont Departments of Fish & Wildlife and Forests, Parks and Recreation. Waterbury, VT.
- Regan, R. J. 1998. Position paper on the handling of nuisance, orphaned, and sick or injured wildlife. Vermont Department of Fish & Wildlife. Waterbury, VT.
- Scharf, C.M. and D.H. Hirth. 2000. Impact of bark stripping on mountain ash in Vermont. *Alces* 36:41-52.
- Silliker, Bill Jr. 2001. Moose watchers guide. R.L. Lemke Corp. Islesboro, ME.
- Williams, J. 1707. The redeemed captive returning to Zion. Boston. As cited in John Demos, The unredeemed captive. Alfred A. Knopf, Inc. New York. 1994.



JOHN HALL

BLACK BEAR

I. Management History

The black bear is a native species in Vermont. It is the smallest of the three North American bear species, grizzly and polar bear comprising the other two, and the only one of the three found in the eastern United States.

To survive in Vermont, black bears require large tracts of forestland. As a result, historical accounts suggest that the state had a fairly abundant bear population when the first settlers arrived. The influx of settlers into Vermont significantly changed habitat conditions for bears. With their axes, the settlers literally cut their farms out of the forests and progressively whittled away the black bear's habitat – confining bears to those mountainous areas too steep or rocky to farm. It was then that Vermont's bear population reached its lowest point.

Loss of habitat was not the only reason for the decline of the bear population. The rapidly expanding human population used their fat, flesh, and hide to sustain themselves. Not being held in high esteem, bears were treated as vermin, readily associated with crop loss and livestock depredation. In 1831 the Vermont Legislature imposed a bounty on bears. Over the next 110 years, 1,295 bounty claims were paid out.

It was habitat change, however, not changes in the bounty laws that saved the Vermont black bear from extinction. Decades of farmers leaving the land following the Civil War led to a pattern of reforestation that provided great benefit to the bear population. Sentiment towards black bears began to change as well. Perhaps echoing the conservation views championed by President Teddy Roosevelt, Vermonters began to view bears and other wildlife as an important natural resource. In 1941 not only was the bounty on bears repealed, but they could only be hunted between June 1 and December 31 each year. Not insensitive to the potential bear damage farmers could incur, Vermont's Legislature obligated the Department, then known as the "Fish and Game Service," to reimburse persons for damages to "livestock." This is still the law.

Laws and regulations affecting the management of black bears during the twentieth century became more frequent as Vermont's human population



VICKI COMBS

continued to grow. Beginning in 1955 the reporting of harvested bears was required. In 1961, the season was shortened to the 91 days between September 1 and November 30. Other changes regulating the harvest of bears occurred over the next three decades, including prohibiting trapping (1967), limiting the harvest to one bear per season (1968), a prohibition on baiting and requiring bear houndsmen to hold a special permit (1972), and reducing the season length twice (1974 and again in 1990).

During this time of changing management and reforestation, the bear population has grown from an estimated 2,000 bears in 1975 to approximately 5,000 in 2008. Today bears are found in approximately 80% of Vermont from the Massachusetts border to Canada. Compared with their status 100 years ago, black bears are in a secure position. The greatest threat to the survival of black bears is in the form of fragmentation of their habitat (for example, roads and mountainside homes). This situation presents new management challenges for the twenty-first century. Vermonters have indicated they are satisfied with current population levels and wish to see them maintained during the next ten-year management period.

Ensuring the existence of a viable bear population and meeting public expectations for an abundant bear population while, at the same time, not having so many bears that they become a nuisance to agriculture and home owners will be the focus of the management actions contained in this plan.

1997-2006 Plan Accomplishments

- **Recommendation 1. *Revise black bear population objectives to reflect public interest in slightly increasing bear populations and repopulate suitable areas currently unoccupied by a breeding bear population.***

- **Strategy 1.1 *Analyze population data to determine current population levels and establish revised population objectives.***

❖ **Action:** The Department monitored growth of the state's black bear population. Population models indicated that Vermont's black bear population was relatively stable between 1985 and 1990 with about 3,000-3,400 bears existing in the state. Estimates indicate that the steady growth in the bear population occurred over the next ten years with about 4,800-5,200 bears existing by 2000.

- **Strategy 1.2 *Reduce black bear harvests by establishing a bear license or regional management zone.***

❖ **Action:** Reduced Vermont bear harvests from 1996 through 1998 resulted from a combination of widely distributed food supplies and the shortening of the length of the bear season beginning in 1990 that contributed to an increase in the statewide bear population. The plan's population goals were met without establishing a bear license or regional bear management zones. Another reason, however, that these actions were not taken was an increasing level of nuisance bear activity. As nuisance bear complaints increased, Department staff became concerned that a black bear license might reduce hunter participation to the point where harvests would

no longer be an effective bear management "tool." When a bear license was proposed, initial legislative language proposed a fee that the Department felt would discourage hunter participation. For these reasons, the Department abandoned efforts to establish a black bear license.

- **Recommendation 2. *Continue bear habitat conservation strategies such as Act 250, land acquisition, review of wood-to-energy harvest operations, and town and regional planning. The Department should pursue regulated logging and explore instituting a habitat stamp.***

- **Strategy 2.1 *Continue Department efforts on Act 250, land acquisition, review of wood-to-energy harvest operations, and town and regional planning.***

❖ **Action:** Between 1997 and 2006, Department staff reviewed 283 Act 250 projects that could potentially affect an estimated 1,000 acres of critical black bear habitat. As a result of subsequent revisions in these projects, a total of 12,621 acres of black bear habitat were protected during this ten-year period. The Department also published *Conserving Vermont's Natural Heritage*, a book to guide town planning for wildlife habitat, including black bear habitat. A new Department employee was assigned to work with town and regional planning agencies to guide conservation of wildlife habitat.

- **Strategy 2.2 *Pursue regulations on logging in critical bear habitat.***

❖ **Action:** The Department participated on the Heavy Cutting Committee that directed legislation on heavy cutting in Vermont. Department recommendations

to include critical bear habitat in this legislation were not incorporated into the law.

- **Strategy 2.3 *Investigate establishing a habitat stamp.***

❖ **Action:** Various funding "stamps" were discussed with a legislative committee but no action on a habitat stamp occurred.

- **Recommendation 3. *The Department will propose establishing a black bear license.***

❖ **Action:** As described in Recommendation 1, efforts to establish a black bear license were abandoned due to concerns over an increasing bear harvest, increased nuisance bear complaints, and potential for decrease in bear hunter participation. The concern was that this action might result in an increase in the bear population to a point where it exceeded the target population objective established by the plan.

- **Recommendation 4. *Regional management zones may be used to adjust bear harvests to meet higher population objectives.***

❖ **Action:** Regional management zones were also considered as a management tool to increase bear numbers in areas where suitable habitat remained unoccupied. Expansion of the bear population during the previous planning period eliminated the need to adopt management zones.

- **Recommendation 5. *No changes to season length or structure will be initiated until after it is determined if a black bear license will be established.***

❖ **Action:** Bear population goals were achieved without the implementation of a bear license, regional management

1997-2006 Plan Accomplishments

zones, or changes in the season structure. Changes in length of the season could be needed in the future to meet bear population objectives.

➤ **Recommendation 6. The Department will propose hunting hours for bears be changed to correspond to those for deer.**

❖ **Action:** These changes were established in state statute.

➤ **Recommendation 7. Work closely with the Vermont Bear Hound Association to discuss issues of concern.**

❖ **Action:** Department staff participated in many meetings with the Bear Hound Association to discuss bear issues, such as length of training season, nonresident dogs, procedures for addressing public perception/landowner conflicts, and public education. This

cooperative effort has resulted in regulatory changes in bear hound permits related to the ownership and residency requirements of dogs listed on permits. It has also led to successful dealings with negative human-bear interactions. The Department worked to modify state statutes related to black bear causing property damage.

➤ **Recommendation 8. The Department will establish a monitoring program on the sale of all bear parts through a mandatory tagging program.**

➤ **Strategy 8.1 Evaluate level and nature of sale of bear parts.**

❖ **Action:** The Department conducted a survey of successful bear hunters to determine the nature of using harvested bears and bear parts, including whether parts were being sold. Survey results

indicated that bear hunters fully utilized harvested bears. The sale of gall bladders and other parts was found to be insignificant and no threat to the sustainability of Vermont's bear population.

➤ **Strategy 8.2 Department will establish a monitoring program through mandatory tagging for the sale of bear parts.**

❖ **Action:** Results from the Vermont hunter survey indicated that a mandatory tagging program was not necessary to protect Vermont's bear population. It was determined that costs associated with mandatory tagging would not provide a cost-effective benefit in management of the already growing bear population. Department staff continued to monitor the global and national markets for bear parts.

II. 2010-2020 Black Bear Management Issues, Goals, and Strategies

ISSUE 1. Bear Population Size and Distribution

GOAL: Identify an appropriate bear population objective that ensures the viability of a wild, free-ranging bear population, provides for hunting opportunities, and satisfies human social expectations and tolerances for nuisance bear occurrences.

Black bears can be found throughout Vermont where preferred food and cover is located (Fig. 4.1). They are secretive animals that prefer to travel among forest and shrub habitat, usually only using fields and large forest openings at night or in low light. Normal bear behavior includes a strong avoidance of humans. Given these bear characteristics, the greatest bear population densities are found along the spine of the Green Mountains and in the Northeast Kingdom counties of Orleans,

Caledonia, and Essex. Because male and female bears lead separate lives, it is important to recognize the differences in the territorial ranges that each sex selects. Males are more solitary and tend to roam further in search of food and shelter. During the breeding season (June) older, more dominant males will search wider areas for receptive females. Females, on the other hand, tend to use smaller home ranges having high quality food sources and security for raising cubs.

Central to the management of a species is the need to accurately estimate the size of its population, the factors that influence growth and decline of the population, and the distribution of the population across the landscape. Based on this information, management goals can be met that satisfy the species' biological needs and human expectations.

Unlike other big game species, estimates of the bear population must be made using five-year averages. There are several reasons for this: bears live longer, they have a low reproductive rate, and harvests vary, depending on food supplies. Although the five-year averages do not pinpoint current bear populations,

they do reflect population trends very well up to the previous year. The data for making population estimates include all known bear mortalities (nonhunting and hunting) and include such factors as age, sex, and location of harvest.

Figure 4.2 illustrates the estimated average Vermont black bear population beginning with the five-year period 1983-1987. The graph shows two periods of population increases – the early 1990s and the late 1990s/early 2000s. The 2003 – 2007 estimated population was between 4,600 and 6,100 bears in 2007. This represents an estimated 27% increase over the 1997 population estimate. These increases in the black bear population are consistent with management goals laid out in the previous plan.

In developing the current management plan, the Department sought Vermonters' opinions on whether bear populations in their county should increase, stay the same, or decrease. The majority of Vermonters surveyed (57%) wanted to see bear populations in their county remain the same, 16% wanted the population to increase, 7% wanted it to become lower, and 20% either didn't have an opinion or didn't know (Fig. 4.3)

In general, Vermonters' opinions on bear populations were consistent across regions of the state. There were two exceptions: in Central Vermont 22% of the respondents supported an increase in the population and in Chittenden County 28% of respondents either had 'No Opinion' or 'Didn't Know' (Table 4.1). Of those Vermonters favoring to increase bear populations, wanting to see more bears and the value of bears to




Figure 4.1 Distribution of female bears from harvest data, 2004-2008.




Figure 4.2 Estimated Black Bear populations by 5-year blocks, with 80% confidence limits, 1987-2008.

the ecosystem were given as the primary reasons. Residents of the Central and Southern Vermont regions who wanted bear population increases were particularly interested in seeing more bears. Statewide, reducing bear-human conflicts was the primary reason given for wanting decreases in local bear populations.

In contrast to Vermonters' general satisfaction with bear populations in their county, bear hunters satisfaction declined significantly from 75% to 54% since the previous survey was conducted in 1996. Dissatisfaction increased from 20% to 32% during the same survey interval. The survey was not able to query the rationale for the decline, but factors other than bear population levels, such as access to unposted land or a low bear harvest the previous year may have influenced respondents' opinions.

Management Strategies

- 1.1 Update and re-evaluate Vermont's black bear population model to reflect the most current harvest and biological parameter data available.
- 1.2 Evaluate and develop hunting season structures that align population estimates with biological data, habitat limitations, and public satisfaction data to sustain a bear population between 4,500 and 6,000 animals.

ISSUE 2. Bear Habitat Conservation

GOAL: Maintain a no net loss of function and value of existing bear habitat.

Historically, black bear management programs concentrated on regulating the legal harvest of the species to ensure that the population was sustainable. Today, management objectives in Vermont revolve around maintaining wild, free-ranging, viable populations of black bear as well as the conservation of bear habitat. Wildlife managers are looking toward conservation of large blocks of interconnected forestland and protection of the most critical areas of black bear habitat as the best




Figure 4.3 .Vermont residents' opinion regarding future bear population change

Table 4.1 Vermont Residents' Opinions on Desired Bear Populations by Region.

Region	Increase	Same	Decrease	No Opinion	Don't Know
Chittenden	10 %	56 %	5 %	14 %	14 %
Northeast Kingdom	17 %	60 %	9 %	8 %	6 %
Central Vermont	22 %	54 %	7 %	7 %	10 %
Southern Vermont	18 %	60 %	7 %	7 %	9 %

long-term strategy for sustaining Vermont's bear population.

In Vermont, black bears require large forested areas that have a variety of food resources, particularly hard mast such as acorns and beechnuts, and provide core habitat for successful reproduction and allow them to avoid humans (Hugie 1982; Hammond 2002). Black bears rely on concentrated stands of American beech trees located at least one kilometer from roads and houses as an essential fall source of high nutrition food needed to build fat reserves prior to denning for the winter (Hammond 2002; McLaughlin 1998; McLaughlin et al. 1994; Wolfson 1992; Hugie 1982; Beeman et al. 1977). Researchers have found during years that beechnuts are in short supply, bears travel great distances to find alternative food sources and incur heavier mortality rates (McLaughlin et al. 1994). The availability of hard mast in the fall affects the minimum reproductive age and rate and cub survival. Simply put, concentrated stands of beech trees used by black bears are critical to the survival and reproduction of bears in Vermont.

Other important hard and soft mast food resources in Vermont include acorns, cherries, berries, apples, and mountain ash.

Vermont's bear population has increased in recent decades as forests have increased over the landscape and recent bear management strategies have encouraged population growth. Bears are now found throughout much of the state, yet the greatest concentrations of Vermont bears are found in "core" habitats that tend to be remote from roads, human developments, and human activity. Vermont black bears need large forested blocks of sufficient size to meet the home range and food requirements of female bears and cubs. The existing range, although becoming increasingly more fragmented in some parts of the state, has been sufficient to support an increasing bear population. Large public and private forest land holdings play an important habitat conservation role in this regard.

A recent study at the University of Vermont, however, indicates that increases in human development will diminish bear habitat (Donovan et al. 2007). The study projects that between the years 2000 and 2020, the number of housing units in Vermont will increase by at least 12,107 and that most of these units will occur in what are now relatively undeveloped locations. Under this scenario, the occurrence of black bear would likely decline in some areas of the state in the next 12 years (Fig. 4.4).

In the mid-1980s, the Department recognized the negative impact that housing developments were having on key black bear feeding and travel areas and began recommending through Act 250, the state's land use and development law, protection of

critical bear habitat. For six years during the 1990s, Department biologists studied the movements and behavior of radio-collared bears in relation to roads, houses, ski trails, and various recreational activities. The findings from this study have helped the Department in its efforts to work with developers to include the habitat needs of bears into their long-term planning processes (Hammond 2002).

Today, the Vermont Fish & Wildlife Department provides advice and technical assistance for the protection of critical bear habitat, such as beech and oak stands, wetlands, and travel corridors. For example, the Department has been working with the Vermont Agency of Transportation to address the issue of habitat connectivity by developing wildlife suitability maps identifying areas that support animals that require large areas, such as black bear and moose. These maps help identify areas that should be conserved and managed so that animals can safely cross roads that bisect their habitat. The map also provides towns and regional planning commissions with a focus for land use planning (Fig. 4.5).

Public opinion surveys suggest that Vermonters continue to strongly support many forms of habitat conservation. Surveys also found that the public supports land conservation efforts in order to maintain the existing habitat base. In addition, 89% of the respondents said it was important to them to know that species like the black bear exist in Vermont, even though they are seldom seen. Eighty percent of Vermonters support using Act 250 as an important habitat protection tool (Duda et al. 2007). Although Act 250 is unique and effective legislation, it does not apply to development involving all critical bear habitat. A survey conducted by Responsive Management (Duda et al. 2007) found that 92% of the general public supported the Department working with town or regional planning commissions to design town plans that address and preserve important wildlife habitat. As a result, the Department has increased its efforts to work with towns and regional planners by providing technical assistance and on the ground assistance for related issues involving conservation of wildlife habitat.




Figure 4.4. Probability of occupancy of a site by black bear in Vermont in relation to the percent of development.

Management Strategies

- 2.1 Maintain and enhance habitat protection efforts through Act 250, wood-to-energy harvest review, work with town and regional planning commissions, land acquisition, and other conservation methods.
- 2.2 Provide technical assistance in managing for critical bear habitat in the Use Value Appraisal program.
- 2.3 Revise and update “A Landowner’s Guide, Wildlife Habitat Management for Vermont Woodlands” to include habitat management recommendations for black bears.

ISSUE 3. Human/Bear Conflicts

GOAL: Minimize the overall number of negative interactions occurring between bears and humans to achieve acceptable levels of human safety and social acceptance.

Bear populations, like all wildlife populations, are normally restricted by two factors — biological carrying capacity and cultural carrying capacity. As described previously, biological carrying capacity is the maximum number of animals an environment will support on a sustained basis. Population density and distribution depends on availability of food, cover, and space. Cultural carrying capacity is the maximum number of bears that can coexist compatibly with local human populations. Bear habitat can often support more animals than the public is willing to tolerate. Bears are large animals capable of causing extensive property damage and even human injury.

Department personnel have documented an increase in the number of people reporting conflicts with bears since the last management plan (1997-2008). This is also reflected in survey data from 2007 that found 14% of wildlife damage incidents were related to nuisance bears. This represents a seven-fold increase from 1996 when only 2% of incidents were related to nuisance bears. In spite of this increase in bear/human conflicts, a large majority of Vermonters (70%) are tolerant of bears on their property while only 18% are not (Duda et al. 2007). Had Vermont residents not had this tolerance for bears, the Department expects that many more might have registered reports of conflict given the increase in both bear numbers and the human population in the past ten years.




Figure 4.5 Wildlife Suitability Map example for black bear and other wide-ranging species.

Hunting plays a significant role in shaping Vermont’s cultural carrying capacity for bears. The Vermont Fish & Wildlife Department uses regulated hunting both to provide harvest and utilization of bears and as a tool to maintain bear numbers at target population levels throughout the state. Hunting also teaches bears to be wary of humans. This reduces the number of bears that might become “nuisance animals” causing damage to livestock or farmers’ crops, raiding dumpsters, or entering buildings in search of food.

The history of hunting and utilizing bears for food in Vermont is a long one. The Department believes that regulated hunting and the training of hunting dogs helps keep Vermont’s bears wild, which in turn has encouraged a higher cultural carrying capacity. The extreme wariness of the Vermont black bear may be related to the bounty system that was in place for 110 years ending half way through the twentieth century. Following the end of the bounty system, liberal hunting seasons and the chasing of bears with hounds has continued to make bears wary of humans. Currently, nuisance bear situations are more likely to occur when there is a shortage of natural food sources that cause them to become bolder in their search for food.

Generally, the wariness of black bears limit their exposure to human-occupied landscapes. A

shortening of the bear hunting season in 1990 resulted in a planned increase in the bear population, resulting in more bears attempting to establish home ranges in less secluded areas that had previously been unoccupied. Vermont now has more bears living in closer proximity to human residences. This situation has increased the likelihood of undesirable human interactions. These situations include, but are not limited to, the destruction of farmers' crops, commercial beehives, and fruit orchards; the killing of livestock; the raiding of garbage barrels and bird feeders; and an increase in the number of bear-motor vehicle collisions.

The Department has developed posters, brochures, and public service announcements designed to increase awareness and to help the general public understand black bear behavior and to live better with black bears (Fig. 4.6). These public outreach efforts advise citizens to remove bird feeders from their yards when bears are not in hibernation and discourage feeding bears through the slogan, "A fed bear is a dead bear." Game wardens also advise and help landowners who report damage from bears.



Management Strategies

- 3.1 Update statewide policy for handling black bear/human conflicts.
- 3.2 Improve and disseminate outreach/education materials and messages for minimizing human/bear conflicts.
- 3.3 Monitor bear/human conflicts and explore new strategies for reducing the number of complaints from the public.
- 3.4 Use permitted houndsmen with trained bear hounds to haze bears and keep them wary of humans.

ISSUE 4. Bear Management Strategies and Season Structure

GOAL: Optimize public hunting opportunity for the utilization of bears for food and other appropriate purposes and ensure hunter satisfaction within biologically sustainable regulations.

People hunt for many different reasons, but over 90% of hunters who were surveyed listed the reason they hunt black bear was "for food." Most hunters also have a deep appreciation of the out-of-doors and love and respect the species they pursue during hunting season and watch during the rest of the year. This appreciation often results in hunters leading efforts for increased harvest regulation, habitat protection, and other conservation initiatives.

There are currently 25 laws and regulations that regulate the harvest, utilization, and sale of bears in Vermont. Black bear season is currently set on a statewide basis with no regulatory differences among wildlife management units. The season length is one of the longest in the nation, running from September 1 to the Wednesday following the opening day of the November deer rifle season. Use of trained hunting dogs to hunt bears is allowed in Vermont by permit only. Baiting for the purpose of taking bears is prohibited. The bag limit is currently set at one bear per licensed hunter per season.

The management of Vermont's black bear population through regulated hunting offers several challenges.

Figure 4.6 Two-foot by three-foot black bear poster developed by the Department for distribution to the public.

Annual black bear harvests are sometimes more reflective of food availability, weather events influencing the timing of denning, and other factors affecting bear movements instead of simply increases or decreases in the population (Fig. 4.7). For this reason, managing a bear population requires reviewing several years of harvest information before proposing regulatory changes to the hunting season. The low reproductive rate and longevity of black bears further complicate management by delaying bear population responses to harvest adjustments (Fig. 4.8).

Black bear hunting participation rates in Vermont are relatively low, remaining significantly below that of white-tailed deer and wild turkey. They have decreased from an estimated 28% of hunters in 1996 to 17% in 2007 probably as a result of shortening the length of the bear season that overlaps with the November rifle deer season. Prior to shortening the season in 1990, bear harvest levels were greatly affected by deer hunters that opportunistically harvested black bear while pursuing deer. Given these facts, it may come as no surprise that bear hunting satisfaction decreased from 75% to 54% between 1996 and 2007.

There are a variety of management strategies available for stabilizing and maintaining existing bear populations in Vermont while providing hunting opportunities. Listed below are management tools that can, individually or in combination, aid in regulating the bear harvest to meet the statewide population goals of 4,500 to 6,000 bears. It must be emphasized that bear season length and structure have historically been adjusted to increase or reduce the statewide bear population.




Figure 4.7. Annual Vermont black bear harvest, 1985-2008.




Figure 4.8 Vermont bear harvest as a proportion of population estimate.

Bear License

A key component to an accurate population measure based on harvest is the parameter of hunter effort. Simply examining harvest differences each year cannot provide a reliable correlation between harvest and total numbers of bears. Bear hunters in Vermont are not required to purchase a separate bear license. A bear tag is included as part of the big game hunting license. This license has a long history (at least 45 years) and has resulted in the expectation that a bear tag is part of the value purchased with the big game license.

A separate black bear license would be one way to determine the number of hunters intending to pursue bears each year. However, the sale of separate bear licenses would not provide other important information such as hunting hours expended in pursuit of bears, the number of bear sightings, or WMU preferences. Collection of all of these data could be accomplished without requiring a separate bear license. Since 2000, surveys indicate that 46% of

all bears harvested were taken incidentally by hunters while hunting another game species. Deer hunters took the largest percent of bears, 29% (Table 4.2).

Survey results from 2007 indicate opposition to the establishment of a separate black bear license has remained at 61%. The percent of hunters supporting a separate bear license had declined to 29% in 2007 from 31% in 1995. These results suggest implementation of a separate license would be difficult. Given that Vermont's bear population has been increasing, it is possible that it will be necessary to extend bear season further into the November deer rifle season in the future. Reducing the number of hunters that may take a bear by requiring a new and separate bear license could jeopardize the Department's future ability to control total bear numbers.

Bag Limit

Vermont's statewide bear season bag limit of one bear was first established in 1968. Bag limits may be effective means of adjusting harvest levels to meet particular population goals. The single bear season bag limit has served Vermont well in initially reducing bear harvests and allowing for population growth. New population goals that require stabilizing or potentially reducing the number of bears could involve re-examining the current bear season bag limit. However, increasing the annual bag limit for bears might call for reducing the length of the bear hunting season, a move that could require eliminating the current overlap of bear season with the first five days of the deer rifle season. Increasing opportunity for one segment of the hunting population will likely decrease opportunity for another.

To date, only Oregon, Washington, and Alaska have fall season bag limits greater than one. Several states and Canadian provinces hold a spring bear season that includes its own bag limit. Although increased fall bag limits are a relatively untested management tool, they may be important in stabilizing bear populations if they can be implemented cautiously and other management tools prove to be ineffective.

Table 4.2 Percent of Vermont bears harvested while hunting for a specific species 2001 – 2008.

SPECIES HUNTED	SEPTEMBER	OCTOBER	NOVEMBER	TOTAL	PERCENT
Bear	1,233	883	264	2,380	54%
Deer	0	473	797	1,270	29%
Birds	4	16	3	23	<1%
Other	15	55	14	84	2%
Unknown	270	230	132	632	14%
Total	1,522	1,657	1,210	4,389	100%

It would be critical to monitor any increased harvest from expanded bag limits to evaluate its effects on regional bear populations and the sex and age of the animals harvested.

The Department believes that revising the bear season bag limit is one possible way to stabilize and control the bear population. Since this method would reallocate the bear resource, any proposal to change bag limits, however, will need to follow a rigorous public outreach effort and significant buy-in from the hunting public.

Regional Management Zones

Management of black bear is currently conducted on a statewide basis because data in measuring hunter effort and distribution are inadequate to inform fine scale regulation of harvest. Although simple to administer, comply with, and enforce, a statewide bear season does limit the flexibility of the Department to adjust the harvest in response to regional issues or variables. In spite of its small size, Vermont does have significant regional differences in bear density, bear habitat use, food supplies, weather patterns, road access, habitat fragmentation, hunting pressure, number of nuisance complaints, and development pressure. As a better understanding of bear population distribution develops, the flexibility to tailor hunting seasons to regional differences may be necessary.

Season Length and Structure

Vermont has regulated the annual bear harvest for the past 40 years by simply adjusting the length of the hunting season. In particular, the number of days that bear season is open during the November deer rifle season has the greatest effect on the total bear harvest especially during years when food supplies are abundant and bears continue to feed instead of going to their dens for the winter.

In 1990, Vermont's bear season was shortened by four days in November (from the second Sunday of deer season to the first Wednesday) in order to reduce bear harvests and increase the population. As previously discussed, this management action was very effective and resulted in a significant increase in Vermont's bear population.

The Department has demonstrated that incremental changes in the number of days that bear season extends into the November rifle deer season is an effective means to regulate the harvest of bears and to adjust the bear population. However, creative ways to achieve population goals on a local or regional basis may be needed in the future. As human and/or bear populations grow, greater flexibility may be needed to address the specific issues to best respond to the needs of bears and people.

Hunting Bears with Hounds

Bear hunting with hounds can be a controversial method to locate and pursue a bear. The Department recognizes and acknowledges that there are issues of public concern such as hounds on private property, the use of telemetry, and the length of the training season. Nevertheless, the Department continues to support bear hunting with hounds as a legitimate and biologically sound hunting method that has advantages in that chasing bears keeps bears wild and minimizes nuisance and other conflicts with humans. Vermont game wardens routinely recommend bear houndsmen to property owners who are dealing with nuisance bears. Houndsmen have come to the rescue of many a farmer who has had problems with bears in their corn, apple orchard, or beehives. Many times, chasing a bear away will prevent its death at the hands of the property owner.

Bear houndsmen are required to have a permit to train and hunt bear with dogs in Vermont. The number of nonresident bear houndsmen permitted to hunt in Vermont is limited to 10% of the resident permit numbers. Recent law changes have placed greater restrictions on the ownership and residency of the dogs permitted to run on the permits of Vermont resident houndsmen. Because the number of bears taken with the aid of hounds is only about 10-15% of the total bear season harvest, bear hunting with dogs is not the most important method for controlling the bear population. The benefits of hunting bears with dogs are significant, however, and the Department will continue to address issues of public concern that would restrict hound hunting in Vermont. The

Department will also continue to work closely with the Vermont Bear Hound Association to discuss and understand the issues of concern and identify actions that can be taken to address them.

Sale of Bear Parts

The Department continues to participate in and monitor national and international assessments of the effect of trade in bear parts on wild bear populations. Recent changes in the market for bear parts along with increasing black bear populations in North America have reduced concerns over this activity in recent years. The 1992 listing of the American black bear as a CITES Appendix I species now also provides significant monitoring of international trade.

The Department will continue to monitor the sale of black bear parts. If trends and activity in the sale of bear parts, particularly gallbladders, is found to be detrimental to Vermont's bear population or pose a threat to bear populations in other parts of the world, it may propose further regulation or prohibition of such sales.

Guided Commercial Bear Hunts

Bear hunters have expressed their concern that commercial guides have been securing exclusive hunting rights to key bear feeding areas such as cornfields near traditional bear travel routes. This allows guides to offer clients a "guarantee" of sorts that they will take a bear because of the high concentration of bears in these areas. Because cornfields may attract bears from several miles, it has been suggested that the cumulative harvest of bears at these sites could have a disproportionate impact on bear populations in several nearby towns. On face value, this seems to be a plausible argument, but it has not been borne out by statistics.

Guiding, when properly administered, can be a quality introduction to hunting and a form of hunter mentoring. These are important components of hunting recruitment and retention. Poorly administered guide services that are purely profit-driven can be very damaging to hunting. Developing some standard for commercial guiding may be a way to assess the effect on the bear population from the guiding industry through analysis of hunter effort and harvest data. A standard might also serve as a marketing tool for guides. Guide programs administered by fish and wildlife departments in other states are not self-supporting and are costly to administer.



Management Strategies

- 4.1 Hunting season management strategies and season structure will be evaluated and adjusted to maintain the population goal of 4,500 to 6,000 bears. Changes in hunting season structure will consider, when necessary, the use of season length, regionalization, or incremental changes to season bag limits to achieve population goals.
- 4.2 Work with partner organizations on issues related to bear management as they are raised throughout the management plan period and develop specific strategies to address them. Such strategies may range from legislative changes to educational efforts.



JOHN HALL

REFERENCES- BLACK BEAR

- Beeman, L.E. and M.R. Pelton. 1977. Homing of black bears in the Great Smoky Mountains National Park. International Conference on Bear Research and Management 3:87-95.
- Duda, M. D., M. Jones, A. Criscione, C. Craun, T. Beppler, T. Winegord, A. Lanier, S. J. Bissell, and J. B. Herrick. 2007. Public opinion on wildlife species management in Vermont. Report by Responsive Management for the Vermont Fish & Wildlife Department, Waterbury, VT.
- Donovan, T.M. A. Troy, and A. Voinov. 2007. Land use change in Northern Forests: Assessments and recommendations for conserving biodiversity. Vermont Cooperative Fish and Wildlife Research Unit. 32 pp.
- Elowe and Dodge. 1989. Factors affecting black bear reproductive success and cub survival. Journal of Wildlife Management. 53:962-968.
- Hammond, F.M. 2002. The effects of resort and residential development on black bears in Vermont. Final Report. Vermont Agency of Natural Resources, Department of Fish & Wildlife.
- Hugie, R.D. 1982. Black bear ecology and management in the northern conifer-deciduous forests of Maine. University of Montana, Missoula, MT.
- McLaughlin, C.R. 1998. Modeling effects of food and harvests on female black bear populations. Ph.D. Dissertation. University of Maine.
- McLaughlin, C.R., G.J. Matula Jr. and R.J. O'Connor. 1994. Synchronous reproduction by Maine black bears. International Conference on Bear Research and Management. (1):471-479.
- Wolfson, D.L. 1992. Development of a quantitative procedure to assign a value rating to beech stands as black bear habitat. Unpublish. M.S. Thesis. Antioch College, Keene, NH.

WILD TURKEY

I. Management History

Records from the late 1700s and early 1800s indicate wild turkeys were present in southern Vermont. Most turkeys in the state seem to have existed along the Taconic Mountain Range in the southwest and along the Connecticut River Valley in the southeast. Massive loss of forestland and unregulated market hunting in the early nineteenth century led to the disappearance of Vermont's wild turkeys by the mid-1800s.

A number of private fish and game clubs attempted to re-establish the birds during the late 1950s at various locations around the state by releasing turkeys raised on game farms. This effort failed, however, because these birds lacked the inherent hardiness and survival skills of wild turkeys. These stocking attempts convinced the Department that successful reintroduction of turkeys into Vermont would require live-trapping of the hardier wild birds from another state.

In the late 1960s, the Department made arrangements with the New York Department of Environmental Conservation to permit Vermont biologists to live-trap wild turkeys and relocate them. Vermont first released 17 wild New York birds in Pawlet, Vermont, in 1969. A second release of 14 wild birds was made in Hubbardton in 1970. Today's wild turkey population of more than 50,000 birds directly descends from this original stock of 31 New York wild turkeys. The Department initially expected the expansion of the wild turkey population to be limited to the part of the state reported to be historical wild turkey range, south of US Route 4. Only 30 years after introduction, turkeys ranged across the entire state. These hardy birds have far exceeded expectations and have successfully exploited Vermont's mosaic of forestland and dairy farms.

The Department began efforts to expand wild turkey range within the state soon after their initial introduction. Over a ten-year period ending in mid-1980s, live-trap and transfer techniques were employed to capture and move wild birds from the original release area in Rutland County to other parts of the state. Birds were released in Bennington, Brattleboro, Bristol, Dummerston, Grand Isle, Halifax, Jericho, Milton, Norwich, Pownal,



Rockingham, Shaftsbury, Springfield, Strafford, and Weybridge. Birds were also restocked in Alburg, Fairfax, Georgia, Grand Isle, and Swanton in the mid-1990s to augment a struggling local population perhaps limited by overharvesting in the fall.

Vermont's first, modern wild turkey hunting season was held in parts of Addison, Bennington, and Rutland Counties in the spring of 1973. A season was held for 12 days (May 9-20) with a limit of one bearded turkey. Twenty-three turkeys were harvested by 579 permitted hunters. The first fall hunt, held in 1975, occurred in a limited area of southwestern Vermont.

Over the past 30 years, wild turkeys have thrived in Vermont and public participation in turkey hunting has continued to increase. Reduction in fall harvest opportunities imposed following the disastrous and extremely severe winter of 1993-94 helped stimulate rapid turkey population growth and expansion. The population is estimated to have increased from approximately 12,000 to 45,000 birds in the period from 1995 through 2001. The increase in spring turkey harvest mirrors the species' population growth (Fig. 5.1).

Turkey hunting opportunities have been expanded to new areas of the state as the population has grown. The entire state of Vermont was opened for the first time to spring turkey hunting in 2004. In addition, relatively liberal fall turkey hunting opportunities, compared to those in other states, are now offered

in most areas of the state. Average annual combined spring and fall harvest of turkeys now totals about 5,800 birds per year. Viewing opportunities have also expanded tremendously for thousands of Vermonters who delight in simply viewing wild turkeys in their natural setting. As a result of the Department's turkey management initiatives, the wild turkey population has risen to the highest level in Vermont history. The wild turkey population is currently estimated to exceed 50,000 birds statewide.

Wild turkey research has found that short-term turkey population fluctuations result from combinations of random environmental conditions such as rainfall and temperature events that affect nesting hatching success, survival of poults and winter survival. Long-term population trends, however, are primarily influenced by changes in the quantity and quality of suitable habitat across the landscape.

Forests now dominate close to 75% of Vermont's land area with only about 15% in an open, nonforested condition. Although the eastern wild turkey is primarily regarded as a forest-dwelling bird, ideal turkey habitat includes a diverse mix of habitat types, forest succession stages and open land, which provides the greatest opportunity for feeding, nesting, and brood rearing. Research shows that turkey nesting rates are consistently higher and turkey populations more stable in habitats consisting of a mosaic of forests and fields than in those areas composed mainly of either mostly forest or mostly open land. The highest densities of turkeys in Vermont follow this pattern, occurring in areas where the available habitat is closer to the ideal mosaic mix of conditions.

Forests are an important habitat component for turkeys especially when forests consist of oak, beech, and pine stands that produce abundant hard mast crops (acorns, beechnuts, and other seeds) that are consumed by the birds in the fall and winter months. Forests also provide the large, dominant trees used by turkeys for roosting. These types of forests are relatively common in Vermont, especially in the




Figure 5.1. Spring Turkey Harvests 1972 - 2008

Champlain Valley, Connecticut River Valley, and the foothills of the Green Mountains and Taconic Mountains. Forest management practices can be used to insure availability of adequate mast crops and roost trees through time.

Clearings and openings in the forest are also a vital habitat component for Vermont's wild turkey population. Whether created as farm pastures, hay fields, or openings within the forest, herbaceous plants such as grasses and clover provide critical habitat for turkey broods. These open areas have abundant insect populations on which young turkeys rely during early growth. The most beneficial clearings are old pastures, dominated by a mix of forbs, weeds, and fruit-bearing shrubs, as opposed to monocultures of grasses. Management can be used to enhance openings through periodic, selective mowing, liming, and fertilizing to favor desirable plant species.

Turkey brood range is a very important habitat consideration. Hen turkeys nest on the ground and prefer nesting in locations having lateral cover of herbaceous plants and shrubs. Recent clear cuts, thinned timber stands, fields, and croplands provide cover suitable for nesting and brooding turkeys. The best management practices for enhancing forest nesting habitat are conventional forest regeneration practices, especially even-aged timber management and group selection methods.

Trends in agriculture may affect the future distribution and abundance of turkeys statewide. As the number of Vermont farms continues to decline and the trend toward increasing forestation and forest age progresses, availability of open land may limit wild turkey production in Vermont.

II. 2010-2020 Turkey Management Issues, Goals, and Strategies

The overall goal of wild turkey management in Vermont is to manage the state's wild turkeys to sustain healthy, abundant populations that will provide hunting and viewing opportunities and will satisfy social expectations and tolerances for turkeys. This management goal aims to sustain an abundant wild turkey population that is truly wild and that is below both the biological carrying capacity of its habitat and the cultural carrying capacity.

ISSUE 1. Turkey Population

GOAL: To adequately assess Vermont's wild turkey populations and trends.

Sustaining healthy wild turkey populations in each wildlife management unit (WMU) that is consistent with Department goals requires an ability to accurately estimate abundance of turkeys or at least trends in relative abundance. Harvest data, nesting success indices as provided by brood surveys, and hunter pressure through license sales are critical elements of the Department's ability to monitor turkey population trends.

Management Strategies

- 1.1 Annually collect and assess turkey harvest data to determine trends as well as summer/fall turkey sighting survey data in order to direct future management decisions.
- 1.2 Conduct the public annual Internet turkey brood survey

1999-2008 Plan Accomplishments

Vermont's wild turkey population was estimated to be approximately 23,000 to 25,000 birds in 1998 at the time the previous turkey management plan was written. The population had risen to 50,000 to 55,000 birds by 2008, a doubling of the population during the ten-year management period. New spring turkey harvest records have been set in eight of the past 11 years. Wild turkey enthusiasts throughout our state have enjoyed countless hours of harvest and viewing opportunities of this highly regarded native wildlife resource.

The following is a list of specific issues and recommendations that were originally proposed in the 1999 plan with the resulting action taken to address them during the past ten-year planning cycle.

► **Recommendation 1. Adjust spring hunting hours to begin one-half hour before sunrise to noon (previous hours were 5:00 a.m. to 11:00 a.m.).**

► **Recommendation 2. Adopt turkey hunting season zones based on existing wildlife management units (WMUs).**

❖ **Action:** Both of these recommendations were adopted by regulation in 2000.

► **Recommendation 3. Change (expand) existing spring turkey hunting zones.**

❖ **Action:** The expansion of spring turkey hunting was adopted by numerous regulation changes throughout the planning period.

2000 – Expanded spring hunting to include all of WMUs H1, H2, and G

2001 – Expanded spring hunting to include Zones B, D1, and D2; WMU A opened to spring hunting by

permit, 75 permits issued

2002 – Held the first spring youth season first weekend prior to start of May season; 80 permits issued in WMU A

2003 – Opened WMU A to all licensed hunters, no WMU A permits required

2004 – Expanded spring season to include all of Zone C and E, resulting in entire state of Vermont open to spring hunting for all licensed hunters

► **Recommendation 4. Change fall hunting zones and season length, and establish threshold guidelines for initiating, liberalizing, or curtailing fall hunting seasons.**

❖ **Action:** Changes to the fall wild turkey season were adopted by regulations established in 2000 and 2003.

2000 – Expanded the fall season in J1 to include the entire zone with seven-day shotgun season

– Expanded fall hunting in Zones H2 and J2 to include the regular bow and arrow season and a new archery-only season during the current seven-day shotgun season in adjacent zones

– Reduced fall shotgun season in WMUs G, I, L, M1, M2, O1, O2, P, Q from 16 days to 7 days in length

2003 – Expanded fall bow hunting season length in H1 to include regular bow and arrow season and through regular seven-day shotgun season

– Expanded fall seven-day shotgun hunting in J2 and H2

along with the Department staff summer turkey survey.

- 1.3 Continue the turkey project's investigation into the genetic variability and structure of the statewide population.
- 1.4 Evaluate new wild turkey population estimation methods and models for use in Vermont.
- 1.5 Evaluate the use of a public Internet survey to assess winter flock sightings.

ISSUE 2. Public Satisfaction with Current Population Levels

GOAL: Assess public and hunter satisfaction with current turkey population levels and management program.

Respondents to a 2007 public opinion survey were asked their opinion about wild turkey population levels around the state. The majority of Vermonters (60%) were satisfied with the turkey population in their county; 15% wanted more turkeys, 10% wanted fewer turkeys, and 15% had no opinion. In an effort to gauge the current level of satisfaction among Vermont's turkey hunters, the Department asked survey participants: "How satisfied are you with your wild turkey hunting experience in Vermont over the past five years?" Ninety-two percent (92%) of the respondents indicated that they were either "Very Satisfied" or "Somewhat Satisfied" with their hunting experience in Vermont. This level of satisfaction for turkey hunting was higher than any other big game species and 16% higher than the opinion survey taken in 1998. The complete results of this question are reported in Table 5.1.

Management Strategies

- 2.1 Provide statewide spring bearded-bird-only seasons (including the Youth and regular May season) and limited fall either-sex hunting seasons in WMUs that can sustain a fall harvest so as to provide for population stability.
- 2.2 Prioritize high quality spring hunting over additional fall harvest opportunity.
- 2.3 Manage fall turkey harvests through changes in fall hunting season length within WMUs depending upon stability or growth of three-year average spring harvest densities, except in WMU A Champlain Islands where inadequate forest cover exists to sustain a fall firearm harvest.

ISSUE 3. Fall Turkey Hunting

GOAL: To provide appropriate opportunity for sustainable fall hunting while maintaining current levels of high quality spring turkey hunting.

The topic of fall turkey hunting is perhaps one of the most misunderstood facets of turkey management in Vermont and generates the most comments from the hunting community. While there is inherent variation in both annual production and survival of wild turkeys, fall either-sex hunting can play a pivotal role in regulating population size. Research on wild turkeys and population modeling in several other states indicates that significant fall hunting pressure can suppress turkey population growth and reduce spring population densities. Although the effects of spring turkey hunting may not be entirely benign due to potential nest disturbance, illegal harvesting of hens, and effects on age structure of male turkeys, fall turkey hunting can have a much more profound impact on turkey populations. Vermont's experience with fall turkey hunting in Grand Isle and Franklin counties in the mid-1980s demonstrated how quickly heavy fall harvests can reduce turkey populations. This is the principal reason that the current fall season bag limit and season length is less in zones with lower turkey densities. In some cases these limits are more conservative than some hunters would prefer.

The Department supports the management practice of using fall hunting zones to regulate turkey populations in areas having the best turkey habitat and highest densities for the following reasons:

- *Fall wild turkey seasons impact turkey populations by primarily removing female turkeys.* Sixty-five to 70% of Vermont's fall harvest consists of female turkeys. This large

Table 5.1 2007 Hunter Satisfaction Survey Results

Satisfaction level	Percent of respondents
Very satisfied	57%
Somewhat satisfied	35%
Neither satisfied nor dissatisfied	0%
Somewhat dissatisfied	7%
Very dissatisfied	1%

female fall harvest reduces the number of hens nesting the following spring. Fall harvests of hens followed by a severe winter and/or poor reproduction the following spring can quickly change the abundance of turkeys — thus fall hunting in this situation can only add to the mortality rate.

- ***The illegal harvest of female turkeys may be more of a factor in the fall season when entire broods are vulnerable to harvest.*** Several states have documented higher illegal take during the fall season.
- ***Turkeys in Vermont are living at the northern fringe of their continental range and are more vulnerable to natural mortality from severe winters and cold, wet springs.*** Severe winters can result in substantial population losses and depress spring reproductive success. While Vermont can experience severe winter conditions throughout the state, this factor is especially significant in the more northerly and higher elevation wildlife management units. The harvest of female turkeys can be additive to these natural mortalities.
- ***The regulatory process does not allow for timely changes to the fall hunting season in response to annual fluctuations in turkey productivity.*** Changing harvest regulations via the mandatory, Administrative Procedures Act is a very deliberative, lengthy process. The time frame for developing a change in turkey regulations, from preparing a proposed rule until final adoption of the rule, requires approximately 18 to 22 weeks. This severely limits the Department's ability to respond quickly to significant increases or decreases in poult production or survival. This also requires the Department to be conservative when proposing harvest regulations.

When asked in the 2007 opinion survey, the majority of turkey hunters (68%) support the current management strategy to limit fall turkey hunting for the purpose of maximizing spring turkey

harvests. Although this indicates strong support for the current approach, there may be opportunities to systematically enhance/expand fall turkey hunting without compromising the goal of providing quality spring turkey hunting in Vermont.

Management Strategies

- 3.1 Provide public opportunity to harvest wild turkey for food and other utilitarian purposes.
- 3.2 Facilitate healthy, abundant spring turkey populations that are stable using modest, fall hunting seasons/bag limits to control the population. When the three-year spring average harvest density reaches the specific threshold value, liberalization of fall hunting in a WMU may be called for (initiate shotgun seasons, extend gun seasons). See Table 5.2.
- 3.3 Consider reducing the current guideline for the threshold as to when fall gun hunting opportunities could be initiated in a new WMU, from the three-year average spring harvest density of one bird per square mile, to an average harvest density of .75 bird per square mile.
- 3.4 Lengthen the current fall seven-day shotgun season to a nine-day season.
- 3.5 Expand the fall shotgun season to include WMUs H1, D1, and B with a nine-day shotgun season.
- 3.6 Expand the fall archery turkey season, coinciding with the opening of the deer archery season, to allow archery hunting statewide.
- 3.7 Investigate establishing a new separate "Fall Gun Season Only" tag.



NWTF.ORG

Table 5.2 Spring Wild Turkey Harvest 1998-2008

WMU	Spring Harvest per Square Mile											3-year Average
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2006-08
A	1.10	1.57	1.86	1.90	1.62	3.33	2.76	2.38	2.86	3.19	3.10	3.05
B				0.28	0.37	0.25	0.43	0.59	0.83	0.84	1.32	1.00
C							0.20	0.25	0.39	0.29	0.49	0.39
D-1				0.34	0.37	0.23	0.39	0.56	0.70	0.54	0.99	0.74
D-2				0.36	0.44	0.25	0.35	0.53	0.60	0.53	0.65	0.59
E	0.01	0.03	0.02	0.04	0.08	0.05	0.17	0.25	0.27	0.20	0.28	0.25
F-1	2.14	2.90	3.59	3.92	4.14	3.50	3.40	3.36	3.14	3.78	2.61	3.18
F-2	1.73	1.55	2.76	2.64	2.34	2.02	2.12	2.04	1.99	1.80	1.57	1.79
G	0.10	0.18	0.26	0.21	0.25	0.16	0.32	0.29	0.35	0.37	0.40	0.37
H-1	0.11	0.41	0.60	0.63	0.80	0.52	0.60	0.79	0.77	0.68	0.85	0.77
H-2	0.58	0.78	1.24	1.03	1.64	1.37	1.37	1.56	1.16	1.40	1.78	1.45
I	0.17	0.19	0.24	0.24	0.30	0.20	0.23	0.25	0.24	0.24	0.20	0.23
J-1	0.49	0.46	0.74	0.85	0.84	0.64	0.78	0.81	0.60	0.90	0.86	0.79
J-2	0.41	0.67	0.63	0.83	0.78	0.54	0.74	0.78	0.67	0.83	0.66	0.72
K-1	1.29	1.96	1.95	2.00	1.85	1.70	1.09	1.08	1.42	1.34	1.28	1.35
K-2	1.43	1.66	1.74	2.02	1.35	1.50	1.33	1.49	1.34	1.50	1.30	1.38
L	0.11	0.21	0.25	0.33	0.25	0.21	0.15	0.28	0.26	0.28	0.30	0.28
M-1	0.23	0.31	0.41	0.34	0.23	0.26	0.17	0.36	0.27	0.47	0.42	0.39
M-2	0.53	0.62	0.87	0.86	0.65	0.65	0.51	0.77	0.75	0.88	1.08	0.90
N	0.92	1.01	1.24	1.47	0.92	0.91	0.67	0.79	0.91	0.97	1.09	0.99
O-1	0.21	0.19	0.34	0.44	0.27	0.33	0.21	0.41	0.31	0.44	0.45	0.40
O-2	0.58	0.77	0.61	0.91	0.75	0.75	0.49	0.72	0.75	0.84	0.93	0.84
P	0.12	0.14	0.15	0.16	0.17	0.13	0.10	0.17	0.15	0.21	0.25	0.20
Q	0.40	0.53	0.34	0.40	0.44	0.44	0.43	0.50	0.50	0.59	0.50	0.53

ISSUE 4. Wild Turkey/Human Conflicts

GOAL: To minimize and manage agricultural damage and nuisance turkey incidents.

Some wild turkey nuisance complaints and/or negative interactions with the public are unavoidable. While complaints have increased in recent years as the turkey population has grown, the annual number of complaints is relatively low compared with those for black bear and white-tailed deer.

The majority of the nuisance wild turkey complaints stem from turkeys' consumption or spoilage of silage or other stored crops. This situation often occurs when deep snow limits turkey mobility and restricts the birds' access to natural foods. Extreme weather creates intense stress on wild turkey populations whose fall food supplies become buried under snow at the same time that cold temperatures cause fat reserves to dwindle. Wild turkeys have a strong survival instinct that leads large winter flocks to exploit a convenient high calorie agricultural crop. Given this natural survival instinct, it is difficult to discourage them, especially once birds have established a pattern of regular feeding around a farm. While fall hunting may reduce the numbers of

offending birds to some degree, it will not solve this problem.

Farmers can protect silage in exposed bunkers by periodically placing waste silage close to the forest where turkeys are taking shelter. By starting this early before turkeys become accustomed to going to the bunker, the birds may be diverted to a food source that has little value to the farmer.

Dairy farmers have expressed concerns regarding potential transmission of diseases to their livestock from turkey feeding/defecation in feed bunkers. Vermont, New Hampshire, and Minnesota have conducted numerous investigations involving disease testing of local wild turkey flocks and, to date, have found no evidence of the presence of Salmonella DT 104 bacteria in these birds. Thus, farmers' concerns for disease transmission between wild turkeys and dairy cows appear to be unwarranted.

A turkey damage control regulation has been promulgated as another method to help address the nuisance issue. Under the "turkey damage rule," a landowner under game warden supervision may take a pre-approved number of offending turkeys that have been determined to have caused repeated or substantial damage to cultivated crops. The use of lethal control by shooting is normally considered to be the last option. With the assistance of a game warden, Department biologists, and U.S. Department of Agriculture's Wildlife Services personnel, other control methods such as hazing and/or fencing are attempted first. Frequently, complaints can be handled simply by providing technical or management assistance over the phone to educate landowners regarding turkey behavior and methods to change problem behavior.

Local National Wild Turkey Federation (NWTF) members can assist in quelling wild turkey/human conflicts. It is possible, given the demographic shift from farming to increased rural development, that nonagricultural nuisance complaints may increase in the near future. With this in mind, the Department will continue to adapt its approach to fit the issue.

Management Strategies

- 4.1 Provide property owners with access to coordinated services of personnel trained to deal with nuisance turkey issues including wildlife biologists, game wardens, and USDA Wildlife Services staff to assist with nuisance complaints

via technical guidance/assistance on techniques to minimize/discourage damage.

- 4.2 Conduct follow-up site visits to nuisance complaint sites when necessary and provide hazing equipment to help ameliorate persistent nuisance situations.
- 4.3 Solicit assistance from local volunteers through the Vermont Chapter of the National Wildlife Turkey Federation (NWTF) to help provide on-the-ground assistance to landowners via hazing and behavior modification efforts.
- 4.4 Assist USDA Wildlife Services staff with development of educational materials to inform and educate farmers about techniques for minimizing conflicts.
- 4.5 Compile and evaluate wild turkey damage complaint reports from farmers, state game wardens, biologists and wildlife service personnel to document problems, management approaches and results.
- 4.6 Develop/modify a standard set of protocols/guidelines/solutions to perceived and actual conflicts caused by wild turkeys (nuisance animals, agricultural damage).

ISSUE 5. Turkey Habitat Management and Conservation

GOAL: To encourage conservation and appropriate habitat management practices to support and sustain Vermont's wild turkey population.

Habitat quality and quantity are necessary to achieve wild turkey management goals. Land use changes that convert habitat to a lower quality or result in permanent habitat loss diminish its ability to sustain healthy, abundant turkey populations.

Management Strategies

- 5.1 Continue efforts on wildlife management areas and other public lands to provide habitat demonstration areas to promote appropriate commercial and noncommercial vegetation management practices beneficial to turkeys and other wildlife. This includes the use of prescribed fire and other management practices to establish and maintain long-term mast production areas.
- 5.2 Provide technical information and assistance regarding turkey habitat management to private



landowners and other land managers, town planning commissions via staff biologists, habitat demonstration projects, LIP and WHIP program lands, etc.

- 5.3 Update the “A Landowner’s Guide, Wildlife Habitat Management for Vermont Woodlands” and make it available on the Department’s website and in published form.
- 5.4 Work with the NWTF regional biologists and chapter volunteers on development of the North American Wild Turkey Management Plan.
- 5.5 Work with partnering organizations on high priority projects and issues.

ISSUE 6. Perception Regarding the Interaction Between Deer and Wild Turkeys, Ruffed Grouse and Wild Turkeys, and Various Predators and Wild Turkeys

GOAL: To improve the public’s knowledge, awareness, and understanding of the role of the wild turkey and its interactions within the ecosystem.

While the number of wild turkeys has increased dramatically throughout Vermont over the last decade, ruffed grouse and deer have at times declined in abundance. This leads some hunters to assume that turkeys could somehow be having a negative impact upon these other popular species. Some hunters believe that turkeys are eating more and more of the available food. However, biologists throughout the range where these species overlap believe that changes in deer or grouse numbers have nothing to do with the size of the turkey population. The factors limiting survival and populations of deer and grouse

are different than those limiting turkeys. Although the effects of winter severity can limit all three species, their effects vary by species. In winter, deer require softwood cover and woody browse. Turkeys don’t eat woody browse. While the formation of crusts on the snow surface can trap or prevent grouse from burrowing below the snow’s surface, crusts make it easy for turkeys to get around in search of food.

Through all the restoration efforts and the tremendous population growth there have been no documented reports of wild turkeys having any negative impact on other wildlife or threatened or endangered species. Because of their general and opportunistic feeding habits and adaptability, the wild turkey seems to be able to find a noncompetitive niche in which to survive regardless of the other species found in the area. One researcher noted that turkeys “usually have filled a vacant environmental niche wherever they have been introduced and no significant environmental problem has been attributed to them.” (Wunz 1992, National Wild Turkey Federation 2001).

DEER AND WILD TURKEYS

The most common concern expressed regarding turkeys competing with deer is that they out-compete deer for hard mast such as acorns or beech nuts. While it’s true that both deer and turkeys feed more on mast during years of mast abundance, but so do bears, squirrels, grouse, blue jays, and numerous small mammal species. Of these, turkeys may leave the most obvious evidence of feeding due to their scratching, but it’s highly unlikely that the birds consume mast to the detriment of deer. Autumn is the period of greatest wild food abundance. Wild apples, corn and other agricultural crops, grasses and forbs, berries and seeds of all kinds are used by both turkeys and deer and many other animals. In fact, a Pennsylvanian researcher used fencing to determine that of all species feeding on red oak acorns, deer actually obtain the greatest proportion of mast. Regarding beech mast, a Michigan researcher (Rosemier et al. 2005) found that in non-mast years, rodents actually consume most of the beechnuts. Considering the fact that only two 150 pound deer (300 lbs) equal the biomass of a flock of about 30 juvenile turkeys (or 15 large adults), it is easy to see how a few deer could easily consume considerably most of the mast crop.

While high turkey densities are believed to have no significant negative impact upon deer populations,

high deer densities do have a harmful impact upon turkeys, ruffed grouse, and other forest birds because excessive browsing of shrubs reduces protective cover, food sources, and nesting sites (Witmer and DeCalesta 1991).

RUFFED GROUSE AND WILD TURKEYS

Dan Dessecker, a forest biologist with the Ruffed Grouse Society, points out that “In order for increasing wild turkey populations to be able to exert a direct negative influence on local ruffed grouse, there would have to be some form of competition” (Dessecker 1996). This competition would be expected to focus on some limited resource, such as space, food, breeding areas, nesting sites, or some other resource. Dessecker notes that it’s highly doubtful that these two species compete for any limited resource. Ruffed grouse thrive in dense young forest stands. Turkeys prefer relatively open mature forests.

Regarding breeding areas, grouse drum on logs surrounded by dense shrubs, and turkey gobblers display in fields or forest openings. Although hens of both species nest in middle-aged or mature forest, their nest site requirements are quite general with both species using a wide range of sites that are found throughout forests. Again, it is very unlikely that there is limited space and competition for nest sites.

Research has shown that wild turkeys do not affect other bird species by eating young birds or destroying nests. Dr. Bill Palmer, a game bird biologist in Florida, used micro-video cameras and radio-tagged hens to monitor more than 400 quail nests and broods in an area with very high turkey populations (30-60 turkeys per square mile). The research did not record a single turkey destroying a nest or eating or killing a quail chick (Zimmer 2002). Gary Zimmer, another Ruffed Grouse Society forest biologist, points out that young grouse can fly well at only three weeks of age. When threatened the brood flushes in all directions to find cover and hide. That would make it nearly impossible for a turkey, with its poor sense of smell, to locate and harm grouse chicks (Zimmer 2002).

Competition for food is also not likely to be significant between turkeys and ruffed grouse. Both birds are generalists, in that they feed on an extremely wide variety of foods throughout the year. During winter, the most stressful period when food resources are most scarce, turkeys and grouse typically use different food sources (Whitaker 1998).



NWTF.ORG

Ruffed grouse feed on the buds of trees and shrubs. Dessecker notes that turkeys are heavy birds that can only stand on stout tree limbs, so they prefer to feed on the ground on waste grains, acorns and beechnuts, and residual fruits and seeds, such as highbush cranberry, burdock, and ash seeds.

Dave Neu, a regional biologist for the National Wild Turkey Federation, states, “Ruffed grouse and wild turkey are two species that have evolved together for thousands of years and their habitats slightly overlap. There is no documented evidence that either species directly impacts populations of the other” (Zimmer 2002). Although turkey populations have increased while grouse have decreased in some portions of Vermont, the population changes are mainly due to changes in agriculture and forest habitat. As young forests mature, the habitat becomes more suitable for turkeys and less attractive to grouse. Thus, populations of both species birds respond to changing habitat conditions rather than turkeys displacing grouse. Gary Zimmer stated it best when he said that “Any impact caused to ruffed grouse populations by turkeys is insignificant compared to declines in young-forest habitats...” (Zimmer 2002).

PREDATORS AND WILD TURKEYS

Department personnel are often asked why they don’t promote a bounty on coyotes or some other form of predator control to “protect” wild turkeys or other game species. It is well established that predator “control” will not protect wild turkeys. Predator/prey relationships are extremely dynamic and complex. These relationships involve a variety of factors that defy a simple, quick fix. Wild turkeys are prey to a long list of predators including coyotes, bobcats, foxes, fisher, weasels, skunks, opossum, raccoons, snakes, hawks, owls, domestic dogs,

and humans. In the case of implementing “coyote control,” for example, assuming that this could be effective, removal of coyotes would only reduce competition among the remaining host of predators that would continue to prey on turkeys. Coyotes, in fact, prey upon weasels, opossums, raccoons, foxes, and rarely skunks. All of these species are effective predators of nests, chicks, and nesting turkey hens. For this reason, it is possible that removal of coyotes could allow the populations of these other predators to increase resulting in more, not less, turkey predation and an overall decrease in a turkey population. Complex species relationships are common in nature. In fact, the rapid growth in Vermont’s turkey population has occurred during a time when the coyote population has been abundant.

Many of the qualities that hunters admire so much about these birds, such as their incredible eyesight, ability to detect movement and wariness, are products of the turkey’s long evolutionary history that they share with their predators. As wild turkey populations increase, the potential role of this species as a significant source of prey for other Vermont animals may now be greater than ever before.

Management Strategies

- 6.1 Promote sound scientific principles regarding inter-species competition and predator-prey relationships through a variety of outreach methods including public speaking events, web-based information and links, and print and broadcast media.

ISSUE 7. Developing and Maintaining an Informed Public is Crucial to the Management Success of the Wild Turkey Project.

GOAL: To ensure continued information exchange and program acceptance by keeping the general public, state and federal agencies informed on the status of the wild turkey resource in Vermont.


Habitat conservation and public use of the turkey resource are best accomplished when citizens are well-informed. Understanding the public’s opinion regarding turkey biology, habitat, and management issues is important in making acceptable management decisions.

Management Strategies

- 7.1 Disseminate wild turkey project information to the public/media professionals via biological reporting stations, teacher workshops, private and public landowner visits/conferences, slide/video presentations, mail correspondence, popular and technical reports, etc.
- 7.2 Use the Department’s library to fill all public requests for its video production “The Wild Turkey in Vermont” as well as its wildlife study guide “The Wild Turkey Education Kit.”
- 7.3 Continue involvement with standing professional committees, regulatory bodies and cooperative agreements with nongovernmental organizations to assist the Department with meeting the goals and objectives of this plan.

REFERENCES - WILD TURKEY

- Dessecker, D.R. 1996. The turkey question. Ruffed Grouse Society Magazine April-June 1996, pp. 35-36.
- National Wild Turkey Federation. 2001. A proposal for the introduction of wild turkeys in Nova Scotia. 44 pages.
- Rosemier, J.N., A.J. Storer, and D.J. Flaspohler. 2005. “Disruption of mast production in American Beech (*fagus grandifolia*) and effects on small mammal communities resulting from beech bark disease” in Evans, C.A., J.A. Lucas, and M.J. Twery. 2005. Beech Bark Disease: Proceedings of the Beech Bark Disease Symposium. Gen. Tech. Rep. NE-331. Newtown Square PA, US. Department of Agriculture Forest Service, Northern Research Station. 149 p.
- Whitaker, D.M. 1998. Turkeys: agents of mayhem? Ruffed Grouse Society Magazine 10(4): 10-26 (discontinuous pages).
- Witmer, G.W., and D.S. DeCalesta. 1991. The need and difficulty of bringing the Pennsylvania deer herd under control. Fifth Eastern Wildlife Damage Control Conference (1991). <http://digitalcommons.unl.edu/ewdcc5/45/> [2007].
- Wunz, G.A. 1992. “Wild turkeys outside their historic range” Pages 361-384 in J.G. Dickson, ed., The wild turkey: Biology and management. Stackpole Books, Mechanicsburg, PA.
- Zimmer, G. 2002. Are turkeys impacting ruffed grouse? Northwest Management, Inc., www.consulting-foresters.com [2007].




Vermont's Big Game Management Plan 2010-2020

Prepared by:

*Shawn Haskell, Cedric Alexander, Forrest Hammond, John Buck, Chris Conner,
John Austin, Kim Royar, Scott Darling, Curtis Smiley, Doug Blodgett, John
Gobeille, Don Isabelle, Paul Hamelin, Tom Decker, Wayne Laroche, Cathy Merrill,
Rebecca Davison, and Lilla Lumbra*

Approved by:

 12/8/2009
Wayne Laroche, Commissioner Date
Vermont Fish & Wildlife Department

